

Lawrence Berkeley National Laboratory

LBL Publications

Title

Impact of Genetic Modifications on the Magnetic and Electronic Structure of Nanoparticles
Particles Synthesized by Magnetotactic Bacteria

Permalink

<https://escholarship.org/uc/item/3k7323gt>

Authors

Liberati, Marco
Murat, D.
Komeili, A.
et al.

Publication Date

2010-04-16

Impact of Genetic Modifications on the Magnetic and Electronic Structure of Nanoparticles Synthesized by Magnetotactic Bacteria.

M. Liberati¹; D. Murat²; A. Komeili²; E. Arenholz¹

1. Advanced Light Source, Lawrence Berkeley National Laboratory, Berkeley, CA, United States.

2. Department of Plant and Microbial Biology, UC Berkeley, Berkeley, CA, United States.

To optimize magnetic nanoparticles (MNP) for applications such as cancer therapy and drug delivery, the ability to tailor their size, morphology, surface chemistry as well as their magnetic and electronic properties is of critical importance. Moreover, the appropriate functionalization of their surface to ensure biocompatibility with the human body is necessary.

Magnetotactic bacteria (MTB) represent a promising approach for the synthesis of MNP for biomedical applications. Once placed in iron rich and oxygen poor environments [3,4,5], they synthesize single crystals of magnetite (i.e. Fe_3O_4) which form of linear chains within the bacteria. Compared to chemically engineered MNP, the magnetosomes single crystals display numerous advantages:

- 1) narrow size distribution and uniform morphologies
 - 2) typical sizes (10-130 nm) not easily achievable through chemical synthesis
 - 3) superparamagnetic properties
 - 4) genetic control with access to morphologies inaccessible with chemical methods
 - 5) lower toxicity compared to some artificial MNP
 - 6) enveloped naturally by a biological membrane that can be specifically functionalized
- To control and tune the particle properties through genetic modifications of the MTB is expected to be possible [5] but needs to be confirmed through the detailed characterization of their magnetic, structural and electronic properties.

We have employed X-ray absorption (XA) spectroscopy and X-ray magnetic circular dichroism (XMCD) to determine the magnetic and electronic state of MNP synthesized by different genetic modifications of the *Magnetospirillum magneticum* strain AMB-1. The XA data are indicative of Fe ions in Fe^{2+} and Fe^{3+} in octahedral and tetrahedral sites as well as metallic Fe. In all samples a Fe XMCD was observed at ambient temperatures in external fields of 0.2 T. From a detailed analysis of the spectral shape of the XMCD signal we conclude that the pure strain AMB-1 forms Fe_3O_4 single crystals [6], while the genetically modified strain forms MNP with Fe in different states of oxidation.

Our results indicate that the characteristics of MNP produced by magnetotactic bacteria can be tuned through genetic modification of bacteria and that soft x-ray absorption spectroscopies provide valuable information about the resulting MNP.

The Advanced Light Source is supported by the Director, Office of Science, Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231.

References:

1. O. Salata, J. Nanobiotech. 2, 1 (2004)
2. C. Sun et al., Adv. Drug Deliv. Rev. 60, 1252 (2008)

3. D. Faivre and D. Schuler, Chem. Rev. 108, 4875 (2008)
4. C. Lang and D. Schuler, J. Phys. Cond. Matter. 18, S2815 (2006)
5. A. Komeili, Annu. Rev. Biochem. 76, 351 (2007)
6. S. Staniland et al., Nat. Nanotech. 3, 158 (2008)