

Lawrence Berkeley National Laboratory

Recent Work

Title

INDUCED NUCLEOPHILIC SUBSTITUTION BENZO[a]PYRENE

Permalink

<https://escholarship.org/uc/item/3k77h2jb>

Authors

Johnson, Milton D.
Calvin, Melvin.

Publication Date

1972-09-01

INDUCED NUCLEOPHILIC SUBSTITUTION.
BENZO[a]PYRENE

Milton D. Johnson and Melvin Calvin

September 1972

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*



DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Induced Nucleophilic Substitution in Benzo[a]pyrene

Cavalieri and Calvin have proposed^{1,2} that the carcinogenic action of benzo[a]pyrene (BaP) may arise from aryl hydroxylase induced binding to nucleophilic cellular components. The theory proposes that attack by electrophilic oxygen, produced in the hydroxylase system, occurs at the 6 position of the hydrocarbon to give a carbonium ion, localized primarily at the 1 and 3 positions, which can undergo attack with nucleophilic cellular components. Alternatively, initial attack could occur at 1 or 3, followed by nucleophilic reaction at 6.

Some evidence for the theory has been obtained by using iodonium ion as a model for the hydroxylase system. Benzo[a]pyrene and iodine dipyridine nitrate react to give either the 6-iodo or 6-pyridinium derivative or a mixture depending upon reaction conditions. Effects of solvent, concentrations and ratios of reactants were investigated and are summarized in the following table. Equimolar quantities of the reagents react within ten minutes in chloroform to give 6-iodo-benzo[a]pyrene in nearly quantitative yield. All other reactions listed require up to 24 hours to reach completion. Formation of the pyridinium derivative requires a 2:1 molar ratio of iodonium reagent to BaP.


Effect of Reaction Variables on Product Formation

Concentration of BaP	Molar Ratios		Solvent	Yield, %	
	py/BaP*	(py-I-py)**/BaP		I-BaP	py-BaP
		1	CHCl ₃	99	trace
		1	MeOH	95	
		1	DMF	95	
			pyridine	no reaction	
1 mg/ml	32	2	1% py/CHCl ₃	50**	50**
0.5 mg/ml	64	2	"		95
5 mg/ml	30	2	5% py/CHCl ₃		95
"	30	1	"		50
1 mg/ml		2	CH ₃ CN		10

* added pyridine

** approximate yields


The formation of the 6-pyridinium derivative may be interpreted as arising via the proposed ionic mechanism with initial electrophilic attack occurring at position 1 or 3 rather than at position 6 as occurs in the formation of 6-iodobenzo[a]pyrene. The position of attack might change for steric reasons since different reagents could be involved as indicated by the following equilibria:



If formation of 6-iodobenzo[a]pyrene arises from attack by I^+ , the equilibrium in pyridine-chloroform will be displaced to the left and a bulkier reagent might attack at position 1 or 3 to result in formation of the 6-pyridinium derivative. Steric factors are important since acylation with succinic anhydride and aluminum chloride gives the product with substitution in the one position.⁴ The excess pyridine present in the reaction mixture may also provide conditions where the addition of pyridine to the sigma complex formed by electrophilic attack at 1 or 3 is faster than the rate of elimination of a proton which would give the 1- or 3-iodo derivative. The dihydro-disubstituted compound thus formed would then undergo elimination of HI to give the 6-pyridinium derivative.

The reaction can also be postulated to proceed via the radical cation of BaP. Although evidence has been found³ that the radical cation of BaP, produced by oxidation with iodine, reacts with pyridine to give the 6-pyridinium derivative, no radical was detected under the conditions of these reactions. The simple iodine reaction does not proceed under the conditions employed in this study.

Both the ionic and radical modes are shown in the following reaction scheme.



The structural assignments for the iodo and pyridinium derivatives are based on the ultraviolet and 220 MHz nuclear magnetic resonance spectra. The UV spectra match reported spectra for 6-iodobenzo[a]pyrene⁵ and 6-pyridinium benzo[a]pyrene perchlorate.³ The NMR assignments are based on comparison with the spectrum of the parent hydrocarbon^{6,7}, coupling constants, splitting patterns, and double resonance decoupling experiments. The NMR spectra of the pyridinium derivative establishes the fact that the plane of the pyridine ring is perpendicular to the plane of the benzo[a]pyrene ring since protons 5 and 7 are shielded, relative to their positions in the parent compound, by the pyridine ring current. This is in contrast to the deshielding which all the protons, including 5 and 7, exhibit in the 6-iodo compound.

This work was supported, in part, by the U.S. Atomic Energy Commission, and, in part, through National Cancer Institute Contract NCI-FS-(71)-58.


Milton D. Johnson

Melvin Calvin

Laboratory of Chemical Biodynamics
University of California
Berkeley, California 94720

References

1. Cavaliere, E., and Calvin, M., Proc. Natl. Acad. Sci. USA, 68, 1251 (1971).
2. Cavaliere, E., and Calvin, M., Photochem. Photobiol., 14, 641 (1971).
3. Rochlitz, J., Tetrahedron, 23, 3048 (1967).
4. Buu-Hoi, N.P., and Lavit, D., Tetrahedron, 8, 1 (1960).
5. Tye, R., Graf, M.J., and Horton, A.W., Anal. Chem., 27, 248 (1955).
6. Bartelle, K.D., Jones, D.W., and Matthews, R.S., Spectrochim. Acta, 25A, 1603 (1969).
7. Haigh, C.W., and Mallion, R.B., J. Mol. Spec. 29, 418 (1969).



XBL 726-4661

The 220 MHz NMR spectra of A. Benzo[a]pyrene in $CDCl_3$,
B, 6-iodobenzo[a]pyrene in tetrahydrofuran, and C. the
6-pyridinium- d_5 derivative of benzo[a]pyrene in methanol.

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720