

UC Davis

UC Davis Previously Published Works

Title

Patterson-Sullivan theory for Anosov subgroups

Permalink

<https://escholarship.org/uc/item/3k9311pz>

Journal

Transactions of the American Mathematical Society, 375(12)

ISSN

0002-9947

Authors

Dey, Subhadip
Kapovich, Michael

Publication Date

2022-12-01

DOI

10.1090/tran/8713

Peer reviewed

PATTERSON–SULLIVAN THEORY FOR ANOSOV SUBGROUPS

SUBHADIP DEY AND MICHAEL KAPOVICH

ABSTRACT. We extend several notions and results from the classical Patterson–Sullivan theory to the setting of Anosov subgroups of higher rank semisimple Lie groups, working primarily with invariant Finsler metrics on associated symmetric spaces. In particular, we prove the equality between the Hausdorff dimensions of flag limit sets, computed with respect to a suitable Gromov (pre-)metric on the flag manifold, and the *Finsler critical exponents* of Anosov subgroups.

CONTENTS

Notations	5
1. Geometric preliminaries	6
2. Critical exponent	12
3. $\bar{\theta}$ -Conformal densities	16
4. Hyperbolicity of Morse image	20
5. Gromov distance at infinity	22
6. Shadow lemma	26
7. Dimension of a $\bar{\theta}$ -conformal density	30
8. Uniqueness of $\bar{\theta}$ -conformal density	33
9. Hausdorff density	38
10. Examples	40
Appendix A. Hausdorff measures on premetric spaces	44
Appendix B. A brief overview of the history of the Patterson–Sullivan theory in higher rank	45
Appendix C. A discussion of the main results without the ι -invariance condition on the type $\bar{\theta}$	47
Acknowledgments	49
References	49

Consider a discrete group Γ of isometries of the n -dimensional hyperbolic space \mathbb{H}^n . The *critical exponent* δ is a fundamental numerical invariant associated with Γ which measures the asymptotic growth rates of Γ -orbits in \mathbb{H}^n . The relation between the Hausdorff dimension of the limit set $\Lambda(\Gamma)$ of Γ and its critical exponent is now a

Received by the editors July 6, 2020, and, in revised form, December 5, 2021, and March 20, 2022.

2020 *Mathematics Subject Classification*. Primary 20F65, 22E40, 53C35.

Key words and phrases. Anosov representations, Patterson–Sullivan theory, critical exponent, Hausdorff dimension.

The second author was partly supported by the NSF grant DMS-16-04241.

classical result. In an influential paper [47], Sullivan proved the following theorem extending pioneering work by Patterson [38] on Fuchsian groups:

Theorem ([47, Thm. 8]). *Let Γ be a convex-cocompact subgroup of the isometry group of \mathbb{H}^n . Then the critical exponent δ of Γ equals to the Hausdorff dimension of $\Lambda(\Gamma)$.*

Later Sullivan generalized this theorem for geometrically finite Kleinian groups [48]. An important ingredient of Sullivan’s proof of this theorem is the existence of a finite, non-null Borel measure on $\Lambda(\Gamma)$ that changes *conformally* under the Γ -action. The construction of such measure goes back to Patterson’s original idea in [38]. Measures of this type (resp. a class of “well-behaved” measures) are commonly referred to as *Patterson–Sullivan measures* (resp. *densities*). We refer to Nicholls’ book [37] for a self-contained exposition on these results.

Since its introduction, the theory of Patterson and Sullivan has attracted a lot of attention. Further developments have been made by various people who analyzed more general classes of discrete groups and their limit sets. We list some of these developments here. Corlette [11] and Corlette–Iozzi [12] proved the above theorem for geometrically finite groups of isometries of rank-one symmetric spaces, and Bishop–Jones [5] extended these results to arbitrary discrete isometry groups of rank-one symmetric spaces. Yue [51] and Ledrappier [33] studied the case of Hadamard spaces of negative curvature.

There has been a considerable amount of development to understand the Patterson–Sullivan theory for discrete subgroups of higher rank semisimple Lie groups acting on its symmetric space, starting with Bishop–Steger [4] and Burger [8] in the rank-two case. Later, Albuquerque [1], Quint [42, 43], and Link [34] considered the case of Zariski-dense discrete subgroups in the isometry groups of general higher rank symmetric spaces. Link [36] also studied the case of products of rank-one symmetric spaces. In Appendix B we discuss these papers in relation to our work in more detail.

In the more abstract setting of Gromov hyperbolic spaces, much of Sullivan’s work in [47] was generalized by Coornaert [10] to the class of quasiconvex-cocompact groups. See also work of Paulin [39] on actions of subgroups of Gromov hyperbolic groups. Recent developments by Das–Simmons–Urbański [13] achieved greater generalizations of the Patterson–Sullivan theory (e.g., a generalization of Bishop–Jones’ theorem) in the case of “infinite-dimensional” Gromov hyperbolic spaces.

The goal of this paper is to study the Patterson–Sullivan theory for *Anosov subgroups*. The notion of Anosov subgroups was first introduced by Labourie [32] to study $\mathrm{PSL}(n, \mathbb{R})$ -Hitchin representations [25] of closed surface groups. This was further developed by Guichard–Wienhard [23] and Kapovich–Leeb–Porti [29–31]. Notably, Anosov subgroups extend the class of convex-cocompact subgroups of rank-one semisimple Lie groups to higher rank.

In this paper, we primarily work with some of the Kapovich–Leeb–Porti’s characterizations of Anosov subgroups. We briefly review these characterizations, the ones which we will need for this paper (namely, $\tau_{\mathrm{mod}}\text{-}URU$, $\tau_{\mathrm{mod}}\text{-}Morse$, and $\tau_{\mathrm{mod}}\text{-}RCA$), in Subsection 1.6. Since we do not use the original notion of Anosov representations as introduced by Labourie [32], and this notion has become classical at the time of writing the paper, we do not include Labourie’s definition. Readers who are interested to understand the connection between Labourie’s definition and Kapovich–Leeb–Porti’s characterizations are encouraged to read [30, Subsec. 5.11].

For instance, it is shown in [30, Subsec. 5.8] that Anosov and Morse properties are equivalent. In the same paper [30], the authors show that Morse (or any other equivalent notion, including RCA) implies the URU property. Finally, in [31], it is shown that URU implies Morse.

Main results. Let G be a noncompact real semisimple Lie group, $X = G/K$ be the associated symmetric space and Γ be a τ_{mod} -Anosov subgroup of G . We will be assuming several conditions on G and X ; they are labeled as “assumption” in Section 1. We consider two types of G -invariant (pseudo-)metrics on X , namely, one is a G -invariant Riemannian metric d_{Riem} of the symmetric space X , and the other one is a G -invariant Finsler distance $d_{\bar{\theta}}$ which depends on the choice of a direction $\bar{\theta}$ in the Weyl chamber (see Section 2). The *critical exponents* of Γ with respect to these two metrics, denoted by δ_{Riem} and $\delta_{\bar{\theta}}$, respectively, are defined in the usual fashion, i.e., as the exponents of convergence of an associated *Poincaré series* (see Section 2). Using the classical construction of Patterson, we define a Γ -invariant $\bar{\theta}$ -conformal density on the flag limit set of Γ (see Section 3).

Throughout this paper, the Finsler metric $d_{\bar{\theta}}$ is given more emphasis than its Riemannian counterpart. For example, the construction of the above mentioned Patterson–Sullivan density is carried out in terms of the Finsler metric. The main reason for this choice is that Finsler metrics reflect the asymptotic geometry of Γ better than the Riemannian metric. In fact, an interesting feature of $d_{\bar{\theta}}$ (for suitably chosen $\bar{\theta}$) is that the orbits of Anosov subgroups in X are Gromov-hyperbolic spaces when equipped with $d_{\bar{\theta}}$. See Corollary 4.8.

Let σ_{mod} be a maximal simplex in the Tits building of X , $\iota : \sigma_{\text{mod}} \rightarrow \sigma_{\text{mod}}$ be the opposition involution, τ_{mod} be an ι -invariant face of σ_{mod} , P be the maximal parabolic subgroup of G that stabilizes τ_{mod} , and $\text{Flag}(\tau_{\text{mod}}) = G/P$ be the partial flag manifold associated to the face τ_{mod} . We fix an ι -invariant type¹ $\bar{\theta} \in \text{int}(\tau_{\text{mod}})$.

Theorem A. *Let Γ be a nonelementary τ_{mod} -Anosov subgroup of G and $\delta_{\bar{\theta}}$ be the Finsler critical exponent for the action of Γ on the symmetric space $(X, d_{\bar{\theta}})$. Then the Patterson–Sullivan density² $\mu^{\bar{\theta}}$ on the flag limit set $\Lambda_{\tau_{\text{mod}}}(\Gamma) \subset \text{Flag}(\tau_{\text{mod}})$ is the unique (up to scaling) Γ -invariant $\bar{\theta}$ -conformal density. Moreover,*

- (i) *The density $\mu^{\bar{\theta}}$ is non-atomic and its dimension equals to $\delta_{\bar{\theta}}$.*
- (ii) *The support of $\mu^{\bar{\theta}}$ is $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ and the action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is ergodic with respect to $\mu^{\bar{\theta}}$.*
- (iii) *The critical exponent $\delta_{\bar{\theta}}$ (as well as the Riemannian critical exponent δ_{Riem}) is positive and finite.*
- (iv) *The $\bar{\theta}$ -Poincaré series of Γ diverges at the critical exponent $\delta_{\bar{\theta}}$. In other words, Γ has $\bar{\theta}$ -divergence type.*
- (v) *The $\delta_{\bar{\theta}}$ -dimensional Hausdorff measure on $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ with respect to a Gromov (pre-)metric³ is a member of a Γ -invariant conformal density (called the Hausdorff density). In particular, the Hausdorff dimension of $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ is $\delta_{\bar{\theta}}$.*

While constructions of conformal densities for discrete subgroups of semisimple Lie groups were done earlier (in the Zariski-dense case) by Albuquerque [1] and

¹Most of our main results are still valid without the assumption that $\bar{\theta}$ is ι -invariant; see the remark after Theorem B.

²See Definition 3.4.

³See Section 5.

Quint [42],⁴ most of the results of our Theorem A are not contained in their work, even in the Zariski-dense setting. Note that the theorem is false in general for nonelementary discrete subgroups of rank-one Lie groups which are not convex-cocompact. The proof of the theorem strongly relies on the τ_{mod} -Anosov condition.

The fact that $\mu^{\bar{\theta}}$ is the unique Γ -invariant $\bar{\theta}$ -conformal density is proven in Corollary 8.4. The main ingredients in the proof are a generalization of Sullivan's *shadow lemma* proven in Theorem 6.1, and an ergodicity argument (see Theorem 8.1) due to Sullivan. The proof of part (i) of the theorem is given in Corollaries 6.2 and 7.5. The second half of part (ii) follows from Theorem 8.3 while the first half follows from the facts that the support of $\mu^{\bar{\theta}}$ is a closed Γ -invariant subset of $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ and the action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is minimal. The part (iii) is proven in Propositions 2.6 and 3.5. See also the remarks following these propositions where δ_{Riem} is analyzed. The part (iv) follows from Corollary 6.5. The Hausdorff density in part (v) is studied in Section 9 (cf. Theorem 9.5). The background Gromov (pre-)metric is introduced in Section 5 where we also prove that the action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}(\Gamma)$ with respect to this metric is *conformal* (see Corollary 5.8).

We should note that some of the results in this paper are proven for more general classes of discrete subgroups of G with the hope that the results may be useful, for instance, in the study of *relatively Anosov subgroups*.⁵

For a much wider class of (uniformly) τ_{mod} -RA⁶ subgroups we prove the following results.

Theorem B. *Let $\Gamma < G$ be a nonelementary τ_{mod} -RA subgroup of G , and let $\delta_{\bar{\theta}}$ be its $\bar{\theta}$ -critical exponent. Then $\delta_{\bar{\theta}} \in (0, \infty]$. Let μ be a β -dimensional Γ -invariant $\bar{\theta}$ -conformal density (if exists).*

- (i) (*Shadow Lemma*) *Fix $x, x_0 \in X$. There exists $r_0 > 0$ such that for all $r \geq r_0$ and all $\gamma \in \Gamma$ satisfying $d_{\text{Riem}}(x, \gamma x_0) > r$,*

$$\mu_x(S(x : B(\gamma x_0, r))) \asymp \exp(-\beta d_{\bar{\theta}}(x, \gamma x_0))$$

(*Theorem 6.1*).

- (ii) $\beta \geq \delta_{\bar{\theta}} - \delta_{\bar{\theta}}^{\text{con}}$ (*Theorem 7.1*).

If we further assume that Γ is uniformly τ_{mod} -regular, then

- (iii) $\delta_{\bar{\theta}}$ *is finite* (*Proposition 2.6*), and

- (iv) *the density μ cannot have atoms at conical limit points* (*Corollary 6.2*).

Remark. Theorem A, with the exception of the item (v), and Theorem B remain valid without the assumption that the type $\bar{\theta} \in \text{int}(\tau_{\text{mod}})$ is ι -invariant. In Appendix C, we show how to generalize these statements without this assumption.

Some historical remarks. The early work on critical exponent of the Patterson–Sullivan theory in the higher rank was mostly developed for general (but, typically, Zariski dense) discrete subgroups of higher rank Lie groups; we discuss this early work (in relation to our paper) in Appendix B. Since the introduction of Hitchin representations of surface groups (and proof of their Anosov property by Labourie) and, more generally, Anosov representations of hyperbolic groups, a substantial work was done investigating different versions of critical exponent and the

⁴See Appendix B where we give a brief discussion about these papers.

⁵Relatively Anosov subgroups defined by Kapovich–Leeb [27] are an extension of the class of geometrically finite groups into the higher rank.

⁶See Subsection 1.6 for this definition.

Patterson–Sullivan theory, and their applications. Below is a brief discussion of this work.

For certain classes of Anosov subgroups, the Patterson–Sullivan theory was used by Sambarino in [45, 46] to solve certain counting problems, while in [7] Bridgeman–Canary–Labourie–Sambarino used related thermodynamic formalism to construct pressure metrics on spaces of Hitchin representations. Moreover, Glorieux–Monclair [20] studied the Patterson–Sullivan theory in the case of convex-cocompact subgroups of the isometry group of $\mathbb{H}^{p,q}$ equipped with the pseudo-Riemannian metric. In their work [40], Potrie and Sambarino prove some interesting inequalities for critical exponents of Hitchin representations and a beautiful rigidity theorem characterizing “Fuchsian” representations in the Hitchin component, which are reminiscent of the earlier inequalities for critical exponents and rigidity theorems (going back to the work of R. Bowen) for Kleinian groups, except that the inequalities are in the opposite direction.

While working on this article, we came to know about two very recent developments by Pozzetti–Sambarino–Wienhard [41] and Glorieux–Monclair–Tholozan [21], which are related to our work. Independently, the authors of these articles proved that the Hausdorff dimension of the limit set of a *projective* Anosov subgroup Γ in the real projective space with respect to the Riemannian metric is bounded above by a certain critical exponent, called the “simple root critical exponent” in the second article. The main result of [41] is stronger than this inequality for a special class of *(1,1,2)-hyperconvex* representations, in which case the Hausdorff dimension equals to the simple root critical exponent. They went further to prove that for hyperconvex subgroups Γ having $\partial\Gamma$ homeomorphic to a sphere, the limit set of Γ in the projective space is a C^1 sphere. As a corollary of these two results, they obtained an earlier result of Potrie–Sambarino [40] on the entropy of Hitchin representations. In [21], the authors also aimed to get a lower bound for the Hausdorff dimension of the limit set of general projective Anosov subgroups Γ . As it is mentioned in [21], initially, the authors aimed to obtain such a lower bound with respect to the Riemannian metric; eventually, they proved such a lower bound for a certain Gromov metric on the limit set. Using our Theorem A and relying on previously known computations of Busemann functions (see Example 5.10), we obtain a lower bound for this Hausdorff dimension with respect to the Riemannian metric (see Theorem 10.1).

After this work was completed, Andrés Sambarino informed us that Ledrappier’s methods from [33] (in conjunction with results of [7, Sec. 3.2]) can be used to obtain some of the results of our paper; we refer the reader to [45, 46] for similar applications of Ledrappier’s work.

NOTATIONS

Here we list some commonly used notations.

- $\mathfrak{B}(Y)$: Class of Borel subsets of a topological space Y
- $B(x, r)$: (Closed) ball of radius r centered at x
- $d_{\bar{g}}, d_{\text{Riem}}$: Finsler and Riemannian metrics, respectively, on X (see Section 2)
- $\widehat{xy}, \overline{xy}$: Finsler⁷ and Riemannian geodesic segments, respectively, connecting $x, y \in X$ (see Section 2)

⁷Note that Finsler geodesic segments connecting two points in X are usually non-unique.

- $\delta_{\bar{g}}, \delta_{\text{Riem}}$: Finsler and Riemannian critical exponents, respectively, of Γ (see Section 2)
- $D_x^{\bar{g}, \epsilon}$: Gromov premetric (see Definition 5.2)
- $\mathcal{B}_T^{\bar{g}}$: Busemann cocycle (see (3.1))

1. GEOMETRIC PRELIMINARIES

In this section, we briefly present some background material needed for the paper.

1.1. Symmetric spaces. A *symmetric space* X is a Riemannian manifold that has an *inversion symmetry* or *point-reflection* with respect to each point $x \in X$: This is an isometric involution $s_x : X \rightarrow X$ fixing x and sending each tangent vector at x to its negative. In this paper we only consider symmetric spaces which are simply-connected and have *noncompact type*. The latter means that X has no flat deRham factor and the sectional curvature of X is non-positive. In particular, X is a Hadamard manifold and, hence, is diffeomorphic to a euclidean space. We refer to Eberlein's book [16] for a detailed discussion of symmetric spaces.

Assumption 1. The symmetric space X is simply-connected and of noncompact type.

A symmetric space X can be written as G/K where G is a semisimple Lie group whose Lie algebra does not have compact and abelian factors, and K is a maximal compact subgroup of G . Moreover, this group G can be chosen to have finite center and be commensurable with the isometry group $\text{Isom}(X)$ of X . For example, one can choose G to be the identity component of $\text{Isom}(X)$.

Assumption 2. The semisimple Lie group G has finite center and is commensurable with the isometry group $\text{Isom}(X)$ of the symmetric space X .

Each point $x \in X$ determines a canonical decomposition of the Lie algebra \mathfrak{g} of G called the *Cartan decomposition*,

$$\mathfrak{g} = \mathfrak{k} + \mathfrak{p},$$

where \mathfrak{k} is tangent to the stabilizer of a point $x \in X = G/K$, and \mathfrak{p} can be realized as the tangent space of the symmetric space X at x . The dimension of a maximal abelian subalgebra $\mathfrak{a} \subset \mathfrak{p}$ is called the *rank* of X . The exponential map $\exp_x : \mathfrak{p} \rightarrow X$ identifies \mathfrak{a} with a maximal flat $F \subset X$ through x and, hence, the rank of X can also be defined as the dimension of a maximal totally geodesic flat subspace in X . A chosen maximal flat $F_{\text{mod}} \subset X$ is called the *model flat* which we isometrically identify with \mathbb{R}^k where $k = \text{rank}(X)$. The image in $\text{Isom}(F)$ of the G -stabilizer of F_{mod} is isomorphic to $\mathbb{R}^k \rtimes W$, where the first factor acts on $F_{\text{mod}} \cong \mathbb{R}^k$ by translations while the second factor W , called the *Weyl group*, is finite, fixes the origin, and is generated by hyperplane reflections. The closures of the connected components of the complement of the reflecting hyperplanes (for hyperplane reflections in W) in F_{mod} are called *chambers*. A chosen chamber is called the *model Weyl chamber*; we denote it by Δ .

1.2. Boundary at infinity. For a symmetric space X , there are multiple notions of (partial) boundary at infinity. The space of equivalence classes of asymptotic rays is called the *visual boundary* of X and denoted $\partial_\infty X$. The visual boundary is naturally identified with the unit tangent sphere $T_x^1 X$ at any point $x \in X$. The

topology it gets from this identification is called the *visual topology*. Attaching the visual boundary to X provides a compactification of X .

Another (strictly finer) topology on $\partial_\infty X$ is given by the G -invariant *Tits angle metric*:

$$\angle_{\text{Tits}}(\zeta, \eta) = \sup_{x \in X} \angle_x(\zeta, \eta),$$

where $\angle_x(\zeta, \eta)$ denotes the angle between the rays emanating from x and asymptotic to ζ and η . The boundary $\partial_\infty X$ with this topology is called the *Tits boundary* $\partial_{\text{Tits}} X$.

The Tits boundary $\partial_{\text{Tits}} X$ carries a canonical G -invariant structure of a spherical simplicial complex called the *Tits building* of X . This can be understood as follows: Consider the ideal boundary $\partial_\infty F_{\text{mod}}$ of F_{mod} where $k = \text{rank}(X)$. This is identified with the unit sphere \mathfrak{a}^1 of \mathfrak{a} and thus, we have an action of the Weyl group $W \curvearrowright \partial_{\text{Tits}} F_{\text{mod}}$. The pair $(\partial_{\text{Tits}} F_{\text{mod}}, W)$ is a spherical Coxeter complex which generates a spherical simplicial complex structure on entire $\partial_{\text{Tits}} X$ by the G -action.

Assumption 3. We assume that the Tits building is *thick*, i.e., every simplex of codimension one is a face of three maximal simplices.⁸

We do not have to worry about this assumption when X is an irreducible symmetric space. The Tits building of X , in that case, is thick. Nevertheless, we impose this assumption to avoid situations like in Example 1.1.

Example 1.1. Let $X = \mathbb{H}^2 \times \mathbb{H}^2$, and $G = \text{PSL}(2, \mathbb{R}) \times \text{PSL}(2, \mathbb{R}) \times (\mathbb{Z}/2\mathbb{Z})$, where the nontrivial element in $\mathbb{Z}/2\mathbb{Z}$ acts by swapping the factors of X . The corresponding Tits building of X is not thick.

We denote the intersection of Δ with the unit sphere in F_{mod} centered at the origin by σ_{mod} . This is a fundamental domain for the action $W \curvearrowright \partial_{\text{Tits}} F_{\text{mod}}$ where $\partial_{\text{Tits}} F_{\text{mod}}$ is identified with the unit sphere in F_{mod} centered at the origin. We call σ_{mod} the *model chamber*. Any other chamber (i.e., a top-dimensional simplex) in the Tits building is naturally identified with σ_{mod} via a G -equivariant map, called the *type map*,

$$\theta : \partial_{\text{Tits}} X \rightarrow \sigma_{\text{mod}}.$$

We reserve the notation τ_{mod} for the faces of σ_{mod} . An ideal point $\zeta \in \partial_{\text{Tits}} X$ (resp. a simplex $\tau \subset \partial_{\text{Tits}} X$) is called *type* $\bar{\theta} \in \sigma_{\text{mod}}$ (resp. of *type* $\tau_{\text{mod}} \subset \sigma_{\text{mod}}$) if $\theta(\zeta) = \bar{\theta}$ (resp. $\theta(\tau) = \tau_{\text{mod}}$). For $\bar{\theta} \in \tau_{\text{mod}}$ and a simplex τ of type τ_{mod} , we use the notation $\bar{\theta}(\tau)$ to denote the unique point in τ of type $\bar{\theta}$. The *opposition involution* ι is an automorphism of σ_{mod} which is defined as the negative of the longest element in the Weyl group.

Two simplices τ_1, τ_2 in the Tits building are called *antipodal* if there exists a point-reflection s_x swapping these two. Their types are related by $\theta(\tau_1) = \iota\theta(\tau_2)$. In particular, when τ_1 has an ι -invariant type τ_{mod} , then any antipodal simplex τ_2 also has type τ_{mod} . In this paper, we only consider types that are ι -invariant.

We now describe an important class of partial boundaries of X which are central to our study. Consider the action of G on the Tits building. The stabilizer of a face τ_{mod} of σ_{mod} is a parabolic subgroup $P_{\tau_{\text{mod}}}$ of G and we identify the quotient $G/P_{\tau_{\text{mod}}}$ with the set of all simplices of type τ_{mod} in the Tits building. This quotient

⁸This is a standing assumption on spherical buildings in the papers by Kapovich, Leeb and Porti we rely upon in our work. See for instance, [30, Lemma 2.4] and its usage elsewhere in that paper.

$G/P_{\tau_{\text{mod}}}$ is a smooth compact manifold, called the *partial flag manifold* of type τ_{mod} and is denoted $\text{Flag}(\tau_{\text{mod}})$. The partial compactification of X by attaching $\text{Flag}(\tau_{\text{mod}})$ is denoted

$$\bar{X}^{\tau_{\text{mod}}} = X \cup \text{Flag}(\tau_{\text{mod}})$$

which is topologized via the topology of *flag convergence* (see Subsection 1.6). In the special case when $\tau_{\text{mod}} = \sigma_{\text{mod}}$, the associated parabolic subgroup $P_{\sigma_{\text{mod}}}$ is minimal and $\text{Flag}(\sigma_{\text{mod}}) = G/P_{\sigma_{\text{mod}}}$ is the *full flag manifold*, also called the *Furstenberg boundary* of X .

A subset $A \subset \text{Flag}(\tau_{\text{mod}})$ is called *antipodal* if any two distinct simplices in A are antipodal.

1.3. Δ -Valued distances and a generalized triangle inequality. There is a canonical map $d_{\Delta} : X \times X \rightarrow \Delta$ which is defined as follows: For a pair of points (x, y) in X , there is an element $g \in G$ which maps x to the origin in Δ and y to a point $v \in \Delta$. We define $d_{\Delta}(x, y) = v$. Note that the norm $\|d_{\Delta}(x, y)\|$ (induced by the euclidean inner product on $F_{\text{mod}} \cong \mathbb{R}^k$) equals $d_{\text{Riem}}(x, y)$ where d_{Riem} denotes the distance function induced by the Riemannian metric on X .

For a pair $(x, y) \in X \times X$, the value $d_{\Delta}(x, y)$ is called the Δ -valued distance between x and y . This is a complete G -congruence invariant for oriented line segments in X . The Δ -valued distances satisfy generalized triangle inequalities (see [28]). In the paper we will need the following triangle inequality. For $x, y, z \in X$,

$$(1.1) \quad \|d_{\Delta}(x, y) - d_{\Delta}(x, z)\| \leq d_{\text{Riem}}(y, z).$$

1.4. Parallel sets, cones, and diamonds. For a detailed discussion on this subsection, we refer to [29, Subsec. 2.4], [30, Subsec. 2.5].

Let τ_{\pm} be a pair of antipodal simplices in the Tits building of X . The *parallel set* $P(\tau_{+}, \tau_{-})$ is the union of all maximal flats in X whose ideal boundary contains $\tau_{+} \cup \tau_{-}$ as a subset. This is a totally geodesic submanifold of X .

For a simplex τ , the *star* $\text{st}(\tau)$ of τ is the union of all chambers in the Tits building containing τ . The *open star* $\text{ost}(\tau)$ of τ is the union of all the open simplices whose closures contains τ . For a face τ_{mod} of σ_{mod} (viewed as a complex), define the open star $\text{ost}(\tau_{\text{mod}})$ similarly. The *boundary* $\partial\text{st}(\tau_{\text{mod}})$ is the complement of $\text{ost}(\tau_{\text{mod}})$ in σ_{mod} .

Let τ_{mod} be an ι -invariant face of σ_{mod} . An ideal point $\xi \in \partial_{\infty}X$ is called τ_{mod} -regular if its type is contained in $\text{ost}(\tau_{\text{mod}})$. Moreover, given an ι -invariant compact subset $\Theta \subset \text{ost}(\tau_{\text{mod}})$, an ideal point $\xi \in \partial_{\infty}X$ is called Θ -regular if its type is contained in Θ . A nondegenerate geodesic segment (or line or ray) in X is called τ_{mod} -regular (resp. Θ -regular) if the ideal endpoints of its line extension are τ_{mod} -regular (resp. Θ -regular).

For a simplex τ in the Tits building and a point $x \in X$, the τ_{mod} -cone $V(x, \text{st}(\tau))$ with apex x is the union of all rays emanating from x asymptotic to a point $\xi \in \text{st}(\tau)$. For a τ_{mod} -regular geodesic segment $\overline{xy} \subset X$, the τ_{mod} -diamond $\diamond_{\tau_{\text{mod}}}(x, y)$ is the intersection of the opposite cones $V(x, \text{st}(\tau_{+}))$ and $V(y, \text{st}(\tau_{-}))$ containing it. The points x and y are called the *endpoints* of $\diamond_{\tau_{\text{mod}}}(x, y)$. The cones and parallel sets can be interpreted as limits of diamonds where, respectively, one or both endpoints diverge to infinity. All of these are convex subsets of X (see [29, Prop. 2.14], [30, Prop. 2.10]). In particular, the cones are *nested*: For every $y \in V(x, \text{st}(\tau))$, $V(y, \text{st}(\tau)) \subset V(x, \text{st}(\tau))$.

Let Θ be an ι -invariant compact subset of $\text{ost}(\tau_{\text{mod}})$. In a similar way as above, the Θ -cone $V(x, \text{ost}_\Theta(\tau))$ with apex x is the union of all rays emanating from x asymptotic to a point $\xi \in \text{st}(\tau)$ of type Θ . Note that $V(x, \text{ost}_\Theta(\tau))$ is strictly contained inside $V(x, \text{st}(\tau))$.

1.5. Morse embeddings. The *Morse property* in higher rank was introduced by Kapovich-Leeb-Porti in [29].

Recall that a *quasigeodesic* in X is a quasiisometric embedding $\phi : I \rightarrow X$ of an interval $I \subset \mathbb{R}$. We say that ϕ is τ_{mod} -*regular quasigeodesic* if for all sufficiently separated points $t_1, t_2 \in I$, the segment $\overline{\phi(t_1)\phi(t_2)}$ is τ_{mod} -regular. We say that ϕ is a τ_{mod} -*Morse quasigeodesic* if it is τ_{mod} -regular and for all sufficiently separated points $t_1, t_2 \in I$, the image $\phi([t_1, t_2])$ is uniformly close to $\diamond_{\tau_{\text{mod}}}(\phi(t_1), \phi(t_2))$.

Let Z be a geodesic Gromov-hyperbolic metric space (cf. Definition 4.2).

Definition 1.2 (Morse embeddings). A quasiisometric map $\phi : Z \rightarrow X$ is called a τ_{mod} -*Morse embedding* if the image of every geodesic is a τ_{mod} -Morse quasigeodesic with uniformly controlled coarse-geometric quantifiers: There exist a constant $D > 0$ and an ι -invariant compact subset $\Theta \subset \text{ost}(\tau_{\text{mod}})$ such that if $z_1 z_2$ is a geodesic segment in Z of length $\geq D$, then $\overline{\phi(z_1)\phi(z_2)}$ is a Θ -regular geodesic in X and the image $\phi([z_1, z_2])$ is D -close to $\diamond_{\tau_{\text{mod}}}(\phi(z_1), \phi(z_2))$.

1.6. Discrete subgroups of G and their limit sets. We consider discrete subgroups with various levels of regularity and their flag limit sets. Most of these notions first appear in the work of Benoist [2]; our discussion follows [29] and [30].

We first recall the notion of *regular sequences* in X . Let τ_{mod} be an ι -invariant face of σ_{mod} . Let $V(0, \partial\text{st}(\tau_{\text{mod}}))$ denote the union of all rays in Δ emanating from 0 asymptotic to points $\xi \in \partial\text{st}(\tau_{\text{mod}})$. A sequence (x_n) on X diverging to infinity is τ_{mod} -*regular* if for all $x \in X$, the sequence $(d_\Delta(x, x_n))_{n \in \mathbb{N}}$ in Δ diverges away from $V(0, \partial\text{st}(\tau_{\text{mod}}))$. Furthermore, a τ_{mod} -regular sequence (x_n) is called *uniformly τ_{mod} -regular* if the sequence $(d_\Delta(x, x_n))_{n \in \mathbb{N}}$ in Δ diverges away from $V(0, \partial\text{st}(\tau_{\text{mod}}))$ at a linear rate,

$$\liminf_{n \rightarrow \infty} \frac{d(d_\Delta(x, x_n), V(0, \partial\text{st}(\tau_{\text{mod}})))}{d(0, d_\Delta(x, x_n))} > 0,$$

where d denotes the euclidean distance on Δ . Accordingly, a sequence (g_n) in G is τ_{mod} -regular (resp. uniformly τ_{mod} -regular) if for some (equivalently, every) $x \in X$, the sequence $(g_n(x))$ is τ_{mod} -regular (resp. uniformly τ_{mod} -regular).

Recall from Subsection 1.2 that we have identified $\text{Flag}(\tau_{\text{mod}})$ with the set of all simplices of type τ_{mod} in the Tits building of X . Also, recall the notion of the stars $\text{st}(\tau)$, and cones $V(x, \text{st}(\tau))$ from Subsection 1.4.

Definition 1.3 (Shadows). For $x \in X$ and $A \subset X$, the *shadow of A in $\text{Flag}(\tau_{\text{mod}})$ from x* is

$$(1.2) \quad S(x : A) = \{\tau \in \text{Flag}(\tau_{\text{mod}}) \mid A \cap V(x, \text{st}(\tau)) \neq \emptyset\}.$$

Remark 1.4. The notion of shadows is used in [31, Subsec. 3.8] to produce a topology, called the *shadow topology*, in $\text{Flag}(\tau_{\text{mod}})$. This topology is generated by the following basic subsets:

$$S(x : B(y, r)), \quad \text{where } x, y \in X, \text{ and } r > 0.$$

Moreover, the shadow topology coincides with the standard topology (i.e., the underlying topological space of a K -invariant Riemannian metric) on $\text{Flag}(\tau_{\text{mod}})$. See [31, Lem. 3.82].

Let (g_n) be a τ_{mod} -regular sequence in G . A sequence (τ_n) in $\text{Flag}(\tau_{\text{mod}})$ is called a *shadow sequence* of (g_n) if there exists $x \in X$ such that, for every $n \in \mathbb{N}$, $\tau_n = S(x : \{g_n x\})$. A τ_{mod} -regular sequence (g_n) is said to be τ_{mod} -*flag-convergent* to $\tau \in \text{Flag}(\tau_{\text{mod}})$ if a(ny) shadow sequence (τ_n) of (g_n) converges to τ . This notion of flag-convergence is the same as the one originally introduced in [31], where the shadow topology was not yet defined.

The notion of flag-convergence leads to the definition of *flag limit sets* of discrete subgroups $\Gamma < G$.

Definition 1.5 (Limit sets). The τ_{mod} -*flag limit set* of a discrete subgroup Γ of G , denoted by $\Lambda_{\tau_{\text{mod}}}(\Gamma)$, is the subset of $\text{Flag}(\tau_{\text{mod}})$ which consists of all limit simplices of τ_{mod} -flag-convergent sequences on Γ .

Remark 1.6. The flag limit set $\Lambda_{\tau_{\text{mod}}}$ is Γ -invariant.

More generally, one defines τ_{mod} -flag-limit sets of a subset $Z \subset X$ as the accumulation subset of Z in $\text{Flag}(\tau_{\text{mod}})$ with respect to the topology of flag-convergence.

Now, we review definitions of several classes of discrete subgroups of G with various levels of regularities:

R: A discrete subgroup $\Gamma < G$ is τ_{mod} -*regular* if for all $x \in X$ and all sequences of distinct elements (γ_n) in Γ , the sequence $(\gamma_n x)$ is τ_{mod} -regular. For τ_{mod} -regular subgroups Γ , the flag limit set $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ provides a compactification of the orbit $\Gamma x \subset X$, i.e., $\Gamma x \sqcup \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is compact.

RA: A τ_{mod} -regular subgroup Γ is τ_{mod} -*RA* (*regular antipodal*) if its limit set $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ is antipodal, i.e., every two distinct elements of $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ are antipodal to each other. For τ_{mod} -RA subgroups Γ , the action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is a *convergence action*⁹ (see [29, Prop. 5.38]). A τ_{mod} -RA subgroup Γ is called *nonelementary* if $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ consists of at least three (hence infinitely many) points; otherwise Γ is called *elementary*. If Γ is nonelementary then the action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is *minimal*, i.e., every orbit of Γ is dense, and $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ is perfect.¹⁰

RC: For a τ_{mod} -regular subgroup Γ , a limit simplex $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is a *conical limit point* if there exist $x \in X$, $c > 0$ and a sequence (γ_n) of pairwise distinct isometries on Γ such that

$$d_{\text{Riem}}(\gamma_n x, V(x, \text{st}(\tau))) \leq c,$$

where d_{Riem} denotes the Riemannian distance on X . The set of all conical limit simplices is denoted by $\Lambda_{\tau_{\text{mod}}}^{\text{con}}(\Gamma)$. A subgroup $\Gamma < G$ is called τ_{mod} -*RC* if $\Lambda_{\tau_{\text{mod}}}(\Gamma) = \Lambda_{\tau_{\text{mod}}}^{\text{con}}(\Gamma)$.

RCA: A subgroup Γ is τ_{mod} -*RCA* if it is both τ_{mod} -RA and τ_{mod} -RC.

⁹Recall that an action $\Gamma \curvearrowright Z$ is called a *convergence action* if the induced action $\Gamma \curvearrowright Z^{(3)}$ is properly discontinuous. Here $Z^{(3)}$ denotes the space of all triples of pairwise distinct points in Z . The action $\Gamma \curvearrowright Z$ is called *uniform convergence action* if, in addition, $\Gamma \curvearrowright Z^{(3)}$ is a cocompact action.

¹⁰This follows from a general result for convergence actions by Gehring–Martin [19] and Tukia [49]. See also [29, Subsec. 3.2] or [30, Subsec. 3.3].

U: A finitely generated subgroup $\Gamma < G$ (equipped with the word metric) is said to be *undistorted* if one (equivalently, every) orbit map $\Gamma \rightarrow \Gamma x \subset X$ is a quasiisometric embedding.

UR: A discrete subgroup $\Gamma < G$ is *uniformly τ_{mod} -regular* if for all $x \in X$ and all sequences of distinct elements (γ_n) in Γ , the sequence $(\gamma_n x)$ is uniformly τ_{mod} -regular.

URU: A subgroup $\Gamma < G$ is said to be τ_{mod} -URU if it is both τ_{mod} -uniformly regular and undistorted.

Morse: A discrete finitely generated subgroup (equipped with a word metric) $\Gamma < G$ is called τ_{mod} -Morse if it is word-hyperbolic and, for an(y) $x \in X$, the orbit map $\Gamma \rightarrow \Gamma x$ is a τ_{mod} -Morse embedding. See Subsection 1.5.

In [30, Equiv. Thm. 1.1] and [31], the properties Morse, RCA and URU are proven to be equivalent to the Anosov property defined by Labourie [32] and Guichard-Wienhard [23].

Theorem 1.7 ([30, Equiv. Thm. 1.1]). *The following classes of nonelementary discrete subgroups of G are equal:*

- (i) τ_{mod} -RCA,
- (ii) τ_{mod} -Morse,
- (iii) $P_{\tau_{\text{mod}}}$ -Anosov,
- (iv) τ_{mod} -URU.

In the sequel, any discrete subgroup that satisfies any of equivalent conditions in the theorem will be called a τ_{mod} -Anosov subgroup.

1.7. Illustrating examples. In this paper, we consider the following two classes of examples.

Example 1.8 (Product of rank-one symmetric spaces). Let X be a product of k rank-one symmetric spaces (X_i, d_i) ,

$$X = X_1 \times \cdots \times X_k.$$

The rank of X is k . Let G be a semisimple Lie group commensurable with the isometry group of X . (For example, we may take $G = \text{Isom}(X_1) \times \cdots \times \text{Isom}(X_k)$.) Assumption 3 on page 7 amounts to the requirement that G preserves the factors of the direct product decomposition of X .

The model maximal flat F_{mod} can be viewed as the product of some chosen geodesic lines (coordinate axes), one for each deRham factor. The Weyl group W is generated by reflections along the coordinate hyperplanes and the longest element in it is the reflection about the origin. The model Weyl chamber Δ can be realized as the nonnegative orthant. Here is a formula for the Δ -valued distances: For $(x_1, \dots, x_k), (y_1, \dots, y_k) \in X_1 \times \cdots \times X_k$,

$$(1.3) \quad d_{\Delta}((x_1, \dots, x_k), (y_1, \dots, y_k)) = (d_1(x_1, y_1), \dots, d_k(x_k, y_k)) \in \mathbb{R}_{\geq 0}^k.$$

It follows that the opposition involution ι acts on Δ trivially.

Recall that the Tits boundary of a product of two symmetric spaces is the simplicial join of their individual Tits buildings and, for rank-one symmetric spaces, the Tits boundary is discrete. These two facts imply that the $(p-1)$ -simplices in the Tits building of X for $1 \leq p \leq k$ can be parametrized by p -tuples $(\xi_{r_1}, \dots, \xi_{r_p}) \in \partial_{\infty} X_{r_1} \times \cdots \times \partial_{\infty} X_{r_p}$, $1 \leq r_1 < \cdots < r_p \leq k$,

$$(\xi_{r_1}, \dots, \xi_{r_p}) \leftrightarrow \tau = \text{span}\{\xi_{r_1}, \dots, \xi_{r_k}\}.$$

We say that such a simplex τ has type $\tau_{\text{mod}} = (r_1, \dots, r_p)$. The incidence structure can be understood as follows: Two simplices have a common q -face if and only if they have q equal coordinates.

The star $\text{st}(\tau) = (\xi_{r_1}, \dots, \xi_{r_p})$ is the minimal subcomplex of the Tits building containing all chambers $(\zeta_1, \dots, \zeta_k)$ satisfying $\zeta_{r_i} = \xi_{r_i}$, for all $i \in \{1, \dots, p\}$.

Since the opposition involution ι fixes each chamber point-wise, every face τ_{mod} of σ_{mod} and every type is ι -invariant. Every two chambers (resp. faces of the same type) in $\partial_{\text{Tits}} X$ are antipodal to each other unless they have a common face (resp. sub-face).

Example 1.9 ($X = \text{SL}(k+1, \mathbb{R})/\text{SO}(k+1, \mathbb{R})$). We take $G = \text{SL}(k+1, \mathbb{R})$, $K = \text{SO}(k+1, \mathbb{R})$; the symmetric space $X = G/K$ is identified with the set of all positive definite, symmetric matrices in $\text{SL}(k+1, \mathbb{R})$. In this case $\text{rank}(X) = k$ and X is irreducible. The standard choice of a model flat F_{mod} is the subset of all diagonal matrices $a = \text{diag}(a_1, \dots, a_{k+1}) \in \text{SL}(k+1, \mathbb{R})$ with positive diagonal entries. We identify the model flat with \mathfrak{a} via the logarithm map

$$\log : a = \text{diag}(a_1, \dots, a_{k+1}) \mapsto (\log a_1, \dots, \log a_{k+1}),$$

where \mathfrak{a} is viewed as the hyperplane in \mathbb{R}^{k+1} consisting of all points with zero sum of coordinates.

The Weyl group $W = \text{Sym}_{k+1}$ acts on \mathfrak{a} by permuting the coordinates. The standard choice for the model Weyl chamber $\Delta = \mathfrak{a}_+$ consists of all the points in \mathfrak{a} with decreasing coordinate entries. The Cartan projection¹¹ $\rho : \text{SL}(k+1, \mathbb{R}) \rightarrow \mathfrak{a}_+$ can be written as $g \mapsto \log a$ where a is associated to g via the *singular value decomposition* $g = uav$, $u, v \in \text{SO}(k+1, \mathbb{R})$. The logarithm of i -th singular value of g will be denoted by $\sigma_i(g)$. The opposition involution ι maps $(\sigma_1, \dots, \sigma_{k+1}) \in \mathfrak{a}_+$ to $(-\sigma_{k+1}, \dots, -\sigma_1)$.

The Tits building of X can be identified with the incidence geometry of flags in \mathbb{R}^{k+1} . The Furstenberg boundary consists of full flags

$$V_1 \subset \dots \subset V_{k+1} = \mathbb{R}^{k+1}, \quad \dim(V_i) = i.$$

The partial flags are

$$V : V_{r_1} \subset \dots \subset V_{r_p} \subset V_{r_{p+1}} = \mathbb{R}^{k+1}, \quad \dim(V_{r_i}) = r_i,$$

$1 \leq r_1 < \dots < r_p < r_{p+1} = k+1$, which are elements of $\text{Flag}(\tau_{\text{mod}})$ where $\tau_{\text{mod}} = (r_1, \dots, r_p)$. The opposition involution maps τ_{mod} to $\iota\tau_{\text{mod}} = (k+1-r_p, \dots, k+1-r_1)$. It follows that τ_{mod} is ι -invariant if and only if $r_i + r_{p+1-i} = k+1$, for each $i = 1, \dots, p$. The partial flag manifold $\text{Flag}(\tau_{\text{mod}})$ consisting of all partial flags V of type $\tau_{\text{mod}} = (r_1, \dots, r_p)$ naturally embeds into the product of Grassmannians $\text{Gr}_{r_1}(\mathbb{R}^{k+1}) \times \dots \times \text{Gr}_{r_p}(\mathbb{R}^{k+1})$.

Suppose that $\tau_{\text{mod}} = (r_1, \dots, r_p)$ is ι -invariant. A pair $V^\pm \in \text{Flag}(\tau_{\text{mod}})$ is antipodal if and only if $V_{r_i}^+ + V_{r_{p+1-i}}^- = \mathbb{R}^{k+1}$ for each $i = 1, \dots, p$.

2. CRITICAL EXPONENT

On a symmetric space $X = G/K$, we consider two natural (pseudo-)metrics. Let $d_{\text{Riem}}(\cdot, \cdot)$ denote the distance function on X of the (fixed) G -invariant Riemannian metric on X . Throughout the paper, we fix an ι -invariant face τ_{mod} of σ_{mod} , and fix an ι -invariant type θ in the interior of τ_{mod} .

¹¹Or the Δ -valued distance in the sense that $d_\Delta(x, gx) = \rho(g)$.

Let $d_{\bar{\theta}}$ denote the polyhedral Finsler (pseudo-)metric¹² on X :

$$(2.1) \quad d_{\bar{\theta}}(x, y) = \langle d_{\Delta}(x, y) | \bar{\theta} \rangle.$$

The inner product above is the euclidean inner product on F_{mod} coming from the Riemannian metric on X . Since $\bar{\theta}$ is in the unit sphere of F_{mod} , and since the diameter of a Weyl chamber for the spherical metric is at most $\pi/2$,¹³ we have

$$(2.2) \quad 0 \leq d_{\bar{\theta}}(x, y) \leq d_{\text{Riem}}(x, y).$$

Remark 2.1. The distance function $d_{\bar{\theta}}$ depends on the choice of $\bar{\theta} \in \tau_{\text{mod}}$.

The metric space (X, d_{Riem}) is a complete Riemannian manifold and, in particular, it is *geodesic*: Any two points in X can be connected by a geodesic segment. The (pseudo-)metric space $(X, d_{\bar{\theta}})$ is also a geodesic space. The geodesics in $(X, d_{\bar{\theta}})$ are called *Finsler geodesics*. All the Riemannian geodesics are also Finsler, however, the converse is generally false: There are non-Riemannian Finsler geodesics when $\text{rank}(X) \geq 2$. The precise description of all Finsler geodesics is given in [26, Subsec. 5.1.3]. We merely use this description as a definition of Finsler geodesics.

Definition 2.2 (Finsler geodesics). Let $I \subset \mathbb{R}$. A path $\ell : I \rightarrow X$ is called a *Finsler geodesic* if there exists a pair of antipodal flags $\tau_{\pm} \in \text{Flag}(\tau_{\text{mod}})$ such that $\ell(I) \subset P(\tau_+, \tau_-)$ and

$$\ell(t_2) \in V(\ell(t_1), \text{st}(\tau_+)), \quad \forall t_1 \leq t_2.$$

Moreover, given an ι -invariant compact subset $\Theta \subset \text{ost}(\tau_{\text{mod}})$, a Finsler geodesic $\ell : I \rightarrow X$ is called a Θ -*Finsler geodesic* if, in addition to the above, it satisfies the following stronger condition:

$$\ell(t_2) \in V(\ell(t_1), \text{ost}_{\Theta}(\tau_+)), \quad \forall t_1 \leq t_2.$$

Remark 2.3. Finsler geodesics give alternative description of diamonds, namely, the τ_{mod} -diamond $\diamond_{\tau_{\text{mod}}}(x, y)$ is the union of all Finsler geodesics connecting the endpoints x and y . See [26, Subsec. 5.1.3].

Notation. In this paper, we use the notation \overline{xy} to denote the Riemannian geodesic segment connecting a pair of points $x, y \in X$. To denote a Finsler geodesic segment connecting x and y , we use the notation \widehat{xy} .

Below we let $*$ be either “Riem” or $\bar{\theta}$. Let $\Gamma < G$ be a subgroup, and $x, x_0 \in X$. Define the *orbital counting function* $N_*(r, x, x_0) : [0, \infty) \rightarrow [0, \infty]$,

$$N_*(r) = N_*(r, x, x_0) = \text{card}\{\gamma \in \Gamma \mid d_*(x, \gamma x_0) < r\}.$$

Using $N_*(r)$, following [1] and [43], we define the *critical exponent* δ_* of Γ by

$$(2.3) \quad \delta_* = \limsup_{r \rightarrow \infty} \frac{\log N_*(r)}{r} \in [0, \infty].$$

The critical exponents $\delta_{\bar{\theta}}$ and δ_{Riem} will be called the $\bar{\theta}$ -*critical exponent* and *Riemannian critical exponent*, respectively.

¹²Our definition is same as the one in [26, Subsec. 5.1.2]. It is remarked in the second paragraph of [26, p. 2571] that $d_{\bar{\theta}}(x, y) = -b^{\bar{\theta}} \circ d_{\Delta}(x, y)$, where $b^{\bar{\theta}} = -\langle \cdot | \bar{\theta} \rangle$ is the Busemann function on the flat $F_{\tau_{\text{mod}}} \supset \Delta$, with the gradient $\bar{\theta}$ and normalized at the origin.

¹³This follows from the fact that the action $W \curvearrowright \mathfrak{a}$ is *essential*, i.e., W does not fix any proper subspace of \mathfrak{a} .

Remark 2.4. The discussion in [1] and [43] is mostly limited to the case when $\bar{\theta}$ is regular, i.e., belongs to the interior of σ_{mod} .

The critical exponent is independent of the chosen points x and x_0 . The proof is standard: Consider the Poincaré series

$$(2.4) \quad g_s^*(x, x_0) = \sum_{\gamma \in \Gamma} \exp(-sd_*(x, \gamma x_0)).$$

It is a well-known fact that $g_s^*(x, x_0)$ converges if $s > \delta_*(x, x_0)$ and diverges if $s < \delta_*(x, x_0)$ where $\delta_*(x, x_0)$ denotes the right side of (2.3). Using the triangle inequality, we obtain

$$\exp(-sd_*(x, x_0)) g_s^*(x_0, x_0) \leq g_s^*(x, x_0) \leq \exp(sd_*(x, x_0)) g_s^*(x_0, x_0).$$

Hence, convergence or divergence of $g_s^*(x, x_0)$ is independent of the choice of x and so is $\delta_*(x, x_0)$. For a similar reason, it is also independent of the choice of x_0 .

Definition 2.5. A discrete subgroup Γ of G is of $\bar{\theta}$ -convergence type if the $\bar{\theta}$ -Poincaré series $g_{\bar{\theta}}^*(x, x_0)$ converges at the critical exponent $\delta_{\bar{\theta}}$. Otherwise, we say that Γ has $\bar{\theta}$ -divergence type.

Since the action $\Gamma \curvearrowright X$ is properly discontinuous, δ_{Riem} is bounded above by the volume entropy of X which is finite.¹⁴ For the $\bar{\theta}$ -critical exponent, (2.2) implies the following lower bound,

$$(2.5) \quad \delta_{\text{Riem}} \leq \delta_{\bar{\theta}}.$$

Finiteness of $\delta_{\bar{\theta}}$ is more subtle because, in general, $d_{\bar{\theta}}$ is only a pseudo-metric and therefore, the orbital counting function $N_{\bar{\theta}}$ may take infinity as a value. However, if the angular radius of the model Weyl chamber σ_{mod} with respect to $\bar{\theta}$ is $< \pi/2$, then $d_{\bar{\theta}}$ is a metric equivalent to d_{Riem} and, consequently, $\delta_{\bar{\theta}}$ is finite in this case. In particular, when G is simple, then diameter of σ_{mod} is $< \pi/2$ and therefore, $\delta_{\bar{\theta}}$ is finite.

The following finiteness result holds in the general pseudo-metric case.

Proposition 2.6. *For a uniformly τ_{mod} -regular subgroup $\Gamma < G$, the $\bar{\theta}$ -critical exponent $\delta_{\bar{\theta}}$ is finite.*

Proof. When Γ is uniformly τ_{mod} -regular, the distance functions d_{Riem} and $d_{\bar{\theta}}$ restricted to an orbit Γx are coarsely equivalent: There exist $L \geq 1, A \geq 0$ such that, for all $x_1, x_2 \in \Gamma x$,

$$(2.6) \quad L^{-1}d_{\text{Riem}}(x_1, x_2) - A \leq d_{\bar{\theta}}(x_1, x_2) \leq d_{\text{Riem}}(x_1, x_2).$$

The right side of this inequality comes from (2.2). From this we get $\delta_{\text{Riem}} \leq \delta_{\bar{\theta}} \leq L\delta_{\text{Riem}}$. Since δ_{Riem} is finite, $\delta_{\bar{\theta}}$ is also finite. \square

Remark 2.7.

- (1) It is clear from the proof of Proposition 2.6 that when Γ is uniformly τ_{mod} -regular, then $\delta_{\bar{\theta}}$ is positive if and only if δ_{Riem} is positive.
- (2) As Anosov subgroups are uniformly regular (see Theorem 1.7), Proposition 2.6 applies to the class of Anosov subgroups.

¹⁴Finiteness of the volume entropy of a symmetric space follows, for instance, from the fact that X has curvature bounded below combined with the Bishop–Günter volume comparison theorem, see e.g. [6, Sec. 11.10, Cor. 4].

Before closing this section, we compute Finsler distances $d_{\bar{\theta}}$ in two examples.

Example 2.8 (Product of rank-one symmetric spaces). We continue with the discussion from Example 1.8. Let $\tau_{\text{mod}} = (r_1, \dots, r_p)$ be a face of the model chamber, let $\bar{\theta} = (1/\sqrt{p}, \dots, 1/\sqrt{p})$ be its barycenter, and let $d_{\bar{\theta}}$ be the corresponding metric on X . Using the formula for d_{Δ} from (1.3), we get

$$(2.7) \quad d_{\bar{\theta}}(x, y) = \frac{1}{\sqrt{p}} \sum_{j=1}^p d_{r_j}(x_{r_j}, y_{r_j}).$$

Example 2.9 ($X = \text{SL}(k+1, \mathbb{R})/\text{SO}(k+1, \mathbb{R})$). We continue with the discussion from Example 1.9. The Riemannian metric on X is given by the restriction of the Killing form B of $\mathfrak{g} = \mathfrak{sl}(k+1, \mathbb{R})$ to \mathfrak{p} ,

$$(2.8) \quad B(P, Q) = 2(k+1) \text{tr}(PQ), \quad P, Q \in \mathfrak{p}.$$

Note that the inner product B on \mathfrak{a} (which we identify with F_{mod}) can be written as

$$(2.9) \quad \langle (\sigma_1, \dots, \sigma_{k+1}) | (\sigma'_1, \dots, \sigma'_{k+1}) \rangle = 2(k+1) \sum_{i=1}^{k+1} \sigma_i \sigma'_i.$$

Let $\tau_{\text{mod}} = (r_1, \dots, r_p)$ be an ι -invariant face of the model chamber σ_{mod} and let $\Delta_{\tau_{\text{mod}}}$ be the corresponding face of the model euclidean Weyl chamber Δ ,

$$\Delta_{\tau_{\text{mod}}} = \left\{ \sigma \in \mathfrak{a}_+ \mid \sigma = \underbrace{(\sigma_1, \dots, \sigma_1)}_{r_1\text{-times}}, \dots, \underbrace{(\sigma_i, \dots, \sigma_i)}_{(r_i - r_{i-1})\text{-times}}, \dots, \underbrace{(\sigma_{p+1}, \dots, \sigma_{p+1})}_{(k+1-r_p)\text{-times}} \right\}.$$

For notational convenience we denote σ in the above expression simply by the $(p+1)$ -vector $(\sigma_1, \dots, \sigma_{p+1})$ (by identifying the repeated entries). With this convention, the opposition involution acts by

$$\iota(\sigma_1, \dots, \sigma_{p+1}) = (-\sigma_{p+1}, \dots, -\sigma_1).$$

We identify τ_{mod} with the unit sphere (w.r.t. the metric in (2.9)) in $\Delta_{\tau_{\text{mod}}}$ centered at the origin, i.e., τ_{mod} consists of all elements $(\sigma_1, \dots, \sigma_{p+1}) \in \Delta_{\tau_{\text{mod}}}$ satisfying $2(k+1) \sum_{i=1}^{p+1} (r_i - r_{i-1}) \sigma_i^2 = 1$. An element $\bar{\theta} = (\sigma_1, \dots, \sigma_{p+1}) \in \tau_{\text{mod}}$ lies in the interior of τ_{mod} if and only if $\sigma_1 > \dots > \sigma_{p+1}$. Moreover, $\bar{\theta}$ is ι -invariant if and only if $\sigma_i + \sigma_{p+2-i} = 0$ for all $i = 1, \dots, p+1$.

The Finsler distance $d_{\bar{\theta}}$ can be calculated explicitly in terms of the above formulas. In the special case¹⁵ when $\tau_{\text{mod}} = (1, k)$ and

$$\bar{\theta} = (1/2\sqrt{k+1}, 0, -1/2\sqrt{k+1}),$$

the unique ι -invariant type in the interior of τ_{mod} , we have a simple formula for the Finsler distance: For all $g \in \text{SL}(k+1, \mathbb{R})$,

$$(2.10) \quad d_{\bar{\theta}}(x_0, gx_0) = \sqrt{k+1} (\sigma_1(g) - \sigma_{k+1}(g)),$$

where the point x_0 is the identity matrix.

¹⁵We will only focus on this case from now on.

3. $\bar{\theta}$ -CONFORMAL DENSITIES

Recall that Busemann functions define the notion of “distance from infinity.” For $\tau \in \text{Flag}(\tau_{\text{mod}})$, let $b_\tau^{\bar{\theta}} : X \rightarrow \mathbb{R}$ denote the Busemann function based at the ideal point $\bar{\theta}(\tau) \in \partial_\infty X$ normalized at x_0 , i.e., $b_\tau^{\bar{\theta}}(x_0) = 0$. Using Busemann functions, one defines the *Busemann cocycle* as

$$(3.1) \quad \mathcal{B}_\tau^{\bar{\theta}}(x, y) = b_\tau^{\bar{\theta}}(x) - b_\tau^{\bar{\theta}}(y).$$

$\mathcal{B}_\tau^{\bar{\theta}}$ satisfies the *cocycle condition*: For each triple $x, y, z \in X$,

$$(3.2) \quad \mathcal{B}_\tau^{\bar{\theta}}(x, y) + \mathcal{B}_\tau^{\bar{\theta}}(y, z) = \mathcal{B}_\tau^{\bar{\theta}}(x, z).$$

These functions are related to the Finsler distance functions by

$$(3.3) \quad \mathcal{B}_\tau^{\bar{\theta}}(x, y) = \lim_{n \rightarrow \infty} (d_{\bar{\theta}}(x, z_n) - d_{\bar{\theta}}(y, z_n))$$

whenever (z_n) is a sequence in X flag-converging to τ , cf. [26, Prop. 5.43]. Note that

$$-d_{\bar{\theta}}(x, y) \leq \mathcal{B}_\tau^{\bar{\theta}}(x, y) \leq d_{\bar{\theta}}(x, y).$$

Remark 3.1. As usual, the Busemann functions and cocycles depend on the choice of $\bar{\theta}$. Also, note that these functions can take negative values. However, $|\mathcal{B}_\tau^{\bar{\theta}}(x, y)|$ satisfy the triangle inequality and, hence, are pseudo-metrics on X .

We define our notion of “conformal densities” on $\text{Flag}(\tau_{\text{mod}})$ using these Busemann cocycles.

For a topological space S , we let $M_+(S)$ denote the set of Borel probability measures on S . Recall that a group H of self-homeomorphisms of S acts on $M_+(S)$ by *pull-back*: For every $B \in \mathfrak{B}(S)$, $h \in H$,

$$\mu \mapsto h^* \mu, \quad h^* \mu(B) = \mu(h^{-1}(B)).$$

Definition 3.2 ($\bar{\theta}$ -Conformal density). Let $\Gamma < G$ be a discrete subgroup and let $A \subset X$ be a nonempty Γ -invariant subset. By a Γ -invariant $\bar{\theta}$ -conformal A -density μ of dimension $\beta \in [0, \infty)$ on $\text{Flag}(\tau_{\text{mod}})$, we mean a Γ -equivariant map

$$\mu : A \rightarrow M_+(\text{Flag}(\tau_{\text{mod}})), \quad a \mapsto \mu_a,$$

satisfying the following properties:

- (i) For each $a \in A$, $\text{supp}(\mu_a) \subset \Lambda_{\tau_{\text{mod}}}(\Gamma)$.
- (ii) Γ -invariance: μ is Γ -invariant, i.e., $\gamma^* \mu_a = \mu_{\gamma a}$ for each $\gamma \in \Gamma$ and each $a \in A$.
- (iii) $\bar{\theta}$ -conformality: For every pair $a, b \in A$, $\mu_a \ll \mu_b$, i.e., μ_a is absolutely continuous with respect to μ_b , and the Radon–Nikodym derivative $d\mu_a/d\mu_b$ can be expressed as

$$(3.4) \quad \frac{d\mu_a}{d\mu_b}(\tau) = \exp\left(-\beta \mathcal{B}_\tau^{\bar{\theta}}(a, b)\right), \quad \forall \tau \in \text{Flag}(\tau_{\text{mod}}).$$

Remark 3.3.

- (1) In the literature, the item (i) in Definition 3.2 is not required to define Γ -invariant conformal densities. Nevertheless, the uniqueness of such conformal densities for τ_{mod} -Anosov subgroups, which we establish later in the paper (see Corollary 8.4), is *possibly* false unless we impose this extra condition.

- (2) Though we define $\bar{\theta}$ -conformal densities for general discrete subgroups of G , for the purpose of this paper we restrict our discussion only to τ_{mod} -regular subgroups.

A $\bar{\theta}$ -conformal X -density is simply called a $\bar{\theta}$ -conformal density. Note that $\bar{\theta}$ -conformal X -densities and $\bar{\theta}$ -conformal A -densities are in a one-to-one correspondence:

$$(3.5) \quad \{\bar{\theta}\text{-conformal } X\text{-densities}\} \longleftrightarrow \{\bar{\theta}\text{-conformal } A\text{-densities}\}.$$

From an X -density μ , define an A -density by restricting the family. On the other hand, given an A -density μ , extend it to an X -density $\{\mu_x\}_{x \in X}$ by

$$d\mu_x(B) = \int_B \exp\left(-\beta \mathcal{B}_\tau^{\bar{\theta}}(x, a)\right) d\mu_a(\tau), \quad B \in \mathfrak{B}(\text{Flag}(\tau_{\text{mod}})),$$

where μ_a is a density in the family μ . Note that this extension is unique because μ_x and μ_a are absolutely continuous with respect to each other. To check Γ -invariance, note that

$$\begin{aligned} \gamma^* \mu_x(B) &= \int_{\gamma^{-1}B} \frac{d\mu_x}{d\mu_a}(\tau) d\mu_a(\tau) = \int_B \exp\left(-\beta \mathcal{B}_{\gamma^{-1}\tau}^{\bar{\theta}}(x, a)\right) d\mu_a(\gamma^{-1}\tau) \\ &= \int_B \exp\left(-\beta \mathcal{B}_\tau^{\bar{\theta}}(\gamma x, \gamma a)\right) d\mu_{\gamma a}(\tau) = \int_B \frac{d\mu_{\gamma x}}{d\mu_{\gamma a}}(\tau) d\mu_{\gamma a}(\tau) = \mu_{\gamma x}(B), \end{aligned}$$

for every $B \in \mathfrak{B}(\text{Flag}(\tau_{\text{mod}}))$. The other two defining properties are also satisfied.

3.1. The Patterson–Sullivan construction. For every τ_{mod} -regular group Γ with finite $\bar{\theta}$ -critical exponent $\delta_{\bar{\theta}}$, we follow the Patterson–Sullivan construction to construct a $\bar{\theta}$ -conformal density. This construction is standard and already appeared in the work of Albuquerque and Quint, although only in the setting of Zariski dense subgroups $\Gamma < G$ and regular vectors $\bar{\theta}$; we present it here for the sake of completeness. We let $\Gamma < G$ be a τ_{mod} -regular subgroup and let Z denote the Γ -orbit of a point $x_0 \in X$. The union

$$\bar{Z} = Z \cup \Lambda_{\tau_{\text{mod}}}(\Gamma) \subset \bar{X}^{\tau_{\text{mod}}},$$

equipped with the topology of flag-convergence, is a compactification of Z .

For $s > \delta_{\bar{\theta}}$, the $\bar{\theta}$ -critical exponent of Γ , we define a family of positive measures $\mu_s^{\bar{\theta}} = \{\mu_{x,s}^{\bar{\theta}}\}_{x \in X}$ on \bar{Z} by

$$(3.6) \quad \mu_{x,s}^{\bar{\theta}} = \frac{1}{g_s^{\bar{\theta}}(x_0, x_0)} \sum_{\gamma \in \Gamma} \exp(-s d_{\bar{\theta}}(x, \gamma x_0)) D(\gamma x_0),$$

where $D(\gamma x_0)$ denotes the Dirac point mass of weight one at γx_0 . Note that $\mu_{x,s}^{\bar{\theta}}$ is a probability measure when $x \in Z$. Also, note that $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ is a null set for these measures. For $\gamma \in \Gamma$ a straightforward computation shows that

$$(3.7) \quad \gamma^* \mu_{x,s}^{\bar{\theta}} = \mu_{\gamma x, s}^{\bar{\theta}}.$$

Moreover, it is easy to see that $\mu_s^{\bar{\theta}}$ is an absolutely continuous family of measures. Using (3.6) we compute the Radon–Nikodym derivatives $d\mu_{x,s}^{\bar{\theta}}/d\mu_{x_0,s}^{\bar{\theta}}$,

$$(3.8) \quad \psi_{x,x_0,s}^{\bar{\theta}}(z) = \frac{d\mu_{x,s}^{\bar{\theta}}}{d\mu_{x_0,s}^{\bar{\theta}}}(z),$$

where for $s \geq 0$,

$$\psi_{x,x_0,s}^{\bar{\theta}}(z) := \exp(-s(d_{\bar{\theta}}(z,x) - d_{\bar{\theta}}(z,x_0))).$$

The formula for $\psi_{x,x_0,s}^{\bar{\theta}}$ above only makes sense when $z \in Z$. Since $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ is a null set, we extend $\psi_{x,x_0,s}^{\bar{\theta}}$ continuously to $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ by setting

$$\psi_{x,x_0,s}^{\bar{\theta}}(\tau) = \exp\left(-s\mathcal{B}_{\tau}^{\bar{\theta}}(x,x_0)\right).$$

The continuity of this function can be verified using properties of $d_{\bar{\theta}}$ (e.g., see [26, Sec. 5.1.2] and (3.3)).

Next we prove that $\psi_{x,x_0,s}^{\bar{\theta}} \rightarrow \psi_{x,x_0,\delta_{\bar{\theta}}}^{\bar{\theta}}$ uniformly as $s \rightarrow \delta_{\bar{\theta}}$. For $S \geq s, s' > \delta_{\bar{\theta}}$ and $z \in Z$,

$$\begin{aligned} & |\psi_{x,x_0,s'}^{\bar{\theta}}(z) - \psi_{x,x_0,s}^{\bar{\theta}}(z)| \\ &= |\exp(-s'(d_{\bar{\theta}}(z,x) - d_{\bar{\theta}}(z,x_0))) - \exp(-s(d_{\bar{\theta}}(z,x) - d_{\bar{\theta}}(z,x_0)))| \\ &= \exp(-s(d_{\bar{\theta}}(z,x) - d_{\bar{\theta}}(z,x_0))) |\exp((s-s')(d_{\bar{\theta}}(z,x) - d_{\bar{\theta}}(z,x_0))) - 1| \\ &\leq \exp(Sd_{\text{Riem}}(x,x_0)) |\exp((s-s')(d_{\bar{\theta}}(z,x) - d_{\bar{\theta}}(z,x_0))) - 1|. \end{aligned}$$

Switching s and s' in the above, we also get

$$\begin{aligned} & |\psi_{x,x_0,s'}^{\bar{\theta}}(z) - \psi_{x,x_0,s}^{\bar{\theta}}(z)| \\ &\leq \exp(Sd_{\text{Riem}}(x,x_0)) |\exp((s'-s)(d_{\bar{\theta}}(z,x) - d_{\bar{\theta}}(z,x_0))) - 1|. \end{aligned}$$

Combining the above two inequalities, we get

$$\begin{aligned} & |\psi_{x,x_0,s'}^{\bar{\theta}}(z) - \psi_{x,x_0,s}^{\bar{\theta}}(z)| \\ &\leq \exp(Sd_{\text{Riem}}(x,x_0)) (\exp(|s'-s| \cdot |d_{\bar{\theta}}(z,x) - d_{\bar{\theta}}(z,x_0)|) - 1) \\ &\leq \exp(Sd_{\text{Riem}}(x,x_0)) (\exp(|s'-s|d_{\text{Riem}}(x,x_0)) - 1). \end{aligned}$$

Since Z is dense in \bar{Z} , the above yields

$$\|\psi_{x,x_0,s'}^{\bar{\theta}} - \psi_{x,x_0,s}^{\bar{\theta}}\|_{\infty} \leq \exp(Sd_{\text{Riem}}(x,x_0)) (\exp(|s'-s|d_{\text{Riem}}(x,x_0)) - 1).$$

Therefore, $\psi_{x,x_0,s}^{\bar{\theta}} \rightarrow \psi_{x,x_0,\delta_{\bar{\theta}}}^{\bar{\theta}}$ uniformly as $s \rightarrow \delta_{\bar{\theta}}$.

Now we construct a $\bar{\theta}$ -conformal density as a limit of the family of densities $\{\mu_s^{\bar{\theta}}\}_{s>\delta_{\bar{\theta}}}$. We first assume that Γ has $\bar{\theta}$ -divergence type.¹⁶ Then, as s decreases to $\delta_{\bar{\theta}}$, the family $\mu_s^{\bar{\theta}} = \{\mu_{x,s}^{\bar{\theta}}\}_{x \in X}$ weakly accumulates to a density $\mu^{\bar{\theta}}$ supported on some subset of $\Lambda_{\tau_{\text{mod}}}(\Gamma)$. By (3.7) we have the Γ -invariance of $\mu^{\bar{\theta}}$, namely, for $\gamma \in \Gamma$,

$$(3.9) \quad \gamma^* \mu_x^{\bar{\theta}} = \mu_{\gamma x}^{\bar{\theta}}.$$

Moreover, since $\mu_x^{\bar{\theta}}$ is obtained as a weak limit of the measures $\mu_{x,s}^{\bar{\theta}}$ and the derivatives $\psi_{x,x_0,s}^{\bar{\theta}} = d\mu_{x,s}^{\bar{\theta}}/d\mu_{x_0,s}^{\bar{\theta}}$ converge uniformly to $\psi_{x,x_0,\delta_{\bar{\theta}}}^{\bar{\theta}}$, it follows that the Radon-Nikodym derivative $d\mu_x^{\bar{\theta}}/d\mu_{x_0}^{\bar{\theta}}$ exists and equals to the limit

$$\lim_{s \rightarrow \delta_{\bar{\theta}}} \frac{d\mu_{x,s}^{\bar{\theta}}}{d\mu_{x_0,s}^{\bar{\theta}}} = \psi_{x,x_0,\delta_{\bar{\theta}}}^{\bar{\theta}},$$

¹⁶This will be the case for Anosov subgroups. See Corollary 6.5.

or more explicitly,

$$(3.10) \quad \frac{d\mu_x^{\bar{\theta}}}{d\mu_{x_0}^{\bar{\theta}}}(\tau) = \exp\left(-\delta_{\bar{\theta}}\mathcal{B}_{\tau}^{\bar{\theta}}(x, x_0)\right).$$

Note that in general weak limits are not unique. In Corollary 8.4 we will prove that for Anosov subgroups Γ we get a unique density in this limiting process.

When Γ has $\bar{\theta}$ -convergence type, we change weights of the Dirac masses by a small amount ([38, Lem. 3.1], see also [37, Sec. 3.1]) in the definition (3.6) to force the Poincaré series to diverge. Define

$$\mu_{x,s}^{\bar{\theta}} = \frac{1}{\bar{g}_s^{\bar{\theta}}(x_0, x_0)} \sum_{\gamma \in \Gamma} \exp(-sd_{\bar{\theta}}(x, \gamma x_0)) h(d_{\bar{\theta}}(x, \gamma x_0)) D(\gamma x_0),$$

where $h : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a subexponential function such that the following modified Poincaré series

$$\bar{g}_s^{\bar{\theta}}(x, x_0) = \sum_{\gamma \in \Gamma} \exp(-sd_{\bar{\theta}}(x, \gamma x_0)) h(d_{\bar{\theta}}(\gamma x, x_0))$$

diverges at $s = \delta_{\bar{\theta}}$. In this case also, a limit density $\mu^{\bar{\theta}}$ has the properties (3.9) and (3.10).

Definition 3.4 (Patterson-Sullivan density). Let Γ be a τ_{mod} -regular subgroup of G such that $\delta_{\bar{\theta}}(\Gamma) < 0$, and let $\bar{\theta} \in \tau_{\text{mod}}$ be an ι -invariant interior point. Any weak limit $\mu^{\bar{\theta}}$ appearing from the construction above is called a *Patterson-Sullivan density of type $\bar{\theta}$* .

3.2. Positivity of the $\bar{\theta}$ -critical exponent. The existence of a $\bar{\theta}$ -conformal density implies that the $\bar{\theta}$ -critical exponent of Γ is positive.

Proposition 3.5. *Suppose that Γ is a nonelementary τ_{mod} -regular antipodal subgroup and $\delta_{\bar{\theta}}$ is finite. Then, $\delta_{\bar{\theta}}$ is also positive.*

Proof. Suppose to the contrary that $\delta_{\bar{\theta}} = 0$. Let $\mu^{\bar{\theta}}$ be a Patterson–Sullivan density constructed above. It follows from the Γ -invariance and $\bar{\theta}$ -conformality that for all $\gamma \in \Gamma$,

$$(3.11) \quad \mu_x^{\bar{\theta}}(\gamma A) = \mu_{\gamma^{-1}x}^{\bar{\theta}}(A) = \mu_x^{\bar{\theta}}(A), \quad \forall A \in \mathfrak{B}(\Lambda_{\tau_{\text{mod}}}(\Gamma)).$$

We use the convergence action property of a τ_{mod} -RA subgroup (see Subsection 1.6).

We first show that $\mu^{\bar{\theta}}$ is atom-free. For if $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$ were an atom, then, since $\Gamma\tau$ is an infinite orbit, $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ would have infinite $\mu_x^{\bar{\theta}}$ -mass by (3.11).

Moreover, using the converge action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}(\Gamma)$, we have an infinite sequence $\gamma_n \in \Gamma$ and points $\tau_{\pm} \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$ such that on $\Lambda_{\tau_{\text{mod}}}(\Gamma) - \{\tau_{-}\}$,

$$\gamma_n |_{\Lambda_{\tau_{\text{mod}}}(\Gamma) - \{\tau_{-}\}} \rightarrow \text{const}_{\tau_{+}}$$

uniformly on compact sets.

Now, pick a compact set $A \subset \Lambda_{\tau_{\text{mod}}}(\Gamma)$ not containing τ_{\pm} such that $\mu_x^{\bar{\theta}}(A) \geq (2/3)\mu_x^{\bar{\theta}}(\Lambda_{\tau_{\text{mod}}}(\Gamma))$ (this is possible because $\mu^{\bar{\theta}}$ is atom-free). Pick a large enough n so that $\gamma_n(A) \cap A = \emptyset$. Then,

$$\mu_x^{\bar{\theta}}(\Lambda_{\tau_{\text{mod}}}(\Gamma)) \geq \mu_x^{\bar{\theta}}(A \cup \gamma_n A) = \mu_x^{\bar{\theta}}(A) + \mu_x^{\bar{\theta}}(\gamma_n A) = 2\mu_x^{\bar{\theta}}(A) \geq \frac{4}{3}\mu_x^{\bar{\theta}}(\Lambda_{\tau_{\text{mod}}}(\Gamma))$$

yields a contradiction. \square

Remark 3.6. As a corollary to Proposition 3.5, the Riemannian critical exponent δ_{Riem} of a nonelementary uniformly τ_{mod} -regular antipodal subgroup is also non-zero. See the remark after Proposition 2.6.

Since Anosov subgroups are uniformly regular and antipodal, we have the following result.

Corollary 3.7. *Let Γ be a nonelementary τ_{mod} -Anosov subgroup of G . Then,*

$$0 < \delta_{\text{Riem}} \leq \delta_{\bar{\theta}} < \infty.$$

4. HYPERBOLICITY OF MORSE IMAGE

In this section we prove that the image of a τ_{mod} -Morse map is Gromov-hyperbolic with respect to the Finsler pseudo-metric $d_{\bar{\theta}}$. As a corollary, we prove that each orbit of an Anosov subgroup is also Gromov-hyperbolic with respect to $d_{\bar{\theta}}$.

We first recall two notions of hyperbolicity. We shall use the symbol δ to denote the hyperbolicity constant in this and the next sections only. We hope that the reader will not confuse this δ with the critical exponent.

Definition 4.1 (Rips hyperbolic). Let (Z, d) be a geodesic metric space. Then, (Z, d) is called $\delta(\geq 0)$ -hyperbolic in the sense of Rips (or Rips hyperbolic) if every geodesic triangle Δ is δ -thin, i.e., each side of Δ lies in the δ -neighborhood of the union of the other two sides.

Definition 4.2 (Gromov hyperbolic). Let (Z, d) be a metric space. For any three points $z, z_1, z_2 \in Z$, the Gromov product is defined as

$$\langle z_1 | z_2 \rangle_z = \frac{1}{2}[d(z, z_1) + d(z_2, z) - d(z_1, z_2)].$$

Then (Z, d) is called $\delta(\geq 0)$ -hyperbolic in the sense of Gromov (or Gromov hyperbolic) if the Gromov product satisfies the following *ultrametric inequality*: For all $z, z_1, z_2, z_3 \in Z$,

$$\langle z_1 | z_2 \rangle_z \geq \min\{\langle z_1 | z_3 \rangle_z, \langle z_2 | z_3 \rangle_z\} - \delta.$$

It should be noted that Gromov's definition applies to all metric spaces whereas Rips' definition works only for geodesic metric spaces. Moreover, Gromov hyperbolicity is not quasiisometric invariant whereas Rips hyperbolicity is (as a consequence of Morse lemma, cf. [15, Cor. 11.43])). For geodesic metric spaces, these two notions of hyperbolicity are equivalent (e.g., see [15, Lemma 11.27]).

Let (Z', d') be Rips hyperbolic and $f : (Z', d') \rightarrow (X, d_{\text{Riem}})$ be a τ_{mod} -Morse map. We denote the image $f(Z')$ by Z . Recall that the $d_{\bar{\theta}}$ is coarsely equivalent to d_{Riem} on Z .¹⁷ Therefore, since f is a quasiisometric embedding with respect to d_{Riem} , it is also a quasiisometric embedding with respect to $d_{\bar{\theta}}$. Moreover, the image of a geodesic (of length bounded below by a constant) in Z' stays within a uniformly bounded Riemannian distance, say $\lambda_0 \geq 0$, from a τ_{mod} -regular Finsler geodesic connecting the images of the endpoints. This is a consequence of the Morse property [31, Thm. 1.1], see also [26, Prop. 12.2]. A consequence of this is that Z is λ_0 -quasiconvex in $(X, d_{\bar{\theta}})$.

¹⁷This is also true for any finite Riemannian tubular neighborhood of Z .

For $\lambda \geq \lambda_0$, let $Y = Y_\lambda$ be the Riemannian λ -neighborhood of Z in X . From the discussion above, it is clear that any two points $z_1, z_2 \in Z$ (with $d_{\text{Riem}}(z_1, z_2)$ sufficiently large) can be connected by a Finsler geodesic $\widehat{z_1 z_2}$ in Y .

Proposition 4.3. *Let c and c' be two Finsler geodesics in Y connecting two points z_1, z_2 . Then they are uniformly Hausdorff close. Here the Hausdorff distance is induced by either d_{Riem} or $d_{\bar{\theta}}$.*

Proof. Since d_{Riem} or $d_{\bar{\theta}}$ are comparable on Y , it is enough to prove the proposition for the Riemannian metric d_{Riem} .

Let \bar{c} and \bar{c}' be the respective nearest point projections of c and c' to Z . Applying the coarse inverse of f , \bar{c} and \bar{c}' map to uniform quasigeodesics \tilde{c} and \tilde{c}' , respectively, in Z' . Since Z' is Rips hyperbolic, \tilde{c} and \tilde{c}' are uniformly close. Applying f to \tilde{c} and \tilde{c}' , we see that \bar{c} and \bar{c}' are uniformly close. Hence c and c' are also uniformly close. \square

Next we observe that geodesic triangles in $(Y, d_{\bar{\theta}})$ with vertices on Z are uniformly thin.

Proposition 4.4. *There exists $\delta \geq 0$ such that every Finsler geodesic triangle $\Delta = \Delta(z_1, z_2, z_3)$ in Y is δ -thin both in Riemannian and Finsler sense.*

Proof. Since Z' is Rips hyperbolic, geodesic triangles in Z' are δ' -thin, for some $\delta' \geq 0$. We map Δ to a uniformly quasigeodesic triangle $\Delta' \subset Z'$ via the coarse inverse map $Y \rightarrow Z'$ of the map f . Since Z' is Rips-hyperbolic, the Morse quasigeodesic triangle Δ' is uniformly thin. Therefore, Δ is also uniformly thin as well. \square

Imitating the proof of [15, Lem. 11.27], we prove that $(Z, d_{\bar{\theta}})$ is Gromov-hyperbolic.

Theorem 4.5 (Hyperbolicity of Morse maps). *Let $Z \subset X$ be the image of a τ_{mod} -Morse map $f : (Z', d') \rightarrow (X, d_{\text{Riem}})$. Then $(Z, d_{\bar{\theta}})$ is Gromov-hyperbolic.*

Proof. Let δ be as in Proposition 4.4. Then the following holds.

Lemma 4.6. *Let $z, z_1, z_2 \in Z$, and let $\widehat{z_1 z_2}$ be any Finsler geodesic in Y connecting z_1 and z_2 . Then,*

$$\langle z_1 | z_2 \rangle_z \leq d_{\bar{\theta}}(z, \widehat{z_1 z_2}) \leq \langle z_1 | z_2 \rangle_z + 2\delta.$$

Proof. The proof is exactly same as [15, Lem. 11.22]. Note that the proof uses δ -thinness of a triangle with vertices z, z_1, z_2 . \square

Let z, z_1, z_2, z_3 be any four points in Z , and let Δ be a Finsler geodesic triangle in Y with the vertices z_1, z_2, z_3 . Let m be a point on the side $\widehat{z_1 z_2}$ nearest to z . By Proposition 4.4, since Δ is δ -thin, $d_{\bar{\theta}}(m, \widehat{z_2 z_3} \cup \widehat{z_1 z_3}) \leq \delta$. Without loss of generality, assume that there is a point n on z_2, z_3 which is δ -close to m . Then, using Lemma 4.6, we get

$$\langle z_2 | z_3 \rangle_z \leq d_{\bar{\theta}}(z, \widehat{z_2 z_3}) \leq d_{\bar{\theta}}(z, \widehat{z_1 z_2}) + \delta,$$

and

$$d_{\bar{\theta}}(z, \widehat{z_1 z_2}) \leq \langle z_1 | z_2 \rangle_z + 2\delta.$$

The theorem follows from this. \square

Quasiisometry of hyperbolic metric spaces extends to a homeomorphism of their Gromov boundaries. At the same time, it is proven in [31, Thm. 1.4] that each τ_{mod} -Morse map

$$f : Z' \rightarrow Z = f(Z') \subset X$$

extends continuously (with respect to the topology of flag-convergence) to a homeomorphism

$$\partial_\infty f : \partial_\infty Z' \rightarrow \Lambda \subset \text{Flag}(\tau_{\text{mod}}).$$

Thus, we obtain

Corollary 4.7. *The Gromov boundary $\partial_\infty Z$ of $(Z, d_{\bar{\theta}})$ is naturally identified with the flag-limit set $\Lambda \subset \text{Flag}(\tau_{\text{mod}})$ of Z : A sequence (z_n) in Z converges to a point in $\partial_\infty Z$ if and only if (z_n) flag-converges to some $\tau \in \Lambda$.*

For a τ_{mod} -Anosov subgroup Γ we know that the orbit map $\Gamma \rightarrow \Gamma x_0$ is a τ_{mod} -Morse embedding (see Subsection 1.6). Then, using Theorem 4.5 we obtain:

Corollary 4.8 (Hyperbolicity of Anosov orbits). *For $x_0 \in X$, let $Z = \Gamma x_0$ where Γ is a τ_{mod} -Anosov subgroup. Then $(Z, d_{\bar{\theta}})$ is Gromov hyperbolic. The Gromov boundary of $(Z, d_{\bar{\theta}})$ is naturally identified with the τ_{mod} -limit set $\Lambda_{\tau_{\text{mod}}}(\Gamma)$.*

5. GROMOV DISTANCE AT INFINITY

For a pair of antipodal simplices $\tau_\pm \in \text{Flag}(\tau_{\text{mod}})$, the *Gromov product* with respect to a base point $x \in X$ is defined as

$$(5.1) \quad \langle \tau_+ | \tau_- \rangle_x^{\bar{\theta}} = \frac{1}{2} \left(\mathcal{B}_{\tau_+}^{\bar{\theta}}(x, z) + \mathcal{B}_{\tau_-}^{\bar{\theta}}(x, z) \right),$$

where z is some point on the parallel set $P(\tau_+, \tau_-)$ spanned by τ_\pm .

The definition Busemann cocycles $\mathcal{B}_\tau^{\bar{\theta}}$ given in (3.1) is free of choice of any particular normalization for the Busemann functions. We use this observation in the proof of Lemma 5.1 which shows that the Gromov products do not depend on the chosen $z \in P(\tau_+, \tau_-)$.

Lemma 5.1. *For $z_1, z_2 \in P(\tau_+, \tau_-)$, one has $b_{\tau_+}^{\bar{\theta}}(z_1) + b_{\tau_-}^{\bar{\theta}}(z_1) = b_{\tau_+}^{\bar{\theta}}(z_2) + b_{\tau_-}^{\bar{\theta}}(z_2)$.*

Proof. Let z be the midpoint of $\overline{z_1 z_2}$ and let $s_z : X \rightarrow X$ be the point reflection about z . Assuming that Busemann functions are normalized at z , s_z transforms $b_{\tau_+}^{\bar{\theta}}(z_1) + b_{\tau_-}^{\bar{\theta}}(z_1)$ into $b_{\tau_-}^{\bar{\theta}}(z_2) + b_{\tau_+}^{\bar{\theta}}(z_2)$. Hence the quantities are equal. \square

Using the Gromov product, we define a *premetric*¹⁸ on $\text{Flag}(\tau_{\text{mod}})$.

Definition 5.2 (Gromov premetric). Given fixed $x \in X$, $\epsilon > 0$, define the *Gromov premetric* $D_x^{\bar{\theta}, \epsilon}$ on $\text{Flag}(\tau_{\text{mod}})$ as

$$D_x^{\bar{\theta}, \epsilon}(\tau_1, \tau_2) = \begin{cases} \exp\left(-\epsilon \langle \tau_1 | \tau_2 \rangle_x^{\bar{\theta}}\right), & \text{if } \tau_1, \tau_2 \text{ are antipodal,} \\ 0, & \text{otherwise.} \end{cases}$$

Remark 5.3. A pair of points $\tau_\pm \in \text{Flag}(\tau_{\text{mod}})$ is antipodal if and only if

$$D_x^{\bar{\theta}, \epsilon}(\tau_+, \tau_-) \neq 0.$$

Lemma 5.4. *$D_x^{\bar{\theta}, \epsilon}$ is a continuous function.*

¹⁸A *premetric* on X is a symmetric, continuous function $d : X \times X \rightarrow [0, \infty)$ such that $d(x, x) = 0$ for all $x \in X$.

Proof. The claim follows from [3, Lem. 3.8]. \square

Definition 5.5 (Conformal maps). Let (Z, d) be a premetric space. A self-homeomorphism $f : (Z, d) \rightarrow (Z, d)$ is called *K-quasiconformal* if, for every $z \in Z$, $\limsup_{r \rightarrow 0} H_f(z, r) \leq K$, where

$$H_f(z, r) := \frac{\sup\{d(f(y), f(z)) \mid d(y, z) \leq r\}}{\inf\{d(f(y), f(z)) \mid d(y, z) \geq r\}}.$$

The map f is called *conformal* if it is 1-quasiconformal.

Lemma 5.6. *Let $\gamma \in G$ and $\Lambda \subset \text{Flag}(\tau_{\text{mod}})$ be a γ -invariant antipodal subset. Then the map $\gamma : \Lambda \rightarrow \Lambda$ is conformal with respect to the premetric $D_x^{\bar{\theta}, \epsilon}$.*

Proof. Given distinct points $\tau_{\pm} \in \Lambda$,

$$\begin{aligned} D_x^{\bar{\theta}, \epsilon}(\gamma\tau_+, \gamma\tau_-) &= \exp\left(-\epsilon\langle\gamma\tau_+ | \gamma\tau_-\rangle_x^{\bar{\theta}}\right) \\ &= \exp\left(-\frac{\epsilon}{2}\left(\mathcal{B}_{\gamma\tau_+}^{\bar{\theta}}(x, z) + \mathcal{B}_{\gamma\tau_-}^{\bar{\theta}}(x, z)\right)\right) \\ &= \exp\left(-\frac{\epsilon}{2}\left(\mathcal{B}_{\tau_+}^{\bar{\theta}}(\gamma^{-1}x, \gamma^{-1}z) + \mathcal{B}_{\tau_-}^{\bar{\theta}}(\gamma^{-1}x, \gamma^{-1}z)\right)\right) \\ &= \exp\left(-\frac{\epsilon}{2}\left(\mathcal{B}_{\tau_+}^{\bar{\theta}}(\gamma^{-1}x, x) + \mathcal{B}_{\tau_-}^{\bar{\theta}}(\gamma^{-1}x, x)\right)\right) D_x^{\bar{\theta}, \epsilon}(\tau_+, \tau_-), \end{aligned}$$

where the last equality follows from the cocycle condition (3.2). Moreover, the continuity of Busemann functions $b_{\tau}^{\bar{\theta}}$ as a function of τ implies that

$$\lim_{\tau_- \rightarrow \tau_+} \mathcal{B}_{\tau_-}^{\bar{\theta}}(\gamma^{-1}x, x) = \mathcal{B}_{\tau_+}^{\bar{\theta}}(\gamma^{-1}x, x).$$

Therefore,

$$(5.2) \quad \lim_{\tau_- \rightarrow \tau_+} \frac{D_x^{\bar{\theta}, \epsilon}(\gamma\tau_+, \gamma\tau_-)}{D_x^{\bar{\theta}, \epsilon}(\tau_+, \tau_-)} = E(\gamma, \tau_+) := \exp\left(-\epsilon\mathcal{B}_{\tau_+}^{\bar{\theta}}(\gamma^{-1}x, x)\right).$$

From this, it can be checked that $\limsup_{r \rightarrow 0} H_{\gamma}(\tau_+, r) = 1$. \square

The premetric $D_x^{\bar{\theta}, \epsilon}$ is not a metric in general since:

- (i) Pairs of distinct non-antipodal points have zero distance.
- (ii) The triangle inequality may fail.

However, as we shall see below, for all sufficiently small $\epsilon > 0$, $D_x^{\bar{\theta}, \epsilon}$ is bilipschitz equivalent to an actual distance function when restricted to “nice” antipodal subsets $\Lambda \subset \text{Flag}(\tau_{\text{mod}})$.

Theorem 5.7. *Let $Z \subset X$ be the image of a τ_{mod} -Morse map $f : (Z', d') \rightarrow (X, d)$, and let $\Lambda \subset \text{Flag}(\tau_{\text{mod}})$ be the flag limit set of Z . There exists $\epsilon_0 > 0$ such that, for all $0 < \epsilon \leq \epsilon_0$ and all $x \in Z$, the premetric $D_x^{\bar{\theta}, \epsilon}$ is 2-bilipschitz equivalent to a metric on Λ . Moreover, the topology induced by $D_x^{\bar{\theta}, \epsilon}$ on Λ coincides with the subspace topology of $\Lambda \subset \text{Flag}(\tau_{\text{mod}})$.*

Proof. The idea of the proof of the first part is due to Gromov [22]: We show that the Gromov product defined in (5.1) restricted to Λ satisfies an ultrametric inequality (see (5.7)).

Let $Y \subset X$ be a Riemannian λ -neighborhood of Z . We assume that λ here is so large such that $x \in Y$ and the image of any complete geodesic l in Z' lies within distance λ from the parallel set spanned by the images of the ideal endpoints of l

under $\bar{f} : \partial_\infty Z' \rightarrow \text{Flag}(\tau_{\text{mod}})$. Note that λ satisfying the last condition exists as a consequence of the Morse property.

Observe that $(Y, d_{\bar{\theta}})$ is a Gromov δ -hyperbolic¹⁹ metric space for some $\delta \geq 0$. This follows from the Gromov hyperbolicity of $(Z, d_{\bar{\theta}})$ (cf. Theorem 4.5) and the fact that Z and Y are (Hausdorff) λ -close to each other.

We recall from Väisälä [50, Sec. 5] that there are multiple ways to define Gromov products on Λ viewed as the Gromov boundary of $(Z, d_{\bar{\theta}})$ and, hence, of $(Y, d_{\bar{\theta}})$. For a distinct pair $\tau_\pm \in \Lambda$, define using the Gromov product $\langle \cdot | \cdot \rangle_x^{\bar{\theta}}$ on $(Y, d_{\bar{\theta}})$ the following two products:

$$\langle \tau_+ | \tau_- \rangle_x^{\text{inf}} = \inf \left\{ \liminf_{i, j \rightarrow \infty} \langle y_i^+ | y_j^- \rangle_x^{\bar{\theta}} \mid (y_n^\pm) \subset Y, y_n^\pm \rightarrow \tau_\pm \right\}$$

and

$$\langle \tau_+ | \tau_- \rangle_x^{\text{sup}} = \sup \left\{ \limsup_{i, j \rightarrow \infty} \langle y_i^+ | y_j^- \rangle_x^{\bar{\theta}} \mid (y_n^\pm) \subset Y, y_n^\pm \rightarrow \tau_\pm \right\}.$$

Then the difference of the above two quantities is uniformly bounded (see [50, Lemma 5.6]), namely, for all distinct pairs $\tau_\pm \in \Lambda$,

$$(5.3) \quad 0 \leq \langle \tau_+ | \tau_- \rangle_x^{\text{sup}} - \langle \tau_+ | \tau_- \rangle_x^{\text{inf}} \leq 2\delta.$$

Finally, $\langle \cdot | \cdot \rangle_x^{\text{inf}}$ satisfies the ultrametric inequality (see [50, 5.12]), i.e., for distinct triples $\tau_1, \tau_2, \tau_3 \in \Lambda$,

$$(5.4) \quad \langle \tau_1 | \tau_2 \rangle_x^{\text{inf}} \geq \min \left\{ \langle \tau_1 | \tau_3 \rangle_x^{\text{inf}}, \langle \tau_2 | \tau_3 \rangle_x^{\text{inf}} \right\} - \delta.$$

By (5.3), $\langle \cdot | \cdot \rangle_x^{\text{sup}}$ also satisfies the ultrametric inequality but with a different constant, 5δ .

Next we compare Väisälä's Gromov products with ours (see (5.1)). Let $\tau_\pm \in \Lambda$ be a pair of antipodal points and let $P = P(\tau_+, \tau_-)$. Note that our assumption on largeness of λ implies that there exist uniformly τ_{mod} -regular sequences (y_n^+) and (y_n^-) on $Y \cap P$ such that $y_n^\pm \rightarrow \tau_\pm$ as $n \rightarrow \infty$. Let $p \in P(\tau_+, \tau_-)$. Then, the additivity of Finsler distances $d_{\bar{\theta}}$ on τ_{mod} -cones (cf. [26, Lem. 5.10]) yields, for large n , $\langle y_n^+ | y_n^- \rangle_p^{\bar{\theta}} = 0$. By definition,

$$\langle y_n^+ | y_n^- \rangle_x^{\bar{\theta}} = \langle y_n^+ | y_n^- \rangle_p^{\bar{\theta}} + \frac{1}{2} \left[(d_{\bar{\theta}}(x, y_n^+) - d_{\bar{\theta}}(p, y_n^+)) + (d_{\bar{\theta}}(y_n^-, x) - d_{\bar{\theta}}(y_n^-, p)) \right],$$

and for large n ,

$$(5.5) \quad \langle y_n^+ | y_n^- \rangle_x^{\bar{\theta}} = \frac{1}{2} \left[(d_{\bar{\theta}}(x, y_n^+) - d_{\bar{\theta}}(p, y_n^+)) + (d_{\bar{\theta}}(y_n^-, x) - d_{\bar{\theta}}(y_n^-, p)) \right].$$

The limit, as $n \rightarrow \infty$, of the right side of this equation equals $\langle \tau_+ | \tau_- \rangle_x^{\bar{\theta}}$ (cf. (3.3)). Therefore,

$$(5.6) \quad \langle \tau_+ | \tau_- \rangle_x^{\text{inf}} \leq \langle \tau_+, \tau_- \rangle_x^{\bar{\theta}} \leq \langle \tau_+ | \tau_- \rangle_x^{\text{sup}}.$$

Hence, by (5.3) and (5.4), $\langle \cdot | \cdot \rangle_x^{\bar{\theta}}$ satisfies the ultrametric inequality with constant 5δ , i.e., for pairwise distinct points $\tau_1, \tau_2, \tau_3 \in \Lambda$,

$$(5.7) \quad \langle \tau_1 | \tau_2 \rangle_x^{\bar{\theta}} \geq \min \left\{ \langle \tau_1 | \tau_3 \rangle_x^{\bar{\theta}}, \langle \tau_2 | \tau_3 \rangle_x^{\bar{\theta}} \right\} - 5\delta.$$

¹⁹Since there is a possibility of confusion, we would like to remind our reader that this δ is *not* a critical exponent.

Applying [50, Prop. 5.16], we get that, for all $0 < \epsilon \leq \epsilon_0$, $D_x^{\bar{\theta}, \epsilon}$ is 2-bilipschitz equivalent to an actual metric²⁰ on Λ . Here the constant ϵ_0 depends only on δ . This completes the proof of the first part of the theorem.

For the second part, note that the inequality (5.6) implies that $D_x^{\bar{\theta}, \epsilon}$ induces the standard topology on Λ as the Gromov boundary of $(Y, d_{\bar{\theta}})$ (see [50, 5.29]). Since, as we noted earlier, this topology is the same as the subspace topology of the flag-manifold $\text{Flag}(\tau_{\text{mod}})$, the second claim of the theorem follows as well. \square

Corollary 5.8 (Conformal metric on Anosov limit set). *Let Γ be a τ_{mod} -Anosov subgroup, $x \in X$. Then there exists $\epsilon_0 > 0$ such that the following holds: Let $0 < \epsilon \leq \epsilon_0$. Then, for all $z \in \Gamma x$, the premetric $D_x^{\bar{\theta}, \epsilon}$ is bilipschitz equivalent to an actual metric on $\Lambda_{\tau_{\text{mod}}}(\Gamma)$.*

Moreover, the action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is conformal with respect to $D_z^{\bar{\theta}, \epsilon}$.

Proof. Since by Theorem 1.7 Anosov subgroups satisfy the Morse property, corollary follows from Theorem 5.7 combined with Lemma 5.6. \square

Example 5.9 (Product of rank-one symmetric spaces). We continue with Example 2.8. Let $\tau = (\xi_{r_1}, \dots, \xi_{r_p})$ be a simplex in the Tits building of type $\tau_{\text{mod}} = (r_1, \dots, r_p)$ and $\bar{\theta} = (1/\sqrt{p}, \dots, 1/\sqrt{p}) \in \tau_{\text{mod}}$. We compute the Busemann cocycle and Gromov distance associated with τ_{mod} and type $\bar{\theta}$.

Let $x = (x_1, \dots, x_k)$, $y = (y_1, \dots, y_k) \in X$. Then

$$\mathcal{B}_{\tau}^{\bar{\theta}}(x, y) = \lim_{t \rightarrow \infty} (d_{\text{Riem}}(\ell(t), x) - t),$$

where $\ell(t)$ is a geodesic ray emanating from y and asymptotic to $\bar{\theta}(\tau)$. A direct computation yields

$$\mathcal{B}_{\tau}^{\bar{\theta}}(x, y) = \frac{1}{\sqrt{p}} \sum_{j=1}^p (b_{\xi_{r_j}}(x_{r_j}) - b_{\xi_{r_j}}(y_{r_j})) = \frac{1}{\sqrt{p}} \sum_{j=1}^p \mathcal{B}_{\xi_{r_j}}(x_{r_j}, y_{r_j}).$$

Hence the Gromov product can be written as

$$\langle \tau_+ | \tau_- \rangle_x^{\bar{\theta}} = \frac{1}{\sqrt{p}} \sum_{j=1}^p \langle \xi_{r_j}^+ | \xi_{r_j}^- \rangle_{x_{r_j}}, \quad \forall \tau_{\pm} = (\xi_{r_1}^{\pm}, \dots, \xi_{r_p}^{\pm}) \in \text{Flag}(\tau_{\text{mod}}),$$

and, finally the Gromov predistance is

$$(5.8) \quad D_x^{\bar{\theta}, \frac{1}{\sqrt{p}}}(\tau_+, \tau_-) = \prod_{j=1}^p D_{x_{r_j}}^{\bar{\theta}, \frac{1}{p}}(\xi_{r_j}^+, \xi_{r_j}^-).$$

Example 5.10 ($X = \text{SL}(k+1, \mathbb{R})/\text{SO}(k+1, \mathbb{R})$). In this case the computations of Busemann functions (see [24]) are explicitly known, and therefore, the Gromov distance can also be computed explicitly. We only give a formula for the Gromov distance in the special case when $\tau_{\text{mod}} = (1, k)$ that corresponds to the partial flags $\{\text{line} \subset \text{hyperplane}\}$ of \mathbb{R}^{k+1} .

We continue with the notations from Example 2.9. The unique ι -invariant type is

$$\bar{\theta} = \left(\frac{1}{2\sqrt{k+1}}, 0, -\frac{1}{2\sqrt{k+1}} \right).$$

²⁰This metric can be constructed as follows: Let $0 < \epsilon \leq \epsilon_0$. On Λ , define $\text{dist}_x^{\bar{\theta}, \epsilon}(\tau, \tau') = \inf \sum_{i=1}^k D_x^{\bar{\theta}, \epsilon}(\tau_{i-1}, \tau_i)$, where the infimum is taken over all finite sequences $\tau = \tau_0, \tau_1, \dots, \tau_k = \tau'$ on Λ . See [50, 5.13].

After equipping \mathbb{R}^{k+1} with the inner product induced by the choice of $x \in X$, the Gromov product (with respect to $x = I_{k+1}$, the identity matrix) can be written as

$$\langle (l_1, h_1) \mid (l_2, h_2) \rangle_x^{\bar{\theta}} = -\frac{\sqrt{k+1}}{2} \log(\sin \angle(l_1, h_2) \cdot \sin \angle(l_2, h_1)),$$

where $\angle(l, h)$ denotes the angle between the line l and the hyperplane h .²¹

Thus, the Gromov predistance formula is

$$(5.9) \quad D_x^{\bar{\theta}, \frac{1}{\sqrt{k+1}}}((l_1, h_1), (l_2, h_2)) = \sqrt{\sin \angle(l_1, h_2)} \sqrt{\sin \angle(l_2, h_1)}.$$

6. SHADOW LEMMA

In this section we prove a generalization Sullivan's shadow lemma in higher rank. The proof we present here is inspired by that of Albuquerque's [1, Thm. 3.3] who treated the case of full flag manifold and Quint [43] who treated general flag-manifolds but only in the case of regular vectors $\bar{\theta}$.

Recall the notion of *shadow* from Definition 1.3. We mainly consider shadows of closed balls (with respect to the Riemannian metric) of non-zero radii in X from a fixed base point $x \in X$. The topology generated by these shadows is the topology of flag convergence. See Remark 1.4.

The main result in this section is the following.

Theorem 6.1 (Shadow lemma). *Let Γ be a nonelementary τ_{mod} -RA subgroup, $x \in X$, and μ a Γ -invariant $\bar{\theta}$ -conformal density of dimension β . There exists $r_0 > 0$ such that for all $r \geq r_0$ and all $\gamma \in \Gamma$ satisfying $d_{\text{Riem}}(x, \gamma x_0) > r$,*

$$\mu_x(S(x : B(\gamma x_0, r))) \asymp \exp(-\beta d_{\bar{\theta}}(x, \gamma x_0)).$$

Here the notation \asymp means that the ratio of the two sides is bounded above and below by positive constants.

It is worth emphasizing that this version of the Shadow lemma is valid for all τ_{mod} -RA subgroups of G which is a much larger class than τ_{mod} -Anosov subgroups. For instance, the relatively τ_{mod} -Morse subgroups of G , a higher rank generalization of the rank-one geometrically finite groups, are also τ_{mod} -RA.

Before presenting the proof, we note two consequences of this theorem.

Corollary 6.2. *Let Γ be a nonelementary uniformly τ_{mod} -RA subgroup. Then any $\bar{\theta}$ -conformal density μ does not have conical limit points as atoms.*

Proof. We observe that any conical limit point $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$ lies in infinitely many shadows $S(x, B(\gamma x_0, r))$ for sufficiently large $r > 0$ (depending on τ). If τ is an atom, then (by Theorem 6.1) the Poincaré series

$$(6.1) \quad g_{\beta}^{\bar{\theta}}(x, x_0) = \sum_{\gamma \in \Gamma} \exp(-\beta d_{\bar{\theta}}(x, \gamma x_0))$$

diverges for every $\beta \geq 0$. Hence $\delta_{\bar{\theta}}$ must be infinite. But this contradicts Proposition 2.6. \square

²¹This formula can be extracted directly from the formula of the Gromov products in Beyrer's paper [3] by applying it to our special case. Note that the determinant of the matrix in the claim of [3, Appendix] is simply the sine of the angle between the line and hyperplane here. Careful readers may notice some discrepancy between the normalizing constant in our formula and the one in Beyrer's work. This has happened due to our choice of the Riemannian metric in X , which was obtained from the Killing form of \mathfrak{g} . In Beyrer's paper, the quantity λ (which is $\bar{\theta}$ in our paper) is not a unit vector of this metric. Also, compare with [24].

The second application of shadow lemma will be given for the following class of subgroups.

Definition 6.3 (Uniformly conical). A τ_{mod} -RA subgroup is called *uniformly conical* if for a given pair of points $x, x_0 \in X$, there is a constant $r > 0$ such that for each conical limit point $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$, there exists a sequence (γ_k) on Γ flag-converging to τ satisfying $d_{\text{Riem}}(\gamma_k x_0, V(x, \text{st}(\tau))) < r, \forall k \in \mathbb{N}$.

We observe that Anosov subgroups satisfy the uniform conicality condition:

Proposition 6.4. *Anosov subgroups are uniformly conical.*

Proof. This follows from the fact that the orbit map $\Gamma \rightarrow \Gamma x_0 \subset X$ is a Morse embedding. Let $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$ be any point and $\xi \in \partial_\infty \Gamma$ be the preimage of τ under the boundary map. Let $(\gamma_k), \gamma_1 = 1_\Gamma$ be a geodesic sequence in Γ asymptotic to ξ . Then the sequence $(\gamma_k x_0)$ is a Morse quasigeodesic in X that is uniformly close to $V(x, \text{st}(\tau))$ (by definition of a Morse embedding). \square

Corollary 6.5. *Let Γ be a nonelementary uniformly conical τ_{mod} -RA subgroup and μ be a $\bar{\theta}$ -conformal density of dimension β . If the conical limit set $\Lambda_{\tau_{\text{mod}}}^{\text{con}}(\Gamma)$ is non-null, then the Poincaré series $g_{\bar{\theta}}^\beta(x, x_0)$ (see (6.1)) diverges.*

For τ_{mod} -Anosov subgroups Γ , Theorem 1.7 implies that $\Lambda_{\tau_{\text{mod}}}^{\text{con}}(\Gamma) = \Lambda_{\tau_{\text{mod}}}(\Gamma)$. Hence the above result applies to all Anosov subgroups with $\Lambda_{\tau_{\text{mod}}}^{\text{con}}(\Gamma)$ replaced by $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ in the statement.

Proof of Corollary 6.5. Writing the elements of Γ in a sequence (γ_n) , define

$$S_N = \sum_{n \geq N} \exp(-\beta d_{\bar{\theta}}(x, \gamma_n x_0)).$$

Convergence of the series (6.1) asserts that $\lim_{N \rightarrow \infty} S_N = 0$. Since Γ is uniformly conical, there exists $r > 0$ such that for all $N \in \mathbb{N}$,

$$\Lambda_{\tau_{\text{mod}}}^{\text{con}}(\Gamma) \subset \bigcup_{n \geq N} S(x : B(\gamma_n x_0, r)).$$

Applying Theorem 6.1, we get

$$\mu_x(\Lambda_{\tau_{\text{mod}}}(\Gamma)) \leq \sum_{n \geq N} \mu_x(S(x : B(\gamma_n x_0, r))) \leq \text{const} \cdot S_N$$

and, the bound above approaches to zero as $N \rightarrow \infty$. Hence we must have $\mu_x(\Lambda_{\tau_{\text{mod}}}^{\text{con}}(\Gamma)) = 0$. \square

The proof of shadow lemma occupies the rest of the section.

Proof of Theorem 6.1. In this proof, we equip $\text{Flag}(\tau_{\text{mod}})$ with a G_x -invariant Riemannian metric. We use the notation $L(\tau)$ to denote the set of all $\tau' \in \text{Flag}(\tau_{\text{mod}})$ which are not antipodal to τ . The complement of $L(\tau)$ in $\text{Flag}(\tau_{\text{mod}})$ is denoted by $C(\tau)$. Note that $L(\tau)$ is closed and hence, compact.

Moreover, if $\tau_n \rightarrow \tau_0$, then the sets $L(\tau_n)$ Hausdorff-converge to $L(\tau_0)$ as $n \rightarrow \infty$. This can be seen as follows. First, we note that $gL(\tau) = L(g\tau)$ for all $g \in G$ and $\tau \in \text{Flag}(\tau_{\text{mod}})$. This is a consequence of the fact that the action $G \curvearrowright \text{Flag}(\tau_{\text{mod}})$ preserves antipodality, i.e., $\tau, \tau' \in \text{Flag}(\tau_{\text{mod}})$ are antipodal if and only if $g\tau$ and

$g\tau'$ are antipodal, $g \in G$. Now, choose a sequence $k_n \in G_x \cong K$ such that $k_n \rightarrow 1_G$ and $k_n\tau_0 = \tau_n$. Since $\text{Flag}(\tau_{\text{mod}})$ is compact and each $L(\tau_n)$, $n \geq 0$, is closed,

$$L(\tau_n) = k_n L(\tau_0) \xrightarrow{\text{Hausdorff}} L(\tau_0).$$

Lemma 6.6. *For every $\varepsilon > 0$, there exists $\delta > 0$ such that, for every $\tau_0 \in \text{Flag}(\tau_{\text{mod}})$ and every $\tau \in B(\tau_0, \delta)$,*

$$N_{\varepsilon/2}(L(\tau)) \subset N_\varepsilon(L(\tau_0)).$$

Proof. We equip the set

$$Y = \{L(\tau) : \tau \in \text{Flag}(\tau_{\text{mod}})\}$$

with the Hausdorff distance d_H . Then, as we noted above, the function $f : \text{Flag}(\tau_{\text{mod}}) \rightarrow Y$, $\tau \mapsto L(\tau)$, is continuous and, hence, uniformly continuous. Therefore, for every $\varepsilon > 0$, there exists $\delta > 0$ such that $d(\tau, \tau_0) < \delta$ implies $d_H(L(\tau), L(\tau_0)) < \varepsilon/2$, which then implies $L(\tau) \subset N_{\varepsilon/2}(L(\tau_0))$. The lemma follows from this. \square

Let $\mathfrak{m} = \mu_x(\Lambda_{\tau_{\text{mod}}}(\Gamma))$ denote the total mass of μ_x , and $\mathfrak{l} = \sup\{\mu_x(\tau) \mid \tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)\}$. Since μ_x is a regular measure and $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ is compact, \mathfrak{l} is realized, i.e., if μ_x has an atomic part, then it has a largest atom. Moreover, since Γ is nonelementary, $\text{supp}(\mu_x)$ is not singleton. In fact, if τ is an atom, then the every point in the orbit $\Gamma\tau$ (which has infinite number of points) is an atom. In particular, $\mathfrak{l} < \mathfrak{m}$.

Lemma 6.7. *Given $\mathfrak{l} < q < \mathfrak{m}$, there exists an $\varepsilon_0 > 0$ such that for all $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$ and all $B \in \mathfrak{B}(\text{Flag}(\tau_{\text{mod}}))$ contained in $N_{\varepsilon_0}(L(\tau))$, $\mu_x(B) \leq q$.*

Proof. If this were false, then we would get a sequence (B_n) of Borel sets, a sequence (ε_n) of positive numbers converging to zero, and a sequence (τ_n) on $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ converging to a point τ_0 such that for every $n \in \mathbb{N}$,

$$B_n \subset N_{\varepsilon_n}(L(\tau_n)), \quad \mu_x(B_n) > q.$$

To get a contradiction, we will show that $\mu_x(\tau_0) \geq q$. Let U be an open neighborhood of $L(\tau_0)$. As $L(\tau_0)$ is compact, there exists $\varepsilon > 0$ such that $N_\varepsilon(L(\tau_0)) \subset U$. Let $\delta > 0$ be a number that corresponds to this ε as in Lemma 6.6. Choose n so large such that $\tau_n \in B(\tau_0, \delta)$ and $\varepsilon_n \leq \varepsilon/2$. By Lemma 6.6, we get $N_{\varepsilon_n}(L(\tau_n)) \subset N_\varepsilon(L(\tau_0))$ and, consequently, $B_n \subset U$. This shows that every open set U containing $L(\tau_0)$ has mass $\mu_x(U) > q$. In particular, $\mu_x(L(\tau_0)) \geq q$.

Finally, as a consequence of the fact that Γ is τ_{mod} -antipodal, we have that $\Lambda_{\tau_{\text{mod}}}(\Gamma) \cap L(\tau_0) = \tau_0$. Since the support of μ_x is contained in $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ (see Definition 3.2),

$$\mu_x(L(\tau_0)) = \mu_x(\Lambda_{\tau_{\text{mod}}}(\Gamma) \cap L(\tau_0)) = \mu_x(\tau_0).$$

The last sentence in the previous paragraph implies that $\mu_x(\tau_0) \geq q$. Hence, $\mathfrak{l} \geq q$. This is a contradiction. \square

Lemma 6.8. *Given $\varepsilon > 0$ there exists $r_1 > 0$ such that for all $r \geq r_1$, the complement of $S(x : B(x_0, r))$ in $\text{Flag}(\tau_{\text{mod}})$ is contained in $N_\varepsilon(L(\tau))$, for some $\tau \in S(x_0 : \{x\})$.*

Proof. For $r > 0$ and $\tau_0 \in \text{Flag}(\tau_{\text{mod}})$, $\tau' \in C(\tau_0)$, consider

$$U(\tau_0, x_0, r) = \{\tau' \in \text{Flag}(\tau_{\text{mod}}) \mid P(\tau_0, \tau') \cap B(x_0, r) \neq \emptyset\}.$$

This is an analogue of shadows (1.2) as viewed from the infinity. It is easy to verify that

$$\bigcup_{r \geq 0} U(\tau_0, x_0, r) = C(\tau_0).$$

Moreover, for $g \in G$, these shadows from infinity transform as $gU(\tau_0, x_0, r) = U(g\tau_0, gx_0, r)$.

If $k \in K = G_{x_0}$, the stabilizer of x_0 , then $kU(\tau_0, x_0, r) = U(k\tau_0, x_0, r)$. Since K is compact, there exists $M \geq 1$ such that the action $k \curvearrowright \text{Flag}(\tau_{\text{mod}})$ is M -Lipschitz for all $k \in K$. Let $r_1 > 0$ be such that $U(\tau_0, x_0, r_1/2)^c \subset N_{\varepsilon/M}(L(\tau_0))$. Here and below, for $A \subset \text{Flag}(\tau_{\text{mod}})$, $A^c = \text{Flag}(\tau_{\text{mod}}) - A$. Then, for any $\tau \in \text{Flag}(\tau_{\text{mod}})$,

$$(6.2) \quad U(\tau, x_0, r/2)^c \subset N_\varepsilon(L(\tau)), \quad \forall r \geq r_1.$$

For $x \in X$, let $\tau \in \text{Flag}(\tau_{\text{mod}})$ be a simplex such that $x \in V(x_0, \text{st}(\tau))$. Then there exists a parameterized geodesic ray x_t starting from x_0 , passing through x and asymptotic to some $\xi \in \text{st}(\tau)$.

Claim. For all $r > 0$, $S(x : B(x_0, 2r)) \supset U(\tau, x_0, r)$.

Proof of claim. Pick $\tau' \in U(\tau, x_0, r)$ and let $\bar{x}_0 \in P(\tau, \tau')$ denote the nearest point projection of x_0 . In addition to the ray x_t , we define another parameterized geodesic ray \bar{x}_t , starting at \bar{x}_0 and asymptotic to ξ . Due to the convexity of the Riemannian distance function on X , the distance $d_{\text{Riem}}(x_t, \bar{x}_t)$ monotonically decreases with t . Moreover, the cones $V(\bar{x}_t, \text{st}(\tau'))$ are nested as t decreases. Then,

$$\begin{aligned} d_{\text{Riem}}(x_0, V(x_t, \text{st}(\tau'))) &\leq d_{\text{Riem}}(x_0, V(\bar{x}_t, \text{st}(\tau'))) + d_{\text{Riem}}(x_t, \bar{x}_t) \\ &\leq d_{\text{Riem}}(x_0, V(\bar{x}_0, \text{st}(\tau'))) + r \leq d_{\text{Riem}}(x_0, \bar{x}_0) + r \leq 2r. \end{aligned}$$

Therefore, $\tau' \in S(x : B(x_0, 2r))$. □

Using (6.2) it follows from the above claim that whenever $r \geq r_1$, the complement of the shadow $S(x : B(x_0, r))$ is contained in $N_\varepsilon(L(\tau))$ for some τ satisfying $x \in V(x_0, \text{st}(\tau))$. □

Lemma 6.9. *For all $r > 0$ and all $\tau \in S(x : B(x_0, r))$,*

$$|d_{\bar{\theta}}(x, x_0) - \mathcal{B}_\tau^{\bar{\theta}}(x, x_0)| \leq 2r.$$

Proof. We recall that $d_{\bar{\theta}}$ can alternatively be defined as

$$d_{\bar{\theta}}(y, z) = \max_{\tau \in \text{Flag}(\tau_{\text{mod}})} \mathcal{B}_\tau^{\bar{\theta}}(y, z),$$

where the maximum above occurs at any point in $S(y : \{z\})$ (see [26, Sec. 5.1.2]). Fix some $\tau_0 \in S(x, \{x_0\})$. Then, for any $\tau \in S(x : B(x_0, r))$,

$$\begin{aligned} |d_{\bar{\theta}}(x, x_0) - \mathcal{B}_\tau^{\bar{\theta}}(x, x_0)| &= |b_{\tau_0}^{\bar{\theta}}(x_0) - b_\tau^{\bar{\theta}}(x_0)| \\ &= |b_{\tau_0}^{\bar{\theta}}(x_0) - b_{\tau_0}^{\bar{\theta}}(k^{-1}x_0)| \\ &\leq d_{\text{Riem}}(x_0, k^{-1}x_0) = d_{\text{Riem}}(x_0, kx_0), \end{aligned}$$

where $k \in K$, stabilizer of x , any element satisfying $\tau = k\tau_0$. In the above we chose the normalizations of the Busemann functions at x .

Let $y \in V(x, \text{st}(\tau)) \cap B(x_0, r)$. Then $y \in V(x, \sigma)$ for some chamber σ in $\text{st}(\tau)$. We identify $V(x, \sigma)$ with the model Weyl chamber Δ . Let $k_1 \in K$ such that $k_1 x_0 \in V(x, \sigma)$. Then, $k_1 \tau_0 = \tau$, and $k_1 x_0 = d_\Delta(x, x_0)$ via the identification above. Moreover, since the map

$$X \rightarrow \Delta, \quad z \mapsto d_\Delta(x, z)$$

is 1-Lipschitz (by the triangle inequality for Δ -distances (1.1)) and $d_\Delta(x, y) = y$, we obtain

$$d_{\text{Riem}}(y, k_1 x_0) \leq d_{\text{Riem}}(y, x_0) \leq r$$

and, in particular, $d_{\text{Riem}}(x_0, k_1 x_0) \leq 2r$. \square

Using the above lemmata, we now complete the proof of Theorem 6.1. We first fix some auxiliary quantities. Let $q \in (\mathfrak{l}, \mathfrak{m})$ and ε_0 be corresponding constant as given in Lemma 6.7. Let δ be a constant given by Lemma 6.6 which corresponds to $\varepsilon = \varepsilon_0$. By Λ we denote the δ -neighborhood of $\Lambda_{\tau_{\text{mod}}}$ and let

$$V = \bigcup_{\tau \in \Lambda} V(x, \text{st}(\tau)) \subset X.$$

Since Γ is discrete, the elements of Γ which send x_0 outside V form a finite set Φ . Let

$$r_0 = \max\{r_1, d_{\text{Riem}}(x, \gamma x_0) \mid \gamma \in \Phi\},$$

where r_1 is a constant that corresponds to $\varepsilon_0/2$ as in Lemma 6.8.

For every $\gamma \in \Gamma$ satisfying $d_{\text{Riem}}(x, \gamma x_0) > r \geq r_0$, we assign an element $\tau_\gamma \in S(x : \{\gamma x_0\}) \cap \Lambda$ (the intersection is nonempty by above). Using Lemma 6.6, for every such τ_γ there exists $\tau_0 \in \Lambda_{\tau_{\text{mod}}}$ so that

$$N_{\varepsilon_0/2}(L(\tau_\gamma)) \subset N_{\varepsilon_0}(L(\tau_0)).$$

By Lemmata 6.7 and 6.8, $\mu_x(S(\gamma^{-1}x : B(x_0, r))) \geq \mathfrak{m} - q$ and by properties of conformal measures,

$$\begin{aligned} \mu_x(S(x : B(\gamma x_0, r))) &= \mu_{\gamma^{-1}x}(S(\gamma^{-1}x : B(x_0, r))) \\ &= \int_{S(\gamma^{-1}x : B(x_0, r))} \exp\left(-\beta \mathcal{B}_\tau^{\bar{\theta}}(\gamma^{-1}x, x)\right) d\mu_x \\ &\asymp \exp\left(-\beta d_{\bar{\theta}}(x, \gamma x_0)\right), \end{aligned}$$

where in the last step we have additionally used Lemma 6.9. This completes the proof. \square

7. DIMENSION OF A $\bar{\theta}$ -CONFORMAL DENSITY

In this section, we establish a lower bound for the dimension of a $\bar{\theta}$ -conformal density. For Anosov subgroups, we prove that the dimension equals the $\bar{\theta}$ -critical exponent (see Corollary 7.5).

Theorem 7.1. *Suppose that Γ is a nonelementary τ_{mod} -RA subgroup. Let μ be a Γ -invariant $\bar{\theta}$ -conformal density of dimension β . Then β has the following lower bound:*

$$(7.1) \quad \beta \geq \delta_{\bar{\theta}} - \delta_{\bar{\theta}}^{\text{con}}.$$

The proof of this theorem is given at the end of this section. The number $\delta_{\bar{\theta}}^{\text{con}}$ above quantifies the maximal exponential growth rate of the orbit Γx_0 in a conical direction. The precise definition is given below.

Definition 7.2 (Critical exponent in conical directions). Suppose that Γ is a τ_{mod} -regular subgroup. For $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$, define

$$N_{\bar{\theta}}^{\text{con}}(r, c, x, x_0, \tau) = \text{card}\{\gamma \in \Gamma \mid d_{\bar{\theta}}(x, \gamma x_0) < r, d_{\text{Riem}}(\gamma x_0, V(x, \text{st}(\tau))) < c\}$$

and

$$(7.2) \quad \delta_{\bar{\theta}}^{\text{con}}(\Gamma) = \sup_{\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)} \left(\lim_{c \rightarrow \infty} \left(\limsup_{r \rightarrow \infty} \frac{\log N_{\bar{\theta}}^{\text{con}}(r, c, x, x_0, \tau)}{r} \right) \right).$$

Note that it is sufficient to take the supremum in the definition of $\delta_{\bar{\theta}}^{\text{con}}(\Gamma)$ over the conical limit set $\Lambda_{\tau_{\text{mod}}}^{\text{con}}(\Gamma)$. For rank-one symmetric spaces, and, more generally, for σ_{mod} -regular subgroups, this number is zero. This can be seen as follows. Let Γ be a σ_{mod} -regular discrete subgroup, and $\sigma \in \text{Flag}(\sigma_{\text{mod}})$ be a limit point. Consider a Riemannian tubular neighborhood T of $V(x, \text{st}(\sigma)) = V(x, \sigma)$. Then the (Riemannian) volume of $T_r := \{y \in T \mid d_{\bar{\theta}}(x, y) < r\}$ grows at most polynomially (of degree equal to the rank of X) with respect to r . Hence the number $N_{\bar{\theta}}^{\text{con}}(r, c, x, x_0, \sigma)$ also grows at most polynomially with r . Therefore, the limit in the innermost bracket of (7.2) is zero.

Below we see that for τ_{mod} -Anosov subgroups also, $\delta_{\bar{\theta}}^{\text{con}}(\Gamma) = 0$. It should be noted that, however, for general discrete subgroups, $\delta_{\bar{\theta}}^{\text{con}}$ could be ∞ .

Proposition 7.3. *Suppose that Γ is a nonelementary τ_{mod} -Anosov subgroup. Then the function $N(r) = N_{\bar{\theta}}^{\text{con}}(r, c, x, x_0, \tau)$ grows linearly with r . In particular, $\delta_{\bar{\theta}}^{\text{con}}(\Gamma) = 0$.*

Proof. Without loss of generality, we can assume that $x = x_0$.²²

Lemma 7.4. *Fix $c > 0$. For any $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$, the set*

$$\{\gamma x_0 \mid \gamma \in \Gamma, d_{\text{Riem}}(\gamma x_0, V(x, \text{st}(\tau))) < c\}$$

is within a uniformly bounded distance from a uniform τ_{mod} -Morse quasiray α emanating from x_0 and asymptotic to τ .

Proof. Pick an arbitrary point $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$. Denote the preimage of τ in $\partial_{\infty}\Gamma$ under the boundary homeomorphism $\partial_{\infty}\Gamma \rightarrow \Lambda_{\tau_{\text{mod}}}(\Gamma)$ by ζ . Since Γ is discrete, we can arrange the elements of $\{\gamma \in \Gamma \mid d_{\text{Riem}}(\gamma x_0, V(x, \text{st}(\tau))) < c\}$ in a sequence (γ_n) . The sequence $x_n = \gamma_n x_0$ converges conically to τ . Let $\alpha : \mathbb{Z}_{\geq 0} \rightarrow X$ be the image (under the orbit map $\Gamma \rightarrow \Gamma x$) of a parametrized geodesic ray $\mathbb{Z}_{\geq 0} \rightarrow \Gamma$ starting at 1_{Γ} and asymptotic to ζ . Then α is a uniform τ_{mod} -Morse quasiray starting at x_0 and asymptotic to τ . Hence α is uniformly close to $V(x_0, \text{st}(\tau))$. Since both sequences (x_n) and $(\alpha(n))$ are uniformly close to $V(x_0, \text{st}(\tau))$, it is enough²³ to understand the simpler case when $\alpha(n)$, $x_n \in V(x_0, \text{st}(\tau))$, for all $n \in \mathbb{N}$.

We claim that the sequence (x_n) is uniformly close to α . Otherwise, after extraction, (x_n) would diverge away from α . Since α is a Morse quasiray, α eventually enters each cone $V(x_n, \text{st}(\tau))$, but further and further away from the tip x_n as n grows. Since the separation between two successive points on α (being a quasi-geodesic) is uniformly bounded, we could find arbitrarily large m 's such that $\alpha(m)$

²²Note that the number $\delta_{\bar{\theta}}^{\text{con}}(\Gamma)$ does not depend on x and x_0 as we have seen in the case of $\delta_{\bar{\theta}}$ in Section 2.

²³For instance, we can consider the nearest point projections of the sequences $\alpha(n)$ and x_n to $V(x_0, \text{st}(\tau))$. The new sequences would be uniformly close to the old ones. Note that the projection sequence of $\alpha(n)$ is also a uniform τ_{mod} -Morse quasiray.

is uniformly close to the boundary of a cone $V(x_n, \text{st}(\tau))$ and is arbitrarily far away from its tip x_n . But this would contradict the uniform τ_{mod} -regularity of the group Γ . \square

We continue with the notations from the proof of the lemma. Since any τ_{mod} -Anosov subgroup $\Gamma < G$ is uniformly τ_{mod} -regular (cf. Theorem 1.7), we may work with the Riemannian metric d_{Riem} in place of the $d_{\bar{\theta}}$. Moreover, we may assume that the sequence (x_n) is sufficiently spaced. Let \bar{x}_n denote the nearest-point projection of x_n to the image of α . Lemma 7.4 implies that $d(x_n, \bar{x}_n)$ is uniformly bounded. Since x_n 's are sufficiently spaced, \bar{x}_n 's are also sufficiently spaced which guarantees that $d_{\text{Riem}}(\bar{x}_n, x_0) \geq \text{const} \cdot n$, for all large n , which in turn implies that $d_{\text{Riem}}(x_n, x_0) \geq \text{const} \cdot n$. The proposition follows from this. \square

As a corollary of the above results, we obtain that any Γ -invariant $\bar{\theta}$ -conformal density must have dimension $\delta_{\bar{\theta}}$ when Γ is τ_{mod} -Anosov. The Patterson–Sullivan densities constructed in Section 3 also had this dimension.

Corollary 7.5. *Suppose that Γ is a nonelementary τ_{mod} -Anosov subgroup. Let μ be a Γ -invariant $\bar{\theta}$ -conformal density of dimension β . Then $\beta = \delta_{\bar{\theta}}$.*

Proof. Recall that Γ -invariant $\bar{\theta}$ -conformal densities are, by definition (see Definition 3.2), supported in the limit set $\Lambda_{\tau_{\text{mod}}}(\Gamma)$. Since for τ_{mod} -Anosov subgroups²⁴ $\Lambda_{\tau_{\text{mod}}}(\Gamma) = \Lambda_{\tau_{\text{mod}}}^{\text{con}}(\Gamma)$, by Corollary 6.5, we know that the Poincaré series $g_{\beta}^{\bar{\theta}}(x, x_0)$ diverges and, consequently, $\beta \leq \delta_{\bar{\theta}}$. The reverse inequality is obtained by a combination of Theorem 7.1 and Proposition 7.3. \square

To close this section, we prove Theorem 7.1.

Proof of Theorem 7.1. We fix some $r \geq r_0$ where r_0 is given by Theorem 6.1.

We first assume that the stabilizer of x_0 in Γ is trivial in which case the function $N_{\bar{\theta}}(R, x, x_0)$ counts the number of orbit points (in Γx_0) within the R -ball in $(X, d_{\bar{\theta}})$ centered at x .

We place a Riemannian ball of radius r at each point in the orbit. In this proof, we reserve the word *ball* to specify these balls. Let

$$c = \min_{1_{\Gamma} \neq \gamma \in \Gamma} \{d_{\text{Riem}}(x_0, \gamma x_0)\}.$$

There exists a number $N \in \mathbb{N}$ that depends only on r , c , and X such that any ball intersects at most N other balls (including itself). Note that the shadows in $\text{Flag}(\tau_{\text{mod}})$ (from x) of two distinct balls are disjoint unless they intersect some common τ_{mod} -cone with tip at x . Also note that, at large distances from x , the balls do not intersect the boundaries of the τ_{mod} -cones because of the τ_{mod} -regularity of the orbit.

Let n_R denote the maximal number of balls in $B_{\bar{\theta}}(x, R)$ that intersect a particular τ_{mod} -cone $V(x, \text{st}(\tau))$. It follows from the definition of $\delta_{\bar{\theta}}^{\text{con}}(\Gamma)$ that

$$(7.3) \quad \limsup_{R \rightarrow \infty} \frac{\log n_R}{R} \leq \delta_{\bar{\theta}}^{\text{con}}(\Gamma).$$

²⁴For τ_{mod} -Anosov subgroups Γ , $\Lambda_{\tau_{\text{mod}}}(\Gamma) = \Lambda_{\tau_{\text{mod}}}^{\text{con}}(\Gamma)$ is a consequence of the fact that τ_{mod} -Anosov subgroups are also τ_{mod} -RCA, see Theorem 1.7.

On the other hand, for each $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$, the maximal number of balls in $B_{\bar{\theta}}(x, R)$ whose shadows intersect τ is n_R . Therefore,

$$(7.4) \quad \frac{N_{\bar{\theta}}(R, x, x_0)}{N \cdot n_R} s(R) \leq \mathbf{m} = \text{total mass of } \mu_x,$$

where $s(R)$ is any lower bound for the measures of the shadows of balls in $B_{\bar{\theta}}(x, R)$. We note that the shadow lemma (Theorem 6.1) produces such a positive lower bound,²⁵ namely, we may take $s(R) = \text{const} \cdot e^{-\beta R}$. Then (7.4) yields

$$N_{\bar{\theta}}(R, x, x_0) \leq \frac{\mathbf{m}N \cdot n_R}{\text{const}} e^{\beta R}.$$

Together with (7.3), the above results in (7.1).

In the general case when Γ_{x_0} is non-trivial, $\Gamma_{x_0} = K \cap \Gamma$ is still at most finite. In this case, $N_{\bar{\theta}}(R, x, x_0)$ will be a constant multiple of the number of Γ -orbit points of x_0 within the R -ball in $(X, d_{\bar{\theta}})$ centered at x . So, we only need to change the constant term in the above inequality. \square

8. UNIQUENESS OF $\bar{\theta}$ -CONFORMAL DENSITY

Recall that an action of a group H on a measure space (S, σ) is said to be *ergodic* if each H -invariant measurable set $B \subset S$ is either null or co-null. In [47], Sullivan proved that for a discrete group Γ of Möbius transformations of the Poincaré ball \mathbb{B}^3 , a Γ -invariant $\bar{\theta}$ -conformal density μ of non-zero dimension is unique (here and henceforth, by “unique” we mean unique up to a constant factor) in the class of all $\bar{\theta}$ -conformal densities of same dimension if and only if the action Γ on the limit set $\Lambda(\Gamma)$ is ergodic with respect to any $\mu_x \in \mu$. See also [37, Thm. 4.2.1]. Generalizing this statement in our setting, we obtain the following result. The proof is essentially same of Sullivan’s theorem, hence we omit the details.

Theorem 8.1. *Suppose that Γ is a nonelementary τ_{mod} -RA subgroup. A Γ -invariant $\bar{\theta}$ -conformal density μ of dimension $\beta > 0$ is unique in the class of all Γ -invariant $\bar{\theta}$ -conformal densities of dimension β if and only if the action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is ergodic with respect to any $\mu_x \in \mu$.*

It is then natural to ask

Question 8.2. For which τ_{mod} -regular subgroups Γ , the action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is ergodic with respect to a conformal measure?

In this section we prove that the Anosov property is a sufficient condition:

Theorem 8.3 (Anosov implies ergodic). *Suppose that Γ is a nonelementary τ_{mod} -Anosov subgroup and μ be a Γ -invariant $\bar{\theta}$ -conformal density. Then the action $\Gamma \curvearrowright \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is ergodic with respect to any $\mu_x \in \mu$.*

As a corollary, we obtain that when Γ is τ_{mod} -Anosov, then, up to a constant factor, there is exactly one Γ -invariant $\bar{\theta}$ -conformal density, namely, the Patterson–Sullivan density.

Corollary 8.4 (Existence and uniqueness of $\bar{\theta}$ -conformal density). *Suppose that Γ is a nonelementary τ_{mod} -Anosov subgroup. Then, up to a constant factor, there exists a unique Γ -invariant $\bar{\theta}$ -conformal density μ , namely, the Patterson–Sullivan density of type $\bar{\theta}$.*

²⁵We may need to disregard a finite number of balls from the picture.

Proof. First of all, by Proposition 3.5, any such density must have a positive dimension. Secondly, by Corollary 7.5 this dimension equals to the critical exponent $\delta_{\bar{\theta}}$. Then the uniqueness follows from the combination of Theorems 8.1 and 8.3. \square

Now we return to the proof of Theorem 8.3.

Proof of Theorem 8.3. Let μ be a Γ -invariant $\bar{\theta}$ -conformal density. Note that the dimension β of μ must be positive (by Proposition 3.5 and Corollary 7.5).

Let B be a Γ -invariant Borel subset of $\Lambda_{\tau_{\text{mod}}}(\Gamma)$. We need to prove that if B is not a null set, then it is co-null. From now on, we assume that B is not a null set, i.e., $\mu_x(B) > 0$.

We need the following lemmata.

Lemma 8.5. *There exists $r_1 > 0$ such that for every $r \geq r_1$ and every $\gamma \in \Gamma$, the shadow $S(x, B(\gamma x_0, r))$ intersects $\Lambda_{\tau_{\text{mod}}}(\Gamma)$.*

Proof. The proof simply follows from the Morse property of the Anosov subgroup Γ . \square

We assume that the r_1 in the lemma also satisfies the ‘‘uniform conicality’’ property for Γ (cf. Proposition 6.4).

Lemma 8.6. *Let $r \geq \max\{r_0, r_1\}$ where r_0 is as in Theorem 6.1. For μ_x -a.e. $\tau \in B$ and every sequence (γ_n) on Γ , $\gamma_n \rightarrow \tau$, satisfying $\tau \in S_n := S(x : B(\gamma_n x_0, r))$, we have*

$$(8.1) \quad \lim_{n \rightarrow \infty} \frac{\mu_x(S_n \cap B)}{\mu_x(S_n)} = 1.$$

Assuming this lemma for a moment, we complete the proof of the theorem. The proof of this lemma is given at the end of this section. Note that Lemma 8.5 is used to ensure that the ratios in Lemma 8.6 are not degenerate.

Let $\tau \in B$ be a *density point*, i.e., τ satisfies (8.1). Such point exists by Lemma 8.6 because B has positive mass. Note that Γ -invariance of B and μ implies that

$$\begin{aligned} \frac{\mu_x(S(\gamma_n^{-1}x : B(x_0, r)) \cap B)}{\mu_x(S(\gamma_n^{-1}x : B(x_0, r)))} &= \frac{\mu_{\gamma_n x}(S_n \cap B)}{\mu_{\gamma_n x}(S_n)} \\ &= 1 - \frac{\mu_{\gamma_n x}(S_n - B)}{\mu_{\gamma_n x}(S_n)} \\ &= 1 - \frac{\int_{S_n - B} \exp(-\beta \mathcal{B}_{\tau}^{\bar{\theta}}(\gamma_n x, x)) d\mu_x}{\int_{S_n} \exp(-\beta \mathcal{B}_{\tau}^{\bar{\theta}}(\gamma_n x, x)) d\mu_x} \\ &\geq 1 - \text{const} \cdot \frac{\mu_x(S_n - B)}{\mu_x(S_n)}, \end{aligned}$$

where the inequality follows by Lemma 6.9. Together with (8.1), we get

$$(8.2) \quad \lim_{n \rightarrow \infty} \frac{\mu_x(S(\gamma_n^{-1}x : B(x_0, r)) \cap B)}{\mu_x(S(\gamma_n^{-1}x : B(x_0, r)))} = 1.$$

Note that by Corollary 6.2, μ is atom-free. Therefore, for every $\varepsilon > 0$ there exists $r > r_1$ such that

$$\mu_x(S(\gamma_n^{-1}x : B(x_0, r))) \geq \mathfrak{m} - \varepsilon,$$

for all large n , where \mathbf{m} denotes the total mass of μ_x . The above follows from the combination of Lemmata 6.7 and 6.8. Therefore, by (8.2),

$$\mu_x(B) \geq \lim_{n \rightarrow \infty} \mu_x(S(\gamma_n^{-1}x : B(x_0, r)) \cap B) \geq \mathbf{m} - \varepsilon,$$

which holds for every $\varepsilon > 0$. Hence $\mu_x(B) = \mathbf{m}$. This completes the proof of the theorem. \square

Now we prove Lemma 8.6. The lemma would have followed from a generalization of the Lebesgue density theorem (cf. [17, Subsec. 2.9.11, 2.9.12]) if we knew that μ_x is, e.g., a *doubling measure*. Since this property is unclear, we adopt a more direct approach. The idea of the proof follows [44, Subsec. 1E] (see also [35, Sec. 3]).

Proof of Lemma 8.6. The proof requires a version of the Lebesgue differentiation theorem.

Sublemma 8.7. *For every bounded measurable function $\Phi : \text{Flag}(\tau_{\text{mod}}) \rightarrow \mathbb{R}_{\geq 0}$,*

$$\Phi(\tau) = \lim_{n \rightarrow \infty} \frac{1}{\mu_x(S(x : B(\gamma_n x_0, r)))} \int_{S(x : B(\gamma_n x_0, r))} \Phi d\mu_x,$$

for μ_x -a.e. $\tau \in \Lambda_{\tau_{\text{mod}}}$ and all $\gamma_n \in \Gamma$ satisfying $\tau \in S(x : B(\gamma_n x_0, r))$.

Proof. For every bounded measurable function $\Psi : \text{Flag}(\tau_{\text{mod}}) \rightarrow \mathbb{R}_{\geq 0}$, define a function Ψ^* on $\text{Flag}(\tau_{\text{mod}})$ which is zero outside $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ and on $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ it is defined by

$$(8.3) \quad \Psi^*(\tau) = \limsup_{N \rightarrow \infty} \frac{1}{\mu_x(S(x : B(\gamma x_0, r)))} \int_{S(x : B(\gamma x_0, r))} \Psi d\mu_x.$$

Here and in the following the limit superior is taken over all $\gamma \in \Gamma$ that satisfy $d_{\text{Riem}}(x, \gamma x_0) \geq N$ and $\tau \in S(x : B(\gamma x_0, r))$.

Let Φ_k be a sequence of continuous functions converging to Φ μ_x -almost surely such that

$$\int_{\text{Flag}(\tau_{\text{mod}})} |\Phi_k - \Phi| d\mu_x < \frac{1}{k}, \quad \forall k \in \mathbb{N}.$$

Then for every $\tau \in \text{Flag}(\tau_{\text{mod}})$ and $\gamma \in \Gamma$, we have

$$(8.4) \quad \limsup_{N \rightarrow \infty} \left| \frac{1}{\mu_x(S(x : B(\gamma x_0, r)))} \int_{S(x : B(\gamma x_0, r))} \Phi d\mu_x - \Phi(\tau) \right| \\ \leq |\Phi - \Phi_k|^*(\tau) + |\Phi_k(\tau) - \Phi(\tau)| \\ + \limsup_{N \rightarrow \infty} \left| \frac{1}{\mu_x(S(x : B(\gamma x_0, r)))} \int_{S(x : B(\gamma x_0, r))} \Phi_k d\mu_x - \Phi_k(\tau) \right|.$$

Since Φ_n are continuous, the last quantity in the right side of the above vanishes. Moreover, the limit of $|\Phi_k(\tau) - \Phi(\tau)|$ as $k \rightarrow \infty$ vanishes at μ_x -a.e. $\tau \in \text{Flag}(\tau_{\text{mod}})$. Therefore, we only need to control the first term of the right side of (8.4): We show that, for all bounded nonnegative measurable functions Ψ on $\text{Flag}(\tau_{\text{mod}})$ and all $\varepsilon > 0$,

$$(8.5) \quad \mu_x(\{\Psi^* > \varepsilon\}) \leq \frac{\text{const}}{\varepsilon} \int_{\text{Flag}(\tau_{\text{mod}})} \Psi d\mu_x,$$

where the constant does not depend on ε or Ψ . The sublemma follows from this as follows: Setting $\Psi = |\Phi - \Phi_k|$ and taking limit as $k \rightarrow \infty$ in (8.5), we see that $|\Phi - \Phi_k|^*$ μ_x -a.s. converges to zero. Hence left-hand side of (8.4) also converges to zero for μ_x -a.e. $\tau \in \Lambda_{\tau_{\text{mod}}}$.

Now we verify (8.5). Let $\varepsilon > 0$ be arbitrary. For $d \geq 0$, let Γ_d be the set of all elements $\gamma \in \Gamma$ such that $d_{\bar{g}}(x, \gamma x_0) \geq d$ and

$$(8.6) \quad \int_{S(x: B(\gamma x_0, r))} \Psi d\mu_x \geq \frac{\varepsilon}{2} \mu_x(S(x : B(\gamma x_0, r))).$$

Claim 1. The union of all shadows $S(x : B(\gamma x_0, r))$ over $\gamma \in \Gamma_d$ covers $\{\Psi^* > \varepsilon\}$.

Proof of claim. The proof is straightforward. \square

We recursively construct a sequence of subsets, $(\Gamma_{d,N})$, of Γ_d in the following way: Let $\Gamma_{d,1} = \{\gamma \in \Gamma_d \mid 0 \leq d_{\bar{g}}(x, \gamma x_0) < 1\}$, and, for $N \geq 2$, define

$$\Gamma_{d,N} = \left\{ \gamma \in \Gamma_d \mid \begin{array}{l} N-1 \leq d_{\bar{g}}(x, \gamma x_0) < N \text{ and } S(x : B(\gamma x_0, r)) \cap \\ S(x : B(\phi x_0, r)) = \emptyset, \forall \phi \in \Gamma_{d,1} \cup \dots \cup \Gamma_{d,N-1} \end{array} \right\}.$$

Set $\Gamma_d^* = \bigcup_{N \geq 1} \Gamma_{d,N}$.

Claim 2. There exists a constant $R \geq r$ such that, for every $d \geq 0$,

$$\{\Psi^* > \varepsilon\} \subset \bigcup_{\phi \in \Gamma_d^*} S(x : B(\phi x_0, R)).$$

Proof of claim. It is enough to prove the claim for very large d . In fact, we assume that d is so large such that $\overline{x(\gamma x_0)}$ is uniformly τ_{mod} -regular for all $\gamma \in \Gamma_d$.

Let $\tau \in \{\Psi^* > \varepsilon\}$ be arbitrary. Then there exists $\gamma \in \Gamma_d$ such that $\tau \in S(x : B(\gamma x_0, r))$. Assume that $\gamma \notin \Gamma_d^*$. By construction of Γ_d^* , there exists $\phi \in \Gamma_d^*$ such that $S(x : B(\gamma x_0, r)) \cap S(x : B(\phi x_0, r)) \neq \emptyset$ and $d_{\bar{g}}(x, \phi x_0) < d_{\bar{g}}(x, \gamma x_0)$.

By Lemma 7.4, both γx_0 and ϕx_0 stay uniformly close to a τ_{mod} -uniform Morse quasigeodesic α with one endpoint at x . Since $d_{\bar{g}}(x, \phi x_0) < d_{\bar{g}}(x, \gamma x_0)$, we may assume that the other endpoint of α is uniformly close to γx_0 . It follows that ϕx_0 is uniformly close to the diamond $\diamond_{\Theta}(x, \gamma x_0)$, since α is, for some $\Theta \subset \tau_{\text{mod}}$. Pick $y \in B(\gamma x_0, r) \cap V(x, \text{st}(\tau))$. Then, by uniform continuity of diamonds (cf. [14, Thm. 3.7]), for some Θ' bigger than Θ , $\diamond_{\Theta}(x, \gamma x_0)$ is contained in a uniform neighborhood of $\diamond_{\Theta'}(x, y)$. Therefore, ϕx_0 is uniformly close to $\diamond_{\Theta'}(x, y)$ and, in particular, to $V(x, \text{st}(\tau))$. We may choose R to be this upper bound. \square

In particular, we get

$$(8.7) \quad \mu_x(\{\Psi^* > \varepsilon\}) \leq \sum_{\phi \in \Gamma_R^*} \mu_x(S(x : B(\phi x_0, R))).$$

Claim 3. If $S(x : B(\gamma x_0, r)) \cap S(x : B(\phi x_0, r)) \neq \emptyset$, for $\gamma, \phi \in \Gamma_d^*$, then $d_{\bar{g}}(\gamma x_0, \phi x_0)$ is uniformly bounded.

Proof of claim. This follows from the Gromov hyperbolicity of $(\Gamma x_0, d_{\bar{g}})$ (see Corollary 4.8) and the fact that both γx_0 and ϕx_0 lie in an annulus $\{x' \in X \mid N-1 \leq d_{\bar{g}}(x, x') < N\}$ in the following way: Let $\tau \in S(x : B(\gamma x_0, r)) \cap S(x : B(\phi x_0, r))$. Let $z \in V(x, \text{st}(\tau))$ be a point uniformly close to Γx_0 . By δ -hyperbolicity,

$$(8.8) \quad \langle \gamma x_0 | \phi x_0 \rangle_x + \delta \geq \min \{ \langle \gamma x_0 | z \rangle_x, \langle \phi x_0 | z \rangle_x \}.$$

Expanding the left side, we get

$$(8.9) \quad \begin{aligned} \langle \gamma x_0 | \phi x_0 \rangle_x + \delta &= \frac{1}{2} (d_{\bar{\theta}}(x, \gamma x_0) + d_{\bar{\theta}}(\phi x_0, x) - d_{\bar{\theta}}(\gamma x_0, \phi x_0)) + \delta \\ &\leq \left(d_{\bar{\theta}}(\phi x_0, x) - \frac{1}{2} d_{\bar{\theta}}(\gamma x_0, \phi x_0) \right) + \delta + \frac{1}{2}, \end{aligned}$$

and expanding the right side, we get

$$\min \{ \langle \gamma x_0 | z \rangle_x, \langle \phi x_0 | z \rangle_x \} = \min \left\{ \frac{1}{2} (d_{\bar{\theta}}(x, \gamma x_0) + d_{\bar{\theta}}(z, x) - d_{\bar{\theta}}(\gamma x_0, z)), \frac{1}{2} (d_{\bar{\theta}}(x, \phi x_0) + d_{\bar{\theta}}(z, x) - d_{\bar{\theta}}(\phi x_0, z)) \right\}.$$

Taking $z \rightarrow \tau$ in the right side of the last one and using (3.3), we get

$$\min \left\{ \frac{1}{2} \left(d_{\bar{\theta}}(\gamma x_0, x) + \mathcal{B}_{\tau}^{\bar{\theta}}(x, \gamma x_0) \right), \frac{1}{2} \left(d_{\bar{\theta}}(\phi x_0, x) + \mathcal{B}_{\tau}^{\bar{\theta}}(x, \phi x_0) \right) \right\},$$

which, by Lemma 6.9, is at least

$$\min \{ d_{\bar{\theta}}(\gamma x_0, x), d_{\bar{\theta}}(\phi x_0, x) \} - r \geq d_{\bar{\theta}}(\gamma x_0, x) - r - 1.$$

Combining this with (8.8) and (8.9), we get

$$d_{\bar{\theta}}(\gamma x_0, \phi x_0) \leq 2r + 2\delta + 3.$$

□

In particular, for each $\tau \in \mu_x(\{\Psi^* > \varepsilon\})$, $\#\{\phi \in \Gamma_d^* \mid \tau \in S(x : B(\phi x_0, r))\}$ is uniformly bounded, say, by $D > 0$. Therefore,

$$(8.10) \quad \sum_{\phi \in \Gamma_r^*} \mu_x(S(x : B(\phi x_0, r))) \leq D \mu_x \left(\bigcup_{\phi \in \Gamma_r^*} S(x : B(\phi x_0, r)) \right).$$

We would like to use the shadow lemma (Theorem 6.1). To this end, we have

$$(8.11) \quad \mu_x(\{\Psi^* > \varepsilon\}) \leq \sum_{\phi \in \Gamma_R^*} \mu_x(S(x : B(\phi x_0, R))) \leq C' \sum_{\phi \in \Gamma_R^*} \exp(-\beta d_{\bar{\theta}}(x, \phi x_0)),$$

where the first inequality is given by (8.7) and the last inequality is given by the shadow lemma with $r_0 \leq r = R$. Note that the necessary condition $d_{\bar{\theta}}(x, \phi x_0) \geq R$ which we needed to apply the shadow lemma in the above follows from the definition of Γ_R^* . Moreover, applying shadow lemma again with $r_0 \leq r = r$, we get another constant $C > 0$ such that

$$(8.12) \quad C^{-1} \sum_{\phi \in \Gamma_R^*} \exp(-\beta d_{\bar{\theta}}(x, \phi x_0)) \leq \sum_{\phi \in \Gamma_R^*} \mu_x(S(x : B(\phi x_0, r))).$$

Combined with (8.10), the inequalities in (8.11) and (8.12) give

$$\mu_x(\{\Psi^* > \varepsilon\}) \leq DC' C \mu_x \left(\bigcup_{\phi \in \Gamma_R^*} S(x : B(\phi x_0, r)) \right).$$

Finally, the above and (8.6) yield

$$\mu_x(\{\Psi^* > \varepsilon\}) \leq \frac{2DC'C}{\varepsilon} \int_{\text{Flag}(\tau_{\text{mod}})} \Psi d\mu_x.$$

This proves (8.5). □

The proof of the lemma follows from the sublemma by taking Φ in the sublemma to be the indicator function for B . \square

9. HAUSDORFF DENSITY

In this section, we restrict our attention to Anosov subgroups. Usually, one defines Hausdorff measures and Hausdorff dimension for metric spaces. In Appendix A, we verify that the theory goes through for premetrics as well. Therefore, we choose to work with premetric spaces. The reader who prefers to work with metrics can assume that $\epsilon > 0$ in the following is chosen so small so that $D_x^{\bar{\theta}, \epsilon}$ is bilipschitz equivalent a metric on $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ (cf. Corollary 5.8).

We fix an $\epsilon > 0$. For $\beta \geq 0$, we let $\mathcal{H}_x^{\bar{\theta}, \epsilon, \beta}$ denote the β -dimensional Hausdorff measure on the premetric space $(\Lambda_{\tau_{\text{mod}}}(\Gamma), D_x^{\bar{\theta}, \epsilon})$. These definitions can be found in Appendix A. The Hausdorff dimension of a Borel subset $B \subset \Lambda_{\tau_{\text{mod}}}(\Gamma)$ is then defined as

$$\dim_{\text{Haus}}^{\bar{\theta}, \epsilon}(B) = \inf\{\beta \mid \mathcal{H}_x^{\bar{\theta}, \epsilon, \beta}(B) = 0\} = \sup\{\beta \mid \mathcal{H}_x^{\bar{\theta}, \epsilon, \beta}(B) = \infty\}.$$

We observe that if for some $\beta \geq 0$, $\mathcal{H}_x^{\bar{\theta}, \epsilon, \beta}(B) \in (0, \infty)$, then $\dim_{\text{Haus}}^{\bar{\theta}, \epsilon}(B) = \beta$.

Remark 9.1. The Hausdorff dimension $\dim_{\text{Haus}}^{\bar{\theta}, \epsilon}(B)$ does not depend on the choice of a base-point $x \in X$, although the definition above involved such a basepoint. This follows from the fact that for any $x, z \in X$, the identity map

$$\text{id} : (\Lambda_{\tau_{\text{mod}}}(\Gamma), D_x^{\bar{\theta}, \epsilon}) \rightarrow (\Lambda_{\tau_{\text{mod}}}(\Gamma), D_z^{\bar{\theta}, \epsilon})$$

and its inverse are locally Lipschitz maps. We show this in the proof of Proposition 9.2. For this reason, we have dropped the basepoint from the notation of the Hausdorff dimension.

Proposition 9.2. *Suppose that for some $\beta \geq 0$*

$$(9.1) \quad \mathcal{H}_x^{\bar{\theta}, \epsilon, \beta}(\Lambda_{\tau_{\text{mod}}}(\Gamma)) \in (0, \infty).$$

Let $Z = \Gamma x$. Then $\mathcal{H}^{\bar{\theta}, \epsilon, \beta} = \{\mathcal{H}_z^{\bar{\theta}, \epsilon, \beta}\}_{z \in Z}$ is a Γ -invariant $\bar{\theta}$ -conformal Z -density of dimension $\beta\epsilon$.

Proof. Let $y, z \in Z$. Define a function $f : \Lambda_{\tau_{\text{mod}}}(\Gamma) \times \Lambda_{\tau_{\text{mod}}}(\Gamma) \rightarrow \mathbb{R}_{\geq 0}$ by

$$f(\tau_1, \tau_2) = \begin{cases} \frac{D_y^{\bar{\theta}, \epsilon}(\tau_1, \tau_2)}{D_z^{\bar{\theta}, \epsilon}(\tau_1, \tau_2)}, & \tau_1 \neq \tau_2, \\ \exp\left(-\epsilon \mathcal{B}_\tau^{\bar{\theta}}(y, z)\right), & \tau_1 = \tau_2 = \tau. \end{cases}$$

By a calculation similar to the proof of Lemma 5.6, we obtain

$$\lim_{\tau_1, \tau_2 \rightarrow \tau} \frac{D_y^{\bar{\theta}, \epsilon}(\tau_1, \tau_2)}{D_z^{\bar{\theta}, \epsilon}(\tau_1, \tau_2)} = \exp\left(-\epsilon \mathcal{B}_\tau^{\bar{\theta}}(y, z)\right)$$

which shows that f is continuous. For $\tau \in \Lambda_{\tau_{\text{mod}}}(\Gamma)$ and sufficiently small $\eta > 0$, let U_η be a neighborhood of τ in $\Lambda_{\tau_{\text{mod}}}(\Gamma)$ such that $\forall \tau_1, \tau_2 \in U_\eta$,

$$D_y^{\bar{\theta}, \epsilon}(\tau_1, \tau_2) \leq \left(\exp\left(-\epsilon \mathcal{B}_\tau^{\bar{\theta}}(y, z)\right) + \eta\right) D_z^{\bar{\theta}, \epsilon}(\tau_1, \tau_2).$$

Hence the identity map $\text{id} : (\Lambda_{\tau_{\text{mod}}}(\Gamma), D_x^{\bar{\theta}, \epsilon}) \rightarrow (\Lambda_{\tau_{\text{mod}}}(\Gamma), D_y^{\bar{\theta}, \epsilon})$ restricted to U_η is L_η -Lipschitz, where $L_\eta := \exp\left(-\epsilon \mathcal{B}_\tau^{\bar{\theta}}(y, z)\right) + \eta$. In particular, the map id is

locally Lipschitz. Therefore, for any $B \in \mathfrak{B}(U)$, $\mathcal{H}_y^{\bar{\theta}, \epsilon, \beta}(B) \leq L_\eta^\beta \mathcal{H}_z^{\bar{\theta}, \epsilon, \beta}(B)$. This also shows that $\mathcal{H}_y^{\bar{\theta}, \epsilon, \beta} \ll \mathcal{H}_z^{\bar{\theta}, \epsilon, \beta}$. Taking limit as $\eta \rightarrow 0$, we obtain

$$\frac{d\mathcal{H}_y^{\bar{\theta}, \epsilon, \beta}}{d\mathcal{H}_z^{\bar{\theta}, \epsilon, \beta}}(\tau) \leq \exp\left(-\beta\epsilon\mathcal{B}_\tau^{\bar{\theta}}(y, z)\right),$$

and by switching the role of y and z in the above we also obtain the reverse inequality. Hence

$$\frac{d\mathcal{H}_y^{\bar{\theta}, \epsilon, \beta}}{d\mathcal{H}_z^{\bar{\theta}, \epsilon, \beta}}(\tau) = \exp\left(-\beta\epsilon\mathcal{B}_\tau^{\bar{\theta}}(y, z)\right)$$

which proves $\bar{\theta}$ -conformality. Suppose that $y = \gamma z$ for some $\gamma \in \Gamma$. Then for any $B \in \mathfrak{B}(\Lambda_{\tau_{\text{mod}}}(\Gamma))$,

$$\mathcal{H}_{\gamma z}^{\bar{\theta}, \epsilon, \beta}(B) = \int_B \exp\left(-\beta\epsilon\mathcal{B}_\tau^{\bar{\theta}}(\gamma z, z)\right) d\mathcal{H}_z^{\bar{\theta}, \epsilon, \beta} = \int_B d\left(\gamma^*\mathcal{H}_z^{\bar{\theta}, \epsilon, \beta}\right) = \gamma^*\mathcal{H}_z^{\bar{\theta}, \epsilon, \beta}(B)$$

and Γ -invariance also follows. Therefore, $\mathcal{H}^{\bar{\theta}, \epsilon, \beta}$ is a conformal Z -density of dimension $\beta\epsilon$. \square

Remark 9.3.

- (1) Note that if such a family $\{\mathcal{H}_z^{\bar{\theta}, \epsilon, \beta} \mid z \in Z\}$ exists, then it may be extended to a *full* $\bar{\theta}$ -conformal density via the correspondence in (3.5).
- (2) By the uniqueness of $\bar{\theta}$ -conformal density (Corollary 8.4), the number β in Proposition 9.2 equals to $\delta_{\bar{\theta}}/\epsilon$.
- (3) In the following we shall see that, indeed, the $\delta_{\bar{\theta}}/\epsilon$ -dimensional Hausdorff measure $\mathcal{H}_x^{\bar{\theta}, \epsilon, \frac{\delta_{\bar{\theta}}}{\epsilon}}$ is finite and non-null (i.e., it satisfies (9.1)).

Next we show that if $\beta = \delta_{\bar{\theta}}/\epsilon$, then the β -dimensional Hausdorff measure $\mathcal{H}_x^{\bar{\theta}, \epsilon, \beta}$ satisfies (9.1). Let us first discuss the simpler case, namely, when the pseudo-metric $d_{\bar{\theta}}$ is a metric. There is an abundance of examples when this occurs, e.g., in the case when $X = G/K$ is an irreducible symmetric space (i.e., G is simple).

Let (Y, d) be a proper, geodesic, Gromov hyperbolic metric space and Γ be a nonelementary discrete group of isometries acting properly discontinuously on Y . Let Λ be the limit set of Γ in $\partial_\infty Y$. Further, assume that Γ is *quasiconvex-cocompact*, i.e., the quasiconvex hull $\text{QCH}(\Lambda)$ is nonempty and $\text{QCH}(\Lambda)/\Gamma$ is compact. In [10], Coornaert proved the following result.

Theorem 9.4 ([10, Cor. 7.6]). *Suppose that the critical exponent δ of Γ is finite. Then the δ -dimensional Hausdorff measure on Λ with respect to a Gromov metric D is finite and non-null.*

To apply this theorem to our case, we need an appropriate setup. In Section 4, we proved that the orbit $Z = \Gamma x$ is a Gromov hyperbolic space with respect to the Finsler metric $d_{\bar{\theta}}$ (cf. Corollary 4.8) and it is also proper. But Z fails to be geodesic. This problem can be remedied by taking a uniform neighborhood Y of Z in X such that Z is quasiconvex in Y , and then putting the intrinsic path-metric d on Y induced by $d_{\bar{\theta}}$ (this requires positivity of $d_{\bar{\theta}}$), and finally by completing Y in this metric. Then (Y, d) is proper, geodesic and Gromov hyperbolic. Moreover, (Y, d) and the isometrically embedded $(Z, d_{\bar{\theta}})$ are Hausdorff-close and, in particular, (Y, d) is quasiisometric to $(Z, d_{\bar{\theta}})$ by a $(1, A)$ -quasiisometry. This implies that there is a bilipschitz homeomorphism from $\partial_\infty Y$ (equipped with the metric D^ϵ defined by

$D^\epsilon(\xi_1, \xi_2) = D(\xi_1, \xi_2)^\epsilon$ where D is a Gromov metric on $\partial_\infty Y$ to $(\Lambda_{\tau_{\text{mod}}}(\Gamma), D_x^{\bar{\theta}, \epsilon})$. Note that the action $\Gamma \curvearrowright (Y, d)$ satisfies all the properties needed to apply Theorem 9.4. Therefore, by this theorem the $\delta_{\bar{\theta}}/\epsilon$ -dimensional Hausdorff measure on $\partial_\infty Y$ (and, consequently, also on $\Lambda_{\tau_{\text{mod}}}(\Gamma)$) is finite and non-null.

In the general case where the positivity of $d_{\bar{\theta}}$ is unknown, the above argument still works after some modifications. Let us go back to our construction in the above paragraph. Let Y be a uniform Riemannian neighborhood of Z in which Z is quasiconvex w.r.t. $d_{\bar{\theta}}$. Define a new Γ -invariant metric $\bar{d}_{\bar{\theta}}$ on Y by

$$\bar{d}_{\bar{\theta}}(y, z) = \max \{d_{\bar{\theta}}(y, z), \varepsilon d_{\text{Riem}}(y, z)\}, \quad \forall y, z \in Y,$$

where $\varepsilon > 0$ is some number that is strictly lesser than L^{-1} given in (2.6). Note that for $y, z \in Z$, if $d_{\bar{\theta}}(y, z)$ is sufficiently large, then $\bar{d}_{\bar{\theta}}(y, z) = d_{\bar{\theta}}(y, z)$. Moreover, for a given ι -invariant compact subset $\Theta \subset \text{ost}(\tau_{\text{mod}})$ and a possibly smaller ε (depending on the choice of Θ), any Θ -Finsler geodesic (see Definition 2.2) connecting these two points remains a geodesic in this new metric. In other words, Z remains quasiconvex in Y with respect to $\bar{d}_{\bar{\theta}}$.

Observe that the identity embedding $(Z, d_{\bar{\theta}}) \rightarrow (Y, \bar{d}_{\bar{\theta}})$ is a $(1, A)$ -quasiisometric embedding for some large enough A and the image is Hausdorff-close to Y . Therefore, in this case also we get a natural identification of the Gromov boundaries of $(Z, d_{\bar{\theta}})$ and $(Y, \bar{d}_{\bar{\theta}})$. Next, considering intrinsic metrics, we complete Y as before to get a proper, geodesic, Gromov hyperbolic space (Y, d) . The rest of the argument works as before.

Using Proposition 9.2 together with the remark after the proposition, we obtain the following result.

Theorem 9.5. *Suppose that Γ is a nonelementary τ_{mod} -Anosov subgroup of G . If $\beta = \delta_{\bar{\theta}}/\epsilon$, then the β -dimensional Hausdorff density $\mathcal{H}^{\bar{\theta}, \epsilon, \beta} = \{\mathcal{H}_z^{\bar{\theta}, \epsilon, \beta} \mid z \in X\}$ is a Γ -invariant $\bar{\theta}$ -conformal density of dimension $\delta_{\bar{\theta}}$. In particular, the Hausdorff dimension with respect to the metric $D_x^{\bar{\theta}, \epsilon}$ satisfies*

$$\dim_{\text{Haus}}^{\bar{\theta}, \epsilon}(\Lambda_{\tau_{\text{mod}}}(\Gamma)) = \frac{\delta_{\bar{\theta}}}{\epsilon}.$$

Moreover, $\mathcal{H}^{\bar{\theta}, \epsilon, \beta}$ equals to a non-zero multiple of the Patterson–Sullivan density of type $\bar{\theta}$ corresponding to Γ .

We have mostly completed the proof of this theorem. The only remaining “moreover” part follows from the uniqueness of the Γ -invariant $\bar{\theta}$ -conformal density (Theorem 8.4).

Corollary 9.6. *Let Γ be a τ_{mod} -Anosov subgroup of G . For all $x \in X$, the Hausdorff dimension of the premetric space $(\Lambda_{\tau_{\text{mod}}}(\Gamma), D_x^{\bar{\theta}, 1})$ equals to $\delta_{\bar{\theta}}$.*

10. EXAMPLES

10.1. Product of two hyperbolic planes. Let Γ_1, Γ_2 be isomorphic discrete cocompact subgroups of $\text{PSL}(2, \mathbb{R})$ where the isomorphism is given by $\phi : \Gamma_1 \rightarrow \Gamma_2$. We let $f : S^1 \rightarrow S^1$ be the equivariant homeomorphism of ideal boundaries of hyperbolic planes determined by ϕ .

The discrete subgroup

$$\Gamma = \{(\gamma_1, \phi\gamma_1) \mid \gamma_1 \in \Gamma_1\} < G = \text{PSL}(2, \mathbb{R}) \times \text{PSL}(2, \mathbb{R})$$

acts on $X = \mathbb{H}^2 \times \mathbb{H}^2$ as a σ_{mod} -Anosov subgroup. (This follows, for instance, from the fact that Γ is an URU subgroup of G .) The σ_{mod} -limit set of Γ in the full flag manifold $S^1 \times S^1$ equals the graph of the map f .

We denote by d_1 (resp. d_2) the distance functions of the constant -1 curvature Riemannian metrics on the first (resp. second) factor of the product $\mathbb{H}^2 \times \mathbb{H}^2$.

Unlike in Section 5, we work with the Finsler metric on $\mathbb{H}^2 \times \mathbb{H}^2$ given by

$$(10.1) \quad d_{\bar{\theta}}((x_1, x_2), (y_1, y_2)) = \frac{d_1(x_1, y_1) + d_2(x_2, y_2)}{2}.$$

Basically, we have multiplied the distance function in (2.7) corresponding to $\bar{\theta} = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$, for $p = 2$, by a factor $1/\sqrt{2}$ in order to avoid cumbersome radical constants.

By the formula of the Gromov predistance (5.8), for $\epsilon = 1$ and $x = (x_1, x_2)$, $D_x^{\bar{\theta}, 1}(\tau_+, \tau_-)$ is bilipschitz equivalent to the product

$$\sqrt{\alpha_1 \alpha_2},$$

where $\tau_{\pm} = (\xi_1^{\pm}, \xi_2^{\pm})$ and α_i is the angle between ξ_i^+, ξ_i^- as measured from x_i , $i = 1, 2$.

By [4, Thm. 2 & 3] we note that the $\bar{\theta}$ -critical exponent $\delta_{\bar{\theta}}$ of Γ is at most 1. This can also be obtained by comparing the Hausdorff dimensions as follows. Note that by the formula of the Gromov predistance, the identity map

$$(S^1 \times S^1, \rho) \rightarrow (\text{Flag}(\sigma_{\text{mod}}), D_x^{\bar{\theta}, 1})$$

is Lipschitz, where ρ is a Riemannian distance function on $S^1 \times S^1 = \partial_{\infty} \mathbb{H}^2 \times \partial_{\infty} \mathbb{H}^2$. Moreover, the limit set of Γ in $S^1 \times S^1$ is the graph of a BV (i.e., bounded variation) function, hence, is a rectifiable curve, and, thus, has Hausdorff dimension 1 with respect to ρ . Consequently, with respect to $D_x^{\bar{\theta}, 1}$, $\dim_{\text{Haus}}(\Lambda_{\sigma_{\text{mod}}}(\Gamma)) \leq 1$. By Theorem 9.5, $\delta_{\bar{\theta}} \leq 1$ as well.

Moreover, by [4, Thm. 2] and Corollary 6.5, $\delta_{\bar{\theta}} = 1$ if and only if ϕ is induced by an isometry of \mathbb{H}^2 , equivalently, f is a Möbius transformation.

We further note that one can use [8] as an alternative argument for both inequality and the equality case.

10.2. Hilbert entropy of projective Anosov representations. A subgroup $\Gamma < \text{SL}(k+1, \mathbb{R})$, $k \geq 2$, is called *projective Anosov* if it is τ_{mod} -Anosov for $\tau_{\text{mod}} = (1, k)$ (see Examples 1.9, 2.9, and 5.10 for notations). The $\bar{\theta}$ -critical exponent associated to the unique ι -invariant type

$$\bar{\theta} = \left(\frac{1}{2\sqrt{k+1}}, 0, -\frac{1}{2\sqrt{k+1}} \right)$$

in τ_{mod} will be denoted, as usual, by $\delta_{\bar{\theta}}$.

Let $\Gamma < \text{SL}(k+1, \mathbb{R})$ be a projective Anosov subgroup. In [21], the authors defined the following two critical exponents of Γ , namely, the *Hilbert critical exponent* (corresponding to the sum of all simple roots)

$$\delta_{1, k+1} = \limsup_{r \rightarrow \infty} \frac{\log \text{card}\{\gamma \in \Gamma \mid \sigma_1(\gamma) - \sigma_{k+1}(\gamma) < r\}}{r}$$

and the *simple root critical exponent* (corresponding to the first simple root)

$$\delta_{1, 2} = \limsup_{r \rightarrow \infty} \frac{\log \text{card}\{\gamma \in \Gamma \mid \sigma_1(\gamma) - \sigma_2(\gamma) < r\}}{r}.$$

A direct computation yields

$$\sqrt{k+1}\delta_{\bar{\theta}} = \delta_{1,k+1},$$

which follows from the formula of $d_{\bar{\theta}}$ given by (2.10). Also note that (by (5.9)) for a pair of partial flags $(l_1, h_1), (l_2, h_2) \in \text{Flag}(\tau_{\text{mod}})$,

$$(10.2) \quad D_x^{\bar{\theta}, 1/\sqrt{k+1}}((l_1, h_1), (l_2, h_2)) = \sqrt{\sin \angle(l_1, h_2)} \sqrt{\sin \angle(l_2, h_1)} \leq \sin \angle(l_1, l_2),$$

where the right side equals the distance (with respect to the constant curvature Riemannian metric on $\mathbb{R}P^k$ determined by $x \in X$) between the points l_1, l_2 in $\mathbb{R}P^k$. This together with Theorem 9.5 implies that

$$(10.3) \quad \delta_{1,k+1} = \sqrt{k+1}\delta_{\bar{\theta}} = \dim_{\text{Haus}}^{\bar{\theta}, 1/\sqrt{k+1}}(\Lambda_{\tau_{\text{mod}}}(\Gamma)) \leq \dim_{\text{Haus}}^{\text{Riem}}(\xi^1(\partial_{\infty}\Gamma)),$$

where $\xi^1 : \partial_{\infty}\Gamma \rightarrow \mathbb{R}P^k$ is the Γ -equivariant embedding²⁶ of $\partial_{\infty}\Gamma$ into $\mathbb{R}P^k$ and $\dim_{\text{Haus}}^{\text{Riem}}$ denotes the Hausdorff dimension with respect to the Riemannian metric.

The critical exponent $\delta_{1,2}$ is known to give an upper bound for $\dim_{\text{Haus}}^{\text{Riem}}(\xi^1(\partial_{\infty}\Gamma))$ (see [41, Prop. 4.1] or [21, Thm. 4.1]). By above, we obtain a lower bound.

Theorem 10.1. *Let $\Gamma < \text{SL}(k+1, \mathbb{R})$ be a projective Anosov subgroup. Then*

$$\delta_{1,k+1} \leq \dim_{\text{Haus}}^{\text{Riem}}(\xi^1(\partial_{\infty}\Gamma)) \leq \dim \mathbb{R}P^k = k.$$

Remark 10.2. If one considers the τ_{mod} -flag limit set $\Lambda_{\tau_{\text{mod}}}$ of Γ in the flag manifold $\text{Flag}(\tau_{\text{mod}})$, then a similar lower bound can be obtained for the Hausdorff dimension corresponding to the Riemannian metric. Note that (10.2) also holds if one replaces $\sin \angle(l_1, l_2)$ by $\sin \angle(h_1, h_2)$. Hence,

$$(10.4) \quad D_x^{\bar{\theta}, 1/\sqrt{k+1}}((l_1, h_1), (l_2, h_2)) \leq \sqrt{2 \sin^2 \angle(l_1, l_2) + 2 \sin^2 \angle(h_1, h_2)}$$

which shows that the identity map $(\Lambda_{\tau_{\text{mod}}}, \rho) \rightarrow (\Lambda_{\tau_{\text{mod}}}, D_x^{\bar{\theta}, 1/\sqrt{k+1}})$ is a Lipschitz map, where ρ is a(ny) Riemannian distance function in $\text{Flag}(\tau_{\text{mod}})$. Therefore,

$$\delta_{1,k+1} \leq \dim_{\text{Haus}}^{\text{Riem}}(\Lambda_{\tau_{\text{mod}}}).$$

This recovers the lower bound obtained in [20, Cor. 1.2].

10.3. Hilbert entropy of $\text{PSL}(3, \mathbb{R})$ -Hitchin representations. In suitable projective Anosov classes, one may be able to improve the inequality in (10.2) to get a better bound for the Hilbert critical exponent $\delta_{1,k+1}$. Here we present an example of such an improvement.

Let $\Gamma = \pi_1(S)$ be a surface²⁷ group. By [9], the $\text{PSL}(3, \mathbb{R})$ -Hitchin representations $\rho : \Gamma \rightarrow \text{PSL}(3, \mathbb{R}) = \text{SL}(3, \mathbb{R})$ consist of holonomies of convex $\mathbb{R}P^2$ -structures in S . In particular, $\rho(\Gamma)$ preserves a properly convex (open) domain Ω in $\mathbb{R}P^2$ with (C^1 -) boundary $\partial\Omega = C$, and the action $\Gamma \curvearrowright \Omega$ is properly discontinuous.

Since Hitchin representations are P_1 -Anosov, Theorem 10.1 shows that

$$\delta_{1,3}(\rho(\Gamma)) \leq 2.$$

However, this *most general* upper bound is weak for the Hitchin representations. In Proposition 10.3 we will obtain a stronger bound.

Let $\tau_{\text{mod}} = (1, 2)$. The τ_{mod} -flag limit set of $\rho(\Gamma)$ can be equivariantly identified with the set of flags $\{(\xi, \xi^*) \mid \xi \in C\}$, where $\xi^* \subset \mathbb{R}P^2$ is the line tangent to C

²⁶Composition of the Γ -equivariant boundary embedding $\partial_{\infty}\Gamma \rightarrow \text{Flag}(\tau_{\text{mod}})$ and the projection map $\text{Flag}(\tau_{\text{mod}}) \rightarrow \mathbb{R}P^k = \text{Gr}_1(\mathbb{R}^{k+1})$.

²⁷More precisely, S is a closed surface of genus $g \geq 2$.




FIGURE 1

through ξ . We denote the dual curve $\{\xi^* \mid \xi \in C\} \subset (\mathbb{R}P^2)^*$ by C^* ; this curve C^* also bounds a properly convex domain in $(\mathbb{R}P^2)^*$.

Claim. There exists a constant L such that, for every $\xi, \eta \in C$,

$$(10.5) \quad D(\xi, \eta) := \sqrt{d_{\mathbb{R}P^2}(\xi, \eta^*)d_{\mathbb{R}P^2}(\eta, \xi^*)} \leq L \cdot d_{\mathbb{R}P^2}(\xi, \eta)d_{(\mathbb{R}P^2)^*}(\xi^*, \eta^*).$$

Proof. The curve C is a simple closed C^1 -curve in an affine chart \mathbb{R}^2 , bounding the convex subset Ω above. Let $d_{\mathbb{R}^2}(\xi, \eta) = \psi$ and $\angle(\eta^*, \xi^*) = \theta$ as in Figure 1. When $\xi \neq \eta$ are uniformly close, $0 < \psi \leq \psi_0$, then the angle θ is acute. See Figure 1. Then,²⁸ $d_{\mathbb{R}^2}(\xi, \eta^*) \asymp \psi\beta$, $d_{\mathbb{R}^2}(\eta, \xi^*) \asymp \psi\alpha$, and $\alpha + \beta = \theta$. Thus,

$$\frac{d_{\mathbb{R}^2}(\xi, \eta^*)d_{\mathbb{R}^2}(\eta, \xi^*)}{d_{\mathbb{R}^2}(\xi, \eta)^2(\angle(\eta^*, \xi^*))^2} \asymp \frac{\psi^2\alpha\beta}{\psi^2(\alpha + \beta)^2} = \frac{\alpha\beta}{(\alpha + \beta)^2} \leq \frac{1}{4}.$$

So,

$$\sqrt{d_{\mathbb{R}^2}(\xi, \eta^*)d_{\mathbb{R}^2}(\eta, \xi^*)} \leq \text{const} \cdot d_{\mathbb{R}^2}(\xi, \eta)(\angle(\eta^*, \xi^*)).$$

Also, since C (resp. C^*) is bounded in the affine chart, the euclidean distances (resp. the angular distance) above are equivalent to the projective distances in (10.5). Hence the above inequality justifies (10.5). \square

Also, note that the premetric D in left side of (10.5) is bilipschitz equivalent to the premetric $D_x^{\theta, 1/\sqrt{3}}$ (see the formula in (10.2)), left side of the inequality in (10.4). But, the square root of the right side of (10.5) is

$$\sqrt{L} \sqrt{d_{\mathbb{R}P^2}(\xi, \eta)d_{(\mathbb{R}P^2)^*}(\xi^*, \eta^*)} \leq \sqrt{\frac{L}{2}} \sqrt{d_{\mathbb{R}P^2}(\xi, \eta)^2 + d_{(\mathbb{R}P^2)^*}(\xi^*, \eta^*)^2}.$$

In the right side of the above inequality, we obtain a multiple of the Riemannian distance in $\text{Flag}(\tau_{\text{mod}})$. Thus, the identity map

$$(\Lambda_{\tau_{\text{mod}}}, \rho) \rightarrow (\Lambda_{\tau_{\text{mod}}}, D_x^{\theta, 1/\sqrt{3}})$$

is a 2-Hölder map. Here ρ denotes the distance function of the Riemannian metric on $\text{Flag}(\tau_{\text{mod}})$. It is known that $\Lambda_{\tau_{\text{mod}}} \subset \text{Flag}(\tau_{\text{mod}})$ is a Lipschitz curve, and hence $\dim_{\text{Haus}}^{\text{Riem}}(\Lambda_{\tau_{\text{mod}}}(\Gamma)) = 1$.

²⁸We remind our reader that the symbol \asymp is used to mean that the ratio of both sides is bounded above and below by some positive constants.

We obtain

$$\delta_{1,3}(\rho(\Gamma)) = \dim_{\text{Haus}}^{\bar{\theta}, 1/\sqrt{3}}(\Lambda_{\tau_{\text{mod}}}(\Gamma)) \leq \frac{\dim_{\text{Haus}}^{\text{Riem}}(\Lambda_{\tau_{\text{mod}}}(\Gamma))}{2} = \frac{1}{2}$$

(cf. (10.3)). We record this result in Proposition 10.3.

Proposition 10.3. *For any Hitchin representation $\rho : \Gamma \rightarrow \text{PSL}(3, \mathbb{R})$,*

$$\delta_{1,3}(\rho(\Gamma)) \leq \frac{1}{2}.$$

Compare with the last paragraph of [40, p. 892].

APPENDIX A. HAUSDORFF MEASURES ON PREMETRIC SPACES

Let X be a metrizable topological space. Recall that an *outer measure* is a function $\mu : \mathcal{P}(X) \rightarrow [0, \infty]$ that satisfies

- (i) $\mu(\emptyset) = 0$,
- (ii) for all $A, B \in \mathcal{P}(X)$ with $A \subset B$, $\mu(A) \leq \mu(B)$, and
- (iii) for all countable collection $\{A_k \mid k \in \mathbb{N}\}$ of subsets of X ,

$$\mu\left(\bigcup_{k \in \mathbb{N}} A_k\right) \leq \sum_{k \in \mathbb{N}} \mu(A_k).$$

A set $A \subset X$ is called μ -*measurable* if for every $E \in \mathcal{P}(X)$, $\mu(A) = \mu(A \cap E) + \mu(A \cap E^c)$. By Carathéodory's theorem (cf. [18, Thm. 1.11]), μ -measurable sets form a σ -algebra to which μ restricts as a complete measure.

Assume now that X is compact. The outer measure μ is called *good* if additionally,

- (iv) for all $A, B \subset X$ with $\bar{A} \cap \bar{B} = \emptyset$, $\mu(A \cup B) = \mu(A) + \mu(B)$.

Lemma A.1 asserts that, for outer measures μ on compact metrizable spaces, the σ -algebra of Borel sets is a subalgebra of the σ -algebra of μ -measurable sets.

Lemma A.1. *Let X be a compact metrizable space. If μ is a good outer measure on X , then every Borel set $B \in \mathfrak{B}(X)$ is measurable.*

Proof. Let d be a metric on X . Then the condition (iv) implies that

- (iv') for all $A, B \subset X$ with $d(A, B) > 0$, $\mu(A \cup B) = \mu(A) + \mu(B)$.

Therefore, μ is a metric outer measure on (X, d) . By [18, Prop. 11.16], Borel subsets of X are measurable. \square

Definition A.2 (Premetric space). Let X be a topological space. A symmetric continuous function $d : X \times X \rightarrow [0, \infty]$ is called a *premetric* on X . A pair (X, d) consisting of a metrizable topological space X and a premetric d on X is called a *premetric space*.

In what follows, we consider only *positive* premetrics, i.e.,

$$d(x, y) > 0 \iff x \neq y, \quad \forall x, y \in X.$$

Let (X, d) be a compact positive premetric space. Then d satisfies the following *separation property*:

$$(A.1) \quad d(A, B) > 0 \iff \bar{A} \cap \bar{B} = \emptyset, \quad \forall A, B \subset X.$$

Let $\varepsilon > 0$, $\beta > 0$. For every $A \subset X$, define

$$\mathcal{H}_\varepsilon^\beta(A) = \inf_{\mathcal{U}} \left\{ \sum_{k \in \mathbb{N}} \text{diam}_d(U_k)^\beta \mid \mathcal{U} = \{U_k \mid k \in \mathbb{N}\} \text{ covers } A, \text{ mesh}(\mathcal{U}) \leq \varepsilon \right\}.$$

In the above, $\text{mesh}(\mathcal{U})$ is the supremum of the d -diameters of the members of \mathcal{U} . Then

$$\mathcal{H}_\varepsilon^\beta : \mathcal{P}(X) \rightarrow [0, \infty]$$

is an outer measure on X (cf. [18, Prop. 1.10]). Define the β -dimensional Hausdorff measure \mathcal{H}^β by

$$\mathcal{H}^\beta(A) = \lim_{\varepsilon \rightarrow 0} \mathcal{H}_\varepsilon^\beta(A).$$

Theorem A.3. *The Hausdorff measure \mathcal{H}^β is a good outer measure.*

Proof. We need to check the properties (i)–(iv). Since, for all $\varepsilon > 0$, $\mathcal{H}_\varepsilon^\beta$ is an outer measure, taking limit $\varepsilon \rightarrow 0$, properties (i)–(iii) are easily verified. Therefore, we only need to check that \mathcal{H}^β satisfies property (iv).

Let $A, B \subset X$ such that $A \cap B = \emptyset$. By (A.1), $d(A, B) = d_0 > 0$. Let $\varepsilon < d_0$ be a positive number and \mathcal{U} be a countable open cover of $A \cup B$ with $\text{mesh}(\mathcal{U}) \leq \varepsilon$. If such open cover does not exist, then $\mathcal{H}_\varepsilon^\beta(A \cup B)$ (and hence, $\mathcal{H}^\beta(A \cup B)$) is infinity. Otherwise, \mathcal{U} can be written as a disjoint union $\mathcal{U}_A \sqcup \mathcal{U}_B$ where \mathcal{U}_A consists of all open sets in \mathcal{U} that intersect A and \mathcal{U}_B consists of the rest. Clearly, \mathcal{U}_A and \mathcal{U}_B are open covers of A and B , respectively. Therefore,

$$\sum_{E \in \mathcal{U}} \text{diam}_d(E)^\beta = \sum_{E \in \mathcal{U}_A} \text{diam}_d(E)^\beta + \sum_{E \in \mathcal{U}_B} \text{diam}_d(E)^\beta \geq \mathcal{H}_\varepsilon^\beta(A) + \mathcal{H}_\varepsilon^\beta(B).$$

Since the above holds for any cover \mathcal{U} with $\text{mesh} \leq \varepsilon$, we have

$$\mathcal{H}_\varepsilon^\beta(A \cup B) \geq \mathcal{H}_\varepsilon^\beta(A) + \mathcal{H}_\varepsilon^\beta(B).$$

Taking limit $\varepsilon \rightarrow 0$, we get $\mathcal{H}^\beta(A \cup B) \geq \mathcal{H}^\beta(A) + \mathcal{H}^\beta(B)$. The reverse inequality follows from property (iii). Therefore, $\mathcal{H}^\beta(A \cup B) = \mathcal{H}^\beta(A) + \mathcal{H}^\beta(B)$. This completes the proof. \square

By Lemma A.1 and Theorem A.3, we obtain the following result.

Corollary A.4. *Every Borel subset of X is \mathcal{H}^β -measurable.*

The Hausdorff dimension of a Borel subset $B \subset (X, d)$ is then defined as

$$\dim_{\text{Haus}}(B) = \inf\{\beta \mid \mathcal{H}^\beta(B) = 0\} = \sup\{\beta \mid \mathcal{H}^\beta(B) = \infty\}.$$

APPENDIX B. A BRIEF OVERVIEW OF THE HISTORY OF THE PATTERSON–SULLIVAN THEORY IN HIGHER RANK

For reader's convenience, in this appendix we discuss connection of our work with some notable earlier papers on the Patterson–Sullivan theory for higher rank symmetric spaces (of noncompact type).

To the best of our knowledge, Bishop–Steger [4] and Burger [8] were the first to investigate Poincaré series associated with a discrete isometry group Γ of a symmetric space of rank ≥ 2 . Precisely, for a pair of Fuchsian subgroups $\Gamma_i < \text{Isom}(\mathbb{H}^2)$, $i = 1, 2$, and an isomorphism $\phi : \Gamma_1 \rightarrow \Gamma_2$, Burger took the subgroup

$\Gamma = \{(\gamma_1, \phi(\gamma_1)) \mid \gamma_1 \in \Gamma_1\} < \text{Isom}(\mathbb{H}^2 \times \mathbb{H}^2)$ and considered the *Finsler Poincaré series*

$$\sum_{(\gamma_1, \gamma_2) \in \Gamma} \exp(-s(ad_1(\gamma_1 x_1, x_1) + bd_2(\gamma_2 x_2, x_2))), \quad (a, b) \in \mathbb{R}_+^2,$$

where d_1 (resp. d_2) and x_1 (resp. x_2) denote the distance function and a fixed point, respectively, in the first (resp. second) factor of $\mathbb{H}^2 \times \mathbb{H}^2$. Then he studied the set of all points $(a, b) \in \mathbb{R}_+^2$ for which the Poincaré series has the critical exponent $s = 1$ which he called the “Manhattan curve.” He proved that each point (a, b) in this set gives rise to a unique “ (a, b) -dimensional” density with respect to which the action $\Gamma \curvearrowright \partial_\infty \mathbb{H}^2 \times \partial_\infty \mathbb{H}^2$ is ergodic, [8, Thm. 4]. Theorem A of our paper illustrates this result, compare Subsection 10.1, although in that example we only consider the case $a = b$, the general case can also be obtained by suitably changing the weights in the formula (10.1) for the Finsler metric. On the other hand, Bishop and Steger considered the closely related Poincaré series

$$\sum_{(\gamma_1, \gamma_2) \in \Gamma} \exp(-sd_1(\gamma_1 x_1, x_1) - (1-s)d_2(\gamma_2 x_2, x_2)), \quad s \in (0, 1),$$

and showed that it diverges if and only if ϕ is induced by an isometry of \mathbb{H}^2 . We discussed the connection of this result with our work in Subsection 10.1 for $s = 1/2$; for a general s , one needs to modify the Finsler metric as above.

Patterson–Sullivan measures for general symmetric spaces X were introduced and studied by Albuquerque [1]. His main result is that for a generic²⁹ Zariski-dense discrete subgroup Γ of $G = \text{Isom}_0(X)$, the support of any Patterson–Sullivan density of dimension $\delta_{\text{Riem}}(\Gamma)$ in the visual boundary of X lies in a single regular G -orbit $G \cdot \xi_0$ where ξ_0 is the direction in the Weyl chamber along which the growth-rate of Γ -orbits in X is “maximal.” More precisely, any Patterson–Sullivan density of dimension $\delta_{\text{Riem}}(\Gamma)$ is supported on $G \cdot \xi_0 \cap \Lambda(\Gamma)$. He also proved that the critical exponent $\delta_{\bar{\theta}}(\Gamma)$ defined in terms of the Finsler pseudometric induced by the linear functional in the Weyl chamber dual to a vector $\xi \in \sigma_{\text{mod}}$ is minimal precisely when $\xi = \xi_0$. Albuquerque further showed that this minimal Finsler critical exponent equals to the Riemannian critical exponent.

Quint [42, 43] generalized Albuquerque’s results to arbitrary Zariski-dense discrete subgroups $\Gamma < G$ and to general partial flag-manifolds $G/P_{\tau_{\text{mod}}}$. For such a group Γ , he defined the *indicator of growth* function

$$\psi_\Gamma : \Delta \rightarrow \mathbb{R} \cup \{-\infty\},$$

which can be regarded as a higher rank analogue of the critical exponent, and showed that it is strictly positive in the interior of the Benoist’s limit cone of Γ . Note, however, that the orbital counting (done in the definition of the function ψ_Γ) in Quint’s paper is different from ours since he takes the infimum of exponents of convergence $\tau_{\mathcal{C}}$ over certain open cones $\mathcal{C} \subset \Delta$. For a positive linear function ϕ on Δ satisfying $\phi \geq \psi_\Gamma$ and $\phi(x) = \psi_\Gamma(x)$ for some $x \in \text{int}(\Delta)$, Quint (using the Patterson–Sullivan construction) defined a “ (Γ, ϕ) -Patterson measure” supported on the flag-limit set of Γ in G/P_θ . Observe that each ϕ defines a G -invariant polyhedral Finsler metric on X by the composition $\phi \circ d_\Delta$. This part of Quint’s

²⁹Here Γ is called generic if the support of any Patterson–Sullivan density lies in the regular part of the visual boundary.

work has nontrivial overlap with ours (specifically, the construction of Patterson–Sullivan densities in Section 3 of our paper). However, neither work subsumes the other since we do not assume Zariski density but require subgroups to be τ_{mod} -RA, while Quint does not assume the τ_{mod} -RA property but requires Zariski density. Quint’s work contains wealth of other results such as a proof of the concavity property of ψ_Γ and its applications: These issues are not discussed at all in our paper.

Taking inspiration from [8] and building on [1], Link [34] considered a class of densities (coming from the Patterson–Sullivan construction) on the visual boundary of X associated with a Zariski-dense discrete subgroup $\Gamma < \text{Isom}_0(X)$ that generalize the conformal densities. After introducing a notion of Hausdorff measures appropriate for her setup, Link showed that for a regular point $\xi \in \partial_\infty X$, the Hausdorff dimension of $G \cdot \xi \cap \Lambda(\Gamma)$ is bounded above by a suitable exponent of growth of Γ -orbits in X ; she also proved the equality for a certain class of groups which she calls “radially cocompact” (see [34, Sec. 6]). In her subsequent work [35], Link proved that the action of Γ on the “ray-limit set” is ergodic³⁰ with respect to these generalized Patterson–Sullivan measures.³¹ These results are parallel to the equality of the Finsler critical exponent and the Hausdorff dimension of flag-limit sets in $G/P_{\tau_{\text{mod}}}$ for τ_{mod} -Anosov subgroups (Theorem 9.5) and the ergodicity theorem (Theorem 8.3) proved in our paper. However, the conicality condition for limit points of Anosov subgroups is a vast relaxation of Link’s notion of radial limit points. Moreover, it follows from the main theorem of [2] and Lemma 7.4 that, say, a Zariski dense τ_{mod} -Anosov subgroup can never be radially cocompact unless G has rank one.

To conclude the discussion, we would like to repeat that Albuquerque, Quint and Link treat general Zariski-dense discrete subgroups, while we study Anosov and, more generally, regular antipodal subgroups, but do not assume Zariski density. While many of our proofs work for general τ_{mod} -RA subgroups, the Anosov property is critical in several places, for instance, in the proof of vanishing of $\delta_\theta^{\text{con}}$ (Proposition 7.3), the proof of ergodicity, the proof that (in a suitable range) the Gromov pre-metric on the limit set is a metric, etc. We would also like to point out that, even in the Zariski-dense case, parts (ii), (iv) and (v) (ergodicity, divergence and relation of Finsler critical exponent and Hausdorff dimension of the limit set) in our Theorem A are new since, to our knowledge, the most general case in this direction was considered by Link [35] but was conditioned on the density of the ray-limit set.

APPENDIX C. A DISCUSSION OF THE MAIN RESULTS WITHOUT THE ι -INVARIANCE CONDITION ON THE TYPE $\bar{\theta}$

In this last appendix, on our referee’s suggestion, we show that Theorem A (except for item (v)), and Theorem B hold true without the ι -invariance condition that we imposed on the type $\bar{\theta} \in \text{int}(\tau_{\text{mod}})$.

Suppose that $\bar{\theta} \in \sigma_{\text{mod}}$ is a type which is not assumed to be ι -invariant. Then the G -invariant pseudo-metric $d_{\bar{\theta}} : X \times X \rightarrow [0, \infty)$ defined in (2.1) is in general

³⁰More precisely, she proves that if A is a measurable Γ -invariant subset of the ray-limit set, then either A is a null set or the complement of A in the full limit set is null. However, this result does not exclude the possibility that the ray-limit set itself is a null set for the generalized Patterson–Sullivan measures.

³¹In fact, Albuquerque [1] also attempted to prove ergodicity, but there was a gap in the proof. This gap was discovered and fixed by Link [35, p. 612].

asymmetric. However, the following still holds: Since $d_\Delta(y, x) = \iota d_\Delta(x, y)$, we have

$$(C.1) \quad d_{\bar{\theta}}(y, x) = d_{\iota\bar{\theta}}(x, y), \quad \forall x, y \in X.$$

The inequality (2.2) still holds. The function $d_{\bar{\theta}}$ satisfies the triangle inequality for the asymmetric distance functions:

$$d_{\bar{\theta}}(x, z) \leq d_{\bar{\theta}}(x, y) + d_{\bar{\theta}}(y, z), \quad \forall x, y, z \in X.$$

See [26, Subsec. 5.1.2] for more details.

Let $\bar{\theta} \in \text{int } \tau_{\text{mod}}$. In the asymmetric case, we define the $\bar{\theta}$ -critical exponent $\delta_{\bar{\theta}}$ of a discrete group $\Gamma < G$ in the same way as it was done for the symmetric case in Section 2, see (2.3). As a consequence of the above triangle inequality, the critical exponent does not depend on the choice of a base-point in X (see the paragraph after Remark 2.4). The definition of $\bar{\theta}$ -convergence/divergence type is also same as in the symmetric case (see Definition 2.5). Proposition 2.6 for uniformly τ_{mod} -regular groups Γ , which only depends on the fact that d_{Riem} and $d_{\bar{\theta}}$ are coarsely equivalent on an Γ -orbit in X , is valid in this case.

All the definitions (in particular, the crucial definition of $\bar{\theta}$ -conformal densities) and results in Section 3 remain valid in the asymmetric case.

We leave the results in Sections 4 and 5 as they are; those results are mostly independent of Sections 6, 7, and 8. We remark in passing that an asymmetric Gromov product on $\text{Flag}(\tau_{\text{mod}})$ can be defined as follows: Define the Gromov product with respect to a base point $x \in X$ by

$$(C.2) \quad \langle \tau_+ | \tau_- \rangle_x^{\bar{\theta}} = \frac{1}{2} \left(\mathcal{B}_{\tau_+}^{\bar{\theta}}(x, z) + \mathcal{B}_{\tau_-}^{\iota\bar{\theta}}(x, z) \right), \quad \tau_{\pm} \in \text{Flag}(\tau_{\text{mod}}) \text{ are antipodal,}$$

where z is some point on the parallel set $P(\tau_+, \tau_-)$ spanned by τ_{\pm} . Following the proof of Lemma 5.1, it can be checked that the definition in (C.2) does not depend on the choice of $z \in P(\tau_+, \tau_-)$. The above leads to an asymmetric Gromov premetric $D^{\bar{\theta}, \epsilon}$ in $\text{Flag}(\tau_{\text{mod}})$ (cf. Definition 5.2) which satisfies

$$D_x^{\bar{\theta}, \epsilon}(\tau_1, \tau_2) = D_x^{\iota\bar{\theta}, \epsilon}(\tau_2, \tau_1), \quad \forall \tau_1, \tau_2 \in \text{Flag}(\tau_{\text{mod}}).$$

The results in Sections 6 and 7 are valid verbatim in the asymmetric case. In the proof of Theorem 8.3 in Section 8, we only need to modify the justification of Claim 3 in the proof of the crucial Sublemma 8.7 in the asymmetric case. The statement in that claim is: *If $S(x : B(\gamma x_0, r)) \cap S(x : B(\phi x_0, r)) \neq \emptyset$, for $\gamma, \phi \in \Gamma_d^*$, then $d_{\bar{\theta}}(\gamma x_0, \phi x_0)$ is uniformly bounded.* The proof of this claim uses the main result of Section 4 and it was given for symmetric functions $d_{\bar{\theta}}$ (note that in that proof, the requirement that $\bar{\theta}$ has unit length is unimportant, only the hyperbolicity of Γ -orbits in X is relevant). When $d_{\bar{\theta}}$ is asymmetric, we consider the *symmetrization* function $d_{\bar{\theta}}^{\text{sym}} = (d_{\bar{\theta}} + d_{\iota\bar{\theta}})/2$,

$$d_{\bar{\theta}}^{\text{sym}}(x, y) = \left\langle d_\Delta(x, y) \left| \frac{\bar{\theta} + \iota\bar{\theta}}{2} \right. \right\rangle, \quad x, y \in X,$$

and the previous proof implies that, under the hypothesis, $d_{\bar{\theta}}^{\text{sym}}(\gamma x_0, \phi x_0)$ is uniformly bounded. Since $d_{\bar{\theta}} \leq 2d_{\bar{\theta}}^{\text{sym}}$, $d_{\bar{\theta}}(\gamma x_0, \phi x_0)$ is also uniformly bounded under the same hypothesis.

ACKNOWLEDGMENTS

This project was a part of the first author’s dissertation work at UC Davis. We are grateful to Olivier Glorieux and Hee Oh for pointing out some mistakes in a previous version of this paper and to Andrés Sambarino for telling us about Ledrappier’s work and related results. Finally, we are thankful to the anonymous referee for carefully reading our paper and making numerous suggestions to improve the text.

REFERENCES

- [1] P. Albuquerque, *Patterson-Sullivan theory in higher rank symmetric spaces*, *Geom. Funct. Anal.* **9** (1999), no. 1, 1–28, DOI 10.1007/s000390050079. MR1675889
- [2] Y. Benoist, *Propriétés asymptotiques des groupes linéaires* (French, with English and French summaries), *Geom. Funct. Anal.* **7** (1997), no. 1, 1–47, DOI 10.1007/PL00001613. MR1437472
- [3] J. Beyrer, *Cross ratios on boundaries of symmetric spaces and Euclidean buildings*, *Transform. Groups* **26** (2021), no. 1, 31–68, DOI 10.1007/s00031-020-09549-5. MR4229658
- [4] Christopher Bishop and Tim Steger, *Representation-theoretic rigidity in $\mathrm{PSL}(2, \mathbf{R})$* , *Acta Math.* **170** (1993), no. 1, 121–149, DOI 10.1007/BF02392456. MR1208564
- [5] Christopher J. Bishop and Peter W. Jones, *Hausdorff dimension and Kleinian groups*, *Acta Math.* **179** (1997), no. 1, 1–39, DOI 10.1007/BF02392718. MR1484767
- [6] Richard L. Bishop and Richard J. Crittenden, *Geometry of manifolds*, AMS Chelsea Publishing, Providence, RI, 2001. Reprint of the 1964 original, DOI 10.1090/chel/344. MR1852066
- [7] Martin Bridgeman, Richard Canary, François Labourie, and Andres Sambarino, *The pressure metric for Anosov representations*, *Geom. Funct. Anal.* **25** (2015), no. 4, 1089–1179, DOI 10.1007/s00039-015-0333-8. MR3385630
- [8] Marc Burger, *Intersection, the Manhattan curve, and Patterson-Sullivan theory in rank 2*, *Internat. Math. Res. Notices* **7** (1993), 217–225, DOI 10.1155/S1073792893000236. MR1230298
- [9] Suhyoung Choi and William M. Goldman, *Convex real projective structures on closed surfaces are closed*, *Proc. Amer. Math. Soc.* **118** (1993), no. 2, 657–661, DOI 10.2307/2160352. MR1145415
- [10] Michel Coornaert, *Mesures de Patterson-Sullivan sur le bord d’un espace hyperbolique au sens de Gromov* (French, with French summary), *Pacific J. Math.* **159** (1993), no. 2, 241–270. MR1214072
- [11] Kevin Corlette, *Hausdorff dimensions of limit sets. I*, *Invent. Math.* **102** (1990), no. 3, 521–541, DOI 10.1007/BF01233439. MR1074486
- [12] Kevin Corlette and Alessandra Iozzi, *Limit sets of discrete groups of isometries of exotic hyperbolic spaces*, *Trans. Amer. Math. Soc.* **351** (1999), no. 4, 1507–1530, DOI 10.1090/S0002-9947-99-02113-3. MR1458321
- [13] Tushar Das, David Simmons, and Mariusz Urbański, *Geometry and dynamics in Gromov hyperbolic metric spaces*, *Mathematical Surveys and Monographs*, vol. 218, American Mathematical Society, Providence, RI, 2017. With an emphasis on non-proper settings, DOI 10.1090/surv/218. MR3558533
- [14] Subhadip Dey, Michael Kapovich, and Bernhard Leeb, *A combination theorem for Anosov subgroups*, *Math. Z.* **293** (2019), no. 1-2, 551–578, DOI 10.1007/s00209-018-2208-9. MR4002289
- [15] Cornelia Druţu and Michael Kapovich, *Geometric group theory*, *American Mathematical Society Colloquium Publications*, vol. 63, American Mathematical Society, Providence, RI, 2018. With an appendix by Bogdan Nica, DOI 10.1090/coll/063. MR3753580
- [16] Patrick B. Eberlein, *Geometry of nonpositively curved manifolds*, *Chicago Lectures in Mathematics*, University of Chicago Press, Chicago, IL, 1996. MR1441541
- [17] Herbert Federer, *Geometric measure theory*, *Die Grundlehren der mathematischen Wissenschaften*, Band 153, Springer-Verlag New York, Inc., New York, 1969. MR0257325
- [18] Gerald B. Folland, *Real analysis*, 2nd ed., *Pure and Applied Mathematics (New York)*, John Wiley & Sons, Inc., New York, 1999. Modern techniques and their applications; A Wiley-Interscience Publication. MR1681462

- [19] F. W. Gehring and G. J. Martin, *Discrete quasiconformal groups. I*, Proc. London Math. Soc. (3) **55** (1987), no. 2, 331–358, DOI 10.1093/plms/s3-55_2.331. MR896224
- [20] Olivier Glorieux and Daniel Monclair, *Critical exponent and Hausdorff dimension in pseudo-Riemannian hyperbolic geometry*, 2016.
- [21] Olivier Glorieux, Daniel Monclair, and Nicolas Tholozan, *Hausdorff dimension of limit sets for projective Anosov representations*, arXiv:1902.01844, 2019.
- [22] M. Gromov, *Hyperbolic groups*, Essays in group theory, Math. Sci. Res. Inst. Publ., vol. 8, Springer, New York, 1987, pp. 75–263, DOI 10.1007/978-1-4613-9586-7_3. MR919829
- [23] Olivier Guichard and Anna Wienhard, *Anosov representations: domains of discontinuity and applications*, Invent. Math. **190** (2012), no. 2, 357–438, DOI 10.1007/s00222-012-0382-7. MR2981818
- [24] Toshiaki Hattori, *Collapsing of quotient spaces of $SO(n) \backslash SL(n, \mathbf{R})$ at infinity*, J. Math. Soc. Japan **47** (1995), no. 2, 193–225, DOI 10.2969/jmsj/04720193. MR1317280
- [25] N. J. Hitchin, *Lie groups and Teichmüller space*, Topology **31** (1992), no. 3, 449–473, DOI 10.1016/0040-9383(92)90044-I. MR1174252
- [26] Michael Kapovich and Bernhard Leeb, *Finsler bordifications of symmetric and certain locally symmetric spaces*, Geom. Topol. **22** (2018), no. 5, 2533–2646, DOI 10.2140/gt.2018.22.2533. MR3811766
- [27] Michael Kapovich and Bernhard Leeb, *Relativizing characterizations of Anosov subgroups, I*, arXiv:1807.00160, 2018.
- [28] Michael Kapovich, Bernhard Leeb, and John Millson, *Convex functions on symmetric spaces, side lengths of polygons and the stability inequalities for weighted configurations at infinity*, J. Differential Geom. **81** (2009), no. 2, 297–354. MR2472176
- [29] Michael Kapovich, Bernhard Leeb, and Joan Porti, *Morse actions of discrete groups on symmetric space*, arXiv:1403.7671, 2014.
- [30] Michael Kapovich, Bernhard Leeb, and Joan Porti, *Anosov subgroups: dynamical and geometric characterizations*, Eur. J. Math. **3** (2017), no. 4, 808–898, DOI 10.1007/s40879-017-0192-y. MR3736790
- [31] Michael Kapovich, Bernhard Leeb, and Joan Porti, *A Morse lemma for quasigeodesics in symmetric spaces and Euclidean buildings*, Geom. Topol. **22** (2018), no. 7, 3827–3923, DOI 10.2140/gt.2018.22.3827. MR3890767
- [32] François Labourie, *Anosov flows, surface groups and curves in projective space*, Invent. Math. **165** (2006), no. 1, 51–114, DOI 10.1007/s00222-005-0487-3. MR2221137
- [33] François Ledrappier, *Structure au bord des variétés à courbure négative* (French), Séminaire de Théorie Spectrale et Géométrie, No. 13, Année 1994–1995, Sémin. Théor. Spectr. Géom., vol. 13, Univ. Grenoble I, Saint-Martin-d’Hères, 1995, pp. 97–122, DOI 10.5802/tsg.155. MR1715960
- [34] G. Link, *Hausdorff dimension of limit sets of discrete subgroups of higher rank Lie groups*, Geom. Funct. Anal. **14** (2004), no. 2, 400–432, DOI 10.1007/s00039-004-0462-y. MR2062761
- [35] Gabriele Link, *Ergodicity of generalised Patterson-Sullivan measures in higher rank symmetric spaces*, Math. Z. **254** (2006), no. 3, 611–625, DOI 10.1007/s00209-006-0962-6. MR2244369
- [36] Gabriele Link, *Asymptotic geometry in products of Hadamard spaces with rank one isometries*, Geom. Topol. **14** (2010), no. 2, 1063–1094, DOI 10.2140/gt.2010.14.1063. MR2629900
- [37] Peter J. Nicholls, *The ergodic theory of discrete groups*, London Mathematical Society Lecture Note Series, vol. 143, Cambridge University Press, Cambridge, 1989, DOI 10.1017/CBO9780511600678. MR1041575
- [38] S. J. Patterson, *The limit set of a Fuchsian group*, Acta Math. **136** (1976), no. 3–4, 241–273, DOI 10.1007/BF02392046. MR450547
- [39] Frédéric Paulin, *On the critical exponent of a discrete group of hyperbolic isometries*, Differential Geom. Appl. **7** (1997), no. 3, 231–236, DOI 10.1016/S0926-2245(96)00051-4. MR1480536
- [40] Rafael Potrie and Andrés Sambarino, *Eigenvalues and entropy of a Hitchin representation*, Invent. Math. **209** (2017), no. 3, 885–925, DOI 10.1007/s00222-017-0721-9. MR3681396
- [41] Maria Beatrice Pozzetti, Andrés Sambarino, and Anna Wienhard, *Conformality for a robust class of non-conformal attractors*, J. Reine Angew. Math. **774** (2021), 1–51, DOI 10.1515/crelle-2020-0029. MR4250471
- [42] Jean-François Quint, *Divergence exponentielle des sous-groupes discrets en rang supérieur* (French, with French summary), Comment. Math. Helv. **77** (2002), no. 3, 563–608, DOI 10.1007/s00014-002-8352-0. MR1933790

- [43] J.-F. Quint, *Mesures de Patterson-Sullivan en rang supérieur* (French, with English and French summaries), *Geom. Funct. Anal.* **12** (2002), no. 4, 776–809, DOI 10.1007/s00039-002-8266-4. MR1935549
- [44] Thomas Roblin, *Ergodicité et équidistribution en courbure négative* (French, with English and French summaries), *Mém. Soc. Math. Fr. (N.S.)* **95** (2003), vi+96, DOI 10.24033/msmf.408. MR2057305
- [45] Andrés Sambarino, *Quantitative properties of convex representations*, *Comment. Math. Helv.* **89** (2014), no. 2, 443–488, DOI 10.4171/CMH/324. MR3229035
- [46] Andrés Sambarino, *The orbital counting problem for hyperconvex representations* (English, with English and French summaries), *Ann. Inst. Fourier (Grenoble)* **65** (2015), no. 4, 1755–1797. MR3449196
- [47] Dennis Sullivan, *The density at infinity of a discrete group of hyperbolic motions*, *Inst. Hautes Études Sci. Publ. Math.* **50** (1979), 171–202. MR556586
- [48] Dennis Sullivan, *Entropy, Hausdorff measures old and new, and limit sets of geometrically finite Kleinian groups*, *Acta Math.* **153** (1984), no. 3-4, 259–277, DOI 10.1007/BF02392379. MR766265
- [49] Pekka Tukia, *Convergence groups and Gromov’s metric hyperbolic spaces*, *New Zealand J. Math.* **23** (1994), no. 2, 157–187. MR1313451
- [50] Jussi Väisälä, *Gromov hyperbolic spaces*, *Expo. Math.* **23** (2005), no. 3, 187–231, DOI 10.1016/j.exmath.2005.01.010. MR2164775
- [51] Chengbo Yue, *The ergodic theory of discrete isometry groups on manifolds of variable negative curvature*, *Trans. Amer. Math. Soc.* **348** (1996), no. 12, 4965–5005, DOI 10.1090/S0002-9947-96-01614-5. MR1348871

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, 10 HILLHOUSE AVE, NEW HAVEN, CONNECTICUT 06511

Email address: subhadip.dey@yale.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, ONE SHIELDS AVE, DAVIS, CALIFORNIA 95616

Email address: kapovich@math.ucdavis.edu