UC Irvine

UC Irvine Previously Published Works

Title

Neutron scattering studies on lanthanum cuprates (invited) (abstract)

Permalink

<https://escholarship.org/uc/item/3m41p2rt>

Journal

Journal of Applied Physics, 69(8)

ISSN

0021-8979

Authors

Hayden, SM
Aeppli, G
Cheong, SW
et al.

Publication Date

1991-04-15

DOI

10.1063/1.348344

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Neutron scattering studies on lanthanum cuprates (invited) (abstract)

S. M. Hayden

Institut Laue-Langevin, 38042 Grenoble, France

G. Aeppli and S-W. Cheong

AT&T Bell Laboratories, Murray Hill, New Jersey 07974

H. A. Mook

Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831

Z. Fisk

Los Alamos National Laboratory, Los Alamos, New Mexico 87545

D. Rytz

Hughes Research Laboratory, Malibu, California 90265

The results of recent inelastic neutron scattering investigations of the magnetic excitations in La_2CuO_4 are reviewed. Experiments were performed using a wide range of incident neutron energies. Our study of the paramagnetic phase is consistent with a picture for the magnetic dynamics which shows conventional spin waves at higher energies and an overdamped response due to spin-wave interactions at lower energies. The evolution of the magnetic dynamics with doping in the $\text{La}_{2-x}(\text{Ba},\text{Sr})_x\text{CuO}_4$ system is discussed.

Hole dynamics in the copper oxide planes (invited) (abstract)

S. A. Trugman

Theoretical Division, B262, Los Alamos National Laboratory, Los Alamos, New Mexico 87545

The physics of a hole doped into the Mott insulating state of the CuO_2 planes is studied in a large variational space. The hole quasiparticle that forms is quite different from a free electron by virtue of the cloud of spin flips (with respect to the background antiferromagnet) that it carries with it. The momentum dependence of the quasiparticle energy is unusual: $\epsilon(\mathbf{k})$ is lowest at nonzero momentum, and has an anisotropic dispersion. At finite doping, the quasiparticles are assumed to fill the rigid band described by the single-hole $\epsilon(\mathbf{k})$, forming a weakly interacting Fermi gas. Normal state properties of the high-temperature superconducting oxides, such as $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$, are calculated in the relaxation time approximation. Good agreement with experiment is obtained for the magnetic susceptibility, Hall coefficient, thermopower, specific heat, and plasma frequency as a function of both doping x and temperature.¹

Work supported by the U.S. Department of Energy.

¹S. A. Trugman, Phys. Rev. Lett. **65**, 500 (1990).