

UC Davis

UC Davis Previously Published Works

Title

Rickettsia africae and Novel Rickettsial Strain in Amblyomma spp. Ticks, Nicaragua, 2013 - Volume 24, Number 2—February 2018 - Emerging Infectious Diseases journal - CDC

Permalink

<https://escholarship.org/uc/item/3m44r0fh>

Journal

Emerging Infectious Diseases, 24(2)

ISSN

1080-6040

Authors

Vogel, Helena
Foley, Janet
Fiorello, Christine V

Publication Date

2018-02-01

DOI

10.3201/eid2402.161901

Peer reviewed

Table. Cysticercosis incidence rates by sex, residence, and age group, Shandong Province, China, 1975–2014

Characteristic	No. (%) patients	Incidence rate, cases/ 1 million population (95% CI)
Sex		
M	1,288 (65.98)	29.1 (27.5–30.7)
F	664 (34.02)	15.5 (14.4–16.7)
Residence		
Rural	1,346 (68.95)	20.6 (19.5–21.7)
Urban	606 (31.05)	27.9 (25.7–30.1)
Age group, y		
<1–9	94 (4.82)	6.7 (5.4–8.1)
10–19	170 (8.71)	12.5 (10.6–14.3)
20–29	410 (21.00)	26.6 (24.0–29.1)
30–39	546 (27.97)	37.2 (34.1–40.3)
40–49	409 (20.95)	32.7 (29.5–35.9)
50–59	185 (9.48)	26.0 (22.2–29.7)
≥60	138 (7.07)	14.3 (11.9–16.7)

and regions than the study by Chen et al., in which the 10–29-year age group and middle regions of the province showed the highest incidence rates (6).

Our study has a few limitations. First, the long, asymptomatic latent period of cysticercosis affects diagnostic efficiency and age-specific incidence estimates. Second, our data were incomplete because of some missing information for cases we identified. Third, independent confirmation might affect incidence estimates from early in the study period. However, our multidagnostic approach substantially reduced misdiagnosis rates and increased the efficiency of diagnosing cysticercosis (9).

In summary, our analyses show that Shandong Province has been a cysticercosis-endemic area for many years. Improved surveillance and control are needed to address the elevated risk for cysticercosis in western regions of this province.

This study was supported by the National Natural Science Fund (no. 31502057) from the National Natural Science Foundation of China and the Innovation Project of Shandong Academy of Medical Sciences.

About the Author

Dr. Gongzhen Liu is an assistant researcher in the field of pathogen biology, Shandong Institute of Parasitic Diseases, Shandong Academy of Medical Sciences, World Health Organization Collaborating Centre on Vector-Borne Diseases and Food-Borne Parasitic Diseases, Jining, China. His current research interest is the role of invasive parasites and interactions with host cells.

References

- Garcia HH, Nash TE, Del Brutto OH. Clinical symptoms, diagnosis, and treatment of neurocysticercosis. *Lancet Neurol.* 2014; 13:1202–15. [http://dx.doi.org/10.1016/S1474-4422\(14\)70094-8](http://dx.doi.org/10.1016/S1474-4422(14)70094-8)
- World Health Organization. First World Health Organization report on neglected tropical diseases. Geneva: The Organization; 2009 [cited 2017 Oct 24]. http://www.who.int/neglected_diseases/2010report/en/

- Yang GJ, Liu L, Zhu HR, Griffiths SM, Tanner M, Bergquist R, et al. China's sustained drive to eliminate neglected tropical diseases. *Lancet Infect Dis.* 2014;14:881–92. [http://dx.doi.org/10.1016/S1473-3099\(14\)70727-3](http://dx.doi.org/10.1016/S1473-3099(14)70727-3)
- Sorvillo FJ, DeGiorgio C, Waterman SH. Deaths from cysticercosis, United States. *Emerg Infect Dis.* 2007;13:230–5. <http://dx.doi.org/10.3201/eid1302.060527>
- Zhen TM, Yang YJ, Li DJ, Ge LY, Liu X, Dai W, et al. Diagnosis of cysticercosis. Diagnostic criteria of the health industry in the People's Republic of China. 2012;WS381:1–14.
- Chen XX, Liu X, Li GP, Fu B, Zhao CL, et al. Seroepidemiological study on the infectious status of cysticercus in Shandong Province [in Chinese]. *China Trop Med.* 2003;3:441–2.
- Wu W, Qian X, Huang Y, Hong Q. A review of the control of clonorchiasis sinensis and Taenia solium taeniasis/cysticercosis in China. *Parasitol Res.* 2012;111:1879–84. <http://dx.doi.org/10.1007/s00436-012-3152-y>
- Ge LY, Zhang JJ, Wang CY. Analysis of 3,596 out-patients with taeniasis and cysticercosis [in Chinese]. *Chin J Parasitic Dis Contr.* 1993;6:23–6.
- Shi FM, Ge LY, Yang SH, Xu WH. Study on immunological diagnosis of cysticercosis [in Chinese]. *Chinese Journal of Parasitic Diseases.* 1989;2:65–6.

Address for correspondence: Zhenhua Yu or Xin Liu, Shandong Institute of Parasitic Diseases, Shandong Academy of Medical Sciences, World Health Organization Collaborating Centre on Vector-Borne Diseases and Food-Borne Parasitic Diseases, Jining 272033, China; email: sfyzyh123@163.com or liux3276@163.com

***Rickettsia africae* and Novel Rickettsial Strain in *Amblyomma* spp. Ticks, Nicaragua, 2013**

Helena Vogel, Janet Foley, Christine V. Fiorello

Author affiliation: University of California, Davis, California, USA

DOI: <https://doi.org/10.3201/eid2402.161901>

We report molecular detection of *Rickettsia africae* in *Amblyomma ovale* ticks from Nicaragua and a novel rickettsial strain in an *A. triste* tick. Of 146 ticks from dogs, 16.4% were *Rickettsia* PCR positive. The presence of *Rickettsia* spp. in human-biting ticks in Nicaragua may pose a public health concern.

Obligately intracellular *Rickettsia* spp., typically transmitted by ticks, cause a multitude of mild to severe rickettsial diseases in humans and other animals. Novel *Rickettsia* species have been identified through molecular

Table. Number of dogs and ticks sampled, tick species, and prevalence of rickettsiae in ticks in 3 indigenous communities, northern Nicaragua

Category	Amak	Raiti	Arang Dak	Total
No. dogs sampled	11	10	19	40
No. ticks collected and tested	25	55	66	146
<i>A. ovale</i> (PCR-positive)	25 (4)	45 (10)	57 (4)	127 (18)
<i>A. sculptum</i> (PCR-positive)	0	4 (3)	8 (2)	12 (5)
<i>A. triste</i> (PCR-positive)	0	6 (1)	1 (0)	7 (1)
Prevalence of rickettsiae in ticks, % (95% CI)	16.0 (5.25–36.9)	25.5 (15.1–39.3)	9.09 (3.75–19.4)	16.4 (11.0–23.7)

techniques (1). Rickettsiae in Central America have primarily been reported in ticks, dogs, and humans, with limited data on tick species and rickettsial prevalence in Nicaragua (1). In an earlier study, 87% of 77 dogs in the Bosawás Biosphere Reserve were seropositive for rickettsiae (2); the ticks in that study were collected from 40 of those dogs.

The Bosawás Reserve in remote northern Nicaragua, part of the second largest tropical rainforest in the Western Hemisphere, is inhabited by 2 rapidly growing populations of indigenous people: the Miskito and the Mayangna. These subsistence-based communities use dogs for hunting in the reserve. Increasing connectivity with outside areas, population growth, and interference of dogs with wildlife pose an increased risk for the emergence of zoonotic rickettsioses. We planned to expand information on zoonotic *Rickettsia* spp. in Nicaragua by surveying ticks from hunting dogs for diversity, number, and presence of rickettsiae.

We collected ticks in 2013 from villages at similar latitude and longitude measured by using global positioning system (GPS): Arang Dak (14.51583, –84.99944), Amak (14.06542, –85.142233), and Raiti (14.59464, –85.02772) (Table). Arang Dak is the smallest of the 3 villages and closest to the densest part of the rainforest; Raiti is the largest and most developed village of the 3 and is situated on a heavily traveled route through the reserve. We obtained owner consent before physical examination and sampling of ticks from dogs and stored ticks in 70% ethanol. In the laboratory, we identified ticks for sex, life stage, and species by using a key (3) and screened tick DNA for *Rickettsia* spp. by real-time PCR (4). *Rickettsia*-positive samples were further tested by conventional PCR targeting the outer membrane protein A gene (*ompA*) (5). We also amplified the *rpmB* and *17kDa* genes of the rickettsia in the *Amblyomma triste* ticks we recovered (4). We sequenced each amplicon by using the forward primer at University of California Davis Sequencing (Davis, CA, USA) and compared sequences to those in the GenBank database by using the BLAST algorithm (<https://blast.ncbi.nlm.nih.gov>).

Of 146 ticks from 40 dogs, 126 (86%) were *A. ovale*, 12 were *A. sculptum*, and 7 were *A. triste*. We detected rickettsial DNA in 24 (16.4%, 95% CI 11.0%–23.7%) of the 146 ticks: 18 *A. ovale*, 5 *A. sculptum*, and 1 *A. triste*. We deposited rickettsial sequences from these ticks into GenBank (accession no. KX530472, KX576685, and KX576686).

By location, the PCR prevalence was 25.5% (95% CI 15.1%–39.3%) in Raiti, 16.0% (95% CI 5.25%–36.9%) in Amak, and 9.09% (95% CI 3.75%–19.4%) in Arang Dak. These differences were statistically significant ($p = 0.05$ by Fisher exact test). The finding of highest prevalence in the most populated community is consistent with peridomestic animals maintaining the infection, and the rainforest and remote wildlife not being significant sources.

For the 576-bp *ompA* sequence, all from *A. ovale* ticks were identical and were 99.6% homologous with sequences from GenBank identified as *R. africae*. *R. africae* has not been reported in *A. ovale* ticks or in North, Central, or South America. *R. africae* causes a mild rickettsiosis known as African tick-bite fever and was first described in a patient in the Western Hemisphere in 1998 (1). *R. africae* has been detected in *A. variegatum* ticks by using PCR and in humans in Guadeloupe by using serology (6) and more recently in *A. loculosum* ticks from New Caledonia (7). In Brazil, adult *A. ovale* ticks bite humans most frequently and are present from the borders of Mexico to those of Argentina (8). *A. ovale* is a common human-biting tick in Central and South America and poses a public health concern.

Sequences of *ompA* in 2 of 5 PCR-positive *A. sculptum* matched 99.6% to *Candidatus R. amblyommii* in GenBank (*ompA* of the other samples did not amplify, likely because they were relatively weak on real-time PCR). *Candidatus R. amblyommii* is common among *Amblyomma* spp. ticks in the New World and was reported in *A. sculptum* ticks in Brazil (9). *Candidatus R. amblyommii* has unknown pathogenicity but has been implicated in rickettsiosis cases in humans (9).

The *ompA* amplicon from *A. triste* ticks matched *Rickettsia* sp. ARAGAOI; sequencing of the *rpmB* and *17kDa* genes was unsuccessful. This rickettsial species was originally described in marsupials in Brazil (10). Further monitoring of tick vectors in this remote area is needed to characterize local risk and detect possibly emerging vector-borne disease.

Acknowledgments

The authors thank members of J.F.'s laboratory for their technical assistance and feedback; and L. Schwartz, J. Liu, J. Koster, F. Diaz-Santos, and U. Coleman for assistance with field logistics. K. Thomas produced the map.

This project was funded by the One Health Institute and the Wildlife Health Center of University of California, Davis.

About the Author

Miss Vogel is a veterinary student at University of California Davis School of Veterinary Medicine and works as a laboratory intern in J.F.'s laboratory. Her research interest is the ecology of emerging vectorborne diseases.

References

1. Parola P, Paddock CD, Socolovschi C, Labruna MB, Mediannikov O, Kernif T, et al. Update on tick-borne rickettsioses around the world: a geographic approach. *Clin Microbiol Rev.* 2013;26:657–702. <http://dx.doi.org/10.1128/CMR.00032-13>
2. Fiorello CV, Straub MH, Schwartz LM, Liu J, Campbell A, Kownacki AK, et al. Multiple-host pathogens in domestic hunting dogs in Nicaragua's Bosawas Biosphere Reserve. *Acta Trop.* 2017;167:183–90. <http://dx.doi.org/10.1016/j.actatropica.2016.12.020>
3. Voltzit OV. A review of neotropical *Amblyomma* species (Acari: Ixodidae). *Acarina.* 2007;15:3–134 [cited 2016 Nov 23]. <http://biotecavirtual.minam.gob.pe/biam/bitstream/handle/minam/890/BIV00727.pdf?sequence=1&isAllowed=y>
4. Stephenson N, Blaney A, Clifford D, Gabriel M, Wengert G, Foley P, et al. Diversity of rickettsiae in a rural community in northern California. *Ticks Tick Borne Dis.* 2017;8:526–31. <http://dx.doi.org/10.1016/j.ttbdis.2017.02.014>
5. Roux V, Fournier P-E, Raoult D. Differentiation of spotted fever group rickettsiae by sequencing and analysis of restriction fragment length polymorphism of PCR-amplified DNA of the gene encoding the protein rOmpA. *J Clin Microbiol.* 1996;34:2058–65.
6. Parola P, Vestris G, Martinez D, Brochier B, Roux V, Raoult D. Tick-borne rickettsiosis in Guadeloupe, the French West Indies: isolation of *Rickettsia africae* from *Amblyomma variegatum* ticks and serosurvey in humans, cattle, and goats. *Am J Trop Med Hyg.* 1999;60:888–93. <http://dx.doi.org/10.4269/ajtmh.1999.60.888>
7. Eldin C, Mediannikov O, Davoust B, Cabre O, Barré N, Raoult D, et al. Emergence of *Rickettsia africae*, Oceania. *Emerg Infect Dis.* 2011;17:100–2. <http://dx.doi.org/10.3201/eid1701.101081>
8. Labruna MB, Camargo LMA, Terrasini FA, Ferreira F, Schumaker TT, Camargo EP. Ticks (Acari: Ixodidae) from the state of Rondônia, western Amazon, Brazil. *Syst Appl Acarol.* 2005;10:17–32. <http://dx.doi.org/10.11158/saa.10.1.4>
9. Nunes EC, Vizzoni VF, Navarro DL, Iani FC, Durães LS, Daemon E, et al. *Rickettsia amblyommii* infecting *Amblyomma sculptum* in endemic spotted fever area from southeastern Brazil. *Mem Inst Oswaldo Cruz.* 2015;110:1058–61. <http://dx.doi.org/10.1590/0074-02760150266>
10. Blanco CM, Teixeira BR, da Silva AG, de Oliveira RC, Strecht L, Ogrzewalska M, et al. Microorganisms in ticks (Acari: Ixodidae) collected on marsupials and rodents from Santa Catarina, Paraná and Mato Grosso do Sul states, Brazil. *Ticks Tick Borne Dis.* 2017;8:90–8. <http://dx.doi.org/10.1016/j.ttbdis.2016.10.003>

Address for correspondence: Janet Foley, University of California, Department of Medicine and Epidemiology, 1320 Tupper Hall, Davis, CA 95616, USA; email: jefoley@ucdavis.edu

Amebaborne *Attilina massiliensis* Keratitis, France

Alexandre Battaini, Bernard La Scola, Gaëlle Ho Wang Yin, Louis Hoffart, Michel Drancourt

Author affiliations: Aix Marseille Université, Marseille, France (A. Battaini, B. La Scola, M. Drancourt); Aix-Marseille University–APHM, Hôpital de la Timone, Marseille (G.H.W. Yin, L. Hoffart)

DOI: <https://doi.org/10.3201/eid2402.170541>

We report a case of *Acanthamoeba castellanii* keratitis in a person who wore contact lenses. The amoebae hosted an amoeba-resistant bacterial symbiont, provisionally named “*Attilina massiliensis*,” a yet undescribed α -Proteobacterium.

Amebal keratitis is an aggressive ocular infection that can lead to blindness (1). It is usually associated with wearing soft contact lenses; Dart et al. documented that in countries with a high prevalence of contact lens wear, 85%–88% of *Acanthamoeba* keratitis cases occurred in contact lens users (1). These amoebae host amoeba-resistant bacteria, and increase their pathogenicity to the host (2). Amoeba hosting intra-amebal microorganisms have been rarely documented in cases originating in contaminated contact lenses (3) and never in mixed keratitis. We report a case of mixed amoeba–amebal-resistant bacterial keratitis.

A 17-year-old woman who wore contact lenses consulted the ophthalmology department of the clinic associated with Hôpital de la Timone, Marseille, France, in July 2016, after experiencing 1 month of keratoconjunctivitis symptoms related to an undocumented clinical diagnosis of herpes virus keratitis of the left eye. The patient had been prescribed a 1-week treatment with valacyclovir (3×/d) and a corneal dressing. Examination of the left eye showed 4/10 visual acuity; the right eye was normal. Slit-lamp examination showed a central radial keratoneuritis, central corneal edema, central diffuse infiltrate, and a punctate superficial keratitis with no predescemetic precipitates and no satellite lesions (Figure). The patient was admitted to the hospital and was administered hourly topical treatments of polyhexamethylene biguanide eye drops, hexamidine, and 1% atropine. The patient, whose diagnosis was early-stage *Acanthamoeba* keratitis infection, was discharged after 5 days of treatment; a corneal swab sample at discharge was negative for herpes virus, varicella zoster virus, adenovirus, enterovirus, cytomegalovirus, and *Chlamydia trachomatis*. Follow-up 7 days later yielded reduced symptoms. We followed up on the patient biweekly and slowly tapered drugs over 4 months; the previously negative