UC San Diego

UC San Diego Previously Published Works

Title

Cost of Prescription Drug-Related Morbidity and Mortality.

Permalink

<https://escholarship.org/uc/item/3n76n4z6>

Journal

The Annals of pharmacotherapy, 52(9)

ISSN

1060-0280

Authors

Watanabe, Jonathan H

McInnis, Terry

Hirsch, Jan D

Publication Date

2018-09-01

DOI

10.1177/1060028018765159

Peer reviewed

Cost of Prescription Drug–Related Morbidity and Mortality

Annals of Pharmacotherapy
2018, Vol. 52(9) 829–837
© The Author(s) 2018
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1060028018765159
journals.sagepub.com/home/aop

Jonathan H. Watanabe, PharmD, MS, PhD¹ ,
Terry McInnis, MD, MPH², and Jan D. Hirsch, PhD¹

Abstract

Background: Public attention and recent US Congressional activity has intensified focus on escalating medication prices. However, the actual cost of medication use extends beyond the up-front cost of purchasing medicines. It also encompasses the additional medical costs of morbidity and mortality resulting from nonoptimized medication regimens, including medication nonadherence. **Objectives:** Applying the most current nationally representative data sources, our goal was to estimate the cost of prescription drug–related morbidity and mortality in the United States. **Methods:** Total costs of nonoptimized prescription drug use and average pathway costs for a patient who experienced a treatment failure (TF), a new medical problem (NMP), or a TF and NMP were modeled in Microsoft Excel (Microsoft Corporation, Redmond, WA) and TreeAge Pro Healthcare, v2014 (TreeAge Software, Inc, Williamstown, MA), respectively. **Results:** The estimated annual cost of prescription drug–related morbidity and mortality resulting from nonoptimized medication therapy was \$528.4 billion in 2016 US dollars, with a plausible range of \$495.3 billion to \$672.7 billion. The average cost of an individual experiencing TF, NMP, or TF and NMP after initial prescription use were \$2481 (range: \$2233, \$2742), \$2610 (range: \$2374, \$2848) and \$2572 (range: \$2408, \$2751), respectively. **Conclusions:** The estimated annual cost of drug-related morbidity and mortality resulting from nonoptimized medication therapy was \$528.4 billion, equivalent to 16% of total US health care expenditures in 2016. We propose expansion of comprehensive medication management programs by clinical pharmacists in collaborative practices with physicians and other prescribers as an effective and scalable approach to mitigate these avoidable costs and improve patient outcomes.

Keywords

medication therapy management, health care policy, pharmacoeconomics, pharmacist/physician issues, drug-related problems, administration, adverse drug reactions, clinical pharmacy, cost, pharmaceutical care

Introduction

Prevention and management of chronic disease is fundamental to improve population-level health outcomes and blunt the upward trend in health care spending in the United States. Close to three-quarters of physician office and hospital outpatient clinic visits involve drug therapy.¹ The cost of drug therapy in the United States has been escalating because of increased utilization and drug prices.² However, the cost associated with drug use reaches beyond the purchase of prescribed medications to encompass additional medical costs of morbidity and mortality resulting from medication regimens that are not optimized to effectively treat the indication resulting in a treatment failure (TF), where the resolvable medical problem is not adequately treated, a new medical problem (NMP), where a newly prescribed medication causes or contributes to an incident clinical symptom or syndrome, or both a TF and NMP. This cost has most recently

been estimated as \$290 billion equating to 13% of total annual US medical costs in 2008.³ Although widely misdescribed in the published literature and policy documents as the cost associated with “patient nonadherence to medications,” this estimate and the preceding estimates (\$76.6 billion in 1995 and \$177.4 billion in 2000) actually reflect medical resource utilization caused by TFs and NMPs that arising from nonoptimized medication use. Nonadherence to the indicated medication regimen is just one of multiple

¹University of California San Diego, La Jolla, CA, USA

²Value Based Care, Laboratory Corporation of America, Burlington, NC, USA

Corresponding Author:

Jonathan H. Watanabe, Skaggs School of Pharmacy and Pharmaceutical Sciences, Division of Clinical Pharmacy, University of California San Diego, 9500 Gilman Drive MC 0764, La Jolla, CA 92093-0764, USA.
Email: jhwatanabe@ucsd.edu

potential causal factors leading to a TF, resulting in downstream health services use.^{4,5}

The need for a more current estimate of the cost of nonoptimized medication use is clear given the significant movements in our health care system since the 2008 estimate.³ Since then, health care reform has expanded medication use in the United States. For example, patients who gained coverage via Medicaid expansion under the Affordable Care Act (ACA) filled 79% more prescriptions.⁶

Escalating prices of medications in the United States have garnered public outcry. However, the limited use of systematic and coordinated approaches that reduce costs attributable to nonoptimized medication regimens demands similar attention. A more current estimate of the cost related to nonoptimized medication utilization can serve as a linchpin in policy, payment, and health care delivery system reform. These are important elements as we contend not only with the cost of medications, but also with the impact of prescription drugs on overall medical costs, patient outcomes, and the valuation of interventions to optimize medication use.

The objective of this article is to update the estimated cost of drug-related morbidity and mortality resulting from nonoptimized medication therapy from a third-party payer perspective.

Methods

Data and Sources

This cost-of-illness (COI) analysis used the original decision analytic model and probabilities used by all prior published estimates (Figure 1), cost data sources most closely matching those used in the most recent update, and the same source for estimated number of physician office visits in the United States.^{3-5,7} All costs were adjusted for medical inflation to 2016.⁸ Our total cost estimate reflects the most current monetary value of medical resources and the frequency of use based on the most current US physician office visit estimates. Costs captured in this analysis were for resource use spending on treatment of a medical problem (ie, direct medical costs). For this COI study, we followed the Consolidated Health Economic Evaluation Reporting Standards (CHEERS) guidelines for reporting health economic evaluations in construction of this manuscript. CHEERS recommends description of the incremental cost per outcome benefit. A priori, our objective was completion of a COI study that quantified the cost of the medical problem. Hence, incremental cost per benefit was not evaluated.⁹

The original modeling included path probabilities based on expert opinion.⁴ The two published updates of the cost analysis retained these probabilities, although Ernst and

Grizzle⁵ did search the literature for evidence to revise the original proportions and found none.^{3,5} Similarly, our literature search did not uncover published data to support modifying the original probabilities in the prior models. Applying PubMed database search terms *drug therapy problem*, *medication related problem*, *drug-related problem*, *new medical problem*, and *treatment failure* with date range January 1, 2000, to November 17, 2017, produced 90 articles which, on review, did not contain updated probabilities for progression to NMPs and TFs or downstream probabilities for health services utilization on review by one of the study authors (JDH). Removing date range criteria yielded the prior published study by Ernst and Grizzle.

Costs. Base case analyses used the weighted average of the mean physician fees for new and established patients of \$222 per visit (2014; \$236, 2016 value).¹⁰ Average prescription cost of \$71 (\$74, 2016 value), calculated as total prescription sales divided by total number of prescriptions sold in 2015 (\$286 797 651 607/4 065 175 064).^{11,12} Average cost of an ED visit was \$1233 (\$1571, 2016 value) based on the 10 most common types of ED visits.¹³ Average cost of a hospital admission was \$13 450 (\$14 324, 2016 value).¹⁴ The average cost of a long-term care (LTC) admission—\$62 178 in 2016—was calculated using data from the 1999 and 2004 National Nursing Home Survey (NNHS) from the Centers for Disease Control and Prevention (CDC) and longtermcare.gov.¹⁵⁻¹⁷ The NNHS is a validated, nationally representative survey that provides estimates of nursing homes, current residents, and discharges in the United States that include explicit capture of duration of stay. Using the NNHS from the CDC resulted in a larger length of stay (LOS) than the reference used in prior estimates.^{4,5} The most recent 2004 NNHS included the average time since admission (ATSA) not average LOS (ALOS) as in previous versions.¹⁶ Therefore, 2004 ALOS was estimated to be 254 days via multiplication of the 2004 ATSA (835 days) by the ratio of ALOS (271.5 days) to ATSA in the 1999 NNHS data ($892.4 / 271.5 = 3.29$). Multiplying the ALOS by the 2010 average semiprivate daily cost (\$205/day) resulted in a value of \$52 078 (\$62 178 in 2016 dollars).¹⁷ Consistent with prior published estimates, death attributable to a TF or NMP was assumed to be preceded by a hospitalization. Hence, although base case cost of a death event was zero dollars, the pathway cost resulting in death resulting from TF or NMP included the cost of a hospitalization, and death events contributed to the total hospitalization count.^{4,5} The number of physician office visits used as the basis for the population scale-up was 928 630 000.¹⁸

Data Analyses

Consistent with prior published studies, total costs of nonoptimized prescription drug use incorporating resource costs

Figure 1. Drug-related morbidity and mortality decision analytic model.
Abbreviation: NMB = net monetary benefit.

and probabilities were modeled in Microsoft Excel (Microsoft Corporation, Redmond, WA).^{4,5} The average pathway costs from initial provider visit for a patient who experienced a TF, NMP, and TF and NMP were calculated within a decision analytic model (Figure 1) using TreeAge Pro Healthcare, v2014

(TreeAge Software, Inc, Williamstown, MA). Probabilistic sensitivity analyses using Monte Carlo simulation were completed to generate a plausible range around the mean estimates. If available, published standard errors were used in the sensitivity analyses. Otherwise, an error margin of 10% of

the mean input estimate was applied for these analyses consistent with the 10% range used in the Ernst and Grizzle analysis.⁵

Population Scale-up to Estimated Cost of Nonoptimized Drug Utilization. The total cost of morbidity and mortality resulting from nonoptimized drug utilization was calculated by multiplying the number of health service events (estimated via proportions in the decision analytic model) by estimated event costs. Sensitivity analyses were completed to investigate the impact of using other nationally representative sources of data for the resource costs. Specifically, cost of ED visits, hospitalizations, LTC, and the number of physician office visits were modified in these analyses. ED visit cost was decreased to the reported average cost across all types of ED visits of \$1116 in 2016 dollars.¹⁹ Hospitalization cost was increased to \$24 600 (net patient revenue [\$813 437 515 663] divided by total number of admissions [33 066 720]) adjusted to \$26 199 in 2016 dollars and decreased to \$10 889 adjusted to \$11 597 in 2016 dollars.^{20,21} The average cost of a LTC admission was decreased to \$57 158 using average daily cost of \$225.²² The number of physician office visits was increased to 1.067 billion to reflect a recent estimate, which may better reflect ACA impact.²³ Total pathway costs beginning with initial visit for each of the specific scenarios delineated in the decision analytic model were tabulated (Table 1).

Results

The estimated annual cost of prescription drug–related morbidity and mortality resulting from nonoptimized medication therapy was \$528.4 billion in 2016 US dollars (Table 2). Sensitivity analyses using alternate cost inputs revealed total costs ranging from a minimum of \$495.3 billion in the scenario that used a lower estimate of hospitalization cost (CDC Healthcare Cost and Utilization Project cost data) to a maximum of \$672.7 billion using the highest estimate of hospitalization cost (American Hospital Association cost data).^{20,21} Using the Department of Health and Human Services Agency for Healthcare Research and Quality estimate for ED visit cost produced a total cost estimate of \$517.7 billion.¹⁹ Applying a larger estimate of physician office visits produced a total cost of \$607.2 billion.²³ Using a lower LTC cost estimate derived from the Genworth Cost of Care Survey, the total cost estimate was \$505.5 billion.²² Hence, the estimated annual cost of prescription drug–related morbidity and mortality yielded a plausible range of \$495.3 to \$672.7 billion.

Based on the output from the decision-analytic model (Figure 1), the average cost of a physician encounter that incorporates costs of patients receiving no drug therapy, those receiving successful drug therapy, and downstream medical costs of nonoptimized drug utilization was

estimated to be \$850 per encounter (plausible range based on sensitivity analyses: \$792, \$908). The average cost of an individual experiencing TF, NMP, or TF and NMP after initial prescription use were \$2481 (plausible range: \$2233, \$2742), \$2610 (plausible range: \$2374, \$2848), and \$2572 (plausible range: \$2408, \$2751).

Discussion

This analysis has updated the annual cost of drug-related morbidity and mortality resulting from nonoptimized medication therapy from a third-party payer perspective and estimated it to be \$528.4 billion. Nonoptimized medication therapy was also estimated to result in 275 689 deaths per year. The cost estimate represents the additional medical resources utilized (eg, additional medications, ED visits, hospitalizations, LTC stays, provider visits) to resolve problems attributable to initial indication and utilization of prescription drugs, including nonadherence to indicated therapy. The potentially avoidable \$528.4 billion estimate is roughly 16% of the total US health care expenditure (\$3.2 trillion in 2015).²⁴ This proportion has doubled over the past 20 years since the original 1995 estimate that represented 8% of total US health care expenditures (2000 and 2008 estimates represented 13% of total health care spending).³⁻⁵

Medical price inflation alone would account for an increase from the 2008 estimate of \$290 billion to \$370 billion in 2016, leaving an excess of \$158 billion attributable to increased numbers of patients with TF, NMP, or a combination of TF and NMP caused by nonoptimized prescription drug utilization. We estimated that a patient experiencing TF, a NMP caused by drug therapy, or a combination of both costs approximately \$2500 considering additional drug therapy and other medical costs needed for resolution.

Our estimate of \$528 billion represents only direct medical costs, which are relevant for a third-party payer perspective. The estimate does not include direct nonmedical costs (eg, transportation or caregiving costs) nor indirect costs related to lost productivity or disability. The estimate is greater than the direct medical costs reported for many prevalent chronic diseases such as heart disease and stroke (\$230 billion), diagnosed diabetes (\$197 billion), cancer (\$187 billion), obesity (\$147 billion), and arthritis and related conditions (\$126 billion)—all reported here in 2016 dollars.²⁵⁻²⁹

Many factors will continue to drive costs associated with nonoptimized medication therapy, not least of which is the aging US population and the concurrent surge in the number of Medicare enrollees to 81.8 million by 2030.³⁰ In addition, the pipeline of new medications has been skewed to expensive specialty medications with more than 700 in development.³¹ In 2016, specialty drugs accounted for more than one-third of total drug expenditures.² The United States will also be challenged by increases in drug utilization rates

Table 1. Drug-Related Morbidity and Mortality Decision Analytic Model Pathway Costs in 2016 US Dollars.^a

Pathway Outcome	Initial Treatment		Cost of Negative Therapeutic Outcome					Total Pathway Cost
	Physician Visit ^b	Drug ^c	Physician Visit	Drug	ED Visit ^d	Hospital Admission ^e	LTC Admission ^f	
Optimal outcome	\$236	\$74						310
Treatment failure (TF) ^f								
No treatment	\$236	\$67 ^g						\$303
Physician visit	\$236	\$67	\$236					\$539
Additional prescription	\$236	\$67	\$236	\$74				\$613
Emergency department (ED) visit	\$236	\$67			\$1571			\$1874
Hospital admission	\$236	\$67				\$14 324		\$14 627
Long-term care (LTC) admission	\$236	\$67					\$62 178	\$62 481
Death	\$236	\$67				\$14 324		\$14 627
New medical problem (NMP)								
No treatment	\$236	\$74						\$310
Physician visit	\$236	\$74	\$236					\$546
Additional prescription	\$236	\$74	\$236	\$74				\$620
ED visit	\$236	\$74			\$1571			\$1881
Hospital admission	\$236	\$74				\$14 324		\$14 634
LTC admission	\$236	\$74					\$62 178	\$62 488
Death	\$236	\$74				\$14 324		\$14 634
TF and NMP								
No treatment	\$236	\$67						\$303
Physician visit	\$236	\$67	\$236					539
Additional prescription	\$236	\$67	\$236	\$74				\$613
ED visit	\$236	\$67			\$1571			\$1874
Hospital admission	\$236	\$67				\$14 324		\$14 627
LTC admission	\$236	\$67					\$62 178	\$62 481
Death	\$236	\$67				\$14 324		\$14 627
No drug therapy	\$236							\$236

^aCosts rounded to nearest dollar.^bMedical Expenditure Panel Survey (MEPS).¹⁰^cSource: Kaiser Family Foundation.^{11,12}^dSource: Caldwell et al.¹³^eSource: Medical Expenditure Panel Survey.¹⁴^fSource: Department of Health and Human Services.^{15,16,17}^gEstimated drug cost reduced by 10% to reflect rate of new prescriptions never filled.

and drug prices for most major disease categories (eg, inflammatory conditions, diabetes, oncology).² Retail prescription drugs along with home health are the 2 sectors of personal health care spending with the highest projected average spending growth through 2025.³² Payers will also likely grapple with increasing prices of medical resources (eg, ED visits, hospitalizations) because the rate of medical price growth is expected to accelerate in the 2018 to 2025 period.³³ Considering the projected increase in up-front costs resulting from additional medications needed for a graying US population and escalation in prescription costs, optimizing medication use is unquestionable.

Comprehensive Medication Management

Reducing these utilization and cost trends will require a systematic approach to medication management to mitigate the proportion of patients experiencing TF, NMP, or TF and NMP resulting from nonoptimized medication therapy. Clinical pharmacists, in collaborative practice with physicians and other prescribers such as nurse practitioners and physician assistants, will increasingly be relied on to assume responsibility for real-time drug therapy decisions in an expanding diversity of care settings by delivering comprehensive medication management (CMM) services. This

Table 2. Summary Cost of Drug-Related Morbidity and Mortality in 2016 Dollars.

	Number of Events (millions)	Cost Per Event	Total Cost (billions) ^a
Total physician visits	160.4	\$236	\$37.8
Total hospital admissions	12.2	\$14 324	\$174.0
Total emergency department visits	23.7	\$1571	\$37.2
Total long-term care admissions	4.4	\$62 178	\$271.6
Total additional prescriptions	105.9	\$74	\$7.8
Total deaths	0.276	—	—
Total	—	—	\$528.4

^aMay not equal events times cost for each resource in this table because of rounding.

transition has already begun in health systems such as Kaiser-Permanente and the Veterans Health Administration where pharmacists provide face-to-face patient care in collaborative-practice clinics.³³⁻³⁵ The approach will directly resolve costly drug therapy problems while addressing a shortage of primary care physicians.³⁶ The importance of reducing NMPs has been illuminated by clinician investigators who have focused on the clinical phenomenon referred to as the “prescribing cascade,” in which an adverse drug reaction (ADR) is misconstrued as a new syndrome, resulting in prescribing of a new medication. The proposed approach to arresting the prescribed cascade is a critical evaluation of the need for the new potential pharmacological agent in context of a comprehensive review of the complete medication profile. This evaluation should include explicit consideration of the current medication regimen as possibly causal on the symptom being addressed. Nonmedication treatment should be considered as well as alternative agents with lower risk of ADRs. If medication treatment is merited, the lowest effective medication dose should be prescribed.³⁷ CMM incorporates consideration of the prescribing cascade as well as the principles to prevent it from occurring.

CMM would also enrich other specialty areas such as behavioral health where medication management is playing a growing role.³⁸ Seminal work published in 2003 from the Asheville Project demonstrated a reduction in days lost to sick time, decreased medical costs, and improved achievement of treatment goals for patients with chronic disease receiving CMM.³⁹ Published evidence of the effectiveness of CMM has continued to build along with increasing calls for pharmacists in the health care team.^{40,41} A 2011 report to the US Surgeon General summarized evidence available for return on investment of medication management programs, reaching a maximum of 12:1 and averaging between 3:1 to 5:1.⁴² One example of a specific CMM program is the team-based medication use system in Minnesota that has lowered per member per month health care costs and increased the percentage of patients achieving treatment goals.⁴³ The benefits of CMM extend to improvements in quality of life,

access to other health care providers, and continuity of care as reported in a recent review of CMM in California.⁴⁴

CMM focused on achieving optimal clinical and patient-centered goals of therapy is becoming recognized as a cornerstone of delivery and payment reform efforts.⁴⁵ Hepler and Strand⁴⁶ proposed eight categories of drug therapy problems (untreated indication, improper drug selection, subtherapeutic dosage, failure to receive drugs, overdosage, ADRs, drug interactions, and drug use without indication). The CMM process is directly related to recognition and resolution of these drug therapy problems (DTPs) that can prevent use of unnecessary medical services (Figure 2). A general framework has been developed in prior studies in which DTPs give rise to TF and NMPs. Johnson and Bootman,⁴ building from the Hepler and Strand work, proposed a causal path where DTPs give rise to TF and NMPs, but did not specify the pathway or the probabilities in this process. Rather, the prior published COI estimates for non-optimized medication use focused on the downstream probability of TFs and NMPs when medication use was indicated.⁴

Drug therapy problems are drivers of medical resource use and are frequently cited as the largest cause of adverse events post-hospital discharge, and it has been estimated that 13% of adverse drug events after discharge from the hospital result in an ED visit or rehospitalization.^{47,48} Johnson and Bootman⁴ categorized this specific scenario as the ADR DTP producing an NMP.

Whereas inappropriate consumption of medications is one important element of DTPs, the existence of untreated indications, subtherapeutic dosing, or drug interactions of medications are common scenarios leading to NMPs and TFs. A large study of commercial insurance patients revealed that the most common issues were related to the need for additional medications (34%) and doses that were too low (20%).⁴⁹ Considering underuse of appropriate, effective, and safe medications as the omission side of DTPs in ambulatory care represents a large opportunity to improve outcomes and reduce the COI.⁵⁰ The majority of accountable care organization and value-based quality

1. Identify patients that have not achieved clinical goals of therapy.
2. Understand the patient's personal medication experience/history and preferences/beliefs.
3. Identify actual use patterns of all medications including OTCs, bioactive supplements, and prescribed medications.
4. Assess each medication (in the following order) for appropriateness, effectiveness, safety (including drug interactions), and adherence, focused on achievement of the clinical goals for each therapy.
5. Identify all drug therapy problems (the gap between current therapy and that needed to achieve optimal clinical outcomes).
6. Develop a care plan addressing recommended steps, including therapeutic changes needed to achieve optimal outcomes.
7. Patient agrees with and understands care plan, which is communicated to the prescriber/provider for his/her consent/support.
8. Document all steps and current clinical status versus goals of therapy.
9. Follow-up evaluations with the patient are critical to determine effects of changes, reassess actual outcomes, and recommend further therapeutic changes to achieve desired clinical goals/outcomes.
10. Comprehensive medication management is a reiterative process—care is coordinated with other team members and personalized (patient-unique) goals of therapy are understood by all team members.

Figure 2. Comprehensive medication management clinical approach.^a

^aPatient-Centered Primary Care Collaborative.⁵³

metrics hinge on optimized medication use to achieve population level goals of therapy.⁵¹

Although CMM represents the most rigorous and structured clinical process to optimize medication therapy, the clinical service referred to as medication therapeutic management (MTM) also entails review of the medication regimen with the goal of optimizing medication regimens. However, MTM can be conducted in the absence of the physical patient and the patient's complete medical record by simple review of the medication list. Whereas in-person, real-time MTM can be pursued, in general, MTM differs from CMM based on the absence of a few general features: CMM requires an evaluation of the patient's clinical status to ensure evidence-based concordance between the patient's syndromes and medication therapy. CMM involves specification of a treatment plan that incorporates follow-up evaluation to assess timely progression to prespecified treatment goals. CMM is designed to operate under a collaborative practice agreement with a physician.⁵²

As described, a systematic categorization scheme for the major types of DTPs related to indication, effectiveness, safety, and adherence is available and accepted.⁵³ Enhanced medication therapy management models are now being tested by the Centers for Medicare and Medicaid Services that utilize new Systematized Nomenclature of Medicine—Clinical Terms (SNOMED CT) codes to report on medication management activities and outcomes.⁵⁴ Lessons learned from these pilot programs will expand documentation efforts across the range of CMM models to facilitate outcomes tracking of pharmacist interventions. The coding system will also supply policy makers with objective data to supplant the current expert opinion–based estimates of TF, NMP, or TF and NMP occurrence. Code use will also furnish valuable data to determine resource utilization estimates after clinical interventions. These measurable outcomes will support production of real-world data models for future estimates of the cost of nonoptimized drug

therapy and provide pathway probabilities to model the chain from DTPs to TFs and NMPs.

There were limitations to this analysis. Medical resource costs used in the models were the most current available. However, not all were from the same year because of varying report dates. All were adjusted to 2016 US dollars. Because the most recent available data for number of physician office visits was from 2014, changes in visit volume caused by ACA deployment after 2014 were not fully accounted for in the base case model. Hence, this estimate is likely conservative. However, our sensitivity analysis did include an estimation that factored in an increase in visits. Model probabilities were based on expert opinion. These were maintained for consistency with prior published methods and the absence of replacement probability values in the published literature. Because of variations in available resource cost estimates, we performed sensitivity analyses that varied these inputs and reported total costs from each of these scenarios providing a range around the base case estimate.

Conclusions

The estimated annual cost of drug-related morbidity and mortality resulting from nonoptimized medication therapy from a third-party payer perspective was \$528.4 billion, equivalent to 16% of total US health care expenditures in 2016. This estimate includes neither direct nonmedical costs (eg, transportation or caregiving costs) nor indirect costs related to lost productivity. On average, a patient experiencing TF, a NMP resulting from nonoptimized drug therapy, or a combination of both costs roughly \$2500 considering additional drug therapy and other medical costs needed for resolution. To reduce these avoidable medical costs, improve patient care, and address the primary care provider shortage, we propose expansion of CMM programs as a cost-effective, scalable policy.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Dr. Watanabe is supported by the National Institutes of Health Loan Repayment Program through the National Institute on Aging. Study author employers and Supporting organizations did not have any role in the design, methods, analysis, or preparation of this manuscript.

ORCID iD

Jonathan H. Watanabe, <https://orcid.org/0000-0002-2543-5305>

References

- Centers for Disease Control and Prevention. Therapeutic drug use. <https://www.cdc.gov/nchs/fastats/drug-use-therapeutic.htm>. Accessed May 28, 2017.
- Express Scripts. Express scripts 2016 drug trend report. <http://lab.express-scripts.com/lab/drug-trend-report>. Accessed May 17, 2017.
- Network for Excellence in Health Innovation. Thinking outside the pillbox. *A system-wide approach to improving patient medication adherence for chronic disease*. <https://www.nehi.net/publications/17-thinking-outside-the-pillbox-a-system-wide-approach-to-improving-patient-medication-adherence-for-chronic-disease/view>. Accessed March 2, 2018.
- Johnson JA, Bootman JL. Drug-related morbidity and mortality: a cost-of-illness model. *Arch Intern Med*. 1995;155:1949-1956.
- Ernst FR, Grizzle AJ. Drug-related morbidity and mortality: updating the cost-of-illness model. *J Am Pharm Assoc*. 2001;41:192-199.
- Owens C. Study finds Obamacare increased prescription drug use, reduced spending. <https://morningconsult.com/2016/08/17/study-finds-obamacare-increased-prescription-drug-use-reduced-spending/>. Accessed March 2, 2018.
- Centers for Disease Control and Prevention. Table 76: Visits to physician offices, hospital outpatient departments, and hospital emergency departments, by age, sex, and race: United States, selected years 2000–2012. <https://www.cdc.gov/nchs/data/abus/abus15.pdf#076>. Accessed March 2, 2018.
- Halfhill TR. Tom's inflation calculator. http://www.halfhill.com/inflation_js.html. Accessed May 30, 2017.
- Husereau D, Drummond M, Petrou S, et al; CHEERS Task Force. Consolidated Health Economic Evaluation Reporting Standards (CHEERS) statement. *BMJ*. 2013;346:f1049. doi:10.1136/bmj.f1049
- Medical Expenditure Panel Survey (MEPS). Table 1: total utilization and mean expenses per visit by type of ambulatory health care service, 2014. https://meps.ahrq.gov/data_stats/summ_tables/hc/mean_expend/2014/table1.htm. Accessed May 30, 2017.
- Kaiser Family Foundation. Total retail sales for prescription drugs filled at pharmacies. <http://kff.org/other/state-indicator/total-sales-for-retail-rx-drugs/>. Accessed May 30, 2017.
- Kaiser Family Foundation. Total number of retail prescription drugs filled at pharmacies <http://kff.org/other/state-indicator/total-retail-rx-drugs/>. Accessed May 30, 2017.
- Caldwell N, Srebotnjak T, Wang T, Hsia R. "How much will I get charged for this?" Patient charges for top ten diagnoses in the emergency department. *PLoS One*. 2013;8:e55491. doi:10.1371/journal.pone.0055491
- Medical Expenditure Panel Survey. Table 2: Total utilization and mean expenses for inpatient stays by length of stay, 2014. https://meps.ahrq.gov/data_stats/summ_tables/hc/mean_expend/2014/table2.htm. Accessed May 30, 2017.
- Department of Health and Human Services. The National Nursing Home Survey: 1999 summary. *Page 44, table 25 and 39*. https://www.cdc.gov/nchs/data/series/sr_13/sr13_152.pdf. Accessed May 30, 2017.
- Department of Health and Human Services. The National Nursing Home Survey (NNHS): 2004 overview. *Table 7, page 19*. https://www.cdc.gov/nchs/data/series/sr_13/sr13_167.pdf. Accessed May 30, 2017.
- Department of Health and Human Services. Costs of care. <https://longtermcare.acl.gov/costs-how-to-pay/costs-of-care.html>. Accessed May 30, 2017.
- Centers for Disease Control and Prevention. Table 76: Visits to physician offices, hospital outpatient departments, and hospital emergency departments by age, sex, and race: United States, selected years 2000–2012. <https://www.cdc.gov/nchs/data/abus/abus15.pdf#076>. Accessed May 30, 2017.
- Medical Expenditure Panel Survey. Table 1: Total utilization and mean expenses per visit by type of ambulatory health care service, 2014. https://meps.ahrq.gov/data_stats/summ_tables/hc/mean_expend/2014/table1.htm. Accessed May 30, 2017.
- Health Forum LLC. *AHA Hospital Statistics*. Chicago, IL: Health Forum LLC; 2016:12-13, table 3.
- Healthcare Cost and Utilization Project (HCUP). Trends in inpatient stays. <https://www.hcup-us.ahrq.gov/faststats/NationalTrendsServlet?measure1=02&characteristic1=01&time1=10&measure2=&characteristic2=01&time2=10&expansionInfoState=hide&dataTablesState=hide&definitionsState=hide&exportState=hide>. Accessed May 30, 2017.
- Genworth. Compare long term care costs across the United States. <https://www.genworth.com/about-us/industry-expertise/cost-of-care.html>. Accessed May 30, 2017.
- Medical Expenditure Panel Survey (MEPS). Table 1: Total utilization and mean expenses per visit by type of ambulatory health care service, 2014. https://meps.ahrq.gov/data_stats/summ_tables/hc/mean_expend/2014/table1.htm. Accessed May 30, 2017.
- Centers for Medicare and Medicaid Services. NHE fact sheet. <https://www.cms.gov/research-statistics-data-and-systems/statistics-trends-and-reports/nationalhealthexpenddata/nhe-fact-sheet.html>. Accessed May 30, 2017.
- Go AS, Mozaffarian D, Roger VL, et al; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2014 update: a report from the American Heart Association. *Circulation*. 2014;129:e28-e292. <http://circ.ahajournals.org/content/early/2013/12/18/01.cir.0000441139.02102.80.full.pdf>. Accessed May 21, 2017. doi:10.1161/01.cir.0000442015.53336.12

26. American Diabetes Association. The cost of diabetes. <http://www.diabetes.org/advocacy/news-events/cost-of-diabetes.html>. Accessed May 21, 2017.
27. National Cancer Institute. Cancer prevalence and cost of care projections. <http://costprojections.cancer.gov/>. Accessed May 21, 2017.
28. Finkelstein EA, Trogon JG, Cohen JW, Dietz W. Annual medical spending attributable to obesity: payer- and service-specific estimates. *Health Aff (Millwood)*. 2009;28:w822-w831. doi:10.1377/hlthaff.28.5.w822
29. Centers for Disease Control and Prevention. The cost of arthritis in US adults. http://www.cdc.gov/arthritis/data_statistics/cost.htm. Accessed May 21, 2017.
30. Blumenthal D, David K, Guterman S. Medicare at 50: moving forward. *N Engl J Med*. 2015;372:671-677. doi:10.1056/NEJMp1414856
31. IMS Health. Overview of the specialty drug trend. <http://docplayer.net/4230764-Overview-of-the-specialty-drug-trend.html>. Accessed March 2, 2018.
32. Keehan SP, Stone DA, Poisal JA, et al. National health expenditure projections, 2016-25: price increases, aging push sector to 20 percent of economy. *Health Aff (Millwood)*. 2017;36:553-563. doi:10.1377/hlthaff.2016.1627
33. Yu J, Shah BM, Ip EJ, Chan J. A Markov model of the cost-effectiveness of pharmacist care for diabetes in prevention of cardiovascular diseases: evidence from Kaiser Permanente Northern California. *J Manag Care Pharm*. 2013;19:102-114. doi:10.18553/jmcp.2013.19.2.102
34. Mazzolini TA, Irons BK, Schell EC, Seifert CF. Lipid levels and use of lipid-lowering drugs for patients in pharmacist-managed lipid clinics versus usual care in 2 VA Medical Centers. *J Manag Care Pharm*. 2005;11:763-771. doi:10.18553/jmcp.2005.11.9.763
35. Hirsch JD, Bounthavong M, Arjmand A, et al. Estimated cost-effectiveness, cost benefit, and risk reduction associated with an endocrinologist-pharmacist diabetes intense medical management "tune-up" clinic. *J Manag Care Spec Pharm*. 2017;23:318-326. doi:10.18553/jmcp.2017.23.3.318
36. US Government Accountability Office. *Primary Care Professionals: Recent Supply Trends, Projections, and Valuation of Services*. Washington, DC: GAO; 2008. Report no. GAO-08-472T.
37. Rochon PA, Gurwitz JH. Optimising drug treatment for elderly people: the prescribing cascade. *BMJ*. 1997;315:1096-1099.
38. Finley PR, Crismon ML, Rush AJ. Evaluating the impact of pharmacists in mental health: a systematic review. *Pharmacotherapy*. 2003;23:1634-1644.
39. Cranor CW, Bunting BA, Christensen DB. The Asheville Project: long-term clinical and economic outcomes of a community pharmacy diabetes care program. *J Am Pharm Assoc (Wash)*. 2003;43:173-184.
40. Smith M, Bates DW, Bodenheimer T, Cleary PD. Why pharmacists belong in the medical home. *Health Aff (Millwood)*. 2010;29:906-913. doi:10.1377/hlthaff.2010.0209
41. Smith M, Bates DW, Bodenheimer TS. Pharmacists belong in accountable care organizations and integrated care teams. *Health Aff (Millwood)*. 2013;32:1963-1970. doi:10.1377/hlthaff.2013.0542
42. Giberson S, Yoder S, Lee MP. *Improving Patient and Health System Outcomes Through Advanced Pharmacy Practice: A Report to the U.S. Surgeon General*. Washington, DC: Office of the Chief Pharmacist, US Public Health Service; 2011.
43. Isetts BJ. Clinical and economic outcomes of medication therapy management services: the Minnesota experience. *J Am Pharm Assoc*. 2008;48:203-211. doi:10.1331/JAPhA.2008.07108
44. Butler A, Dehner M, Gates RJ, et al. Comprehensive medication management programs: 2015 status in Southern California. *Res Social Adm Pharm*. 2017;13:63-87. doi:10.1016/j.sapharm.2016.02.003
45. McInnis T, Capps K. Get the medications right: a nationwide snapshot of expert practices—comprehensive medication management in ambulatory/community pharmacy. <http://static.correofarmaceutico.com/docs/2016/07/15/getthemedicationsright.v22final-5.20.pdf>. Accessed March 2, 2018.
46. Hepler CD, Strand LM. Opportunities and responsibilities in pharmaceutical care. *Am J Hosp Pharm*. 1990;47:533-543.
47. Forster AJ, Clark HD, Menard A, et al. Adverse events among medical patients after discharge from hospital. *CMAJ*. 2004;170:345-349.
48. Kripalani S, Roumie CL, Dalal AK, et al. Effect of a pharmacist intervention on clinically important medication errors after hospital discharge: a randomized controlled trial. *Ann Intern Med*. 2012;157:1-10. doi:10.7326/0003-4819-157-1-201207030-00003
49. Isetts BJ, Brummel AR, Ramalho de Oliveira D, Moen DW. Managing drug-related morbidity and mortality in the patient-centered medical home. *Med Care*. 2012;50:997-1001. doi:10.1097/MLR.0b013e31826ecf9a
50. McInnis T. Medication underuse: the most underappreciated quality of care gap. *Pharmacy Times* <http://www.pharmacytimes.com/publications/directions-in-pharmacy/2015/march2015/medication-underuse-the-most-underappreciated-quality-of-care-gap>. Accessed May 21, 2017.
51. RTI International. Accountable Care Organization 2016 Program quality measure narrative specifications. <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/shared-savingsprogram/Downloads/2016-ACO-NarrativeMeasures-Specs.pdf>. Accessed May 21, 2017.
52. American College of Clinical Pharmacy; McBane SE, Dopp AL, Abe A, et al. Collaborative drug therapy management and comprehensive medication management: 2015. *Pharmacotherapy*. 2015;35:e39-e50. doi:10.1002/phar.1563
53. Patient-Centered Primary Care Collaborative. The patient-centered medical home: integrating comprehensive medication management to optimize patient outcomes resource guide. <https://www.accp.com/docs/positions/misc/CMM%20Resource%20Guide.pdf>. Accessed June 2, 2017.
54. Academy of Managed Care Pharmacy. Standardized framework for cross-walking MTM services to SNOMED CT codes. <http://www.amcp.org/SNOMED/>. Accessed June 2, 2017.