

UC Davis

UC Davis Previously Published Works

Title

VCAM-1 is a TGF- β 1 inducible gene upregulated in idiopathic pulmonary fibrosis

Permalink

<https://escholarship.org/uc/item/3nb577dj>

Journal

Cellular Signalling, 27(12)

ISSN

0898-6568

Authors

Agassandian, Marianna
Tedrow, John R
Sembrat, John
[et al.](#)

Publication Date

2015-12-01

DOI

10.1016/j.cellsig.2015.09.003

Peer reviewed



Published in final edited form as:

Cell Signal. 2015 December ; 27(12): 2467–2473. doi:10.1016/j.cellsig.2015.09.003.

VCAM-1 is a TGF- β 1 Inducible Gene Upregulated in Idiopathic Pulmonary Fibrosis

Marianna Agassandian, PhD^{1,2}, John R. Tedrow, MD^{1,2}, John Sembrat, BS^{1,2}, Daniel J. Kass, MD^{1,2}, Yingze Zhang, PhD^{1,2}, Elena A Goncharova, PhD^{2,3}, Naftali Kaminski, MD⁵, Rama K. Mallampalli, M.D.^{2,4}, and Louis J Vuga, MD, MPH^{1,2,#}

¹Dorothy P and Richard P Simmons Center for Interstitial Lung Diseases, University of Pittsburgh, Pittsburgh, PA

²Division of Pulmonary, Allergy and Critical Care Medicine, Department of Medicine, University of Pittsburgh, Pittsburgh, PA

³Vascular medicine Institute, University of Pittsburgh, Pittsburgh, PA

⁴Medical Specialty Service Line, Veterans Affairs Pittsburgh Healthcare System, Pittsburgh, Pennsylvania 15240

⁵Section of Pulmonary, Critical Care and Sleep Medicine, School of Medicine, Yale University, New Haven, CT

Abstract

Idiopathic pulmonary fibrosis (IPF) is a chronic lethal interstitial lung disease of unknown etiology. We previously reported that high plasma levels of vascular cell adhesion molecule 1 (VCAM-1) predict mortality in IPF subjects. Here we investigated the cellular origin and potential role of VCAM-1 in regulating primary lung fibroblast behavior. VCAM-1 mRNA was significantly increased in lungs of subjects with IPF compared to lungs isolated from control subjects ($p=0.001$), and it negatively correlated with two markers of lung function, forced vital capacity (FVC) and pulmonary diffusion capacity for carbon monoxide (DL_{CO}). VCAM-1 protein levels were highly expressed in IPF subjects where it was detected in fibrotic foci and blood

Correspondence to: Louis J. Vuga, M.D., M.P.H., NW628 Montefiore University Hospital, 3459 Fifth Avenue, Pittsburgh, PA, 15213, vugalj@upmc.edu.

Rama K. Mallampalli, MD., NW628 Montefiore University Hospital, 3459 Fifth Avenue, Pittsburgh, PA, 15213, mallampallirk@upmc.edu

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflicts of interest: None

Portion of these findings have been presented at the American Thoracic Society conference, San Diego, May 2014

Author Contributions

Conceived and designed the experiments: RKM, MA, LJV, NK, and EAG. Performed the experiments: LJV, and MA. Analyzed the data: LJV, MA, JRT and RKM. Contributed reagents/materials/analysis tools: LJV, DJK, and JS. Wrote the paper: MA, LJV, EAG, and RKM. Proofed paper and contributed to text: MA, LJV, NK, DJK, EAG, and RKM. Developed the IHC imaging, analyzed these data, and provided text material describing experiments: MA, LJV, EAG, DJK and JRT. Recruited subjects and organized specimen and clinical data collections, proofed paper and contributed some ideas/manuscript text: DJK and YZ. Edited paper and contributed scientific scrutiny NK, EAG, and RKM. Contributed to data treatment, statistical analyses and scientific proofreading: JRT.

vessels of IPF lung. Treatment of human lung fibroblasts with TGF- β 1 significantly increased steady-state VCAM1 mRNA and protein levels without affecting VCAM1 mRNA stability. Further, cellular depletion of *VCAM-1* inhibited fibroblast cell proliferation and reduced G2/M and S phases of the cell cycle suggestive of cell cycle arrest. These effects on cell cycle progression triggered by *VCAM1* depletion were associated with reductions in levels of phosphorylated extracellular regulated kinase 1/2 and cyclin D1. Thus, these observations suggest that VCAM-1 is a TGF- β 1 responsive mediator that partakes in fibroblast proliferation in subjects with IPF.

Keywords

IPF; VCAM-1; Lung Fibroblasts; TGF- β 1

Introduction

Idiopathic pulmonary fibrosis is a chronic lethal lung disease with unknown cause and no cure [1, 2]. IPF has the worst prognosis among interstitial lung diseases (ILD) [3] with a median survival of 2.5 to 3 years [4]. The principal hypothesis of IPF pathogenesis has been suggested to be aberrant alveolar- re-epithelialization caused by repeated alveolar epithelial injury [5]. The failure of proper wound healing is proposed to be due to a complex array of mechanisms involving modifications of fibroblast-myofibroblast transformation, epithelial cell apoptosis, changes in hemostasis of cytokines, chemokines, and transforming growth factor β 1 (TGF- β 1), all of which play critical roles in IPF pathogenesis via promoting extracellular matrix accumulation and phenotypic changes of fibroblasts and epithelial cells in IPF lungs [6–8].

Vascular cell adhesion molecule 1 (VCAM-1) is an immunoglobulin superfamily member [9] that is expressed in large and small blood vessels after cytokine stimulation and mediates adhesion of lymphocytes, monocytes, eosinophils, and basophils to vascular endothelium. VCAM-1 plays a critical role in the development of atherosclerosis and rheumatoid arthritis [10], facilitates leukocyte-endothelial cell adhesion and plays a role in signal transduction [11]. We previously reported that high VCAM-1 protein in peripheral blood levels of IPF patients predicts mortality [12], but its mode of expression and putative role as a contributor to IPF pathogenesis remains unclear. In some studies there has been limited success in detecting levels of VCAM-1 protein in IPF lungs [13]

In this study, we demonstrate that VCAM-1 expression is elevated in human IPF lung fibroblasts and that its expression is responsive to actions of a critical mediator of disease, transforming growth factor β 1 (TGF- β 1). Interestingly, silencing *VCAM-1* mRNA inhibits fibroblasts proliferation and impairs cell cycle progression through depletion of specific signaling factors implicating in cellular proliferation. In aggregate, these observations provide a foundation for further studies on the mechanistic role of VCAM-1 in IPF pathogenesis.

Materials and Methods

Materials

VCAM1 antibody was obtained from Novus Biologicals (Littleton, CO). Anti-Collagen type 1 antibody was from Rockland (Limerick, PA). β -actin antibody was purchased from Sigma-Aldrich (St. Louis, MO). The cyclin D1, cyclin D2, cyclin D3, cdk2, cdk4 and cdk6 antibodies were from Cell Signaling (Danvers, MA). The anti- ERK1/2, phosphor-ERK1/2, p38 and phosphor-p38 antibodies were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). The miRNA mini Kit was from Qiagen (Louisville, KY). The primers for VCAM1 and qRT-PCR were purchased from ABI (Foster City, CA). Recombinant human TGF- β was obtained from R&D Systems (Minneapolis, MN). VCAM1 shRNA was purchased from Dharmacon (Lafayette, CO). The CytoSelect BrdU cell proliferation ELISA kit was from Cell Biolabs (San Diego, CA). The Cell Cycle Phase Determination Kit was obtained from Cayman (Ann Arbor, MI). Western Lightning Plus ECL was from PerkinElmer (Boston, MA). Actinomycin D was from Sigma (St. Louis, MO).

Microarrays

Primary lung tissues were isolated from normal ($n=109$) and IPF lungs ($n=134$). Total RNA was extracted, labeled and hybridized to Agilent 44k whole human genome microarrays (Agilent Technologies, Wilmington, DE). After cyclic-LOESS normalization, Genomica and SAM were applied for statistical analysis. A 5% false discovery rate was used for significance in microarray data.

RNA extraction and real-time PCR analysis

All patients were evaluated at the University of Pittsburgh Medical Center, Pittsburgh, PA; the study was performed in accordance to the protocols approved by the University of Pittsburgh Institutional Review Board (IRB). The diagnosis of IPF was established on the basis of American Thoracic Society (ATS) and European Respiratory Society (ERS) criteria [14]. Clinical data were available through the Simmons Center Database at the University of Pittsburgh. Smoking status was defined as previously described [15]. All patients signed informed consent to participate in the study. Total cellular RNA was extracted from the lungs of eleven IPF and eleven control subjects obtained from University of Pittsburgh Health Sciences Tissue Bank (Pittsburgh, PA). RT-PCR was performed according to the manufacturer's protocol as we described previously [16].

Immunohistochemical Analysis

Paraffin-embedded IPF and control lung tissue specimens were obtained from the University of Pittsburgh Health Sciences Tissue Bank (Pittsburgh, PA), and immunohistochemical analysis was performed as previously described [17]. Briefly, the slides were deparaffinized in xylene, ethanol, and rehydrated in phosphate-buffered saline (PBS), blocked for 1 h in 10% bovine serum albumin and incubated with primary anti-VCAM1 antibody (Novus biologicals, Littleton, Co) overnight. Slides were then washed with PBS three times and incubated with biotinylated donkey anti-mouse secondary antibody (Jackson ImmunoResearch Laboratories Inc, West Grove, PA) for 1 h. The images were visualized

with an Olympus CH2™ microscope and obtained using aDP25 camera (Olympus America Inc., Melville, NY).

Cell culture, TGF-β1 stimulation and immunoblot analysis

Early passages (1–3) of primary normal human lung fibroblasts from the University of Pittsburgh Tissue bank as described above were cultured in F-12 (1×) medium supplemented with 10% fetal bovine serum (FBS) according to supplier's protocol. All experiments were performed on cells at 70–80% confluence. Treatment with recombinant TGF-β1 (5 ng/ml) was performed for 24 hrs. Immunoblots were performed with antibodies against VCAM-1 (1:1,000), β-actin (1:10,000) and bands quantitated and analyzed using ImageJ software.

Enzyme-Linked Immunosorbent Assay (ELISA)

IPF subjects ($n=48$) and control subjects ($n=50$) were evaluated at the University of Pittsburgh Medical Center and the study was approved by the IRB at the University of Pittsburgh. Recruitment and data collection has been previously described [18]. Smoking status was defined as described in [19]. The demographic and clinical information of subjects involved in this study are shown in Table 1 from the Lung Tissues Research Consortium (LTRC) for the gene expression cohort and Table 2 for the Pittsburgh plasma cohort. VCAM1 levels were detected using a VCAM-1 ELISA kit according to manufacturer's protocol.

Cell proliferation

The proliferation of lung fibroblasts was detected using a CytoSelect BrdU cell proliferation ELISA kit. To determine BrdU incorporation into cellular DNA during cell proliferation, cells on a 96-well cell culture plate were transfected with *VCAM-1* shRNA or a control shRNA using Effecten transfection reagent. After 12 hrs, medium was changed and cells were incubated for an additional 24 hrs. An aliquot of BrdU was then added to the medium and cells were incubated for an additional 3 hrs at 37°C. After washing with PBS, cells were fixed for 30 min and BrdU incorporation into total cellular DNA was determined using anti-BrdU antibody following the manufacturer's instructions.

Cell cycle phase determination

To determine cell cycle progression, the lung fibroblasts were transfected with *VCAM-1* shRNA or control shRNA as described above [20, 21]. After transfection, cells were fixed and then stained with propidium iodide for 30 mins at room temperature in the dark. Cells in individual cycle phases were analyzed by flow cytometry and captured with a 488 nm excitation laser.

Bleomycin murine model of fibrosis

Male and female C57BL/6 mice (6 to 8 weeks old) are deeply anesthetized and bleomycin at 3 U/kg (standard dose) or 1 U/kg (low dose) or saline control was administered intratracheally in a volume of 50 μL. Mice are sacrificed on days 1, 3, 7, 14, or 21 with pentobarbital, and the lungs are excised for determination of VCAM1 content. All

procedures were executed in accordance with approved protocols through the University of Pittsburgh Institutional Animal Care and Use Committee.

Statistical Analysis

The group comparisons between diseased and control subjects were performed using an unpaired two-tailed Student's *t*-test for normally distributed data. A level of $p < 0.05$ was considered statistically significant. Spearman correlation tests were used for computation of VCAM-1 mRNA correlation with pulmonary function tests utilizing Stata software. The bands on immunoblots were quantified by using ImageJ, and then densitometric ratios were calculated. The ratios data were statistically analyzed to discriminate their differences by using an unpaired student *t* test. The immunoblot data were representatives of 3–5 separate experiments.

Results

VCAM-1 mRNA levels in IPF subjects negatively correlate with pulmonary function

Data mining of our previously published LTRC microarray data revealed that VCAM1 is one of the most significantly expressed genes in IPF lung [22, 23]. Indeed, the steady-state VCAM-1 mRNA levels were significantly increased in IPF lungs compared to controls ($P = 0.0001$) (Figure 1A). Analysis of potential correlation of VCAM-1 mRNA levels with pulmonary function parameters revealed that there is a statistically significant negative correlation between VCAM-1 and DLco percent predicted ($\rho = -0.2$, $P = 0.02$) and FVC percent predicted ($\rho = -0.22$, $P = 0.0121$) (Figure 1B and C, respectively) demonstrating that increased VCAM-1 mRNA negatively correlates with pulmonary function. To validate the microarray results, we performed VCAM-1-specific qRT-PCR analysis of the independent cohort consisting of 11 IPF and 11 control lung specimens. In agreement with microarray data, VCAM-1 mRNA was significantly increased in IPF lungs compared to controls (Figure 1D). Taken together, these data demonstrate that VCAM-1 mRNA is increased in IPF lung that correlates with worse pulmonary function.

VCAM-1 protein levels are elevated in IPF lungs

To determine whether increased VCAM-1 mRNA levels result in elevated expression of protein, we next compared VCAM-1 protein levels in the plasma and lungs of IPF and non-diseased (control) subjects. Comparison of plasma from 48 IPF and 50 controls subjects using VCAM-1 ELISA assays demonstrated that VCAM-1 protein levels were significantly higher in the IPF cohort (Figure 2A). Levels of immunoreactive VCAM-1 were almost undetectable at baseline in control lungs. Similarly, analysis of whole lung tissue from four IPF and four control subjects showed a 3-fold increase in VCAM-1 protein levels in IPF lungs (Figures 2B, C). In addition, mice were given bleomycin as a model of lung fibrosis. In these mice, VCAM1 levels were also modestly increased compared to control lungs (Fig. 2D). Collectively, these data demonstrate that the VCAM-1 protein levels are elevated in both the plasma and lungs of IPF subjects and in a murine lung model of fibrosis.

VCAM-1 expression, cellular localization and regulation by TGF- β 1

To determine the cell types responsible for VCAM-1 expression in IPF lungs, we performed immunohistochemical analysis of lung tissue specimens from four IPF subjects and four control lungs. In control lung, VCAM-1 was poorly if at all, detected in distal lung tissue (Figure 3A). Importantly, VCAM-1 protein levels were markedly higher in IPF lungs with predominant localization in fibrotic areas (Figure 3A, arrows) with additional positive staining detected in lung vessels (Figure 3B, right panel) highlighting that lung fibroblasts are one major site of VCAM-1 expression in IPF lungs. To confirm our findings, we examined primary human fibroblasts isolated from IPF and control lungs. As seen in Figures 4A and B, IPF fibroblasts had higher VCAM-1 protein compared to cells from control lung donors. Of note, treatment of control lung fibroblasts with TGF- β 1, a crucial mediator of IPF pathogenesis increased VCAM-1 protein levels 4-fold in lung fibroblasts (Figure 4C, D) suggesting that fibroblast-responsive up-regulation of VCAM-1 in IPF lungs may be caused by TGF- β 1. To determine whether the increased VCAM-1 protein level is indeed induced by mRNA translation, total RNA was extracted from cells and real time PCR was performed. TGF- β 1 triggered a several-fold increase in the VCAM-1 transcript that was maximal at 6 hrs of analysis (Figure 4E). The response of VCAM-1 mRNA was still robust at 12 and 24 hrs of analysis after TGF- β 1 exposure but was of lower magnitude. Additional analysis of VCAM-1 mRNA stability were conducted using actinomycin D (Figure 4F). VCAM-1 transcript exhibited a $t_{1/2}$ of ~ 6 hrs and the stability of the mRNA was not significantly altered by exogenous TGF- β 1 administration. Taken together, these results suggest that TGF- β 1 released in IPF lung robustly increases VCAM-1 gene transcription rather than modulating the lifespan of its transcript in human lung fibroblasts.

VCAM-1 regulates lung fibroblast proliferation

To investigate the physiologic role of VCAM-1 in human fibroblasts, we focused on cell proliferation given the role of TGF- β 1 in cell growth and repair. We focused on endogenous VCAM-1 in fibroblast proliferation using BrdU labeling of cells (Figure 5A). Cells were transfected with a control shRNA or VCAM-1 shRNA and BrdU labeling were quantitated. Compared to lung fibroblasts treated with control shRNA, VCAM-1 shRNA showed ~47% lower BrdU cellular incorporation (Figure 5A). We next assayed effects of VCAM-1 depletion on cell cycle progression and observed that cells transfected with VCAM-1 shRNA exhibited an increase in G0/G1 coupled to reduced G2/ M and S- phase compared to control lung fibroblasts (Figure 5B). To evaluate potential mechanisms of VCAM-1 depletion, we assayed immunoreactive levels of several mediators of cell proliferative signaling (Figure 5C,D). Indeed, cells transfected with VCAM-1 shRNA showed reduced levels of phosphorylated p38, extracellular signal-regulated kinase $\frac{1}{2}$ (ERK $\frac{1}{2}$) (Figure 5C) and reduced mass of cyclin D1 (Figure 5D). The results suggest that VCAM-1 abundance modulates specific regulatory components involved in fibroblast growth.

Discussion

The mechanisms whereby VCAM-1 levels are elevated in IPF and how this adhesion molecule might contribute to the pathobiology of disease are not well understood. The new findings from this study include (i) the demonstration of a negative correlation between lung

VCAM-1 transcripts and pulmonary function, (ii) the identification that VCAM-1 protein expression is elevated in fibroblasts isolated from human IPF lungs, (iii) that VCAM-1 protein mass is increased in response to a pro-fibrotic regulator, TGF- β 1 through a mechanism that is independent of effects on VCAM-1 mRNA stability, and (iv) that silencing *VCAM-1* expression inhibits fibroblast proliferation by inducing G₀/G₁ cell cycle arrest. These observations provide a biologic framework for future studies investigating the molecular regulation of this adhesion molecule at the level of gene transcription in IPF models.

Our previous work indicated that up-regulated VCAM-1 protein levels in the peripheral blood of IPF subjects are associated with increased mortality and is in line with the association of this biomolecule with disease risk, outcome, and severity [12]. However, to date, the source of VCAM-1 in the circulation was unknown in IPF subjects. We found that in addition to peripheral blood, VCAM-1 mRNA is also significantly upregulated in lung tissues from IPF patients compared to a control cohort. Given that IPF subjects may have pulmonary vascular hypertension, our data are consistent with prior observations showing that serum VCAM-1 also predicts the risk for vascular disease [24]. Interestingly, studies in systemic sclerosis report that increased VCAM-1 in serum is also associated with pulmonary involvement and pulmonary hypertension thus supporting a potential mechanistic role for this protein [25, 26]. VCAM-1 is also observed in atopic asthma where it is related to disease activity [27]. While our data show significant differences in VCAM-1 protein and mRNA expression between IPF subjects and control subjects, we are aware of the limitations associated with a relatively small sample size for the group of subjects tested for validation of microarray datasets with IPF and control ($n=11$ /group). Despite this shortcoming, we were able to show consistently that an increase in IPF VCAM-1 protein in plasma using the existing cohort was sufficient to detect significant differences in VCAM-1 lung levels using immunoblot analysis. Given that lung mRNA VCAM-1 levels in IPF subjects negatively correlate with pulmonary function and VCAM-1 is shed during inflammation [28], these data suggest that the lung is one potential reservoir of blood-derived VCAM-1 that may be predictive of disease severity.

Because the cellular origin of VCAM-1 in pulmonary tissue may be linked to IPF disease pathogenesis, we further identified that VCAM-1 protein is expressed predominantly by IPF fibroblasts in fibrotic foci and is sensitive to TGF- β 1 stimulation. Because TGF- β 1 is a well-recognized driver of fibrotic lung remodeling, it is possible that increased VCAM-1 protein or mRNA concentrations in the peripheral blood may serve as an indicator of active fibrotic lung remodeling. Interestingly, high level expression of VCAM-1 in mouse lungs was observed in a murine model of radiation-induced pulmonary fibrosis, and its expression was significantly reduced after pulmonary manganese superoxide dismutase treatment, suggesting that at least in experimental chronic lung injury VCAM-1 levels might serve as an additional biomarker to monitor effects of anti-oxidant therapeutic intervention [29].

Not surprisingly, in addition to fibrotic foci and early-passage lung IPF fibroblasts, we found increased VCAM-1 levels in lung vessels in human IPF lung. Up-regulation of VCAM-1 is a sensitive marker of endothelial responses to oxidative stress. Further, vascular pathology is observed in IPF subjects with pulmonary hypertension [30, 31]. While a mechanistic role of

VCAM-1 in IPF is not yet fully elucidated, its role as a signaling molecule in other cell types is well documented [32, 33]. As an example, hypoxic exposure up-regulates VCAM-1 together with other adhesion molecules, triggering an array of pro-inflammatory signaling events in pulmonary arteries [34]. Many of these signaling events are reversed during normoxia, but may be accentuated by mediators expressed in IPF. In this regard, TGF- β 1, an integral mediator of pulmonary fibrosis, displays cross-talk with VCAM-1 raising a potential signaling mechanism that may be relevant in human IPF linked to increased adhesion molecules. The link between VCAM1 expression and TGF- β 1, identified by us in IPF fibroblasts, raises the possibility that TGF β 1 may be a driving stimulus that alters VCAM-1 expression leading to both fibrosis and vascular pathology [35].

VCAM-1 transcripts are known to be upregulated by pro-inflammatory stimuli [36] [37]. Here we observed a robust increase in both VCAM-1 mRNA and protein levels in response to exogenous TGF- β 1 in human fibroblasts. The VCAM-1 mRNA lifespan was relatively short ($t_{1/2} \sim 6$ hrs) consistent with the half-life of the transcript in synovial fibroblasts [36]. Interestingly, unlike the effects of IL-4 and TNF- α that induce VCAM-1 mRNA by stabilizing its transcripts, our data suggest that TGF- β 1 triggers an increase in mRNA synthesis in human lung fibroblasts [36]. Our data also contrasts with findings of others in astrocytes where TGF- β 1 inhibits TNF- α and IL-1 β induction of VCAM-1 mRNA [37]. The ability of TGF- β 1 that is centrally involved in IPF pathogenesis to elicit adhesion molecule expression in our studies also raises additional questions as to the physiological role of transcriptionally activated VCAM-1 on modulating fibroblast phenotypic behavior. We found that depletion of endogenous *VCAM-1* results in a decrease in fibroblast growth and reduction in levels of key signaling molecules implicated in cell survival and cycle progression. These results are consistent with observations elsewhere showing that VCAM-1 ligation activates ERK 2 resulting in increased expression of cyclin D1 [38]. Whether VCAM-1 is a bona fide molecular target for therapeutic intervention will require additional studies in preclinical models. These results do not exclude small molecule VCAM-1 inhibitors or biologics against adhesion molecules. In fact, studies are underway targeting the VCAM-1 co-receptor, α 4 β 1 integrin, in neoplasia supporting a role of this molecular apparatus in activating cell survival pathways [39]. However, given its promiscuous roles in numerous biological processes, the present data together with findings of others suggest a potential role of VCAM-1 as a biomarker in IPF and an upstream signaling molecule that regulates the fibroblast phenotype rather than a pharmaceutical target.

Acknowledgements

We would like to thank Dr. Mauricio Rojas for his editorial suggestions in the manuscript. This research was supported by the National Institutes of Health/National Heart, Lung, and Blood Institute [HL108869, HL073745, HL097376, R01HL095397, RC2HL101715, U01HL108642, and P01HL114453, and P50HL0894932], and the Dorothy P. and Richard P. Simmons Endowed Chair for Pulmonary Research. This material is also based upon work supported, in part, by the US Department of Veterans Affairs, Veterans Health Administration, Office of Research and Development, Biomedical Laboratory Research and Development and a VA Merit Review Award.

References

1. ATS/ERS International Multidisciplinary Consensus. Classification of the idiopathic interstitial pneumonias. *Am J Respir Crit Care Med.* 2002; 165:277–304. [PubMed: 11790668]

2. Raghu G CH, Egan JJ, Martinez FJ, Behr J, Brown KK, Colby TV, Cordier JF, Flaherty KR, Lasky JA, Lynch DA, Ryu JH, Swigris JJ, Wells AU, Ancochea J, Bouros D, Carvalho C, Costabel U, Ebina M, Hansell DM, Johkoh T, Kim DS, King TE Jr, Kondoh Y, Myers J, Müller NL, Nicholson AG, Richeldi L, Selman M, Dudden RF, Griss BS, Protzko SL, Schünemann HJ. An official ATS/ERS/JRS/ALAT statement: idiopathic pulmonary fibrosis: evidence-based guidelines for diagnosis and management. *Am J Respir Crit Care Med.* 2011; 183:788–824. [PubMed: 21471066]
3. Lynch J III, Saggar R, Weigt S, Zisman D, White E. Usual Interstitial Pneumonia. *Semin Respir Crit Care Med.* 2006; 27:634–651. [PubMed: 17195140]
4. Flaherty K, Travis W, Colby T, Toews G, Kazerooni E, Gross B, Jain A, Strawderman R III, Flint A, Lynch J III, Martinez F. Histopathologic variability in usual and nonspecific interstitial pneumonias. *Am J Respir Crit Care Med.* 2001; 164:1722–1727. [PubMed: 11719316]
5. Selman M, King TE, Pardo A. Idiopathic pulmonary fibrosis: prevailing and evolving hypotheses about its pathogenesis and implications for therapy. *Annals of internal medicine.* 2001; 134:136–151. [PubMed: 11177318]
6. Pottier N MT, Chevalier B, Puisségur MP, Lebrigand K, Robbe-Sermesant K, Bertero T, Lino Cardenas CL, Courcot E, Rios G, Fourre S, Lo-Guidice JM, Marcet B, Cardinaud B, Barbry P, Mari B. Identification of keratinocyte growth factor as a target of microRNA-155 in lung fibroblasts: implication in epithelial-mesenchymal interactions. *PLoS ONE.* 2009; 4:e6718. [PubMed: 19701459]
7. Brigham ZB, Willis C. TGF- β -induced EMT: mechanisms and implications for fibrotic lung disease. *AJP - Lung Physiol.* 2007; 293:L525–L534.
8. Pandit YH, KV CD, Yarlagadda M, Tzouveleki A, Gibson KF, Konishi K, Yousem SA, Singh M, Handley D, et al. Inhibition and role of let-7d in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med.* 2010; 182:220–229. [PubMed: 20395557]
9. Zhang D, Yuan D, Shen J, Yan Y, Gong C, Gu J, Xue H, Qian Y, Zhang W, He X, Yao L, Ji Y, Shen A. Up-regulation of VCAM1 Relates to Neuronal Apoptosis After Intracerebral Hemorrhage in Adult Rats. *Neurochemical research.* 2015
10. Navarro-Hernandez RE, Oregon-Romero E, Vazquez-Del Mercado M, Rangel-Villalobos H, Palafox-Sanchez CA, Munoz-Valle JF. Expression of ICAM1 and VCAM1 serum levels in rheumatoid arthritis clinical activity. Association with genetic polymorphisms, Disease markers. 2009; 26:119–126.
11. Yamada Y, Arao T, Matsumoto K, Gupta V, Tan W, Fedynshyn J, Nakajima TE, Shimada Y, Hamaguchi T, Kato K, Taniguchi H, Saito Y, Matsuda T, Moriya Y, Akasu T, Fujita S, Yamamoto S, Nishio K. Plasma concentrations of VCAM-1 and PAI-1: a predictive biomarker for post-operative recurrence in colorectal cancer. *Cancer science.* 2010; 101:1886–1890. [PubMed: 20491774]
12. Richards TJ, Kaminski N, Baribaud F, Flavin S, Brodmerkel C, Horowitz D, Li K, Choi J, Vuga LJ, Lindell KO, Klesen M, Zhang Y, Gibson KF. Peripheral blood proteins predict mortality in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med.* 2012; 185:67–76. [PubMed: 22016448]
13. Nakao A, Hasegawa Y, Tsuchiya Y, Shimokata K. Expression of cell adhesion molecules in the lungs of patients with idiopathic pulmonary fibrosis. *Chest.* 1995; 108:233–239. [PubMed: 7541743]
14. M.a.U.C. Demedts. ATS/ERS international multidisciplinary consensus classification of the idiopathic interstitial pneumonias. *Eur Respir J.* 2002; 19:794–796. [PubMed: 12030715]
15. King TE Jr, et al. Predicting survival in idiopathic pulmonary fibrosis: scoring system and survival model. *Am J Respir Crit Care Med.* 2001; 164:1171–1181. [PubMed: 11673205]
16. Vuga L, Ben-Yehudah A, Kovkarova-Naumovski E, Oriss T, Gibson KF, Feghali-Bostwick C, Kaminski N. WNT5A is a Regulator of Fibroblasts Proliferation and Resistance to Apoptosis. *Am J Respir Cell Mol Biol.* 2009
17. Xue J, Kass DJ, Bon J, Vuga L, Tan J, Csizmadia E, Otterbein L, Soejima M, Levesque MC, Gibson KF, Kaminski N, Pilewski JM, Donahoe M, Scirba FC, Duncan SR. Plasma B lymphocyte stimulator and B cell differentiation in idiopathic pulmonary fibrosis patients. *Journal of immunology (Baltimore, Md. :1950).* 2013; 191:2089–2095.

18. Vuga LJ, Milosevic J, Pandit K, Ben-Yehudah A, Chu Y, Richards T, Scierba J, Myerburg M, Zhang Y, Parwani AV, Gibson KF, Kaminski N. Cartilage oligomeric matrix protein in idiopathic pulmonary fibrosis. *PLoS one*. 2013; 8:e83120. [PubMed: 24376648]
19. King TE Jr, Toozé JA, Schwarz MI, Brown KR, Cherniack RM. Predicting survival in idiopathic pulmonary fibrosis: scoring system and survival model. *American journal of respiratory and critical care medicine*. 2001; 164:1171–1181. [PubMed: 11673205]
20. Chen BB, Glasser JR, Coon TA, C Z, HL M, Fenton M, JF M, M B, Mallampalli RK. F box protein FBXL2 targets cyclin D2 for ubiquitination and degradation to inhibit leukemic cell proliferation. *Blood*. 2012; 119:3132–3141. [PubMed: 22323446]
21. Chen BB, Glasser JR, Coon TA, Mallampalli RK. FBXL2 is a ubiquitin E3 ligase subunit that triggers mitotic arrest. *Cell Cycle*. 2011; 10
22. [accessed 2013 May 16] Gene Expression Omnibus. Available from: <http://www.ncbi.nlm.nih.gov/geo/>
23. Lung Tissue Research Consortium. <http://www.ltrcpublic.com/>.
24. Meyer O. Prognostic markers for systemic sclerosis. *Joint, bone, spine : revue du rhumatisme*. 2006; 73:490–494.
25. Shahin AA, Anwar S, Elawar AH, Sharaf AE, Hamid MA, Eleinin AA, Eltablawy M. Circulating soluble adhesion molecules in patients with systemic sclerosis: correlation between circulating soluble vascular cell adhesion molecule-1 (sVCAM-1) and impaired left ventricular diastolic function. *Rheumatology international*. 2000; 20:21–24. [PubMed: 11149656]
26. Denton CP, Bickerstaff MC, Shiwen X, Carulli MT, Haskard DO, Dubois RM, Black CM. Serial circulating adhesion molecule levels reflect disease severity in systemic sclerosis. *British journal of rheumatology*. 1995; 34:1048–1054. [PubMed: 8542206]
27. Popper HH, Pailer S, Wurzinger G, Feldner H, Hesse C, Eber E. Expression of adhesion molecules in allergic lung diseases. *Virchows Archiv : an international journal of pathology*. 2002; 440:172–180. [PubMed: 11964048]
28. Zonneveld R, Martinelli R, Shapiro NI, Kuijpers TW, Plotz FB, Carman CV. Soluble adhesion molecules as markers for sepsis and the potential pathophysiological discrepancy in neonates, children and adults. *Crit Care*. 2014; 18:204. [PubMed: 24602331]
29. Epperly MW, Sikora CA, DeFilippi SJ, Gretton JE, Bar-Sagi D, Archer H, Carlos T, Guo H, Greenberger JS. Pulmonary irradiation-induced expression of VCAM-1 and ICAM-1 is decreased by manganese superoxide dismutase-plasmid/liposome (MnSOD-PL) gene therapy. *Biology of blood and marrow transplantation : journal of the American Society for Blood and Marrow Transplantation*. 2002; 8:175–187.
30. Nathan SD, King CS. Treatment of pulmonary hypertension in idiopathic pulmonary fibrosis: shortfall in efficacy or trial design? *Drug design, development and therapy*. 2014; 8:875–885.
31. Pitsiou G, Papakosta D, Bouros D. Pulmonary hypertension in idiopathic pulmonary fibrosis: a review. *Respiration; international review of thoracic diseases*. 2011; 82:294–304.
32. Lee IT, Yang CM. Inflammatory signalings involved in airway and pulmonary diseases. *Mediators of inflammation*. 2013; 2013:791231. [PubMed: 23690670]
33. Cook-Mills JM, Marchese ME, Abdala-Valencia H. Vascular cell adhesion molecule-1 expression and signaling during disease: regulation by reactive oxygen species and antioxidants. *Antioxidants & redox signaling*. 2011; 15:1607–1638. [PubMed: 21050132]
34. Burke DL, Frid MG, Kunrath CL, Karoor V, Anwar A, Wagner BD, Strassheim D, Stenmark KR. Sustained hypoxia promotes the development of a pulmonary artery-specific chronic inflammatory microenvironment. *Am J Physiol Lung Cell Mol Physiol*. 2009; 297:L238–L250. [PubMed: 19465514]
35. Bartolome RA, Sanz-Rodriguez F, Robledo MM, Hidalgo A, Teixido J. Rapid up-regulation of alpha4 integrin-mediated leukocyte adhesion by transforming growth factor-beta1. *Mol Biol Cell*. 2003; 14:54–66. [PubMed: 12529426]
36. Croft D, McIntyre P, Wibulswas A, Kramer I. Sustained elevated levels of VCAM-1 in cultured fibroblast-like synoviocytes can be achieved by TNF-alpha in combination with either IL-4 or IL-13 through increased mRNA stability. *Am J Pathol*. 1999; 154:1149–1158. [PubMed: 10233853]

37. Winkler MK, Beveniste EN. Transforming growth factor-beta inhibition of cytokine-induced vascular cell adhesion molecule-1 expression in human astrocytes. *Glia*. 1998; 22:171–179. [PubMed: 9537837]
38. Lazaar AL, Krymskaya VP, Das SK. VCAM-1 activates phosphatidylinositol 3-kinase and induces p120Cbl phosphorylation in human airway smooth muscle cells. *J Immunol*. 2001; 166:155–161. [PubMed: 11123288]
39. Chen Q, Massague J. Molecular pathways: VCAM-1 as a potential therapeutic target in metastasis. *Clin Cancer Res*. 2012; 18:5520–5525. [PubMed: 22879387]

Highlights

IPF subjects have increased VCAM-1 mRNA linked to lung dysfunction

TGF- β 1 increases VCAM1 transcript levels without affecting VCAM1 mRNA stability

Cellular *VCAM-1* depletion inhibits fibroblast proliferation and cell cycle regulators

VCAM-1 is a TGF- β 1 responsive mediator that controls cell proliferation in human IPF.




Figure 1. VCAM-1 mRNA levels are increased in IPF lungs and negatively correlate with pulmonary function

A. VCAM1 expression levels were extracted from publically available data by microarray profiling of lung tissues from IPF (IPF: $n = 134$) and control subjects (control: $n = 109$). VCAM-1 is up-regulated in IPF subjects compared to control ($*P = 0.0001$) [22, 23]. **B, C:** Associations between continuous variables were established by a Spearman correlation analysis and the regression lines have been fitted in two-way scatterplots by Stata software. For FVC association; $\rho = -0.2186$, $P = 0.0123$ and for DLco; $\rho = -0.2074$, $P = 0.02$. **D:** VCAM-1 mRNA levels in lungs of control subjects ($n = 11$) and IPF subjects ($n = 11$) were measured using qRT-PCR analysis. Total cellular RNA was extracted with Triazol. The mitochondrial ribosomal protein S18A (MRPS18A) was used as an internal standard. Data represents fold change in VCAM-1/S18A ratios with a ratio for control cells taken as one fold. Data are expressed as means \pm SE using an unpaired students t-test $*P < 0.003$ for controls vs IPF groups.

VCAM1 in IPF




Figure 2. Plasma VCAM-1 levels are increased in IPF subjects

A. Plasma VCAM-1 (ng/ml) was assayed in control subjects ($n=50$) and IPF ($n=48$) subjects using an ELISA assay. Approximately equal sample sizes were used in each group and the comparison between each group was computed using an unpaired student t-test. ($P = 0.0003$, mean = 737.2979 and SD = 186.8587 for IPF and mean = 596.735 and SD = 178.8378 for controls). **B, C.** Whole lung tissue lysates were obtained from IPF lungs and control lungs as described in Methods. Protein levels of VCAM-1 and collagen 1 (Col1, positive control) were determined in lungs by immunoblotting. Shown is a representative of immunoblot for VCAM1, Col1, and β -actin as a loading control. The bands were quantified by ImageJ and the ratios are presented in panel C showing fold change in IPF lungs compared to control lungs. **D.** Control mice (C) or bleomycin (BL) treated mice ($n=3$ mice/group) were euthanized and analyzed for VCAM1 protein in whole lung by immunoblotting (inset) and bands quantitated densitometrically (below).




Figure 3. Expression of VCAM-1 protein in IPF lung

A. Paraffin embedded IPF and control lungs were incubated overnight in primary antibody at 4 ° C. An appropriate secondary antibody was used to identify the target protein on the lung tissue slides. The IPF lung shows staining in endothelial cells (black arrows), airway epithelial cells (red arrowheads) and cells in the interstitium (yellow arrowheads). No staining was detected with the non-immune isotype control (magnification $\times 100$, yellow inset bar=100 microns). **B.** Brown staining for VCAM-1 protein in IPF lung vessels (V) (arrow) was detected and lack of staining was observed in control lung vessels.




Figure 4. Expression of VCAM-1 protein in primary lung fibroblasts

A–B. Control and IPF human lung fibroblasts were cultured and harvested at 70–80% confluence. VCAM-1 protein was examined by immunoblotting. Collagen 1 served as positive control for IPF lung fibroblasts and β -actin was used as a loading control of individual samples. The immunoblot shown is a representative of three independent experiments. In **(B)** individual bands were analyzed densitometrically and corrected for loading and densitometric ratios are shown as a bar graph. * $P < 0.05$ for control vs IPF groups. **C–D.** Lung fibroblasts were cultured and stimulated with TGF- β 1 (5 ng/ml) for 24 hrs. The bands on immunoblots were quantified by ImageJ and densitometric ratios shown in Figure **D**. Statistical analysis was performed by using an unpaired student t-test. * $P = 0.0018$ for control vs TGF- β 1 groups. Data shown are representatives of 3–5 independent experiments. **E.** Lung fibroblasts were cultured and stimulated with or without TGF- β 1 (**E**) for 24 hrs. **F.** Lung fibroblasts were cultured in the presence of actinomycin D (ActD) with or without the inclusion of TGF- β 1 (5 ng/ml) in the culture medium for various time points as shown. In both (**E**) and (**F**) total cellular RNA was extracted and real time PCR was performed to assay VCAM-1 mRNA as shown. The data represents three separate experiments except panel (**F**), $n=2$.




Figure 5. VCAM-1 cellular depletion decreases fibroblast proliferation by impairing cell cycle progression

A. Human lung fibroblasts were transfected with control shRNA or *VCAM-1* shRNA and proliferation of cells was assessed using BrdU labeling as described in Methods. Significantly reduced BrdU incorporation was observed in *VCAM-1* shRNA transfected cells (* $P < 0.05$ in shRNA control vs *VCAM-1* shRNA). **B.** *VCAM-1* cellular depletion was conducted as in (A). *VCAM-1* silencing on cell cycle progression was then determined in fibroblasts after depleting the adhesion molecule. Significantly reduced S phase and G2/M phase were observed in *VCAM-1* shRNA transfected cells ($P < 0.05$) versus control shRNA. **C–D.** Human lung fibroblasts were cultured, transfected and harvested at 70–80% confluence as described above using shRNA. Cell lysates were harvested and processed for immunoblot analysis of indicated cell signaling proteins and key proteins involved in cell cycle regulation. The data are representative of $n=2$ separate experiments.

Table 1

LTRC Lung Cohort	Control	IPF	P-Value
Age	63.6 ± 11.4	63.9 ± 8.2	NS
Female	27 (42.3%)	21 (25%)	NA
Non-Caucasian	8 (7.33%)	9 (6.7%)	NA
FEV1%	95.0 ± 12.62	70.4 ± 18.2.6	1.03E-25
FVC%	94.4 ± 13.12	63.38 ± 16.16	3.55E-38
DLCO %	84.05 ± 16.7	46.7 ± 17.8	5.96E-38

¹⁾FVC = Force Vital Capacity

²⁾DLco= Carbon Monoxide Diffusing Capacity

³⁾And FEV1= Forced Expiratory Volume in One Second.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Pittsburgh Plasma Cohort	Characteristics	IPF
Age	Years	64.8 ± 8.2
Female Sex	Absolute/Percent %	17 (35.4%)
Non-Caucasian	Absolute number	3(6.25 %)
PFTs	FEV1 %	77.2 ± 18.8%
	FVC%	68 ± 19.8 %
	DLCO%	48.2 ± 11 %

⁴⁾ FVC = Force Vital Capacity,

⁵⁾ DLco= Carbon Monoxide Diffusing Capacity

⁶⁾ FEV1= Forced Expiratory Volume in One Second.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript