UCLA

UCLA Previously Published Works

Title

Brine-assisted anatexis: Experimental melting in the system haplogranite-H₂O-NaCl-KCl at deep-crustal conditions

Permalink

<https://escholarship.org/uc/item/3nm1r002>

Authors

Aranovich, LY
Newton, RC
Manning, CE

Publication Date

2013-07-05

DOI

10.1016/j.epsl.2013.05.027

Peer reviewed

Corrigendum

Corrigendum to “Zircon solubility and zirconium complexation in $\text{H}_2\text{O}+\text{Na}_2\text{O}+\text{SiO}_2 \pm \text{Al}_2\text{O}_3$ fluids at high pressure and temperature” [Earth Planet. Sci. Lett. 349–350 (2012) 15–25]

Max Wilke^{a,*}, Christian Schmidt^a, Julien Dubrail^a, Karen Appel^b, Manuela Borchert^b, Kristina Kvashnina^c, Craig E. Manning^d

^a Deutsches GeoForschungsZentrum (GFZ), Telegrafenberg, 14473 Potsdam, Germany

^b Deutsches Elektronen Q6 Synchrotron, Notkestr. 85, 22607 Hamburg, Germany

^c European Synchrotron Radiation Facility, 6 rue Horowitz, 38043 Grenoble, France

^d Department of Earth and Space Sciences, University of California, Los Angeles, CA 90025-1567, USA

The author regrets that Fig. 5a and b in the published paper was not correct; correct figures are given below.

DOI of original article: <http://dx.doi.org/10.1016/j.epsl.2012.06.054>

* Corresponding author.

E-mail address: max@gfz-potsdam.de (M. Wilke).

Fig. 5. Comparison of experimental (thick lines) and simulated (thin lines) XANES spectra for Zr in aqueous fluids. (a) Spectra of Zr in basic and acidic solutions. (b) Spectra of Zr in aqueous fluids with Na–Al–Si components as indicated. Spectra shown here are the same as shown in Fig. 3, where experimental conditions are indicated.