UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

Reconciling agriculture, carbon and biodiversity in a savannah transformation frontier

Permalink

https://escholarship.org/uc/item/3p34t1cf

Journal

Philosophical Transactions of the Royal Society B Biological Sciences, 371(1703)

ISSN

0962-8436

Authors

Estes, LD Searchinger, T Spiegel, M <u>et al.</u>

Publication Date 2016-09-19

DOI

10.1098/rstb.2015.0316

Peer reviewed

PHILOSOPHICAL TRANSACTIONS B

rstb.royalsocietypublishing.org

Research



Cite this article: Estes LD *et al.* 2016 Reconciling agriculture, carbon and biodiversity in a savannah transformation frontier. *Phil. Trans. R. Soc. B* **371**: 20150316. http://dx.doi.org/10.1098/rstb.2015.0316

Accepted: 19 May 2016

One contribution of 15 to a theme issue 'Tropical grassy biomes: linking ecology, human use and conservation'.

Subject Areas:

ecology, environmental science

Keywords:

trade-offs, optimization, savannahs, agriculture, biodiversity, carbon

Authors for correspondence:

L. D. Estes e-mail: lestes@princeton.edu T. Searchinger e-mail: tsearchi@princeton.edu

Electronic supplementary material is available at http://dx.doi.org/10.1098/rstb.2015.0316 or via http://rstb.royalsocietypublishing.org.

THE ROYAL SOCIETY PUBLISHING

Reconciling agriculture, carbon and biodiversity in a savannah transformation frontier

L. D. Estes^{1,2}, T. Searchinger², M. Spiegel¹, D. Tian¹, S. Sichinga⁴, M. Mwale⁴, L. Kehoe⁵, T. Kuemmerle^{5,6}, A. Berven¹, N. Chaney³, J. Sheffield¹, E. F. Wood¹ and K. K. Caylor¹

¹Civil and Environmental Engineering, ²Woodrow Wilson School, and ³Program in Atmospheric and Oceanic Sciences, Princeton University, Princeton, NJ 08544, USA

⁴Zambia Agricultural Research Institute, Mt. Makhulu Research Station, Chilanga, Zambia
⁵Geography Department, and ⁶Integrative Research Institute for Transformations in Human-Environment Systems (IRI THESys), Humboldt University, 10099 Berlin, Germany

(D) LDE, 0000-0002-9358-816X

Rapidly rising populations and likely increases in incomes in sub-Saharan Africa make tens of millions of hectares of cropland expansion nearly inevitable, even with large increases in crop yields. Much of that expansion is likely to occur in higher rainfall savannas, with substantial costs to biodiversity and carbon storage. Zambia presents an acute example of this challenge, with an expected tripling of population by 2050, good potential to expand maize and soya bean production, and large areas of relatively undisturbed miombo woodland and associated habitat types of high biodiversity value. Here, we present a new model designed to explore the potential for targeting agricultural expansion in ways that achieve quantitatively optimal trade-offs between competing economic and environmental objectives: total converted land area (the reciprocal of potential yield); carbon loss, biodiversity loss and transportation costs. To allow different interests to find potential compromises, users can apply varying weights to examine the effects of their subjective preferences on the spatial allocation of new cropland and its costs. We find that small compromises from the objective to convert the highest yielding areas permit large savings in transportation costs, and the carbon and biodiversity impacts resulting from savannah conversion. For example, transferring just 30% of weight from a yield-maximizing objective equally between carbon and biodiversity protection objectives would increase total cropland area by just 2.7%, but result in avoided costs of 27-47% for carbon, biodiversity and transportation. Compromise solutions tend to focus agricultural expansion along existing transportation corridors and in already disturbed areas. Used appropriately, this type of model could help countries find agricultural expansion alternatives and related infrastructure and land use policies that help achieve production targets while helping to conserve Africa's rapidly transforming savannahs.

This article is part of the themed issue 'Tropical grassy biomes: linking ecology, human use and conservation'.

1. Introduction

Meeting growing food demands while minimizing ecological losses presents one of the twenty-first century's major challenges. Many global sustainability studies and climate mitigation pathways call for eliminating or rapidly phasing out emissions from land use change [1,2]. Unfortunately, these ambitions do not appear realistically attainable in sub-Saharan Africa, due to a projected doubling of population by 2050 combined with rapidly growing economies, which will combine to treble food demand [1,3]. For example, even with the

2

substantial yield growth (3–5% per year) projected by the UN Food and Agriculture Organization (FAO), the region would probably need to add more than 100 million hectares of cropland by 2050 [1]. Much and probably the great bulk of that new cropland is likely to occur in the 700 million hectares of savannahs and shrublands of the region that receive sufficient rainfall to permit crop production [1].

Savannahs sometimes receive limited respect in global land use assessments and have been a major focus of agricultural expansion, not only in Africa but in places such as the Cerrado of Brazil and northern Thailand [4]. Their carbon contents and biodiversity values may be completely ignored on the grounds that they are not forests [1]. Yet, the wetter woodlandsavannahs and shrublands of Africa have a similar average richness, although not density, for birds and mammals as the world's wetter tropical forests, and 75% of these habitats have higher overall vertebrate diversity than 75% of the rest of the world [1]. Their carbon contents are also substantial, particularly relative to their likely yields, so that their use for food crops is unlikely to result in fewer carbon emissions per ton of crop [1]. Conversion of over 100 million hectares of these lands would release tens of millions of tons of carbon [1].

The high likelihood of some cropland expansion in sub-Saharan Africa raises the prospect of whether this expansion could be located in ways that meet agricultural production goals, but for substantially lower loss of carbon and biodiversity than conventional agricultural development pathways. Analysing this question must be done at national or sub-national scales, because that is where key decisions are made, and accurate results require data that are high in both resolution and quality.

In this paper, we look at Zambia, which is a bellwether for this land use challenge in sub-Saharan Africa. On the one hand, Zambia has great need to boost its food production. Sixty per cent of Zambia's population is rural [5], *per capita* calorie food availability is just 2100 kcal [6] and child stunting rates are 40% [7]. According to the medium estimate of the United Nations, Zambia's population is projected to grow 265% from 16.2 million in 2015 to 43 million in 2050 [8], which will combine with rapid economic growth (more than 6%, [5]) to substantially increase *per capita* calorie demands [3]. Zambia has also been expanding its agricultural exports and has agronomic and situational characteristics that put it in a strong position to continue to do so to other parts of the region.

At the same time, two thirds of Zambia's 738 400 km² land area is dominated by woodland-savannahs, which have sufficiently dense tree cover to be classified as forest under either of the definitions (more than 10% or more than 30% tree cover) used by the United Nations Framework Convention on Climate Change [9]. These ecosystems store substantial amounts of carbon, with 50% of Zambia having a carbon density of at least 130 t ha⁻¹ (vegetative carbon and soil carbon in the top 1 m), while an additional 40% exceeds 80 t ha⁻¹. Meanwhile, Zambia's deforestation rate of 2500–3000 km² y⁻¹ is one of the highest in the world [10].

The biodiversity of these woody savannahs is also high. Most of Zambia is covered by what the World Wildlife Fund has classified as the Central Zambian Miombo Woodland ecoregion [11,12], which contains some 3800 plant species, making it the 17th richest ecoregion in the world in floral diversity (out of 867) [13]. Vegetation is characterized by a mix of tall, frequently evergreen, trees (15–20 m), broadleaf shrubs and grasses, and is heavily interspersed with wetlands [12]. The large expanses of relatively intact habitat give this woodland the third highest richness for mammal species of the world's ecoregions [11,14,15]. It supports rare species including the black rhino, major predators, and highly migratory, wide-ranging species, such as elephants [12]. Zambia overall also has a high bird diversity, with 753 recorded species [16]. Although few bird and mammal species are endemic, reptile and amphibian endemism is high, with 19 endemic reptile species and 13 endemic amphibians [14].

There is a long history of using models to achieve conservation objectives in the face of cost and other constraints (e.g. [17,18]). MARXAN, perhaps the most widely used of these, is a decision-support program that is designed to efficiently select conservation area networks [19,20]. Within the more general land use planning field, an even broader range of models exists for finding optimal spatial configurations that satisfy multiple land use objectives (see Cao et al. [21] for an example and review of alternatives). In the present case, the need is not to select protected areas but to identify areas for cropland expansion that meet the objectives of agricultural development and environmental conservation. Koh & Ghazoul [22] developed one model for examining trade-offs among oil palm development, rice production and carbon and biodiversity production in Indonesia, and a related approach was used to identify sustainable agricultural intensification solutions in southern Tanzania [23]. In both examples, the model assigned equal weight to each land use objective in the compromise scenarios that were analysed. However, one of the great challenges for land use planning is that different objectives can be difficult to weigh against each other ex ante, because the values are often abstract, making them harder to evaluate without first viewing the consequential trade-offs [21,23]. Governments and other stakeholders also tend to have varying, and often conflicting, priorities, and their willingness to compromise may depend on the degree to which their priorities are impacted.

To overcome these problems, we developed a new, publicly available model that allows varying levels of compromise between competing agricultural and environmental interests to be evaluated while still meeting agricultural production targets. The model adopts different methods than the two aforementioned models [22,23] to estimate agricultural potential and conservation objectives, and can allocate land for multiple crops simultaneously. More fundamentally, our model is based on linear programming, which allows users to place varying weights on the different land use objectives [21]. This feature allows different stakeholders to work together to choose the most appealing combination of weights based on their land use preferences, and their own assessments of what are and are not acceptable costs within those objectives.

We apply this model to Zambia so that we can explore its strengths and limitations as a decision-making tool, and also to answer a basic but important question: is it possible to achieve a reasonable balance of different interests by reducing the carbon and biodiversity costs of expansion while still focusing on land use areas with high agricultural potential? Answering this question is becoming increasingly important as savannahs and other grassy biomes receive increasing agricultural development pressure, and may become increasingly targeted as an indirect result of the laudable efforts to conserve tropical forests [1].

rstb.royalsocietypublishing.org Phil. Trans. R. Soc. B 371: 20150316

3

2. Material and methods

The model we developed seeks to find optimal land use configurations that satisfy the production targets for multiple crops while minimizing four costs: the total land area required, transportation costs, carbon released from land conversion and impacts on biodiversity. Although total land area is treated as a cost, minimizing this cost identifies lands with the highest likely yields. By minimizing total land required as well as transportation costs, the model therefore finds areas that reflect two important indicators of agricultural potential. By minimizing carbon and biodiversity costs, the model also finds areas that minimize two important dimensions of environmental impact. The model chooses final areas depending on the weights that a user assigns to each of these four objectives.

To enable this functionality and to identify optimal solutions, the model is structured to evaluate each objective in terms of production efficiency or the ratio of each cost per hectare to crop production per hectare (yield) that results from converting land to agriculture. For land area, cost is measured as the number of square kilometres required to meet the crop production target. For the other objectives, the costs indicators are transportation time, carbon loss or biodiversity impacts.

We focus our model initially on two crops, maize and soya beans, because of their existing and potential significance for Zambia's agricultural sector, and their likely dominance in the future growth of agriculture in Africa's broader savannah regions [1]. Maize is the largest crop by area and consumption in Zambia [6], while soya beans are expanding rapidly as a commodity crop to meet regional and global demands for animal feed and direct human consumption [24]. Because the model is designed to meet production targets for multiple crops, we apply algorithms for allocating each crop to its highest value cells, given the particular combination of objectives being assessed.

Further details on model structure and all model input calculations can be found in the electronic supplementary material.

(a) Model inputs

Our model focuses on the costs of cropland expansion into areas that are currently not farmed but could be. To identify such areas, we calculated the proportion of currently cropped and settled grid cells within each 1 km² grid cell. We excluded the proportion of cells that were either protected or had slopes greater than 20%, which are both generally considered unsuitable for farming and strong actual predictors of cropland presence or absence [25]. We estimated the costs of converting each cell as described in the following sections.

(i) Likely yields

We spatially estimated likely yields for maize and soya beans using a three-step approach. In the first step, we used the decisionsupport system for agrotechnology transfer (DSSAT, [26]) to simulate maize and soya bean yields at the sites of 40 weather stations distributed across Zambia. They are the points where a 31-year, gridded daily meteorological dataset has received the greatest amount of bias-correction and infilling by weather observations [27–29]. We used soil profiles corresponding to those locations drawn from the WISE v. 1.1 gridded soil profile database [30], and simulated for each location maize yields over all 31 years under commercial input practices for three to four cultivars representative of short and medium season length according to the Zambian crop calendar [31].

In the second step, we used an empirical model to map the DSSAT-modelled yields onto a 1 km grid, taking advantage of a finer resolution soil map that provides a common frame for estimating both yield potential and soil carbon stocks (see below), but has insufficient detail in several parameters

(particularly effective rooting depth) required by DSSAT to provide a sound empirical basis for running the model directly. We used a generalized additive model [32] to predict the values of the DSSAT-simulated yields, using best-fitting subsets of the following weather and soil variables: growing season precipitation, growing degree days, mean growing season shortwave solar radiation, per cent organic carbon, pH and per cent clay content. The soil predictors represented mean values in the top 1 m of soil and were extracted from a new 1 km global database of soil properties [33]. This two-step approach allowed us to more soundly capture crop responses to Zambia's climatic variability and potential management practices, and to base potential yields on the same dataset used to estimate carbon costs (see below).

The first two steps resulted in a map of potential yield under high inputs and good management, and assuming no losses due to other limitations such as diseases. These yield values therefore overestimate what farmers are truly likely to achieve. To reflect more realistic estimates of future yields, we therefore rescaled the yield maps to match FAO-projected Zambian yields in 2050 (4.4 t ha⁻¹ for maize and 3.6 t ha⁻¹ for soya bean). We used the FAO-projections for sub-Saharan Africa as a whole to derive an annual growth factor and applied that factor to the average 2009–2014 yields for each crop in Zambia. Overall, the distribution of yields derived from our crop modelling is most important to targeting land use, while this choice of average yield determines the total quantity of land needed to meet production targets.

(ii) Transportation costs

As a second measure related to agricultural potential, we estimated the travel time to the nearest major market town. Travel time serves here as a proxy for differences in output prices and production costs, such as fertilizer and transport costs [34], and, via the density of road networks, is strongly correlated with the suitability of land for farming [25,35]. For this analysis, we defined market towns as the administrative capitals of Zambia's districts, as well as any town outside of Zambia having a population of at least 10 000 within a region bounded by 19.5° to 34.5° longitude and -19.5° to -8° latitude. These neighbouring towns were included because they may be more easily accessible to Zambians living in border regions than the nearest district capital.

We calculated travel time in hours using cost-distance analysis drawing on several spatial datasets. Vector data for Zambian roads obtained from the Government of Zambia (GOZ) and for the broader region from OpenStreetMap (http://openstreet map.org) were merged to create a single dataset of regional roads that distinguished between trunk, primary, secondary and tertiary roads. For market towns, we obtained district capital town locations from the GOZ and for towns outside Zambia we used data from the Gridded Population of the World Version 3 [36] database. For waterbodies, we used the USGS Hydrosheds database [37] to indicate the position and size of rivers, based on their contributing area, and used a lakes vector provided by the GOZ. The roads, rivers and lakes data, as well as Zambia's border, were converted to 1 km grids wherein each feature was assigned a travel time value based on how long it would take to traverse 1 km. The resulting grids were merged into a single 'friction' surface, which was used to compute the number of travel hours along the fastest route to the nearest market town (see the electronic supplementary material, for further details).

Total travel costs for any land use scenario equal the total time needed to transport the tonnage of crops produced in each converted cell to the nearest town. We divided these costs by 20 to produce 'truck hours' on the assumption of a 20-ton truck. Doing so produces a more intuitive unit, but does not alter mapping outcomes.

rstb.royalsocietypublishing.org Phil. Trans. R. Soc. B 371: 20150316

4

(iii) Carbon

To estimate potential carbon costs from cropland conversions, we developed maps of vegetative and soil carbon stocks for Zambia. Vegetative carbon was calculated from a recent map of aboveground biomass developed by Baccini *et al.* [38]. We estimated below-ground biomass using savannah and miombo woodland-specific root-shoot ratios [39,40], and then converted the total biomass to carbon using a ratio of 0.47 [41]. The same soil carbon dataset used for crop yield prediction, together with an accompanying map of bulk density, was used to calculate soil carbon stocks in the top 1 m. We calculated that the potential carbon loss due to agricultural conversion would be 100% of vegetative carbon and 25% of soil carbon [1].

(iv) Biodiversity

There are many ways of evaluating biodiversity value and potential impacts, including species richness of different taxa in absolute terms or based on level of threat or endemism (e.g. [1,24]). Other possibilities include factors designed to address the significance of contiguity or the impact of different types of development on diversity (e.g. [22,42]). The actual measures selected should reflect both conservation priorities and the quality of available data. For a fine-scaled, country-level analysis such as this one, the species range map data that form the basis for many broader impact assessments (e.g. [43]) have an effective resolution that is too coarse. We therefore developed a composite measure of biodiversity value based on several datasets.

First, we used the potential vegetation maps of East Africa [44], a 2010 landcover dataset for Zambia (http://apps.rcmrd.org/ landcoverviewer/), and a shapefile of protected areas to calculate how much of Zambia's 20 unique vegetation types remained after conversion to croplands and settlements as of 2010, and what proportion of that remainder fell within national parks. We used these values to calculate an index of rarity and threat. We calculated rarity as the proportion of Zambia occupied by each vegetation type's remaining area, which we then log-transformed to account for the highly skewed distribution. For threat, we calculated the proportion of each vegetation type falling within protected areas, and then multiplied the threat and rarity values to create the index [45]. We then used the remaining vegetation map to calculate a second measure, the proportion of untransformed vegetation within an 11×11 km neighbourhood centred on each 1 km². By adding this measure to the first index and normalizing, we created a biodiversity score with a range of 0-1 ha⁻¹, which gives equal weight to a habitat based on (i) how rare or threatened and (ii) how undisturbed it is. Summing the biodiversity score for each converted grid cell generates a total cost of biodiversity impacts for each scenario.

As a final modification, we masked out the areas of existing national parks and game management areas to respect the national judgement that they should not be used for expanding farmland. We retained a third protected category, forest reserves, as these tend to be far less stringently protected, having already lost 8.3% of their area to cropland or settlements, but assigned these areas a high biodiversity score (0.927, the inverse of their converted proportion).

(v) Potential farmland and production targets

Future demand for maize and soya beans produced in Zambia is probably difficult to estimate, and our model can easily be run with different production targets. For this paper, we adopted targets of a fourfold increase for maize for 2050 compared with 2009–2014 average, and a 10-fold increase for soya beans. These increases reflect the rate of growth in Zambia from 2000–2014, and a mean production trend for soya bean growth in southern Africa estimated by Gasparri *et al.* [24], with small additional adjustments to account for the fact that Zambia has one of the highest agricultural potentials in sub-Saharan Africa and has high potential to expand exports.

(vi) Yields on current land

The amount of land conversion to meet a production target will depend on the growth of yields on existing croplands, which are well short of their current potentials [46]. For the principal scenarios we present in this paper, we assume that yields on current cropland will achieve the average likely yields for Zambia's *potential* farmland. However, to examine the possible significance of yield gains on existing cropland, we also examined scenarios in which the yield gains on existing cropland (i) only manage to close 50% of the gap between current and our projected future yields and (ii) exceed the gap by 50% (i.e. achieve 150% of the expected yield gain).

(b) Normalization and weighting

Weighting different objectives to some extent depends purely on preferences, but it is important that these preferences be expressed in a numerical form that reasonably reflects them so that a quantitative solution can be computed. To achieve this result, we first express the different objectives as efficiencies, e.g. carbon loss per ton of crop yield, and then normalize the units of each objective based on the range of its scores. Weights representing the percentage of importance are placed on each objective, expressed as a decimal between 0 and 1, with values closer to 1 indicating stronger preferences, and with all weights summing to 1. The final score is the sum of the resulting products for each objective for each grid cell, which is used to rank each cell according to its ability to meet the production target most for the lowest total cost, given those weights. The model then selects cells in descending rank order until the cumulative production of converted cells reaches each crop's corresponding production target.

Although this weighting system treats choices as relative preferences, our model calculates the actual costs of each objective in absolute terms: total area converted, total transportation costs from new croplands, total carbon released from land conversions and total impacts on biodiversity. After reviewing these costs, model users can then adjust the preferences to reflect desirable or politically acceptable trade-offs.

3. Results

(a) Significance of yield gains on existing cropland

In our main scenario, we assumed that the gap between current and projected future yields would be fully closed on existing cropland, but we also ran the model for cases when only 50% or 150% of the gap was closed (with equal weights across all objectives). In the former case, the total land area required to meet the target increased by 18% (14 634–17 202 km²), transport costs increased 22% (5.52–6.61 × 10⁻⁵ truck hours), carbon losses increased by 19% (597–711 Mt), and the total biodiversity index of converted areas increased by 20% ($5.53-6.61 \times 10^{-5}$). In the 150% gap closure scenario, the impacts were reversed, with impacts declining by almost exactly the same percentage.

(b) Overlap and adjacency from optimal individual solutions

To identify the minimum possible costs for each individual objective, and the degree to which their areas of new cropland correspond, we examined simulations giving 100% of the



Figure 1. Areas converted to new maize and soya bean croplands and their associated costs when each individual objective (yield maximization, and carbon cost, biodiversity cost and transport time minimization) is given 100% weight. Cropland converted under each scenario is shown in the maps, colour-coded according to which objective receives 100% weight. Large bar charts, colour-coded to the maps, display the absolute value of costs resulting from each scenario. Small bar charts show the percentage of overlap (top) and adjacency (bottom; nearest neighbour distance) between the map directly above it, and each of the other three maps.

weight to each of the four different objectives. Figure 1 maps the results and shows the costs for each objective in each simulation, as well as the extent of overlap among objectives.

The scenario designed solely to maximize yield (100% weight) converts land with the most promising yield potential for farmers, which should have economic advantages. These areas have little direct overlap with the areas picked by scenarios that minimize carbon costs, biodiversity impacts or travel time (figure 1). They also have little adjacency, as measured by the average distance between each pixel of one map and its nearest neighbour in the other (figure 1).

The areas selected solely for minimizing carbon, biodiversity or transportation costs also have little direct overlap, but the map reveals that they have high adjacency, confirmed by direct estimates of distance (figure 1). All three criteria tend to identify lands along existing road networks and population and agricultural centres, with the carbon minimizing objective converting a larger area of land in the western half of the country. The reasons for this correlation are probably that carbon stocks and biodiversity values are lowest in areas of high human densities, which have correspondingly high levels of wood harvesting for charcoal and fuelwood, and higher levels of habitat loss and fragmentation leading to lower biodiversity scores.

(c) Potential to harmonize different interests

Although the ideal solutions to maximize each objective do not lead to much direct overlap, areas selected by less ideal but still good solutions for each objective may still overlap. The reasons start with the different potential ranges of costs among the scenarios that maximize each objective.

While the best-case scenario for maximizing yield would convert 13 668 km², the worst-case scenario for yields among our single-purpose optimization scenarios would convert 15 302 km² (the carbon-optimizing scenario). This range in converted areas is only 12%. This difference is far lower than

the range in yields across the country's potential cropland $(3.7-6.1 \text{ t/ha/y ha}^{-1} \text{ for maize and } 0.8-5.1 \text{ t/ha/y ha}^{-1} \text{ for soya bean}$, but yield plays such a significant role for each objective (because it is always in the denominator) that the model selects land with better yield potential even when the yield maximization priority is given no weight. By contrast, the differences in travel times among single-objective scenarios range from a total of $3.4-32.3 \times 10^{-5}$ truck hours, a difference of 872%. Carbon costs differ by 156% (462–1184 Mt), and biodiversity by 68% (4.96–8.33 × 10⁻⁵). These results indicate that there is a strong potential for reducing transportation, carbon and biodiversity costs with only a small increase in additional land area needed to meet production targets. In other words, these cost savings can be achieved for relatively little sacrifice of potential yield, which is inversely related to converted area.

This potential is illustrated in figure 2, which plots the 'efficiency frontier' [47] resulting from the range of possible trade-offs between the yield maximization and carbon protection objectives. Each point represents the most efficient solution, in terms of lowest possible carbon loss and land area converted, for meeting the crop production targets under the given combination of weights. Colour-coded maps indicate the associated cropland conversions for five of the 21 weighting combinations. Adding just 5% weight to carbon protection (orange scenario) results in 31% less carbon loss but less than 1% additional converted area relative to the pure yield maximization objective (red scenario), without significantly altering the location of the new cropland (adjacency = 7.4 km). Transferring 25% weight to carbon (yellow scenario) substantially shifts the conversion map (adjacency to pure yield maximization map = 65 km), but results in 54% less carbon loss. These savings nearly equal the 61% reduction in carbon loss achieved under the pure carbon protection scenario (dark blue).

These last examples illustrate that trade-offs can be usefully assessed in terms of avoided cost, or the per cent difference between the cost paid under a given weight



Figure 2. The costs of the optimal (most efficient) trade-offs between the yield maximization and carbon protection objectives under varying combinations of weights (varied in 5% increments). Costs are measured in total carbon loss and converted area. The conversion areas resulting from a subset of the weight combinations are illustrated in the accompanying maps and are colour-coded to their corresponding costs on the scatter plot, which represents the efficiency frontier. The weightings (expressed as proportions) associated with this subset of scenarios are also provided: Y = yield maximization objective; C = carbon protection objective.

combination and the maximum cost generated by all possible weighting scenarios. By expressing avoided cost as a percentage, we can directly compare the impacts of trade-offs between more than two objectives. Figure 3 shows how much carbon, biodiversity and transport cost can be avoided by transferring increasing amounts of weight away from the yield maximization objective. Sequentially adding weights in 5% increments to both the biodiversity and carbon objectives results in little cost in land area, but rapid, large reductions in both biodiversity and carbon costs. It also results in substantial avoided transport costs, even without assigning any weight to this objective, because of correlations between savings in transport, carbon, and biodiversity. For example, adding just 15% weight each to carbon and biodiversity increases the area of new lands (i.e. reduces the average yield) by just 2.7% (avoided area costs drop to 8% from 10.7%), but avoids potential costs equal to 47% for carbon, 27% for biodiversity and 44% for transportation. For comparison, in the best-case scenario for each objective, 61%, 41% and 90% of carbon, biodiversity and transportation cost would be avoided, respectively.

Figure 4 shows the potential for compromise in which a hypothetical land user only interested in a single objective is willing to 'pay' 5% more costs than the lowest possible costs for that objective. For example, the yield-maximizing user would be willing to pay for an extra 683 km² of new cropland, while the person interested in minimizing carbon costs would pay an additional 231 Mt of carbon. Lines in the bar charts



Figure 3. The impact of transferring weights from the yield maximization objective to the carbon and biodiversity objectives in 5% increments, measured in terms of the per cent cost that is avoided in each objective relative to the highest cost paid across all weight combinations. In these results, no weight was assigned to the transport cost objective.

show the minimum and maximum costs across all possible weighting scenarios. Each of the four different 'compromise' scenarios, represented by a different colour, permits solutions that allow the other scenarios to save substantially relative to



Figure 4. The results of scenarios in which each land use objective is willing to pay 5% more cost to compromise with the other three objectives. Cropland converted under each compromise scenario is shown in the maps, colour-coded according to which objective is making the 5% compromise. Large bar charts, colour-coded to the maps, display the absolute value of costs resulting from each compromise scenario, together with the maximum, minimum and median costs across all weighting permutations (grey horizontal lines in bar charts). Small bar charts show the percentage of overlap (top) and adjacency (bottom; nearest neighbour distance) between the map directly above it, and each of the other three maps, with the colour denoting which map is being compared.

maximum potential costs. The carbon, biodiversity and transport cost objectives in all but two cases avoid more than 50% of their maximum potential costs when any of the other objectives is willing to pay 5% more cost (figure 4).

The location of the conversions associated with each of these compromises show substantial convergence along roads and population and agricultural centres. Overlap between the resulting cropland allocations exceeds 30% between the yield and biodiversity and cost compromises, and 25% between the biodiversity and carbon compromises. The carbon and cost compromises overlap least (less than 10%), followed by carbon and yield (approx. 15%), but each pair of maps has high adjacency, with all compromise allocations having an average nearest neighbour distance of less than 20 km.

We also analysed an 'equal compromise' scenario in which each objective receives equal weight, as shown in figure 5. The bar charts show how this scenario, represented in purple, compares with each of the 5% compromise scenarios shown in figure 4, plus the best and worst case options among all weighting scenarios. Relative to the worst case, equal compromise avoids 80% or more of the cost for carbon, biodiversity and transportation, and 41% of the cost for land area. The equal weighting scenario compares favourably with the approximately 95% savings achieved by carbon, biodiversity and transportation costs within their individual cost categories, and to the 58% savings by the yield objective.

As a general rule, the areas selected for conversion under equal compromise lie along existing major transportation corridors, and generally near areas of existing cropland.

All of these scenarios also suggest the value of deliberate and optimized land use planning. Zambia has designated a large number of farm blocks, and added nine more blocks with roughly one million hectares as recently as 2005 [48]. Although these areas have as a whole attracted relatively little development, there is continued interest in developing them [49]. But there is less than 2% overlap between designated farm blocks and either the yield maximization scenario and the equal weighting scenario, and very little adjacency (see electronic supplementary material, results).

4. Discussion

Our results for Zambia suggest at least the potential to plan agricultural expansion such that it limits carbon and biodiversity costs with limited sacrifice of yield potential. In fact, if agricultural production goals alone are best represented by a combination of agronomic potential and existing road access, then solutions that provide a good balance of these objectives alone also correlate strongly with solutions that limit carbon and biodiversity impacts.

This finding demonstrates the potential benefits of targeting land use based on their ability to deliver a particular benefit for the lowest possible cost. The value of such an efficiency-based approach, which we apply here to agricultural land use, is in line with findings from the field of conservation planning, where it has been shown that conservation outcomes can be achieved for substantially lower cost when both the economic and ecological values of land are considered (e.g. [17,47]).

Another result, if more obvious, is the potential benefits of increasing yields on existing cropland if used to spare land. Actions that boost yields on existing cropland also have the potential to have feedback effects that lead to more local expansion, for example, by boosting the competitiveness and therefore quantity of exports [50,51]. Our analysis ignores those potential feedback effects because all options analysed meet the same production targets, but shows how yield gains could be used to limit environmental impacts.

The fact that the most advantageous compromises occur along existing transportation corridors also appears to have several advantages. Road construction has been a primary

8



Figure 5. Areas converted to new maize and soya bean croplands and their associated costs when all objectives (yield maximization, and carbon, biodiversity and transport cost minimization) compromise equally (i.e. receive 25% weight). Cropland converted under this scenario is denoted by purple areas in the map, with its associated cost impacts shown alongside those from the 5% compromise scenarios (figure 4), together with the maximum, minimum and median costs across all weighting permutations (grey horizontal lines in bar charts).

driver of the location of agricultural expansion [52]. Zambia is already engaged in a programme to upgrade existing major roads, but there is some evidence in other parts of Africa that agriculture responds even more to improvements in the quality of smaller, rural roads than to improvements in larger paved roadways [53,54]. Improving rural roads would be one way of focusing Zambia's agriculture development on existing road networks. Because agricultural development tends to build on the private and public infrastructure that accompanies initial development, the focus on existing areas would also seem likely to be a more robust strategy for avoiding carbon and biodiversity loss over the long-term [35].

Beyond road infrastructure, our results also indicate the importance of assessing current agricultural development plans because of the low overlap between Zambia's agricultural development blocks and the areas selected for any objective.

We believe this type of model would be particularly useful for planning agricultural development in any politically heterogeneous environment. Optimization models that require *ex ante* specification of preferences in mathematical terms work less well for decision-making that must weigh very different objectives in unclear ways. Our model makes an iterative optimization process possible, which allows people to realize their preferences in the face of real information about options. It is particularly suitable for making decisions that will inevitably have strong political elements, and to consensus-building among stakeholders with different preferences.

For this kind of tool to be truly legitimate and useful on a practical level, we believe several elements are required. First, it should be further developed and applied in an iterative way with the government and other stakeholders so that the costs and benefits of different weighting decisions are fully explored. We are developing plans to work with governments, various stakeholders, and international collaborators to further develop and use this model as an operational decision-support tool in Zambia and other countries in the region. Second, development planning must reflect and incorporate other real-world factors beyond those included in our model. For example, large, multi-lane roads can lead to high rates of wildlife mortality, and serve as effective barriers to migration for large mammals, regardless of other land uses [55]. Although our 'equal weight' scenario identifies some target areas in the heavily wooded northeastern part of the country, these concerns would be one reason to explore whether areas closer to existing cropland concentrations in the south might be reasonable substitutes. Such consolidation might also make it easier to support agricultural development with lower attendant infrastructure development costs. And this biodiversity concern is only one example of other issues, many purely practical, which should, and inevitably will, influence where agriculture develops.

Third, data quality is critical. This kind of model focuses on particular hectares, not broad averages, so data errors have little or no chance of averaging out through aggregation. Predicted potential yields and most of our other scores can be highly sensitive to errors in the inputs used to calculate them. These facts mean that long-term use of the model should be accompanied by steady data improvement. It also means that the model should mostly serve as a guide to likely values. Before governments make any important and potentially irreversible decisions, there should be site-specific ground-truthing of soil properties and yield potential, as well as confirmation of biodiversity and carbon characteristics.

Fourth, some elements of the model should also be further developed. Analysis of agricultural potential could incorporate more economic factors, differentiate between smallholder and commercial farming practices and incorporate irrigation potential (e.g. [23]). Additional information on land ownership and title would allow a finer delineation of potential agricultural land, while the carbon estimates would be improved by rootshoot ratios specific to each vegetation type. For biodiversity, defining rarity/threat in relation to national borders may provide a biased view of conservation importance, as habitats that are rare in Zambia might be common across the border, while a relatively common habitat in Zambia might be regionally rare. A regional approach to defining this measure might therefore be more appropriate (e.g. [45]). This index could also be adjusted to factor in additional indicators of biodiversity value, such as Important Bird Areas [56] or Key Biodiversity Areas [57].

Analysis of biodiversity impacts could also be improved by more direct quantification of biodiversity responses to land use. One potentially promising approach would be to incorporate the biodiversity impact model developed by Newbold *et al.* [42], which estimates changes in species diversity and abundance in response to landcover and land-use intensity. With sufficient data, this approach could also allow comparison of the biodiversity impacts of different agricultural practices, including land-sparing approaches that maintain trees, vegetated field borders, or other habitats within the agricultural landscape [58]. The model can be easily adjusted to simulate varying intensities of agricultural management.

The model could also be used to evaluate the temporal persistence of trade-offs in response to key uncertainties, such as the future effects of climate change on agricultural potential. Robustness to such uncertainty should be a key factor in determining new agricultural development areas. Assessment of robustness can be achieved by identifying stable areas of conversion that emerge over many iterations within the bounds of uncertainty, which is a feature of the model that we intend to develop. Despite the current limitations of the model and this initial analysis, our results are promising. They suggest that efficiency-based land use planning approaches such as ours, if operationalized and incorporated into decision-making processes, can help to substantially minimize the ecological cost of cropland expansion in sub-Saharan Africa's savannahs, the twenty-first century's emerging hotspot of agricultural land use change.

Data accessibility. The model source and inputs data are available online at https://github.com/PrincetonUniversity/agroEcoTradeoff.

Authors' contributions. L.D.E. and T.S. conceived and designed the research, together with K.K.C. L.D.E. and M.S. designed and coded the model. L.D.E. developed the input datasets, and ran and analysed simulations, together with T.S. D.T. and L.D.E. developed the crop yield potential surfaces. L.K. and T.K. helped design the biodiversity impact metric. S.S. and M.M. collected and processed spatial datasets used in developing inputs. A.B. processed infrastructure data and helped develop the travel cost surface. J.S., N.C. and E.F.W. developed the bias-corrected weather data needed to estimate yield potential. L.D.E. and T.S. wrote the manuscript, with input from K.K.C., T.K. and L.K.

Competing interests. We declare we have no competing interests.

Funding. This work was supported by funds from: the Norwegian Aid Agency (NORAD) under the Agricultural Synergies Project, the Princeton Environmental Institute Grand Challenges program, the NASA New Investigator Program (NNX15AC64G) and the National Science Foundation (SES-1360463 and SES-1534544). L.K. and T.K. gratefully acknowledge funding by the Einstein Foundation Berlin.

Acknowledgements. We thank Paulo van Breugel for providing the potential vegetation map of East Africa, and Philip Thornton for advice regarding modelling of yield potential.

References

- Searchinger TD, Estes L, Thornton PK, Beringer T, Notenbaert A, Rubenstein D, Heimlich R, Licker R, Herrero M. 2015 High carbon and biodiversity costs from converting Africa's wet savannahs to cropland. *Nat. Clim. Change* 5, 481–486. (doi:10.1038/nclimate2584)
- 2. Field CB *et al.* 2014 *Summary for policymakers*. Cambridge, UK: Cambridge University Press.
- Tilman D, Balzer C, Hill J, Befort BL. 2011 Global food demand and the sustainable intensification of agriculture. *Proc. Natl Acad. Sci. USA* **108**, 1091–6490. (doi:10.1073/pnas.1116437108)
- Morris M, Byerlee D. 2009 Awakening Africa's sleeping giant. Washington, DC: Technical report, World Bank and FAO.
- The World Bank. 2016 World development indicators. See http://data.worldbank.org/indicator/ SP.RUR.TOTL.ZS.
- 6. FAO. 2016 FAOStat: Production. See http://faostat. fao.org.
- International Food Policy Research Institute. 2015 Global Nutrition Report 2015: Actions and Accountability to Advance Nutrition and Sustainable Development. Washington, DC.
- United Nations. 2015 World population prospects: the 2015 revision. New York, NY: United Nations, Department of Economic and Social Affairs.
- Sexton JO *et al.* 2015 Conservation policy and the measurement of forests. *Nat. Clim. Change* 6, 192–196. (doi:10.1038/nclimate2816)

- Vinya R, Syampungani S, Kasumu EC, Monde C, Kasubika R. 2011 Preliminary study on the drivers of deforestation and potential for REDD+ in Zambia. Lusaka, Zambia: FAO/Zambian Ministry of Lands and Natural Resources.
- Olson DM *et al.* 2001 Terrestrial ecoregions of the world: a new map of life on Earth. *BioScience* **51**, 933 – 938. (doi:10.1641/0006-3568(2001)0510933: TEOTWA]2.0.C0;2)
- Hogan CM. 2004 Central Zambezian miombo woodland. The terrestrial ecoregions of Africa and Madagascar: a conservation assessment. Washington, DC: Island Press.
- Kier G, Mutke J, Dinerstein E, Ricketts TH, Küper W, Kreft H, Barthlott W. 2005 Global patterns of plant diversity and floristic knowledge. *J. Biogeogr.* 32, 1107 – 1116. (doi:10.1111/j.1365-2699.2005. 01272.x)
- Fund W. 2011 Ecoregions of Zambia (WWF). See http://www.eoearth.org/view/article/ 152238.
- Estes LD, Greyling L. 2004 Zambezian and mopane woodlands. The terrestrial ecoregions of Africa and Madagascar: a conservation assessment. Washington, DC: Island Press.
- Dowsett JR, Aspinwall RD, Dowsett-Lemaire F, Francoise H. 2008 *The birds of Zambia: an atlas and handbook*. Liège, Belgium: Tauraco & Aves.

- Naidoo R, Balmford A, Ferraro P, Polasky S, Ricketts T, Rouget M. 2006 Integrating economic costs into conservation planning. *Trends Ecol. Evol.* 21, 681–687. (doi:10.1016/j.tree.2006.10.003)
- Margules CR, Pressey RL. 2000 Systematic conservation planning. *Nature* 405, 243-253. (doi:10.1038/35012251)
- Ball IR, Possingham HP, Watts M. 2009 Marxan and relatives: software for spatial conservation prioritisation. In *Spatial conservation prioritisation: quantitative methods and computational tools*, pp. 185–195. Oxford, UK: Oxford University Press.
- Watts ME, Ball IR, Stewart RS, Klein CJ, Wilson K, Steinback C, Lourival R, Kircher L, Possingham HP. 2009 Marxan with zones: software for optimal conservation based land- and sea-use zoning. *Environ. Model. Softw.* 24, 1513–1521. (doi:10. 1016/j.envsoft.2009.06.005)
- Cao K, Batty M, Huang B, Liu Y, Yu L, Chen J. 2011 Spatial multi-objective land use optimization: extensions to the non-dominated sorting genetic algorithm-II. *Int. J. Geogr. Inform. Sci.* 25, 1949–1969. (doi:10.1080/13658816.2011.570269)
- Koh LP, Ghazoul J. 2010 Spatially explicit scenario analysis for reconciling agricultural expansion, forest protection, and carbon conservation in Indonesia. *Proc. Natl Acad. Sci. USA* **107**, 11 140–11 144. (doi:10.1073/pnas.1000530107)

10

- Nijbroek RP, Andelman SJ. 2016 Regional suitability for agricultural intensification: a spatial analysis of the Southern Agricultural Growth Corridor of Tanzania. *Int. J. Agric. Sustainability* 14, 231–247. (doi:10.1080/14735903.2015.1071548)
- Gasparri NI, Kuemmerle T, Meyfroidt P, le Polain de Waroux Y, Kreft H. 2015 The emerging soybean production frontier in Southern Africa: conservation challenges and the role of south-south telecouplings. *Conserv. Lett.* 9, 21–31. (doi:10. 1111/conl.12173)
- Estes LD, Paroz L-L, Bradley BA, Green JM, Hole DG, Holness S, Ziv G, Oppenheimer MG, Wilcove DS. 2014 Using changes in agricultural utility to quantify future climate-induced risk to conservation. *Conserv. Biol.* 28, 427–437. (doi:10.1111/cobi.12205)
- Jones J *et al.* 2003 Modelling cropping systems: science, software and applications. *Eur. J. Agron.* 18, 235–265. (doi:10.1016/S1161-0301(02)00107-7)
- Sheffield J, Goteti G, Wood EF. 2006 Development of a 50-year high-resolution global dataset of meteorological forcings for land surface modeling. J. Clim. 19, 3088–3111. (doi:10.1175/JCLI3790.1)
- Chaney NW, Sheffield J, Villarini G, Wood EF. 2014 Spatial analysis of trends in climatic extremes with a high resolution gridded daily meteorological data set over sub-saharan Africa. J. Clim. 27, 5815– 5835.
- Estes LD, Chaney NW, Herrera-Estrada J, Sheffield J, Caylor KK, Wood EF. 2014 Changing water availability during the African maize-growing season, 1979–2010. *Environ. Res. Lett.* 9, 075005. (doi:10.1088/1748-9326/9/7/075005)
- HarvestChoice. 2010 Converting WISE 1.1 Soil Profile Database for Crop Models. Technical report. Washington, DC: IFPRI.
- Grassini P et al. 2015 How good is good enough? Data requirements for reliable crop yield simulations and yield-gap analysis. *Field Crops Res.* 177, 49-63. (doi:10.1016/j.fcr.2015.03.004)
- 32. Wood SN. 2001 mgcv: GAMs and generalized ridge regression for R. *R News* **1**, 20–25.
- Hengl T *et al.* 2014 SoilGrids 1 km—global soil information based on automated mapping. *PLoS ONE* 9, e105992. (doi:10.1371/journal.pone. 0105992)
- Stifel D, Minten B. 2008 Isolation and agricultural productivity. *Agric. Econ.* **39**, 1–15. (doi:10.1111/j. 1574-0862.2008.00310.x)

- Laurance WF, Sloan S, Weng L, Sayer JA. 2015 Estimating the environmental costs of africa's massive 'development corridors'. *Curr. Biol.* 25, 3202–3208. (doi:10.1016/j.cub.2015.10.046)
- CIESIN C. 2005 Gridded population of the world version 3 (GPWV3): population density grids.
 Palisades, NY: Socioeconomic Data and Applications Center (SEDAC), Columbia University.
- Lehner B, Verdin K, Jarvis A. 2008 New global hydrography derived from spaceborne elevation data. *Eos, Trans. Am. Geophys. Union* 89, 93-94. (doi:10.1029/2008E0100001)
- Baccini A *et al.* 2012 Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps. *Nat. Clim. Change* 2, 182–185. (doi:10.1038/nclimate1354)
- Mokany K, Raison RJ, Prokushkin AS. 2006 Critical analysis of root: shoot ratios in terrestrial biomes. *Glob. Change Biol.* 12, 84–96. (doi:10.1111/j.1365-2486.2005.001043.x)
- Mugasha WA, Eid T, Bollandsås OM, Malimbwi RE, Chamshama SAO, Zahabu E, Katani JZ. 2013 Allometric models for prediction of above- and belowground biomass of trees in the miombo woodlands of Tanzania. *Forest Ecol. Manage.* **310**, 87–101. (doi:10.1016/j.foreco.2013.08.003)
- 41. IPCC. 1996 Land use change and forestry. In *Revised* 1996 IPCC guidelines for national greenhouse gas inventories: reference manual. Geneva, Switzerland: Intergovernmental Panel on Climate Change.
- Newbold T *et al.* 2015 Global effects of land use on local terrestrial biodiversity. *Nature* 520, 45–50. (doi:10.1038/nature14324)
- Kehoe L, Kuemmerle T, Meyer C, Levers C, Václavk T, Kreft H. 2015 Global patterns of agricultural landuse intensity and vertebrate diversity. *Divers. Distrib.* 21, 1308–1318. (doi:10.1111/ddi.12359)
- van Breugel P *et al.* 2011 Potential natural vegetation of eastern Africa (Ethiopia, Kenya, Malawi, Rwanda, Tanzania, Uganda and Zambia), Volume 6: an overview of the methods and material used to develop the map. Forest & Landscape Working Paper 68-2011.
- van Breugel P, Kindt R, Lillesø J-PB, van Breugel M. 2015 Environmental gap analysis to prioritize conservation efforts in eastern Africa. *PLoS ONE* **10**, e0121444. (doi:10.1371/journal.pone.0121444)
- 46. Waddington SR, Li X, Dixon J, Hyman G, de Vicente MC. 2010 Getting the focus right: production

constraints for six major food crops in Asian and African farming systems. *Food Security* **2**, 27–48. (doi:10.1007/s12571-010-0053-8)

- Polasky S *et al.* 2008 Where to put things? Spatial land management to sustain biodiversity and economic returns. *Biol. Conserv.* 141, 1505–1524. (doi:10.1016/j.biocon.2008.03.022)
- Anonymous. 2005 Farm development block plan 2005 – 2007. Technical report. Lusaka, Zambia: Ministry of Finance and National Planning, Republic of Zambia.
- Anonymous. 2015 Undeveloped farm blocks worry Agriculture minister. *Zambia Daily Mail*. https:// www.daily-mail.co.zm/?p=29505.
- Ewers RM, Scharlemann JPW, Balmford A, Green RE. 2009 Do increases in agricultural yield spare land for nature? *Glob. Change Biol.* **15**, 1716–1726. (doi:10.1111/J.1365-2486.2009.01849.X)
- Angelsen A. 2010 Policies for reduced deforestation and their impact on agricultural production. *Proc. Natl Acad. Sci. USA* **107**, 19 639 – 19 644. (doi:10. 1073/pnas.0912014107)
- Laurance WF, Goosem M, Laurance SGW. 2009 Impacts of roads and linear clearings on tropical forests. *Trends Ecol. Evol.* 24, 659–669. (doi:10. 1016/j.tree.2009.06.009)
- Pinstrup-Andersen P, Shimokawa S. 2008 Rural infrastructure and agricultural development. In *Rethinking infrastructure for development* (eds F Bourguignon, B Pleskovic). World Bank Publications.
- Blimpo MP, Harding R, Wantchekon L. 2013 Public investment in rural infrastructure: some political economy considerations. *J. Afr. Econ.* 22, 1464–3723. (doi:10.1093/jae/ejt015)
- Holdo RM, Fryxell JM, Sinclair ARE, Dobson A, Holt RD. 2011 Predicted impact of barriers to migration on the Serengeti Wildebeest population. *PLoS ONE* 6, e16370. (doi:10.1371/journal.pone.0016370)
- Butchart SHM *et al.* 2012 Protecting important sites for biodiversity contributes to meeting global conservation targets. *PLoS ONE* 7, e32529. (doi:10. 1371/journal.pone.0032529)
- Eken G *et al.* 2004 Key biodiversity areas as site conservation targets. *BioScience* 54, 1110–1118. (doi:10.1641/0006-3568(2004)0541110:KBAASC]2.0. (0;2)
- Kremen C. 2015 reframing the land-sparing/landsharing debate for biodiversity conservation: reframing the land-sparing/land-sharing debate. *Annu. N. Y. Acad. Sci.* **1355**, 52–76. (doi:10.1111/nyas.12845)