

UCSF

UC San Francisco Previously Published Works

Title

Rho GEFs and GAPs: Emerging integrators of extracellular matrix signaling

Permalink

<https://escholarship.org/uc/item/3p75m7g7>

Journal

Small GTPases, 6(1)

ISSN

2154-1248

Authors

Kutys, Matthew L
Yamada, Kenneth M

Publication Date

2015-01-02

DOI

10.4161/21541248.2014.989792

Peer reviewed

Rho GEFs and GAPs: Emerging integrators of extracellular matrix signaling

Matthew L Kutys* and Kenneth M Yamada*

Laboratory of Cell and Developmental Biology; National Institute of Dental and Craniofacial Research; National Institutes of Health

Investigating cell migration in 3D settings has revealed that specific extracellular matrix environments require differential activities of the Rho GTPases for efficient migration. However, it is largely unknown how the activities of specific Rho GTPases are modulated to direct cell migration in response to different extracellular matrix cues. We have recently reported that extracellular matrix-dependent regulation of a specific Rho GEF is a fundamental mechanism governing cell migration in different microenvironments, providing a direct mechanism for extracellular matrix-specific regulation of Rho GTPase activity directing cell motility. We discovered that the Rho GEF β Pix has a unique function during cell migration in fibrillar collagen environments by restraining RhoA signaling through a conserved signaling axis involving Cdc42 and the Rho GAP srGAP1. In this Commentary, we expand upon this new pathway and discuss potential mechanotransductive and therapeutic applications. Additionally, we speculate on a generalized role for Rho GEFs and GAPs in providing localized, context-dependent responses to the cellular microenvironment during cell migration and other cellular processes.

During cell migration, cells utilize cell-matrix adhesion in concert with dynamic cytoskeletal rearrangements and myosin II-generated contractility to sense and respond to physical and chemical cues from the extracellular matrix (ECM).^{1–3} Cellular responses to these extracellular signals are frequently orchestrated through the activity and function of the Rho family GTPases Cdc42, Rac1, and RhoA, as well as their guanine nucleotide exchange

factor (GEF) and GTPase activating protein (GAP) regulators. However, recent investigations in 3D ECM and observations of the plasticity of 3D cell migratory modes have revealed that a cell can exhibit variable dependence on the activity of a particular Rho GTPase, which can be dictated by distinct features of the composition and structure of the surrounding ECM.^{4–6} Two key contemporary questions are: What are the major extracellular determinants of cell migration in different 3D ECMs, and how are these distinct environments able to elicit changes in Rho GTPase activity?

Recently, we initiated a study with the hypothesis that cell adhesion to different matrix molecules, such as collagen and fibronectin, would trigger differential regulation of guanine nucleotide exchange factors to regulate migration.⁷ To address this question, we developed an unbiased, ECM-based affinity purification screen to isolate and identify novel GEFs uniquely active under collagen or fibronectin-rich ECM conditions.^{7,8} We discovered that the GEF β Pix has a conserved, extracellular matrix-specific migratory function and is critical for efficient migration in fibrillar collagen environments by restraining the signaling activity of RhoA. Unexpectedly, this suppression occurs through a mechanism of Rho GTPase crosstalk between Cdc42 and RhoA that is mediated by a collagen-specific interaction between the GEF/GAP pair, β Pix and srGAP1.

In addition, our model (Fig. 1) suggested that the collagen-specific β Pix function is dictated by tight phosphorylation of threonine residue 526 (T526) on β Pix. Binding of $\alpha_2\beta_1$ integrin to fibrillar collagen leads, through PP2A, to loss of phosphorylation at T526 on β Pix

Keywords: extracellular matrix, rho GEF, rho GAP, rho GTPase, 3D migration

Abbreviations: GEF, Guanine nucleotide exchange factor; GAP, GTPase activating protein; ECM, extracellular matrix; T526, Threonine 526; FRET, Fluorescence resonance energy transfer; HUVEC, Human umbilical cord vein endothelial cell.

This article not subject to US copyright law.

*Correspondence to: Matthew L. Kutys; Email: mkutys10@gmail.com; Kenneth M. Yamada; Email: kyamada@mail.nih.gov

Submitted: 10/31/2014

Revised: 11/12/2014

Accepted: 11/14/2014

<http://dx.doi.org/10.4161/21541248.2014.989792>




Figure 1. Summary model of the extracellular matrix-specific regulation and function of β Pix during cell migration. During migration in fibronectin-rich environments, β Pix is phosphorylated at threonine 526 and localizes to focal adhesions through associations with GIT1 and paxillin. Conversely, binding of $\alpha_2\beta_1$ to fibrillar collagen leads, through PP2A, to loss of phosphorylation at T526 on β Pix, β Pix localization to the plasma membrane, and associations with srGAP1 and Cdc42. β Pix/Cdc42/srGAP1 act to locally suppress RhoA activity at the leading edge of the cell, establishing inversely polarized gradients of Cdc42 and RhoA activity that are required for efficient cell migration in fibrillar collagen environments.

and promotes association with srGAP1. Taken together, we defined a conserved migratory signaling cascade involving PP2A/ β Pix/srGAP1 that coordinates suppressive crosstalk between Cdc42 and RhoA and is critical for cell migration in fibrillar collagen environments.

The existence of an extracellular matrix-specific pathway directing cell migration may at first seem puzzling physiologically, considering the heterogeneous nature of most environments found *in vivo*.⁹⁻¹¹ However, this mechanism provides an additional tier of control over the plasticity of cell migratory behavior in response to cues from each cell's extracellular environment. It provides the capacity for local changes in extracellular matrix composition and organization to elicit precise spatial control over the movement of cells during tightly regulated processes such as epithelial morphogenesis, angiogenesis, and wound healing. For example, gradients of type I collagen fibers have been reported to be sufficient to drive cytokine-independent endothelial morphogenesis and migration.¹² In fact, the increased angiogenic migratory response downstream of endothelial cell adhesion to fibrillar collagen was attributed to suppression of PKA activity, a reported kinase for T526 on β Pix.^{12,13} In our recent study, we observed that the collagen-

specific function of β Pix was indeed conserved in human umbilical vein endothelial cells (HUVECs). This suggests a potential anti-angiogenic application for the inhibition of β Pix that is dependent on the ECM context.

Another exciting potential application can be surmised from recent concepts concerning the role of fibrillar type I collagen during tumor progression and metastasis. The linearization and perpendicular reorganization of fibrillar type I collagen at the tumor front is a marker of malignant transformation and metastatic potential.¹⁴⁻¹⁶ In certain settings, these fibrillar collagen "tracks" serve as metastatic highways for transformed cells, facilitating dissemination away from the primary tumor.^{17,18} We observed that loss of β Pix in the metastatic breast adenocarcinoma MDA-MB-231 line blocks nearly all motility of these normally motile cells in 3D collagen, with no perturbation of their migration in fibronectin-rich 3D cell-derived matrices. Therefore, inhibition of β Pix may have potential as an anti-metastatic therapeutic, particularly in collagen-dense tumor environments, such as the skin, breast, and pancreas,^{19,20} by preventing primary tumor dissemination along collagen fibrils.

One intriguing element of this study is the identification of a pathway of active

RhoA suppression, a Rho GTPase classically associated with controlling the contractile or mechanotransductive response of the cell to the microenvironment.^{21,22} Traditionally, when we consider cell mechanotransduction, tension-sensitive proteins are thought to be activated through conformational changes in response to increased rigidity or other physical inputs from the surrounding microenvironment.^{23,24} The newly identified collagen-specific β Pix pathway of Cdc42-RhoA crosstalk surprisingly contrasts with this canonical view, in that the activity of RhoA is inherently high, yet is actively suppressed by β Pix/Cdc42/srGAP1 in response to fibrillar collagen. It is not yet clear which environmental factors may be influencing RhoA activity, whether it is induced by growth-factor signaling from serum, cell-matrix adhesion between $\alpha_2\beta_1$ integrin and fibrillar collagen, or a physical aspect of the fibrillar collagen matrix itself. However, this pathway potentially presents an alternative to the classical view of mechanotransduction, i.e., a tunable RhoA responsive mechanism that adapts to changes in collagen matrix rigidity. In essence, increases in matrix rigidity could modulate the phosphorylation of β Pix at T526, which would decrease the activity of Cdc42/srGAP1, remove the suppressive check on RhoA, and tailor appropriate local levels of contractile force required for efficient migration. Preliminary observations from our laboratory support this hypothesis: increasing the rigidity of 2D fibrillar collagen matrices through crosslinking is able to partially rescue the morphology and migratory defects of β Pix knockdown. This observation not only delineates an alternative to traditional mechanotransduction views, but akin to the concept of distinct migratory signaling mechanisms, highlights the importance of elucidating mechanotransduction pathways that are specific to different cell-surface receptors and ECM environments.

It is surprising that no single GEF was identified as specifically active in response to fibronectin. Live-cell timelapse imaging and immunohistochemistry revealed that fibroblasts in 3D collagen and 3D cell-derived matrices migrate with no obvious differences in cell morphology and

polarization, leading edge/adhesive dynamics, or cytoskeletal assembly⁷ (Fig. 2). Additionally, we observed through Cdc42 and RhoA live-cell FRET imaging that fibronectin is able to promote migration through similar mechanisms as fibrillar collagen, yet loss of β Pix is not detrimental to migration in this condition. This finding highlights the importance of identifying matrix-specific migratory pathways and may indicate that the migratory machinery activated in response to fibronectin is driven by a distinct GEF/GAP mechanism involving Cdc42, RhoA, non-canonical GTPases such as RhoC/E/G and Rac2/3, or through some complex concerted action of multiple GTPases. This view is further supported by the observation that while srGAP1 directs GAP activity toward RhoA specifically during fibrillar collagen migration, we observed that loss of srGAP1 leads to increased Rac1 activity during migration on fibronectin. We speculate that this and other potential extracellular matrix-specific GEF/GAP mechanisms will emerge that provide local contextual regulation of cell migration in different microenvironments.

How the cell is able to regulate signaling spatially and temporally across many aspects of cell physiology is currently a major area of investigation. When one considers that the approximate 80 Rho GEFs and 70 Rho GAPs in the human genome outnumber their target GTPases by 3- to 4-fold and that many GEF/GAPs control the same processes,²⁵⁻²⁷ it becomes clear that this apparent redundancy raises questions about the larger role served by these proteins. An emerging paradigm suggests the explanation that Rho GEFs and GAPs serve to specifically transduce the diverse number of extracellular matrix signals to the cell by restricting their exchange/hydrolysis activity to appropriate subcellular locations and the associated downstream targets of the extracellular stimuli.^{25,28} Along with our recent discovery of the collagen-specific roles of β Pix/srGAP1 during cell migration, other examples of contextual GEF activity dictating migratory responses include the role of the RhoA GEF PDZ-RhoGEF during breast cancer motility in response to the chemokine CXCL12²⁹ and the

Rac1 GEF Tiam1 during keratinocyte spreading and migration downstream of $\alpha_3\beta_1$ adhesion to laminin-5.³⁰

Evidence also exists for ECM regulation of GEFs/GAPs in other, non-migration related processes. The GEF Sos regulates cell cycle progression through Rac1 in endothelial cells through $\alpha_5\beta_1$ attachment to fibronectin, but not laminin or charge-mediated attachments.^{31,32} Application of force on different integrin subtypes has revealed that the RhoA GEFs GEF-H1 and LARG are critical for mediating cell-matrix mechanotransduction and the cell stiffening response in collagen and fibronectin environments, respectively.^{33,34} Considering the diverse range of roles played by the Rho GTPases in regulating cell physiology, the complexities of the ECM-responsive GEF/GAP system could be quite vast. Therefore, it is clear that it will be important not only to delin-

eat the molecular functions of the many uncharacterized GEFs and GAPs, but also to elucidate the precise contexts in which they are physiologically relevant.

There are many complex interactions and forms of signaling crosstalk that occur at the leading edge of cells during migration, and the β Pix/srGAP1 complex provides an elegant mechanism for restricting RhoA and concentrating Cdc42 activity toward the leading edge in collagen microenvironments. It will be of interest to expand this approach to GEFs and GAPs toward other Rho family GTPases and different cellular processes to determine whether modulation of GEF/GAP activity directs cell physiology in response to specific chemical and physical properties of the extracellular matrix. By doing so, we will gain a greater understanding of how cells respond to, and navigate within, the complex extracellular matrix environment




Figure 2. Primary human foreskin fibroblasts immunostained for actin (green, phalloidin) during migration in 3D collagen matrix (left, magenta collagen fibers) and 3D cell-derived matrix (right, magenta fibronectin fibers). Cells in both 3D matrix environments can migrate with similar morphologies and dynamics, yet a requirement for β Pix/Cdc42/srGAP1 signaling does not exist in the fibronectin-rich cell-derived matrix. This highlights the importance of understanding how cells respond to diverse ECM environments and appropriately regulate their migratory signaling. Nuclei (blue) visualized by DAPI.

found *in vivo* and potentially how to intervene when these processes become deregulated in disease.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Funding

Supported by the Intramural Research Program of the NIDCR, NIH.

References

- Doyle AD, Petrie RJ, Kutys ML, Yamada KM. Dimensions in cell migration. *Curr Opin Cell Biol* 2013; 25:642-9; PMID:23850350; <http://dx.doi.org/10.1016/j.ccb.2013.06.004>
- Baker BM, Chen CS. Deconstructing the third dimension: how 3D culture microenvironments alter cellular cues. *J Cell Sci* 2012; 125:3015-24; PMID:22797912; <http://dx.doi.org/10.1242/jcs.079509>
- Petrie RJ, Yamada KM. At the leading edge of three-dimensional cell migration. *J Cell Sci* 2012; 125:5917-26; PMID:23378019; <http://dx.doi.org/10.1242/jcs.093732>
- Petrie RJ, Gavara N, Chadwick RS, Yamada KM. Non-polarized signaling reveals two distinct modes of 3D cell migration. *J Cell Biol* 2012; 197:439-55; PMID:22547408; <http://dx.doi.org/10.1083/jcb.201201124>
- Sahai E, Marshall CJ. Differing modes of tumour cell invasion have distinct requirements for rho/ROCK signalling and extracellular proteolysis. *Nat Cell Biol* 2003; 5:711-9; PMID:12844144; <http://dx.doi.org/10.1038/ncb1019>
- Wolf K, Te Lindert M, Krause M, Alexander S, Te Riet J, Willis AL, Hoffman RM, Figdor CG, Weiss SJ, Friedl P. Physical limits of cell migration: control by ECM space and nuclear deformation and tuning by proteolysis and traction force. *J Cell Biol* 2013; 201:1069-84; PMID:23798731; <http://dx.doi.org/10.1083/jcb.201210152>
- Kutys ML, Yamada KM. An extracellular-matrix-specific GEF-GAP interaction regulates rho GTPase crosstalk for 3D collagen migration. *Nat Cell Biol* 2014; 16:909-17; PMID:25150978; <http://dx.doi.org/10.1038/ncb3026>
- Garcia-Mata R, Wennerberg K, Arthur WT, Noren NK, Ellerbrog SM, Burridge K. Analysis of activated GAPs and GEFs in cell lysates. *Methods Enzymol* 2006; 406:425-37; PMID:16472675; [http://dx.doi.org/10.1016/S0076-6879\(06\)06031-9](http://dx.doi.org/10.1016/S0076-6879(06)06031-9)
- Kutys ML, Doyle AD, Yamada KM. Regulation of cell adhesion and migration by cell-derived matrices. *Exp Cell Res* 2013; 319:2434-9; PMID:23751565; <http://dx.doi.org/10.1016/j.yexcr.2013.05.030>
- Frantz C, Stewart KM, Weaver VM. The extracellular matrix at a glance. *J Cell Sci* 2010; 123:4195-200; PMID:21123617; <http://dx.doi.org/10.1242/jcs.023820>
- Byron A, Humphries JD, Humphries MJ. Defining the extracellular matrix using proteomics. *Int J Exp Pathol* 2013; PMID:23419153
- Whelan MC, Senger DR. Collagen I initiates endothelial cell morphogenesis by inducing actin polymerization through suppression of cyclic AMP and protein kinase A. *J Biol Chem* 2003; 278:327-34; PMID:12399469; <http://dx.doi.org/10.1074/jbc.M20754200>
- Howe AK. Regulation of actin-based cell migration by cAMP/PKA. *Biochim Biophys Acta* 2004; 1692:159-74; PMID:15246685; <http://dx.doi.org/10.1016/j.bbamcr.2004.03.005>
- Provenzano PP, Eliceiri KW, Campbell JM, Inman DR, White JG, Keely PJ. Collagen reorganization at the tumor-stromal interface facilitates local invasion. *BMC Med* 2006; 4:38; PMID:17190588; <http://dx.doi.org/10.1186/1741-7015-4-38>
- Levental KR, Yu H, Kass L, Lakins JN, Egeblad M, Erler JT, Fong SF, Csizsar K, Giaccia A, Weninger W, et al. Matrix crosslinking forces tumor progression by enhancing integrin signaling. *Cell* 2009; 139:891-906; PMID:19931152; <http://dx.doi.org/10.1016/j.cell.2009.10.027>
- Provenzano PP, Inman DR, Eliceiri KW, Knittel JG, Yan L, Rueden CT, White JG, Keely PJ. Collagen density promotes mammary tumor initiation and progression. *BMC Med* 2008; 6:11; PMID:18442412; <http://dx.doi.org/10.1186/1741-7015-6-11>
- Condeelis JS, Pollard JW. Macrophages: obligate partners for tumor cell migration, invasion, and metastasis. *Cell* 2006; 124:263-6; PMID:16439202; <http://dx.doi.org/10.1016/j.cell.2006.01.007>
- Shi Q, Ghosh RP, Engelke H, Rycroft CH, Cassereau L, Sethian JA, Weaver VM, Liphardt JT. Rapid disorganization of mechanically interacting systems of mammary acini. *Pro Natl Acad Sci U S A* 2014; 111:658-63; PMID:24379367; <http://dx.doi.org/10.1073/pnas.1311312110>
- Zhang K, Corsa CA, Ponik SM, Prior JL, Pivnicka-Worms D, Eliceiri KW, Keely PJ, Longmore GD. The collagen receptor discoidin domain receptor 2 stabilizes SNAIL1 to facilitate breast cancer metastasis. *Nat Cell Biol* 2013; 15:677-87; PMID:23644467; <http://dx.doi.org/10.1038/ncb2743>
- Shintani Y, Hollingsworth MA, Wheelock MJ, Johnson KR. Collagen I promotes metastasis in pancreatic cancer by activating c-Jun NH(2)-terminal kinase 1 and up-regulating N-cadherin expression. *Cancer Res* 2006; 66:11745-53; PMID:17178870; <http://dx.doi.org/10.1158/0008-5472.CAN-06-2322>
- Marjoram RJ, Lessey EC, Burridge K. Regulation of rhoa activity by adhesion molecules and mechanotransduction. *Curr Mol Med* 2014; 14:199-208; PMID:24467208; <http://dx.doi.org/10.2174/1566524014666140128104541>
- Chrzanoska-Wodnicka M, Burridge K. Rho-stimulated contractility drives the formation of stress fibers and focal adhesions. *J Cell Biol* 1996; 133:1403-15; PMID:8682874; <http://dx.doi.org/10.1083/jcb.133.6.1403>
- Orr AW, Helmke BP, Blackman BR, Schwartz MA. Mechanisms of mechanotransduction. *Dev Cell* 2006; 10:11-20; PMID:16399074; <http://dx.doi.org/10.1016/j.devcel.2005.12.006>
- Chen CS, Tan J, Tien J. Mechanotransduction at cell-matrix and cell-cell contacts. *Annu Rev Biomed Eng* 2004; 6:275-302; PMID:15255771; <http://dx.doi.org/10.1146/annurev.bioeng.6.040803.140040>
- Goicoechea SM, Awadia S, Garcia-Mata R. I'm coming to GEF you: regulation of rhoGEFs during cell migration. *Cell Adh Migr* 2014; 8:535-49; PMID:24727387; <http://dx.doi.org/10.4161/cam.28721>
- Rossmann KL, Der CJ, Sondel J. GEF means go: turning on RHO GTPases with guanine nucleotide-exchange factors. *Nat Rev Mol Cell Biol* 2005; 6:167-80; PMID:15688002; <http://dx.doi.org/10.1038/nrm1587>
- Tcherkezian J, Lamarche-Vane N. Current knowledge of the large rhoGAP family of proteins. *Biol Cell* 2007; 99:67-86; PMID:17222083; <http://dx.doi.org/10.1042/BC20060086>
- Garcia-Mata R, Burridge K. Catching a GEF by its tail. *Trends Cell Biol* 2007; 17:36-43; PMID:17126549; <http://dx.doi.org/10.1016/j.tcb.2006.11.004>
- Struckhoff AP, Rana MK, Kher SS, Burrow ME, Hagan JL, Del Valle L, Worthylyke RA. PDZ-RhoGEF is essential for CXCR4-driven breast tumor cell motility through spatial regulation of rhoa. *J Cell Sci* 2013; 126:4514-26; PMID:23868972; <http://dx.doi.org/10.1242/jcs.132381>
- Hamelers IH, Olivo C, Mertens AE, Pegtel DM, van der Kammen RA, Sonnenberg A, Collard JG. The rac activator tiam1 is required for (alpha)3(beta)1-mediated laminin-5 deposition, cell spreading, and cell migration. *J Cell Biol* 2005; 171:871-81; PMID:16330714; <http://dx.doi.org/10.1083/jcb.200509172>
- Ridley AJ. Cyclin' round the cell with rac. *Dev Cell* 2001; 1:160-1; PMID:11702775; [http://dx.doi.org/10.1016/S1534-5807\(01\)00029-6](http://dx.doi.org/10.1016/S1534-5807(01)00029-6)
- Mettrouchi A, Klein S, Guo W, Lopez-Lago M, Lemichez E, Westwick JK, Giancotti FG. Integrin-specific activation of rac controls progression through the G(1) phase of the cell cycle. *Mol Cell* 2001; 8:115-27; PMID:11511365; [http://dx.doi.org/10.1016/S1097-2765\(01\)00285-4](http://dx.doi.org/10.1016/S1097-2765(01)00285-4)
- Guilloy C, Swaminathan V, Garcia-Mata R, O'Brien ET, Superfine R, Burridge K. The rho GEFs LARG and GEF-H1 regulate the mechanical response to force on integrins. *Nat Cell Biol* 2011; 13:722-7; PMID:21572419; <http://dx.doi.org/10.1038/ncb2254>
- Heck JN, Ponik SM, Garcia-Mendoza MG, Pehlke CA, Inman DR, Eliceiri KW, Keely PJ. Microtubules regulate GEF-H1 in response to extracellular matrix stiffness. *Mol Biol Cell* 2012; 23:2583-92; PMID:22593214; <http://dx.doi.org/10.1091/mbc.E11-10-0876>