Lawrence Berkeley National Laboratory

Recent Work

Title

AN IMPROVED METHOD FOR THE ISOLATION OF CARBOXYDISMUTASE. PROBABLE IDENTITY WITH FRACTION I PROTEIN AND PROTEIN MOIETY OF PROTOCHLOROPHYLL HOLOCHROME

Permalink

<https://escholarship.org/uc/item/3pd9150n>

Author

Trown, P.W.

Publication Date

1964-12-01

University of California

Ernest O. Lawrence Radiation Laboratory

AN IMPROVED METHOD FOR THE ISOLATION OF
CARBOXYDISMUTASE. PROBABLE IDENTITY WITH FRACTION I
PROTEIN AND THE PROTEIN MOIETY OF PROTOCHLOROPHYLL
HOLOCHROME

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Biochemistry

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

AN IMPROVED METHOD FOR THE ISOLATION OF CARBOXYDISMUTASE.
PROBABLE IDENTITY WITH FRACTION I PROTEIN AND THE PROTEIN
MOIETY OF PROTOCHLOROPHYLL HOLOCHROME

P. W. Trown

December 1964

AN IMPROVED METHOD FOR THE ISOLATION OF CARBOXYDISMUTASE. PROBABLE
IDENTITY WITH FRACTION I PROTEIN AND THE PROTEIN MOIETY OF
PROTOCHLOROPHYLL HOLOCHROME

P. W. Trown

From the Laboratory of Chemical Biodynamics, Lawrence Radiation Laboratory,
University of California, Berkeley, California

ABSTRACT

An essentially homogeneous preparation of carboxydismutase has been obtained from spinach leaves by ammonium sulfate fractionation and repeated gel filtration on Sephadex G-200. The enzyme was stabilized during the purification by the presence of sulfate ions at a concentration of 0.1 M or greater, which reversibly inhibits its catalytic activity. Chloride ions at similar concentration caused no inhibition, but irreversibly deactivated the enzyme over a period of several days. The molecular weight of carboxydismutase was determined to be 515,000 by sedimentation equilibrium. Its sedimentation coefficient ($S_{w,20}$) was 18.57 Svedbergs. These and other physical properties correspond closely with those of so-called fraction I protein present in many plants, and also with those of the protein moiety of protochlorophyll holochrome obtained from some etiolated plants. It is concluded that fraction I protein is crude carboxydismutase which may also make up the protein moiety of protochlorophyll holochrome. The purified enzyme showed no significant phosphoriboisomerase or phosphoribulokinase activities, these enzymes being readily separated from carboxydismutase by the procedure employed.

INTRODUCTION

The isolation from Chlorella of a cell-free extract capable of catalysing the carboxylation of RuDP¹ to give PGA was first reported by Quayle et al. (1954). Soon afterwards, an enzyme, now named carboxydismutase or RuDP carboxylase, was isolated from spinach by Weissbach et al. (1956), and independently by Jakoby et al. (1956). The enzyme was purified about 10-fold from the crude extracts by Weissbach et al. It behaved as an almost homogeneous protein on ultracentrifugation and electrophoresis, and was estimated to have a molecular weight of about 300,000. Its sedimentation coefficient was 17 Svedberg units under the conditions used.

The similarity of these physical properties to those of an apparently homogeneous soluble protein obtained from many higher plants and known as "fraction I protein" (Wildman and Bonner, 1947) was noted by Dornier et al. (1957). Subsequent workers (Lyttleton and T'so, 1958; Park and Pon, 1961) confirmed this observation and demonstrated that carboxydismutase activity was associated with "fraction I protein". More recently a highly purified protein-protochlorophyll complex was isolated from etiolated leaves (Boardman, 1962a) which had very similar physical properties to those of fraction I protein and carboxydismutase. It therefore became of interest to determine the physical properties of highly purified carboxydismutase in order to test its possible identity with fraction I protein and also the protein moiety of protochlorophyll holochrome.

The mechanism of action of carboxydismutase has also been the subject of much investigation and speculation (Calvin, 1954b; Hurwitz et al., 1956a; Pon, 1960; Pon et al., 1963; Akoyunoglou and Calvin, 1963). Enzyme purified

¹Abbreviations used are: RuDP, ribulose-1,5-diphosphate; PGA, 3-phosphoglyceric acid; R-5-P, ribose-5-phosphate; Ru-5-P, ribulose-5-phosphate; ATP, adenosine-5'-triphosphate; EDTA, ethylenediaminetetraacetic acid; GSH, reduced glutathione; SAS, saturated ammonium sulfate [70 g/100 ml original volume at 0°].

by the methods described in this paper was used in more recent studies in which the mechanism was partially elucidated (Rabin and Trown, 1964a; Rabin and Trown, 1964b; Trown and Rabin, 1964).

For investigation of its physical properties and the mechanism of its action, a large quantity of very pure, active and, if possible, stable enzyme was required. The isolation procedure of Weissbach et al. was unsuitable because it was lengthy and low-yielding. An improved procedure was therefore sought.

Carboxydismutase has been shown to be located mainly in the chloroplasts of spinach (Heber et al., 1963), and at least 90% of the carboxydismutase activity of chloroplasts was previously shown to be associated with the colourless supernatant protein obtained by high speed centrifugation of sonicated or lysed chloroplasts (Park and Pon, 1961). An extract of lysed chloroplasts was therefore used as the starting material for the isolation rather than a whole-leaf homogenate which contains many more impurities (Pon, 1960; Pon et al., 1963). This communication describes the subsequent purification of the enzyme (by ammonium sulfate fractionation and repeated gel filtration under special conditions) and the determination of some of its physical properties.

EXPERIMENTAL PROCEDURES

Materials.—Enzyme grade ammonium sulfate was purchased from Gallard-Schlesinger Chemical Corp., N. Y.; it gave a negative reaction in the dithizone test for metal contaminants. Tris buffers at the desired pH were prepared from Trizma (Sigma Chemical Co., St. Louis, Mo.) and reagent grade hydrochloric acid or sulfuric acid. They were prepared as stock solutions (total tris, 1.0 M) and freed from heavy-metal ion contaminants as described by Rabin and Trown (1964a). The water and dialysis tubing used in the investigations were freed of heavy-metal ion contaminants by standard procedures (Hughes and Klotz,

1956; Thiers, 1957), and polyethylene apparatus was used wherever possible.

RuDP was purchased from Calbiochem, Los Angeles, Calif., as the Ba salt. It was stated to contain 26% RuDP and was either used without purification in some of the earlier work or was purified as described by Rabin et al. (1958), or by continuous electrophoresis in pyridine acetate buffer (0.25 M in pyridine, pH 5.0) using a Beckman Spinco Model CP continuous flow paper electrophoresis cell (Trown, 1964). Where necessary, the Ba salt of RuDP was converted to the Na salt as previously described (Rabin and Trown, 1964a).

R-5-P was purchased from Calbiochem, Los Angeles, Calif. as the Ba salt. It was stated to be 98.3% pure, and was converted to the Na salt (0.1 M) as follows: 10 mg were suspended in water (750 μ l) at 0°, then Na₂ SO₄ (0.1 M, 250 μ l) was added. The mixture was left at 0° for 15 min and then centrifuged to remove the precipitated BaSO₄.

ATP was purchased from Pabst Laboratories, Milwaukee, Wis., Sephadex G-200 from Pharmacia, Uppsala, Sweden, and Triton X-100 from Rohme and Haas, Philadelphia, Pa. All other chemicals used were reagent grade.

Protein Determination.—Dry weight measurements on salt-free, highly-purified carboxydismutase showed that a solution of the enzyme containing 0.71 mg/ml has an optical density of 1.00 (1 cm path) at 280 m μ . Protein concentrations were determined throughout, using this relationship.

Enzyme Assays.—Carboxydismutase activity was determined by measurement of the acid-stable radioactivity produced in the reaction between RuDP and NaH¹⁴CO₃ as described by Rabin and Trown (1964a). Approximately 5 μ g of protein were used in each assay. A unit of carboxydismutase activity is defined as the amount that will fix 1 μ mole of carbon in 10 min under the assay conditions used.

Phosphoriboisomerase activity was measured by the procedure of Axelrod and Jang (1954), in which Ru-5-P produced from R-5-P is determined by the cysteine-carbazole test. The modification described by Hurwitz et al. (1956b)

was employed.

The combined activities of phosphoriboisomerase and phosphoribulokinase were determined by an assay using R-5-P, $\text{NaH}^{14}\text{CO}_3$ and ATP as substrates in the presence of an excess of carboxydismutase. The total acid-stable radioactivity present after the reaction was taken to be proportional to the RuDP produced by the two enzymes being assayed. A typical incubation mixture contained the following components in a total volume of 250 μl : tris-chloride buffer, pH 8.0, 30 μM ; MgCl_2 , 4 μM ; $\text{NaH}^{14}\text{CO}_3$, 25 $\mu\text{C}/\mu\text{M}$, 3 μM ; R-5-P, 1.25 μM ; ATP, 5 μM ; isomerase-kinase mixture, approximately 10 μg ; purified carboxydismutase, 10 μg . The mixture was incubated at 25° for 10 min and then acetic acid (6 N, 50 μl) was added to stop the carboxydismutase reaction. Aliquots (25 μl) were plated on aluminum planchets for radioactivity counting. For preparations with very low activities, longer incubation periods (up to 6 hr) were employed. A unit of "isomerase-kinase" activity is defined as the amount of the combined enzymes phosphoriboisomerase and phosphoribulokinase required to produce 1 μmole of RuDP in 10 min under the conditions described above. It is assumed that all of the RuDP produced by the isomerase-kinase mixture is converted to PGA by the excess carboxydismutase. The carboxydismutase used in the assay was a highly purified preparation with insignificant isomerase-kinase activity.

Isolation of Carboxydismutase. 1. Preparation of the Chloroplast Extract.—

Chloroplasts were isolated by differential centrifugation from 10 kg of spinach leaves using a phosphate buffered isotonic sucrose solution (pH 7.4) as described by Park and Pon (1961). The chloroplasts were lysed with tris-sulfate buffer (0.002 M, pH 7.4) and subsequently removed by centrifugation at 40,000 x g for 10 min. The resulting supernatant (the "chloroplast extract") was made 90% SAS by the addition of solid ammonium sulfate (the pH being maintained at approx 7.5 by the addition of a concentrated solution of ammonia) and left overnight at 0°.

2. Fractional Precipitation of the Chloroplast Extract.—The material precipitated at 90% SAS from the chloroplast extract was collected by centrifugation at 40,000 x g for 15 min and then re-dissolved in tris-sulfate buffer (0.01 M, pH 7.4). The final volume of the solution was 90 ml, and the concentration of ammonium sulfate, as determined by conductivity measurements, was 27.8% SAS. The concentration of protein in the solution was 28 mg/ml. Solid ammonium sulfate was then added to the solution to obtain consecutively 30, 35, 40, 45, 50, 60 and 70% SAS. After each addition the solution was stirred at 2° for at least 1 hr and was then centrifuged at 40,000 x g for 10 min to collect the precipitate which had formed. Each precipitated fraction was re-dissolved in tris-sulfate buffer (0.01 M, pH 7.6), and dialysed against the same buffer at 0° overnight, by the method of Hospelhorn (1961).

3. Gel Filtration on Sephadex G-200.—After removal of the fines by decantation, Sephadex G-200 was equilibrated with a solution containing tris-sulfate buffer, 0.1 M, pH 7.4; ammonium sulfate, 0.2 M; GSH, 0.001 M; EDTA, Na salt, pH 7.4, 0.001 M. A gel column 44 x 4.7 cm was prepared from this material and cooled to 2°, at which temperature all subsequent operations were performed. The 35-40% SAS and 40-45% SAS cuts from the fractional precipitation were combined and divided into two parts, one of which was retained for subsequent purification. The other half was dialysed overnight against a solution containing tris-sulfate buffer, ammonium sulfate, GSH and EDTA as above. The dialysed solution (7.5 ml containing 580 mg protein) was applied to the top of the Sephadex gel bed (which was protected by a disc of filter paper) through a glass capillary tube as suggested by Porath (1964). Elution of the column was achieved with the same tris-sulfate buffer/ammonium sulfate/GSH/EDTA mixture described above. After analysis of the column effluent by optical density and enzymic activity measurements, fractions containing material with

the highest carboxydismutase activity were combined. The solution was concentrated by precipitation with ammonium sulfate followed by centrifugation and re-solution of the precipitate in tris-buffer (0.1 M, pH 7.4). After dialysis against more of the solution used as effluent of the first column, the enzyme solution was re-applied to an identical Sephadex G-200 column to that already described. The column was eluted in a similar manner and fractions containing the highest specific carboxydismutase activity were again combined. Carboxydismutase was stored until use as a suspension in 50% SAS, at 2°.

Effects of Various Compounds Upon the Stability of Carboxydismutase.—

The compounds listed below were added at the stated concentrations to solutions of partially purified carboxydismutase (35-40% SAS cut) in tris-chloride buffer (0.001 M, pH 7.6) containing EDTA (Na salt, 0.001 M, pH 7.4). The solutions were stored at 0°. Aliquots were withdrawn at various times and assayed for carboxydismutase activity. The compounds studied were: GSH, 0.001 M; sodium ascorbate, 0.001 M; cysteine, 0.001 M; sucrose, 1.0 M, 0.5 M, 0.25 M; Triton X-100, 0.05%, 0.02%, 0.01%; sodium dodecyl sulfate, 0.001 M; sodium chloride, 0.25 M, 0.083 M; ammonium sulfate, 0.25 M, 0.083 M; ammonium chloride, 0.25 M; sodium sulfate, 0.25 M; and potassium phosphate buffer at pH 7.6, 0.2 M, 0.05 M, and 0.01 M. In the case of the inorganic salts, a sample was withdrawn from each solution between 48 and 96 hr after the start and dialysed overnight against tris-chloride buffer, 0.001 M at 0°. The next day, the dialysed and undialysed solutions were assayed at the same time for carboxydismutase activity.

Inhibition of Carboxydismutase by Sulfate and Phosphate.—The total fixation of CO₂ by carboxydismutase in 10 min at 25° was measured at six different concentrations of RuDP in the absence and presence of (a) ammonium sulfate, 0.02 M, (b) ammonium sulfate, 0.1 M, (c) potassium phosphate buffer,

pH 7.6, 0.08 M, and (d) potassium phosphate buffer, 0.04 M. It was assumed that the values for the total fixation of CO_2 were proportional to the initial rates of reaction under the various conditions.

Ultracentrifugal Analysis of Purified Carboxydismutase.—Sedimentation coefficients of purified carboxydismutase at various concentrations of protein ranging from 1 mg/ml to 10 mg/ml were determined in a solution containing ammonium sulfate, 0.1 M, tris-sulfate buffer, 0.01 M, pH 7.4, and EDTA, 0.001 M, at 4.75° in a Beckman Spinco Model E Ultracentrifuge. Diffusion coefficients of carboxydismutase were determined in the same solvent at 4.75° and at protein concentrations within the same range by use of a synthetic boundary cell in the ultracentrifuge. The Rayleigh interference optical system was employed and the data were computed by the method of Longworth (1952).

The molecular weight of carboxydismutase was determined by the method of Archibald (1947), at a protein concentration of 7.3 mg/ml in the solvent already described. Data were obtained from the schlieren patterns and calculations made as described by Ehrenberg (1957). A more accurate value of the molecular weight was obtained from a sedimentation equilibrium experiment at 2095 rpm and 4.75° with a solution of carboxydismutase (3.97 mg/ml) in the solvent described above.

Miscellaneous.— The partial specific volume of carboxydismutase was calculated from the densities of solutions of the enzyme at known concentrations and of the solvent. Densities were measured using a Gay-Lussac 5 ml density bottle at $25^\circ \pm 0.05^\circ$. Purified carboxydismutase was subjected to electrophoresis on 2.5% (final conc.) polyacrylamide gel at pH 7.9. Protein bands were detected by staining the gel columns with Nigrosin in 5% acetic acid.

RESULTS

Stabilization of Carboxydismutase. (a) Organic compounds.---Experiments with partially purified carboxydismutase showed that, of a number of organic compounds tested, only GSH, 0.0 M sucrose and Triton X-100 at a concentration of 0.05% protected the enzyme against deactivation to any significant extent. The results of these experiments are summarized in Table I. Although values of remaining activity are only given for one time (160 hrs), assays were performed approximately every 24 hrs, and indicated a steady trend towards the final values given in Table I. The variation in the activity remaining in the control solutions varied considerably between the various experiments. The cause of this variation is unknown. The ionic detergent, sodium dodecyl sulfate, completely destroyed the carboxydismutase in less than 1 min in contrast with the stabilizing effect shown by the non-ionic detergent Triton X-100.

(b) Inorganic compounds.---The comparable effects upon the stability of carboxydismutase of sodium chloride, sodium sulfate, ammonium chloride, and ammonium sulfate at a concentration of 0.25 M are summarized in Fig. 1. Sodium sulfate and ammonium sulfate had similar effects in that they inhibited carboxydismutase approximately 80% immediately upon addition to the enzyme solution. This inhibition was maintained with time but could be completely reversed by dialysis with restoration of essentially all the initial activity. In contrast, ammonium chloride and sodium chloride caused an apparent activation of about 25% above that of the control solution immediately upon their addition to the enzyme solutions. However, this activation was not maintained since the activity decreased with time and could not be recovered by dialysis. Potassium phosphate buffer at pH 7.6 protected carboxydismutase activity in a similar manner to sodium and ammonium sulfates, except that only 80% of the initial activity was recovered after 120 hrs on removal of the phosphate by

TABLE I

Compound added	% initial activity remaining after 160 hrs	Compound added	% initial activity remaining after 160 hrs
GSH, 0.001 <u>M</u>	95	Triton X-100, 0.05%	75*
Sodium Ascorbate, 0.001 <u>M</u>	70	Triton X-100, 0.01%	40
		Triton X-100, 0.002%	40
Cysteine, 0.001 <u>M</u>	70	Sodium dodecyl sulfate, 0.001 <u>M</u>	0†
Control - no additions	70	Control - no additions	40
Sucrose, 1.0 <u>M</u>	60		
Sucrose, 0.5 <u>M</u>	45		
Sucrose, 0.75 <u>M</u>	45		
Control - no additions	40		

* No loss of activity during first 72 hrs

† No carboxydismutase activity remained after 1 min

dialysis. Phosphate also inhibited carboxydismutase activity: at zero time, solutions containing 0.01, 0.05, and 0.2 M potassium phosphate at pH 7.6 had respectively 80, 40, and 5% of the control solution activity.

Inhibition of Carboxydismutase by Sulfate and Phosphate.— Figures 2 and 3 show the Lineweaver-Burk plots for the inhibition of carboxydismutase by sulfate and phosphate respectively, each at two different concentrations.

Purification of Carboxydismutase.—The results of the fractionation of the 90% SAS-precipitate from the chloroplast extract are summarized in Table II. More than 75% of the total protein was precipitated in the range 35-45% SAS where the highest specific carboxydismutase activity was also found. A 60% increase in the specific carboxydismutase activity was achieved during the fractionation. The highest kinase-isomerase activity was found in the 40-50% SAS range. The ratio of carboxydismutase to kinase-isomerase activities decreased with increasing ammonium sulfate concentrations from a value of 6.5 to 1 in the 30% SAS cut to 0.45 to 1 in the supernatant from the 60% SAS precipitation.

The elution of carboxydismutase from the first of the two columns of Sephadex G-200 used is shown in Fig. 4. Material with an ultraviolet absorption maximum at 260 mμ was separated from carboxydismutase and was eluted both before the main protein peak and also after it in a long "tail" (not shown in Fig. 4). The specific carboxydismutase activity for fractions 75-100 was constant at 1.26 units/mg, but values up to 2.15 units/mg have been obtained in subsequent similar purifications. Fractions 75-100 were combined, concentrated, and then applied to the second Sephadex G-200 column as described in the Experimental Section. The elution pattern of the second column was similar to that of the first except that the first peak was very small and there was almost no "tail". Material from the peak fractions of the second column had

TABLE II

& SAS	O.D.280 units recovered (% total)	λ_{max} (m μ)	Specific Carboxy- dismutase activity (units/mg)*	Specific "Kinase- Isomerase" activity (units/mg)	Carboxydismutase/ kinase-isomerase ratio
30	4.0	260	0.013	0.02	6.5
35	7.7	257	0.23	0.06	3.8
40	54.5	276	0.81	0.22	3.7
45	21.6	277	1.20	0.36	3.3
50	5.9	278	0.35	0.45	0.78
60	2.4	275	0.08	0.12	0.66
Super- natant from 60% SAS	3.6	273	0.009	0.02	0.45
Chloro- plast Extract	—	269	0.74	0.21	3.5

* Units as defined in Experimental Section. This unit is not the same as that defined by Weissbach et al. (1956), since very different conditions of incubation with RuDP were employed in the assay.

a specific carboxydismutase activity of 1.54 units/mg, but activities up to 2.58 units/mg have been obtained in subsequent similar purifications. This activity remained constant over a period of several months when the enzyme was stored as a precipitate in 50% SAS.

The eluate from the first Sephadex G-200 column of a different isolation from that shown in Fig. 4 was also assayed for phosphoriboisomerase and "isomerase-kinase" activities. The results of these assays are shown in Fig. 5. While some phosphoriboisomerase and phosphoribulokinase activity was eluted with the carboxydismutase, the two former enzymes reached their peak specific activities at fractions 180 and 190 respectively, where no carboxydismutase activity could be detected. The elution of phosphoriboisomerase and isomerase-kinase activities from the second Sephadex G-200 column in this experiment was similar to that shown in Fig. 5, except that none of either activity was detected until fraction 150 had been eluted. The maximum specific activities of both enzymes again occurred at about fractions 180-190.

The purified carboxydismutase from the peak fractions of the second Sephadex G-200 column was assayed for isomerase-kinase activity for 6 hrs at 25° in order to obtain enough radioactivity to count. The equivalent of 1 unit of kinase-isomerase activity was detected per 20,000 units of carboxydismutase activity.

Properties of Carboxydismutase.—The sedimentation coefficient of carboxydismutase was proportional to the concentration of the enzyme. A plot of $1/S$ versus concentration was used for the extrapolation to zero concentration, and a value of 11.56 was obtained for S^0 under the conditions employed. Corrections for temperature, viscosity and density, to the standard state gave $S^0_{20,w} = 18.57$. The constant k in the equation $S^0/S = 1 + kc$ was found to have a value of 0.091 (S = sedimentation coefficient at concentration c in grams per 100 ml).

and S^0 = the value of S at infinite dilution). A solution of carboxydismutase (7.3 mg/ml) was frozen and thawed before examining it for sedimentation characteristics. After such treatment three distinct schlieren peaks were visible with sedimentation coefficients of 18S, 26S, and 32S. The areas under the schlieren peaks were in the ratios 6:2:1 respectively. Electron micrographs of such a solution showed the presence of monomers, dimers and trimers in approximately the same ratio. The trimers consisted of monomers arranged in a linear fashion.

The diffusion coefficient of carboxydismutase also varied linearly with concentration and extrapolation of a plot of D versus C gave $D^0 = 1.77 \times 10^{-7}$ cm²/sec. Corrections for temperature, viscosity and density, to the standard state gave $D_{20,w}^0 = 2.93 \times 10^{-7}$ cm²/sec. The partial specific volume of carboxydismutase, as determined from measurements of the densities of enzyme solutions and solvent, was 0.730 ml/g.

The molecular weight of carboxydismutase, as determined by Archibald's method was $560,000 \pm 60,000$. Another value of the molecular weight, 559,000, was obtained using the equation $M = RTS/D(1-\bar{V}\rho)$ where R is the gas constant, T is the absolute temperature, $\bar{V}$ is the partial specific volume, and ρ is the density of the enzyme solution. The most accurate determination of the molecular weight was obtained from a sedimentation equilibrium experiment. A plot of the logarithm of protein concentration versus the square of the distance from the centre of rotation (x^2) derived from this experiment is shown in Fig. 6. The molecular weight derived from these data was $515,000 \pm 10,000$. The slight heterogeneity of the preparation of carboxydismutase is indicated by the departure from linearity of the $\log c$ versus x^2 plot at high values of x^2 , i.e., near the bottom of the centrifuge cell. This heterogeneity was confirmed when the enzyme was subjected to electrophoresis on polyacrylamide gel at pH 7.9.

A small band of contaminant which contained less than 5% of the total fixed Mgrosin colour was observed to move approximately half as fast towards the anode as did the main protein band.

Electron micrographs of the purified enzyme indicated that the molecule is compact and non-linear. Micrographs at 80,000 x magnification in which the protein had been shadowed with Nichrome, showed that many of the particles had a depression in the centre of the upturned face. This finding was confirmed by higher resolution pictures (magnification factor 480,000 x) using sodium phosphotungstate (pH 7.6) in the negative staining technique. Almost every particle had an electron dense centre, i.e., there was a hole or depression in the molecule which could be penetrated by the phosphotungstate. Some particles had an electron dense line across them instead of the hole.

Some of the physical properties described above are presented in Table III, together with values obtained by other workers for "fraction I protein" and protochlorophyll holochrome.

TABLE III

Property	"Fraction I protein"	Carboxydismutase	Protochlorophyll Holochrome
Sedimentation coefficient (Svedbergs)	$S_{w,20} = 17.9-19.5^a$ $S_{20} = 16.2-16.6^b$ $S_{20}^o = 18.5^c$	$S_{\text{observed}} = 17^d$ $S_{w,20} = 18^e$ $S_{w,20}^o = 18.57^f$	$S_{20} = 15.3-16.2^g$ $S_{w,20}^o = 18.0^h$
Diffusion coefficient ($\times 10^7$ cm ² /sec)	$D_{w,20}^o = 2.60-2.75^b$	$D_{\text{observed}} = 5.5^d$ $D_{\text{observed}} = 1.6^e$ $D_{w,20}^o = 2.93^f$	$D_{w,20}^o = 2.7^h$
Partial specific volume (ml/g)	± 0.69	0.73^f	0.73^h
Molecular weight	$\approx 600,000$ (estimate) ⁱ $375,000^a$ $\approx 595,000-620,000^b$	$300,000$ (estimate) ^d $515,000 \pm 10,000^f$	$400,000-700,000^j$ $600,000 \pm 50,000^h$
Mobility on electrophoresis at stated pH	pH 7.2 : towards anode ^a	pH 7.7 towards anode ^d pH 6.9 " " ^e pH 7.9 " " ^f	pH 9.6 towards anode ^j
Particle dimensions (from elec- tron micro- graphs)	$100 \times 200 \text{ \AA}^k$	$100 \times 200 \text{ \AA}^k$ $\sim 100 \text{ \AA}^f$	$100-110 \text{ \AA}^h$

References: a, Eggman et al. (1953); b, Lyttleton (1956); c, Kupke (1962);
d, Weissbach et al. (1956); e, Pon (1960); f, present study; g, Smith
and Kupke (1956); h, Boardman (1962a); i, Singer et al. (1952);
j, Smith (1960); k, Park and Pon (1961).

* Molecular weights calculated assuming $\bar{v} = 0.75$ ml/g.

* Sample contained 11.2% nucleic acid.

DISCUSSION

Carboxydismutase activity has long been known to be inhibited by sulfhydryl blocking agents such as p-chloromercuribenzoate and iodoacetamide (Mayaudon et al. (1957)), and it has been shown recently that a sulfhydryl group is probably an essential part of the active site of the enzyme (Rabin and Trown, 1964a and 1964b; Trown and Rabin, 1964). It was not surprising therefore that reduced glutathione protected the enzymic activity during purification, presumably by inhibition of the oxidation of the essential sulfhydryl group on the enzyme. That cysteine was not able to act in a similar manner may be related to its higher reduction potential (Calvin, 1954a), making it less readily oxidized than the essential enzyme sulfhydryl group.

The protection, though small, afforded the enzyme by 1.0 M sucrose is in agreement with the results of Heitfuss et al. (1959) who found that polyhydroxy compounds in general stabilized cabbage leaf proteins. The stabilization of carboxydismutase by the non-ionic detergent Triton X-100 may be a related phenomenon. Carboxydismutase is evidently a very hydrophilic protein, as is also evident from its very high solubility in aqueous solvents (solutions containing up to 200 mg/ml have been obtained in the present investigation). This hydrophilicity must be due in part to the large number of charged groups the enzyme carries; its mobility on electrophoresis at pH 7.7 (Weissbach et al. (1956)) is remarkable considering its high molecular weight. This high charge may account for the dramatic effect which the ionic detergent sodium dodecyl sulfate has upon the enzyme, contrasting sharply with that of the non-ionic detergent Triton X-100. Deactivation by sodium dodecyl sulfate is immediate and irreversible. Interaction of charged species on the detergent and enzyme could lead to disruption of the tertiary structure and indeed carboxydismutase is broken down into sub-units with sedimentation coefficients of approximately

two S by sodium dodecyl sulfate at the concentration used in the present investigations (Trown, 1964).

Experiments with the four possible salts of the cations Na^+ and NH_4^+ and the anions SO_4^{2-} and Cl^- indicated that the anionic content of the salts controlled their effect on the enzyme. Sulfate ions reversibly inhibit carboxydismutase and also protect it against deactivation. In contrast, chloride ions caused an apparent activation of the enzyme, which, however, was not maintained, and the long-term result was permanent deactivation. Phosphate buffer at pH 7.6 had a similar effect to sulfate; Weissbach et al. (1956) reported both phosphate and arsenate to be inhibitors of carboxydismutase. On the basis of these limited results, it seems that divalent anions are able to inhibit and protect the enzyme, whereas the monovalent anion Cl^- causes deactivation. Such a result is not without precedent; London et al. (1962) found that prostatic acid phosphatase was protected by sulfate ions against thermal denaturation, whereas the ions Cl^- , Br^- and CNS^- accelerated the process. Inhibition of enzymic activity by inorganic anions is also fairly common—for example, transaldolase is inhibited by inorganic phosphate (Bonsignore et al. (1960). In all these cases, it is possible that the anions are able to inhibit and/or protect the enzyme by virtue of their being similar in structure and/or charge distribution to the substrate, which, in the cases cited, is an organic phosphate. Protection of enzymes by substrates or substrate analogues has often been reported (e.g., Delory and King, 1943; Burton, 1951). As previously suggested (Dorner et al. (1957; Lyttleton and Ts'o, 1958; Park and Fon, 1961) and also strongly indicated by results to be discussed later, so-called "fraction I protein" is a crude form of carboxydismutase. We may therefore regard attempts to purify "fraction I protein" and carboxydismutase as synonymous. Such attempts have always been complicated by the

tendency of the enzyme to bind other substances to itself to varying degrees. Among the substances which have at some time been considered to be integral parts of "fraction I protein" are auxin and a phosphatase (Wildman and Bonner, 1947), nucleic acid (Eggman et al., 1961; Mendiola and Akazawa, 1964). However, Weissbach et al. (1956) were able to obtain a preparation of carboxydismutase which did not contain auxin or nucleic acid and which had no other enzymic activity but that of carboxydismutase. More recently, Van Noort and Wildman (1964) were also able to prepare, by precipitation with specific rabbit antibody, "fraction I protein" which contained carboxydismutase activity but not phosphoriboisomerase or phosphoribulokinase activity. The present work supports these findings and furthermore shows that the enzymes phosphoriboisomerase and phosphoribulokinase may be partially separated from carboxydismutase by careful ammonium sulfate fractionation and completely separated by repeated gel filtration on Sephadex G-200. This latter technique was employed by Mendiola and Akazawa (1964) for the purification of "fraction I protein" from rice leaves, but these authors assayed the column eluent for catalysis of the sequence of reactions $R-5-P \rightarrow Ru-5-P \rightarrow RuDP \rightarrow PGA$ using a radioactive substrate (CO_2) in the final stage, and therefore would not have detected phosphoriboisomerase and phosphoribulokinase if they had been separated from carboxydismutase.

The position at which phosphoriboisomerase and phosphoribulokinase were eluted from the Sephadex G-200 column indicates that these enzymes have molecular weights smaller than 200,000, which is normally thought to be the exclusion limit for this particular Sephadex. Since the molecular weight of carboxydismutase is much greater than 200,000 it is excluded from Sephadex G-200, but it nevertheless retarded somewhat on the column compared with a material containing nucleic acid which comes through with the front. The

retardation of carboxydismutase must be due to specific physical interaction between the enzyme and the column material. This may be yet another manifestation of the hydrophilic nature of carboxydismutase, resulting in interaction with the polyhydroxy compound dextran.

The purity of the carboxydismutase prepared by the method described was very high. The slight departure from linearity of the plot of $\log c$ versus x^2 in a sedimentation equilibrium experiment was an indication of the presence in the sample of a small amount of a contaminant of higher molecular weight. This finding was confirmed by electrophoresis on polyacrylamide gel where a very small amount of slower moving (i.e., less charged or of higher molecular weight) contaminant was detected. This contaminant may well be polymerised carboxydismutase which was encountered in preparation of the enzyme which had been stored under non-optimal conditions for extended periods.

The physical properties of carboxydismutase are very similar to those of "fraction I protein" as may be seen in Table III. The sedimentation coefficients of the two substances, particularly those of the more purified preparations, are virtually identical. The properly corrected diffusion coefficient (2.50 – 2.75×10^{-7} cm²/sec) of the "fraction I protein" prepared by Lyttleton (1956) compares well with that determined for purified carboxydismutase (2.93×10^{-7} cm²/sec). The values determined by Weissbach et al. (1956) and Pon (1960) were not corrected to the standard state. Probably the most reliable molecular weight of "fraction I protein", determined by Lyttleton (1956), was calculated for an assumed partial specific volume of 0.75 ml/g. If the value determined for carboxydismutase (0.73 ml/g) is used in the calculation, the molecular weight obtained (550,000–574,000) compares well with that determined for purified carboxydismutase ($515,000 \pm 10,000$). The values obtained by Singer et al. (1952) and Weissbach et al. (1956) were stated by the authors to be estimates, and that obtained by Eggman et al. (1953) was determined for material containing 11.2% nucleic acid. In all cases where it was investigated for both "fraction I

protein" and carboxydismutase, the direction of electrophoretic mobility was towards the anode in the pH range 6.9-7.9. Both "fraction I protein" and carboxydismutase were stated by most of the authors cited under Table III to be unstable below pH 6, both being precipitated and irreversibly denatured. Electron micrographs of both "fraction I protein" and a crude preparation of carboxydismutase showed particles $100 \times 200 \text{ \AA}$ in each preparation which were indistinguishable from each other (Park and Pon, 1961). The preparation appears to contain similar particles with an average diameter of $100 \text{ \AA}$.

The above evidence, in conjunction with the reported finding that material prepared as "fraction I protein" showed carboxydismutase activity (Lyttleton and T'so, 1958; Park and Pon, 1961; Van Noort and Wildman, 1964), strongly indicates that carboxydismutase and "fraction I protein" are identical. Certainly the obvious characteristic of "fraction I protein" -i.e., its high molecular weight and consequent high sedimentation coefficient- is due to carboxydismutase. Any other enzymatic activity present in "fraction I protein" would appear to be due to absorption on carboxydismutase of the enzymes involved. Further evidence to support this hypothesis was recently reported by Trown and Rabin (1964), who found that the equivalent weight of carboxydismutase on the basis of the sulfhydryl group at the active site of the enzyme was 267,000. There are thus two active sites per molecule of 515,000 molecular weight. It is still possible, though not likely on the basis of the present evidence, that a large protein molecule exists in the plant to which the various enzymes of the carbon cycle, including carboxydismutase, are attached. It seems more likely that the large protein molecule is carboxydismutase and that the other enzymes are either bound to it in some kind of organized structure in the plant or become bound, as did auxin and nucleic acid, during the isolation process.

The similarities of the physical properties of carboxydismutase and these

of protochlorophyll holochrome are also striking (see Table III). The most purified preparations of each have virtually identical sedimentation coefficients, diffusion coefficients, partial specific volumes and molecular weights. Although the stability of the holochrome to pH changes has not been studied extensively, a lyophilized preparation of the protein is only soluble above pH 5.0 (below which carboxydismutase is denatured). Electron micrographs of carboxydismutase and protochlorophyll holochrome are indistinguishable from each other. Both show particles with holes or depressions in the centers of the molecules, particularly evident on negative staining with sodium phosphotungstate, and the particle dimensions obtained from the micrographs are the same. Another property reported by Boardman (1962a) for the holochrome is also shown by carboxydismutase. After freezing and thawing a solution of the holochrome, the single schlieren peak corresponding to 18S is split into 3 peaks corresponding to 18S, 26S, and 32S. A similar splitting of the single schlieren peak into 18S, 26S, and 32S peaks was observed in one preparation of carboxydismutase which had been exposed to some unknown factor not normally encountered in isolation or storage of the enzyme. In this case the phenomenon was demonstrated, by electron microscopy, to be due to aggregation into dimers and trimers, a conclusion also reached by Boardman (1962a) for protochlorophyll holochrome.

In reporting the isolation of protochlorophyll holochrome, Boardman (1962a) pointed out the similarity of the holochrome to "fraction I protein" and confirmed that the two substances are indistinguishable in the ultracentrifuge. The present findings, in confirming the identity of "fraction I protein" and carboxydismutase, and taken in conjunction with the striking similarities in physical properties between carboxydismutase and holochrome, make it likely that carboxydismutase makes up the protein moiety of protochlorophyll holochrome. This is not unlikely in view of the ability, already discussed, of carboxydis-

mutase to bind other substances to itself. In this connection it is of interest that the amount of active protochlorophyll per protein molecule varies considerably from one preparation of protochlorophyll holochrome to another and also during the isolation process, Boardman (1962a). In addition, it has been shown that all of the active protochlorophyll in the holochrome preparations is probably not bound to the protein in the same way (Smith and Coomber, 1960; Boardman, 1962b). Ultraviolet light converts only 30% of the transformable protochlorophyll to chlorophyll a; the remainder is converted by visible light (McLeod and Coomber, 1959). These two pieces of evidence do not support the hypothesis that protochlorophyll is bound to a site specifically designed for this purpose in the plant, and it seems that the binding of protochlorophyll to the protein may be fortuitous, possibly occurring during the isolation procedure. Isolation of a protochlorophyll holochrome preparation which showed carboxydismutase activity would shed considerable light on this question, and work is proceeding along these lines.

REFERENCES

- Akoyunoglou, G., and Calvin, M. (1963), *Biochem. Z.* 338, 20.
- Archibald, W. J. (1947), *J. Phys. Chem.* 51, 1204.
- Axelrod, B., and Jang, R. (1954), *J. Biol. Chem.* 209, 847.
- Boardman, N. K. (1962a), *Biochim. Biophys. Acta* 62, 63.
- Boardman, N. K. (1962b), *Biochim. Biophys. Acta* 64, 279.
- Bonsignore, A., Pontremoli, S., Grazi, E., and Horecker, B. L. (1960),
J. Biol. Chem. 235, 1868.
- Burton, K. (1951), *Biochem. J.* 48, 458.
- Calvin, M. (1954a), in "Proc. Symp. on Glutathione, Ridgefield, Conn., 1953",
edit. Colowick et al., Academic Press, N. Y., p. 20.
- Calvin, M. (1954b), *Fed. Proc.* 13, 697.
- Delory, G. G., and King, E. J. (1943), *Biochem. J.* 37, 547.
- Dorner, R. W., Kahn, A., and Wildman, S. G. (1957), *J. Biol. Chem.* 229, 945.
- Eggan, L., Singer, S. J., and Wildman, S. G. (1953), *J. Biol. Chem.* 205, 969.
- Ehrenberg, A. (1957), *Acta Chem. Scand.* 11, 1257.
- Heber, U., Pon, N. G., and Heber, M. (1963), *Plant Physiol.* 38, 355.
- Heitfuss, R., Buchanan-Davidson, D. J., and Stahman, M. A. (1959), *Arch. Bio-
chem. Biophys.* 85, 200.
- Hospelhorn, V. D. (1961), *Analyt. Biochem.* 2, 180.
- Hughes, T. A., and Klotz, J. M. (1956), in "Methods of Biochemical Analysis",
edit. D. Glick, Vol. 3, p. 265.
- Hurwitz, J., Jakoby, W. B., Horecker, B. L. (1956a), *Biochim. Biophys. Acta* 22,
194.
- Hurwitz, J., Weissbach, A., Horecker, B. L., and Smyrniotis, P. Z. (1956b),
J. Biol. Chem. 218, 769.
- Jakoby, W. B., Brummond, D. O., and Cohen, S. (1956), *J. Biol. Chem.* 218, 811.
- Kupke, D. W. (1962), *J. Biol. Chem.* 237, 3287.
- London, M., McHugh, R., and Hudson, P. B. (1962), *J. Gen. Physiol.* 46, 57.

- Longworth, L. G. (1952), J. Am. Chem. Soc. 74, 4155.
- Lyttleton, J. W. (1956), Biochem. J. 64, 70.
- Lyttleton, J. W., and T'so, P. O. P. (1958), Arch. Biochem. Biophys. 73, 120.
- McLeod, G. C., and Coomber, J. (1960), Carnegie Inst. of Washington Year Book, 59, 324.
- Meyaudon, J., Benson, A.A., and Calvin, M. (1957), Biochim. Biophys. Acta 23, 342.
- Mendiola, L., and Akazawa, T. (1964), Biochemistry 3, 175.
- Park, R. B., and Pon, N. G. (1961), J. Mol. Biol. 3, 1.
- Pon, N. G. (1960), Ph.D. Thesis, Univ. of California, Berkeley; Univ. of Calif. Rad. Lab. Report No. 9373.
- Pon, N. G., Rabin, B. R., and Calvin, M. (1963), Biochem. Z. 338, 7.
- Porath, J. (1964), "Advances in Protein Chemistry", Academic Press, New York, Vol. 17, p. 209.
- Quayle, J. R., Fuller, R. C., Benson, A. A., and Calvin, M. (1954), J. Am. Chem. Soc. 76, 3610.
- Rabin, B. R., Shaw, D. F., Pon, N. G., Anderson, J. M., and Calvin, M. (1958), J. Am. Chem. Soc. 80, 2528.
- Rabin, B. R., and Trown, P. W. (1964a), Proc. Nat. Acad. Sci. 51, 497.
- Rabin, B. R., and Trown, P. W. (1964b), Nature 202, 1290.
- Singer, S. J., Eggman, L., Campbell, J. M., and Wildman, S. G. (1952), J. Biol. Chem. 197, 233.
- Smith, J. H. C., and Kupke, D. W. (1956), Nature 178, 751.
- Smith, J. H. C. (1960), Symposia on Comparative Biology, No.1, Academic Press, N. Y., p. 257.
- Smith, J. H. C., and Coomber, J. (1960), Carnegie Inst. of Washington Year Book, 59, 325.
- Thiers, R. C. (1957), in "Methods of Biochemical Analysis", edit. D. Glick, Vol. 5, p. 273.

Trown, P. W., and Rabin, B. R. (1964), Proc. Nat. Acad. Sci. 51.

Trown, P. W. (1964), unpublished results.

Van Noort, G., Hudson, W., and Wildman, S. G. (1961), Plant Physiol. 36,
Suppl. xix.

Van Noort, G., and Wildman, S. G. (1964), Biochim. Biophys. Acta 90, 309.

Weissbach, A., Horecker, B. L., and Hurwitz, J. (1956), J. Biol. Chem. 218,
795.

Wildman, S. G., and Bonner, J. (1947), Arch. Biochem. Biophys. 14, 381.

Fig. 1. Effects of ammonium sulfate, ammonium chloride, sodium sulfate and sodium chloride on the stability and activity of carboxydismutase:

Δ , 0.25 M $(\text{NH}_4)_2 \text{SO}_4$; $\circ$, 0.25 M $\text{NH}_4 \text{Cl}$; Δ , 0.25 M $\text{Na}_2 \text{SO}_4$; $\square$, 0.25 M NaCl ; $\bullet$, control.

Fig. 2. Lineweaver-Burk plot for the inhibition of carboxydismutase by ammonium sulfate. V = reaction velocity ($\mu\text{moles of C fixed per mg protein per 10 min}$), S = concentration of RuDP (mM). $\square$, no ammonium sulfate;

Δ , 0.02 M $(\text{NH}_4)_2 \text{SO}_4$; $\circ$, 0.1 M $(\text{NH}_4)_2 \text{SO}_4$.

Fig. 3. Lineweaver-Burk plot for the inhibition of carboxydismutase by potassium phosphate buffer, pH 7.6. V = reaction velocity ($\mu\text{moles C fixed per mg protein per 10 min}$), S = concentration of RuDP (mM). $\bullet$, no phosphate; $\square$, 0.04 M phosphate; Δ , 0.08 M phosphate.

Fig. 4. Fractionation of carboxydismutase on Sephadex G-200 column (44 x 4.7 cm): $\bullet$, O.D. at 260 m μ ; $\square$, O.D. at 280 m μ ; Δ , carboxydismutase activity [$(\text{dpm/ml}) \times 10^{-6}$].

Fig. 5. Elution of phosphoriboisomerase and phosphoribulokinase from Sephadex G-200 column (44 x 4.7 cm): Δ , phosphoriboisomerase; $\bullet$, "kinase-isomerase" (see Experimental section); line without points, O.D. at 280 m μ .

Fig. 6. Functions of protein concentration (C) and distance from centre of rotation (x) in a sedimentation equilibrium experiment with purified carboxydismutase (3.97 mg/ml) in 0.1 M $(\text{NH}_4)_2 \text{SO}_4$, 0.01 M tris sulfate buffer pH 7.4, 0.001 M EDTA. The rotor was overspeeded at 3397 rpm for 8 hr. and then run at 2095 rpm until equilibrium was attained. The data were obtained from a photograph of the Rayleigh interference pattern taken 120 hr. after reaching 2095 rpm. The rotor temperature was 4.75°.

MU-32792A

Fig. 1

MU-32795A

Fig. 2

MU-32793A

Fig. 3

Fig. 4 MU-32797

Fig. 5

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.