UCSF

UC San Francisco Previously Published Works

Title

Estimating dead-space fraction for secondary analyses of acute respiratory distress syndrome clinical trials.

Permalink

<https://escholarship.org/uc/item/3pv7h3h2>

Journal

Critical care medicine, 43(5)

ISSN

0090-3493

Authors

Beitler, Jeremy R
Thompson, B Taylor
Matthay, Michael A
et al.

Publication Date

2015-05-01

DOI

10.1097/ccm.0000000000000921

Peer reviewed

Estimating Dead-Space Fraction for Secondary Analyses of Acute Respiratory Distress Syndrome Clinical Trials

Jeremy R. Beitler, MD¹; B. Taylor Thompson, MD²; Michael A. Matthay, MD^{3,4};
Daniel Talmor, MD, MPH⁵; Kathleen D. Liu, MD, PhD, MAS^{6,7}; Hanjing Zhuo, MD, MPH⁴;
Douglas Hayden, PhD⁸; Roger G. Spragg, MD¹; Atul Malhotra, MD¹

¹Division of Pulmonary and Critical Care Medicine, University of California, San Diego, San Diego, CA.

²Pulmonary and Critical Care Unit, Massachusetts General Hospital, Boston, MA.

³Departments of Anesthesia and Medicine, University of California, San Francisco, San Francisco, CA.

⁴The Cardiovascular Research Institute, University of California, San Francisco, San Francisco, CA.

⁵Department of Anesthesia, Critical Care, and Pain Medicine, Beth Israel Deaconess Medical Center, Boston, MA.

⁶Division of Nephrology, University of California, San Francisco, San Francisco, CA.

⁷Division of Critical Care Medicine, University of California, San Francisco, San Francisco, CA.

⁸Biostatistics Center, Massachusetts General Hospital, Boston, MA.

Dr. Beitler received support for article research from the National Institutes of Health (NIH). His institution received grant support from the National Heart, Lung, and Blood Institute (NHLBI) (T32-HL007633). Dr. Thompson received support for article research from the NIH. His institution received grant support from the NIH/NHLBI (N01-HR56179). Dr. Matthay received support for article research from the NIH (R37-HL51856 and SCCOR HL-74005), served as a board member for Roche DSMB, and consulted for Cerus and GSK. His institution received grant support from the NHLBI and National Institute of Allergy and Infectious Diseases (NIAID). Dr. Talmor received grant support from the NHLBI and received support for article research from the NIH (UM1-HL108724). Dr. Liu served as a board member for Astute Biomedical (Clinical Events Adjudication Committee), Complexa (Scientific Advisory Board), Cytopheryx (Data Safety Monitoring Board), and Roche DSMB. She received support for travel from the American Thoracic Society (ATS) (Member of publications committee; trip to planning meeting paid for by ATS), American Society of Nephrology (Reimbursement for travel to national meeting for presentation), and Continuous Renal Replacement Therapies Meeting (Reimbursement for travel to national meeting for presentation). She consulted for Chemocentryx, Cerus, and GSK; lectured for the NIH; has stock in Amgen; disclosed other support from Abbott (gift of reagents for biomarker assays) and CMIC (gift of reagents for biomarker assays); and received support for article research from the NIH. Her institution received grant support from the NIH, NHLBI, and NIAID. Dr. Hayden is employed by a contract with the NIH-NHLBI Acute Respiratory Distress Syndrome Network (ARDSNet) (HHSN268200536179C, Fund Number: 219579) and received support for article research from the NIH. His institution received grant support from the NIH-NHLBI as the ARDSNet Clinical Coordinating Center (HHSN268200536179C, Fund Number: 219579). Dr. Malhotra received

support for article research from the NIH (K24-HL093218). His institution consulted for Pfizer, SHC, SGS, Apnex, and Apnicure (relinquished all outside personal income in 2012). The remaining authors have disclosed that they do not have any potential conflicts of interest.

For information regarding this article, E-mail: jbeitler@ucsd.edu

Objectives: Pulmonary dead-space fraction is one of few lung-specific independent predictors of mortality from acute respiratory distress syndrome. However, it is not measured routinely in clinical trials and thus altogether ignored in secondary analyses that shape future research directions and clinical practice. This study sought to validate an estimate of dead-space fraction for use in secondary analyses of clinical trials.

Design: Analysis of patient-level data pooled from acute respiratory distress syndrome clinical trials. Four approaches to estimate dead-space fraction were evaluated: three required estimating metabolic rate; one estimated dead-space fraction directly.

Setting: U.S. academic teaching hospitals.

Patients: Data from 210 patients across three clinical trials were used to compare performance of estimating equations with measured dead-space fraction. A second cohort of 3,135 patients from six clinical trials without measured dead-space fraction was used to confirm whether estimates independently predicted mortality.

Interventions: None.

Measurements and Main Results: Dead-space fraction estimated using the unadjusted Harris-Benedict equation for energy expenditure was unbiased (mean $\pm$ SD Harris-Benedict, 0.59 ± 0.13 ; measured, 0.60 ± 0.12). This estimate predicted measured dead-space fraction to within ± 0.10 in 70% of patients and ± 0.20 in 95% of patients. Measured dead-space fraction independently predicted mortality (odds ratio, 1.36 per 0.05 increase in dead-space fraction; 95% CI, 1.10–1.68; $p < 0.01$). The Harris-Benedict estimate closely approximated this association with mortality in the same cohort (odds ratio, 1.55; 95% CI, 1.21–1.98; $p < 0.01$) and remained independently predictive of death in the larger Acute Respiratory Distress Syndrome Network cohort. Other estimates predicted measured dead-space fraction or its association with mortality less well.

Conclusions: Dead-space fraction should be measured in future acute respiratory distress syndrome clinical trials to facilitate

Copyright © 2015 by the Society of Critical Care Medicine and Wolters Kluwer Health, Inc. All Rights Reserved.

DOI: 10.1097/CCM.0000000000000921

incorporation into secondary analyses. For analyses where dead-space fraction was not measured, the Harris-Benedict estimate can be used to estimate dead-space fraction and adjust for its association with mortality. (*Crit Care Med* 2015; 43:1026–1035)

Key Words: acute lung injury; acute respiratory distress syndrome; capnography; clinical trials as topic; respiratory dead space

Few lung-specific predictors of mortality from acute respiratory distress syndrome (ARDS) exist. Impaired oxygenation assessed by $\text{PaO}_2:\text{FiO}_2$ is a defining feature of ARDS, but severity inconsistently correlates with clinical outcomes (1–3). Oxygenation index, an alternative measure of oxygenation that includes mean airway pressure, may correlate with outcomes more reliably (2). In addition to measures of oxygenation, respiratory system compliance and pulmonary dead-space fraction have been found in multiple studies to predict mortality from ARDS (4–7). Yet while hypoxemia and compliance are commonly reported, dead-space fraction is rarely assessed in clinical trials.

Increased dead-space fraction occurs within hours of ARDS onset and independently predicts mortality, even after accounting for overall illness severity, hypoxemia, and compliance (4). Sustained elevation of dead-space fraction over the first week additionally identifies patients less likely to survive hospitalization (5, 8). Microvascular endothelial injury, microvascular thrombi, and derangements in pulmonary blood flow are characteristic features of ARDS that lead to increased dead-space fraction (9–11). Hyperinflation with excessive applied or intrinsic positive end-expiratory pressure (PEEP) may further increase dead-space fraction (12–14). Lowering tidal volume also increases dead-space fraction (15), an effect that may be offset partially by a brief end-inspiratory pause with each breath (16).

Determination of dead-space fraction requires measurement of expired CO_2 in a volume of expired gas. This volume is either derived by integrating flow or measured directly by collection in a Douglas bag. Expired CO_2 is measured only infrequently in routine clinical care and clinical trials of ARDS, contributing to underreporting of dead-space fraction.

Absent routine measurement or a reliable estimate, most studies do not account for dead-space fraction. Yet, such secondary analyses of clinical trials influence future research directions and, at times, even clinical decisions. Secondary analyses should account for known independent predictors of the outcome of interest to calculate a valid effect estimate, particularly when the analysis is not performed according to the groups to which patients were originally randomized. Otherwise, the calculated effect—benefit or harm—might be attributable to residual confounding, that is, to other differences between groups that would explain the findings.

To address this gap, we evaluated the validity of four approaches to estimating dead-space fraction. Three approaches required predicting energy expenditure: the unadjusted Harris-Benedict estimate, which employs the

eponymous formula for resting energy expenditure (REE); the Siddiki estimate, which adjusts Harris-Benedict to account for hypermetabolic conditions often encountered in critical illness; and the Penn State estimate, which was derived specifically for use in critically ill patients. We also derived a novel approach to estimate dead-space fraction directly without requiring estimation of energy expenditure as an intermediate step. Dead-space fraction estimates were evaluated for their prediction of measured dead-space fraction and prediction of the association between measured dead-space fraction and mortality. We hypothesized that estimating dead-space fraction directly would yield the highest predictive validity.

METHODS

Study Design

De-identified patient-level data pooled from three randomized controlled trials of early ARDS (V_D/V_T cohort) were used to compare directly measured dead-space fraction with four methods for estimating dead-space fraction. To evaluate the association between estimated dead-space fraction and mortality in a larger population, a second cohort was created by pooling data from completed National Heart, Lung, and Blood Institute Acute Respiratory Distress Syndrome Network [ARDSNet] trials in which dead-space fraction was not measured (ARDSNet Cohort). The study was exempt from review by the institutional review board.

Subjects

Patients eligible for the V_D/V_T cohort had baseline measured dead-space fraction obtained within 24 hours of study enrollment and prior to any study interventions. Patients were enrolled in one of three randomized controlled trials testing therapies for early ARDS. These trials were chosen because dead-space fraction was reported in their primary publications. The ARDSNet Albuterol for the Treatment of Acute Lung Injury Trial (ALTA) (17) was a multicenter randomized, placebo-controlled trial in which patients with early ARDS were assigned to receive aerosolized albuterol (5 mg) or saline placebo every 4 hours for up to 10 days. The trial was stopped early for futility, with no significant difference in mortality or ventilator-free days between groups. Esophageal Pressure-Guided Mechanical Ventilation Trial (18) was a single-center randomized, controlled trial in which patients with early ARDS were assigned to undergo mechanical ventilation with PEEP adjustment guided by esophageal pressure or according to the ARDSNet PEEP titration table. The trial was stopped early for reaching the primary endpoint of improved $\text{PaO}_2:\text{FiO}_2$, with no significant difference in mortality or ventilator-free days in the unadjusted primary analysis. The Activated Protein C for Acute Lung Injury Trial (19) was a multicenter randomized, placebo-controlled trial in which patients with early ARDS were assigned to receive activated protein C 24 $\mu\text{g}/\text{kg}/\text{hr}$ or placebo for 96 hours. No significant difference in mortality or ventilator-free days was found.

For the ARDSNet cohort without measured dead-space fraction available, data were pooled from the following ARDSNet trials: ARMA (higher versus lower tidal volumes, low tidal volume group only) (1), ALVEOLI (higher versus lower PEEP) (20), FACTT (liberal versus conservative fluid management and pulmonary artery versus central venous catheter-guided management) (21, 22), ALTA (only patients without measured dead-space fraction were included in the ARDSNet cohort) (17), OMEGA (omega-3 fatty acid and antioxidant supplementation versus placebo) (23), and EDEN (initial trophic versus full enteral feeding) (24). Inclusion and exclusion criteria for each trial are described in the original referenced publications.

Measurement of Physiological Dead-Space Fraction

In the V_D/V_T cohort, physiological dead-space fraction was calculated by measuring mean expired CO_2 using volumetric capnography according to a validated protocol (25). An arterial blood gas was obtained at the time of expired gas analysis. Measured dead-space fraction was calculated using the Eng-hoff modification to the Bohr equation (26):

$$\frac{V_D}{V_T} = \frac{(\text{PaCO}_2 - \text{P}_E\text{CO}_2)}{\text{PaCO}_2}$$

where PaCO_2 and P_ECO_2 represent partial pressure of CO_2 in arterial blood and expired gas, respectively. All measurements were made prior to study interventions.

Estimating Equations for Physiological Dead-Space Fraction

Methods for estimating dead-space fraction that do not require measurement of expired CO_2 typically depend on the alveolar ventilation equation:

$$\text{PaCO}_2 = \frac{\dot{V}\text{CO}_2 \times 0.863}{\dot{V}_A}$$

where PaCO_2 is measured in mm Hg, $\dot{V}\text{CO}_2$ represents CO_2 production (mL/min), and $\dot{V}_A$ represents alveolar minute ventilation (L/min). Because $\dot{V}_A$ is defined as the difference between total minute ventilation and dead-space minute ventilation, this equation can be rewritten and, after solving for V_D/V_T , yields:

$$\frac{V_D}{V_T} = 1 - \frac{(0.863 \times \dot{V}\text{CO}_2)}{(\text{RR} \times V_T \times \text{PaCO}_2)}$$

where RR is the respiratory rate (breaths/min) and V_T is the tidal volume (liters). In this rearranged equation for dead-space fraction, the only variable not routinely available is $\dot{V}\text{CO}_2$, which may be calculated from the REE using the rearranged Weir equation (27):

$$\dot{V}\text{CO}_2 = \frac{\text{REE}}{\left(\frac{5.616}{\text{RQ}} + 1.584 \right)}$$

where RQ is the respiratory quotient, assumed to be 0.8 for this analysis.

In this study, four different strategies for estimating dead-space fraction were considered. All physiological measurements required for dead-space fraction estimates were obtained prior to study interventions associated with the clinical trial.

1. *Unadjusted Harris-Benedict estimate:* The original sex-specific Harris-Benedict equations (28) were used to estimate REE:

Males:

$$\text{REE}_{\text{HB}} = 66.473 + 13.752 (\text{Wt}) + 5.003 (\text{Ht}) - 6.755 (\text{age})$$

Females:

$$\text{REE}_{\text{HB}} = 655.096 + 9.563 (\text{Wt}) + 1.850 (\text{Ht}) - 4.676 (\text{age})$$

with weight (Wt) in kg, height (Ht) in cm, and age in years. The value for REE_{HB} was inserted into the rearranged Weir equation to calculate $\dot{V}\text{CO}_2$, which was then used to calculate dead-space fraction.

2. *Siddiki estimate:* Siddiki et al (29) proposed using a modified Harris-Benedict equation to estimate REE. In this approach, REE_{HB} is adjusted to account for the hypermetabolic state resulting from certain clinical conditions:

$$\text{REE}_{\text{Siddiki}} = \text{REE}_{\text{HB}} \times \text{hf}$$

where hf is a unitless multiplier term for hypermetabolic factors with potential values of 1.13 per °C above 37°C, 1.2 for minor surgery, 1.35 for major trauma, and 1.6 for severe infection. The hypermetabolic factor yielding the highest value for hf is selected to calculate $\text{REE}_{\text{Siddiki}}$. The value for $\text{REE}_{\text{Siddiki}}$ was inserted into the rearranged Weir equation to calculate $\dot{V}\text{CO}_2$, which was then used to calculate dead-space fraction. In the original report of Siddiki et al (29), the rearranged Weir equation for $\dot{V}\text{CO}_2$ differs trivially from that above due to rounding.

3. *Penn State estimate:* An alternative formulation for estimating REE was derived previously by Frankenfield et al (30, 31) specifically for critically ill patients. This approach uses the Mifflin-St. Jeor equation (32) to estimate REE for the patient in good health:

Males:

$$\text{REE}_{\text{MSJ}} = 10(\text{Wt}) + 6.25(\text{Ht}) - 5(\text{age}) + 5$$

Females:

$$\text{REE}_{\text{MSJ}} = 10(\text{Wt}) + 6.25(\text{Ht}) - 5(\text{age}) - 161$$

with weight (Wt) in kg, height (Ht) in cm, and age in years. Additional clinical variables are then incorporated to yield the Penn State equations for REE in critical illness (30, 31): If BMI < 30 kg/m²:

$$\text{REE}_{\text{PSU}} = 0.96 (\text{REE}_{\text{MSJ}}) + 31(\text{RR})(V_T) + 167(T_{\text{max}}) - 6212$$

If $\text{BMI} \geq 30 \text{ kg/m}^2$:

$$\text{REE}_{\text{PSU}} = 0.71(\text{REE}_{\text{MSI}}) + 64(\text{RR})(V_T) + 85(T_{\text{max}}) - 3085$$

where BMI is the body mass index, RR is the respiratory rate (breaths/min), V_T is the tidal volume (liters), and T_{max} is the maximum temperature ($^{\circ}\text{C}$) over the last 24 hours.

4. *Direct estimate from physiological variables:* A novel alternative approach to estimate dead-space fraction directly was developed using least angle regression to derive a prediction model. Only variables with physiological plausibility were considered for inclusion in the model: anthropometrics (height, measured body weight, predicted body weight, body mass index, body surface area, sex, age, and race/ethnicity), respiratory variables (tidal volume, tidal volume per predicted body weight, respiratory rate, minute ventilation, minute ventilation per predicted body weight, PaCO_2 , $\text{PaO}_2/\text{FiO}_2$, respiratory system compliance, PEEP, number of quadrants with infiltrates on chest imaging, and Murray lung injury score) (33), hemodynamic variables that may affect ventilation/perfusion matching (systolic blood pressure, mean arterial pressure, shock as defined by the Brussels criteria (34), heart rate, and rate-pressure product), primary cause of lung injury, and maximum temperature over the previous 24 hours. Clinically relevant multiplicative interaction terms were also considered, consisting of PaCO_2 and each of: minute ventilation, minute ventilation per predicted body weight, body mass index, body surface area, measured body weight, height, sex, age, and temperature.

Model building used least angle regression with five-fold cross-validation to minimize cross-validated mean squared prediction error. For parsimony, only the first five variables were retained in the final model since mean squared prediction error improved minimally with additional variables. The final model was refit using ordinary least squares to derive the reported coefficients:

$$\begin{aligned} V_D / V_T = & 0.1726 + 0.0059(\text{RR}) + 0.0054(\text{PEEP}) + 0.0293(\text{LIS}) \\ & + 0.0036(\text{PaCO}_2 \times \dot{V}_E) + 0.000057(\text{PaCO}_2 \times \text{age}) \end{aligned}$$

where RR is the respiratory rate (breaths/min), PEEP represents set PEEP (cm H_2O) on the mechanical ventilator, LIS is the Murray lung injury score (33), PaCO_2 is measured in mm Hg, and $\dot{V}_E$ represents total minute ventilation (L/min).

Comparison of Approaches

Dead-space fraction estimates were evaluated based on two overarching criteria: prediction of measured dead-space fraction and prediction of the association between measured dead-space fraction and mortality. Measured and estimated dead-space fraction were compared graphically using the Bland-Altman approach for assessing agreement between methods of clinical measurement (35). Quantitatively, methods were compared according to bias and accuracy. Bias describes

whether the estimate systematically underpredicts or overpredicts measured dead-space fraction and was determined by comparing the difference in means between measurement and each estimating equation. A one-sample t test was performed to determine if the mean difference was significantly different from zero. Accuracy describes how close each estimated value for dead-space fraction is to the true measured value and was calculated in two ways. First, the 95th percentile of the absolute difference between measured and estimated values was calculated; the absolute difference was used to avoid canceling effects of negative and positive values. Second, the proportion of estimated dead-space fraction values that fell within ± 0.10 or ± 0.20 of measured dead-space fraction was calculated.

Measured and estimated dead-space fraction were also compared for their ability to predict 28-day mortality. Mortality per dead-space fraction quintile was used to evaluate whether the predictive ability of each dead-space fraction estimate varied by level. Simple logistic regression was used to compare the unadjusted association between each dead-space fraction estimate and mortality. In a sensitivity analysis, measured dead-space fraction was then added to each model to determine whether the association between estimated dead-space fraction and mortality included effects beyond that explained by measured dead-space fraction. Multivariable logistic regression was performed to evaluate the relationship between each dead-space fraction estimate and mortality. First, backward elimination (threshold $p \leq 0.05$ to remain in model) was used to construct the best-fitting model of mortality with measured dead-space fraction, selecting from the following candidate predictors: age, shock, Acute Physiology and Chronic Health Evaluation (APACHE) II, tidal volume per predicted body weight, PEEP, primary cause of lung injury, and respiratory system compliance. Berlin ARDS severity (3) and clinical trial enrolled were forced into the model as categorical variables for face validity. The selected covariates were then used to fit logistic models for each estimate of dead-space fraction. Finally, the area under the receiver operating curve (AUROC) for logistic regression models of mortality was used to determine whether measured dead-space fraction and the best-performing estimate improved predictive validity of the Berlin definition. Statistical analyses were performed using SAS 9.3 (SAS Institute, Cary, NC). A two-sided p value less than or equal to 0.05 was considered statistically significant.

RESULTS

A total of 210 patients in the V_D/V_T cohort were included in the primary analysis. In this cohort, measured dead-space fraction was markedly elevated (mean $\pm$ SD, 0.60 ± 0.12). An additional 3,135 patients enrolled in the ARDSNet trials did not have measured dead-space fraction and were included in the second cohort. Patient characteristics for both cohorts are described in **Table 1**. Overall, mortality at 28 days was significantly lower in the V_D/V_T cohort (16% vs 22%; $p = 0.05$).

TABLE 1. Characteristics of Included Patients With (V_D/V_T Cohort) and Without (ARDSNet Cohort) Measured Dead-Space Fraction

Patient Characteristic	V _D /V _T Cohort				ARDSNet Cohort (n = 3,135)
	ALTA (n = 92)	EPVent (n = 57)	APC (n = 61)	Total (n = 210)	
Demographics					
Age	50 ± 17	54 ± 19	52 ± 18	52 ± 18	51 ± 16
Female (%)	46 (50)	22 (39)	26 (43)	94 (45)	1,439 (46)
Nonwhite race (%)	28 (30)	7 (12)	18 (30)	53 (25)	871 (28)
Body mass index (kg/m²)	29.0 ± 9.0	30.7 ± 9.3	28.3 ± 7.4	29.3 ± 8.7	28.7 ± 7.6
Clinical characteristics					
Acute Physiology and Chronic Health Evaluation II	26 ± 7	26 ± 6	20 ± 7	25 ± 7	27 ± 7
Shock (%)	59 (64)	33 (58)	16 (26)	108 (51)	1,733 (55)
Primary cause of lung injury (%)					
Aspiration or pneumonia	50 (54)	8 (14)	44 (72)	102 (49)	1,952 (62)
Sepsis	24 (26)	14 (25)	9 (15)	47 (22)	659 (21)
Multiple transfusions	2 (2)	0	0	2 (1)	65 (2)
Trauma	8 (9)	15 (26)	0	23 (11)	223 (7)
Other	8 (9)	20 (35)	8 (13)	36 (17)	235 (8)
Respiratory characteristics					
Tidal volume (mL)	390 ± 70	493 ± 96	415 ± 92	425 ± 94	408 ± 94
Tidal volume (mL/kg predicted body weight)	6.3 ± 0.9	7.7 ± 1.4	6.7 ± 1.4	6.8 ± 1.3	6.5 ± 1.4
Respiratory rate (breaths/min)	28 ± 7	25 ± 6	27 ± 7	27 ± 7	27 ± 7
Positive end-expiratory pressure (cm H ₂ O)	9 ± 3	13 ± 4	9 ± 3	10 ± 4	10 ± 4
Pao ₂ /Fio ₂	184 ± 63	148 ± 57	160 ± 63	167 ± 63	172 ± 76
Paco ₂ (mm Hg)	40 ± 8	41 ± 8	41 ± 11	41 ± 9	41 ± 10
Respiratory system compliance (mL/cm H ₂ O)	32 ± 17	36 ± 11	31 ± 22	33 ± 17	32 ± 17
Murray lung injury score	2.4 ± 0.6	2.8 ± 0.6	2.6 ± 0.6	2.6 ± 0.6	2.6 ± 0.7
V _D /V _T measured	0.57 ± 0.12	0.67 ± 0.10	0.59 ± 0.12	0.60 ± 0.12	Not applicable
Clinical outcomes					
Mortality (%)	12 (13)	15 (26)	7 (11)	34 (16)	693 (22)
Vent-free days through day 28	16 ± 10	10 ± 9	15 ± 10	14 ± 10	14 ± 11

V_D/V_T = dead-space fraction, ARDSNet = Acute Respiratory Distress Syndrome Network, ALTA = Albuterol for the Treatment of Acute Lung Injury Trial, EPVent = Esophageal Pressure-Guided Mechanical Ventilation Trial, APC = Activated Protein C for Acute Lung Injury Trial.

Data are reported as mean ± SD or n (%).

Performance of Estimating Equations

Bias was evaluated by comparing mean differences between estimated and measured dead-space fraction (Table 2). Both the Siddiki and Penn State estimates were significantly biased toward underestimation of measured dead-space fraction (mean difference, -0.32 ± 0.35 and -0.08 ± 0.12 , respectively; $p < 0.01$ for both comparisons). The unadjusted Harris-Benedict and direct estimates were unbiased (mean difference, -0.01 ± 0.12 and 0 ± 0.09 , respectively; $p = 0.30$ for unadjusted Harris-Benedict estimate; direct estimate unbiased by design).

Accuracy was evaluated first as the 95th percentile of the absolute difference between measured and estimated dead-space fraction (Table 2). The unadjusted Harris-Benedict and direct estimates were the most accurate estimates of dead-space fraction (95th percentile, 0.20 and 0.17, respectively). The Penn State estimate displayed intermediate accuracy relative to the other formulations (95th percentile, 0.30). The Siddiki estimate was the least accurate (95th percentile, 1.03).

Accuracy was also evaluated as the proportion of dead-space fraction estimates within two prespecified thresholds

TABLE 2. Performance of Estimating Equations Compared to Measured Dead-Space Fraction in V_D/V_T Cohort

Measure of Estimating Equation Performance	Method of Dead-Space Estimation			
	Harris-Benedict	Siddiki	Penn State	Direct
V_D/V_T mean $\pm$ SD, all patients	0.59 $\pm$ 0.13	0.28 $\pm$ 0.38	0.52 $\pm$ 0.14	0.60 $\pm$ 0.08
Survivors	0.58 $\pm$ 0.14	0.25 $\pm$ 0.39	0.50 $\pm$ 0.14	0.59 $\pm$ 0.07
Nonsurvivors	0.67 $\pm$ 0.11	0.47 $\pm$ 0.28	0.59 $\pm$ 0.11	0.66 $\pm$ 0.08
Bias: difference between measured and estimated V_D/V_T mean $\pm$ SD	-0.01 $\pm$ 0.12	-0.32 $\pm$ 0.35 ^a	-0.08 $\pm$ 0.12 ^a	0 $\pm$ 0.09 ^{a,b}
Accuracy: 95th percentile of absolute difference between measured and estimated V_D/V_T	0.20	1.03	0.30	0.17
V_D/V_T accuracy threshold of $\pm$ 0.10				
Accuracy, %	70.1	26.5	52.5	73.0
Values overestimating, %	15.7	3.4	4.9	13.7
Values underestimating, %	14.2	70.1	42.6	13.2
V_D/V_T accuracy threshold of $\pm$ 0.20				
Accuracy, %	95.1	44.6	85.3	98.5
Values overestimating, %	1.5	1.0	0.5	0.5
Values underestimating, %	3.4	54.4	14.2	1.0
Correlation coefficient	0.58 ^a	0.45 ^a	0.56 ^a	0.67 ^a

V_D/V_T = dead-space fraction.

^a $p < 0.01$ between estimated and measured dead-space fraction.

^bDirect estimate was unbiased by design.

of measured dead-space fraction: ± 0.10 and ± 0.20 (Table 2). Again, the unadjusted Harris-Benedict and direct estimates displayed the best performance, with 70.1% and 73.0% of estimates within ± 0.10 of measured dead-space fraction, respectively, and greater than 95% of estimates within ± 0.20 for both formulations. By contrast, the Siddiki and Penn State estimates were considerably less accurate, predicting dead-space fraction to within ± 0.10 of the measured value only 26.5% and 52.5% of the time. The Siddiki estimate alone yielded negative values for dead-space fraction, whereas no estimate produced values greater than or equal to 1.00. Graphical comparison of each estimate with measured dead-space fraction using Bland-Altman plots confirmed these findings (Fig. 1).

Prediction of Mortality

Measured dead-space fraction was significantly higher in nonsurvivors compared with survivors (0.67 ± 0.12 vs 0.59 ± 0.12 ; $p < 0.01$). Estimated dead-space fraction was also significantly higher among nonsurvivors for all estimating equations ($p < 0.01$ for all comparisons) (Table 2).

Mortality by quintile of measured and estimated dead-space fraction is reported in Figure 2. Quintiles were calculated separately for each formulation of dead-space fraction to account for potential differences in scaling. Mortality per unadjusted Harris-Benedict quintile was within $\pm 6.7\%$ of that

per measured dead-space fraction for all quintiles (Fig. 2A). No other estimating equation as closely approximated the measured dead-space fraction per-quintile mortality. In the ARDSNet cohort, mortality similarly increased with successive quintiles of each dead-space fraction estimate (Fig. 2B).

In the unadjusted analysis, higher measured dead-space fraction was significantly associated with increased risk of death (Table 3). The multivariable model-building process identified measured dead-space fraction, APACHE II, and PEEP as statistically significant independent predictors of death, with clinical trial enrolled and Berlin ARDS severity forced into the model for face validity but not reaching statistical significance. In the multivariable analysis, for every 0.05 increase in measured dead-space fraction, odds of death increased by 36% (odds ratio [OR], 1.36; 95% CI, 1.10–1.68; $p < 0.01$).

Each estimate of dead-space fraction was also significantly associated with mortality on unadjusted and multivariable analyses in both cohorts ($p < 0.01$ for all analyses). However, ORs for death varied considerably (Table 3) due to differences in scaling and accuracy of dead-space fraction estimates and variation in predicting the relationship between measured dead-space fraction and mortality. In the V_D/V_T cohort, the Harris-Benedict and Penn State ORs for death were most similar to that of measured dead-space fraction in both the unadjusted and multivariable analyses (multivariable model, OR_{death} per 0.05 increase in dead-space fraction; Harris-Benedict OR,

Figure 1. Bland-Altman analyses comparing measured dead-space fraction with that determined by each estimating equation: Harris-Benedict (A), Siddiki (B), Penn State (C), and direct estimate (D). Plots consist of the difference between estimated and measured dead-space fraction formulations (vertical axis) versus the average value of the two approaches (horizontal axis). Differences greater than zero indicate overestimation.

1.55; 95% CI, 1.21–1.98; Penn State OR, 1.53; 95% CI, 1.21–1.95). The Siddiki OR was considerably lower (OR, 1.19; 95% CI, 1.07–1.31), whereas that of the direct estimate was higher (OR, 1.80; 95% CI, 1.26–2.56). In the ARDSNet cohort, the Harris-Benedict and Penn State ORs for death were similar to each other, whereas the Siddiki and direct estimate ORs differed considerably.

In the sensitivity analysis, only the direct estimate remained significantly associated with mortality after adding measured dead-space fraction to the model, indicating that the direct estimate had an association with mortality that was independent of measured dead-space fraction.

Compared with modeling mortality using the Berlin definition alone, adding measured dead-space fraction significantly improved predictive validity in the V_D/V_T cohort, with an AUROC of 0.689 (95% CI, 0.587–0.791) versus 0.534 (95% CI, 0.440–0.628; $p = 0.02$). Similarly, adding the unadjusted Harris-Benedict estimate to the Berlin definition significantly improved predictive validity for mortality in both the V_D/V_T cohort

(AUROC, 0.714; 95% CI, 0.616–0.813 vs 0.543; 95% CI, 0.449–0.637; $p < 0.01$) and the ARDSNet cohort (AUROC, 0.644; 95% CI, 0.617–0.672 vs 0.592, 95% CI, 0.566–0.618; $p < 0.01$).

Prediction of Outlying Estimates

A post hoc analysis was conducted to identify the subset of patients in whom the unadjusted Harris-Benedict estimate did not predict measured dead-space fraction accurately, to within ± 0.10 . Among baseline characteristics, only $Paco_2$ differed significantly between patients with inaccurate compared with accurate estimates (39 ± 9 vs 41 ± 9 , respectively; $p = 0.04$). Nearly half of the patients (43%) with $Paco_2$ less than 30 mm Hg had an inaccurate Harris-Benedict estimate, compared with just 28% of patients with $Paco_2$ greater than or equal to 30 mm Hg ($p = 0.17$). Considering only underestimates, $Paco_2$ was less than 30 mm Hg in 24% of patients; by contrast, only 8% of patients in whom the Harris-Benedict approach did not underestimate dead-space fraction had $Paco_2$ less than 30 mm Hg ($p < 0.01$). Such marked hypocapnia occurred infrequently

Figure 2. Observed mortality by quintile of measured or estimated dead-space fraction: among patients with measured dead-space fraction (V_D/V_T cohort) (A) and among patients without measured dead-space fraction (Acute Respiratory Distress Syndrome Network cohort) (B). Quintiles were calculated separately for each formulation of dead-space fraction.

in our cohorts (in 10% and 9% of patients in the V_D/V_T and ARDSNet cohorts, respectively), as would be expected during lung-protective ventilation for ARDS. No comparable pattern was seen with overestimation.

DISCUSSION

The present study demonstrates that dead-space fraction is best estimated using the unadjusted Harris-Benedict equation for energy expenditure to predict measured dead-space fraction and its association with mortality. The Harris-Benedict and direct estimates predicted measured dead-space fraction most accurately. However, the direct estimate was associated with death independent of measured dead-space fraction, indicating a relationship with mortality beyond that explained by measured dead-space fraction. Still, even the Harris-Benedict

approach, the best-performing estimate, predicted measured dead-space fraction to within ± 0.10 in only 70% of patients and ± 0.20 in 95% of patients—a considerable range given the scale of measure. Therefore, estimates of dead-space fraction should not replace prospective measurement in future clinical trials of ARDS. For secondary analyses of existing clinical trials data where dead-space fraction was not measured, the unadjusted Harris-Benedict estimate can be used to estimate the independent association between dead-space fraction and mortality.

Increased dead-space fraction is a clinical hallmark of ARDS that independently predicts patient outcomes (3, 4). For this reason, dead-space fraction was considered in formulating the 2012 Berlin definition of ARDS to lend further face validity to the definition (36). However, dead-space fraction was not included in the final Berlin definition because it is not routinely measured (36). The present study found that adjusting for measured dead-space fraction in the Berlin definition significantly improved predictive validity for mortality. Adjusting for estimated dead-space fraction using the Harris-Benedict approach similarly improved the Berlin definition's predictive validity for mortality (3). These findings reinforce the need to adjust for dead-space fraction as a marker of ARDS severity that independently predicts mortality.

Secondary analyses of epidemiological and clinical trials data are a mainstay of ARDS research due to the expense and resources required to conduct clinical trials (37). Because such secondary analyses play a central role in shaping future directions of ARDS research and clinical practice, it is essential they incorporate known independent predictors of ARDS mortality. Oxygenation, respiratory system compliance, and dead-space fraction have been identified repeatedly as predictors of clinical outcomes from ARDS (2, 4, 7). While oxygenation and respiratory system mechanics are frequently considered, dead-space fraction is rarely measured in such studies and thus altogether ignored in secondary analyses.

Few prior reports have attempted to estimate dead-space fraction in ARDS. The Siddiki estimate (29) was shown previously in a study of just 13 patients to substantially underestimate measured dead-space fraction (38), a finding confirmed here in a much larger cohort. In fact, the Siddiki estimate produced negative values for dead-space fraction in some patients because its use of hypermetabolic factors led to overestimating $\dot{V}CO_2$ (39). Frankenfield et al (40) derived a predictive equation for dead-space fraction requiring measurement of end-tidal CO_2 in a heterogeneous population of critically ill patients. Their study excluded patients with FiO_2 greater than 0.60 due to equipment limitations, limiting generalizability to ARDS cohorts. Furthermore, end-tidal CO_2 is not routinely captured in clinical trials or epidemiological data, limiting utility of this approach for the purpose of secondary analyses of existing data.

Alternative surrogates for dead-space fraction have been proposed. The ARDS Berlin Definition Task Force considered total minute ventilation standardized to a $Paco_2$ of 40 mm Hg (3, 41). Post hoc analysis of the Berlin cohort found that stratifying patients with severe ARDS ($PaO_2:FiO_2 \leq 100$) by standardized

TABLE 3. Odds Ratios (95% CIs) for Death per 0.05 Increase in Measured or Estimated Dead-Space Fraction^a

Regression Model	Measured	Harris-Benedict	Siddiki	Penn State	Direct
<i>V_D/V_T</i> cohort					
Unadjusted analysis	1.36 (1.14–1.62)	1.40 (1.14–1.71)	1.12 (1.04–1.22)	1.36 (1.13–1.63)	1.74 (1.34–2.27)
Multivariable analysis ^b	1.36 (1.10–1.68)	1.55 (1.21–1.98)	1.19 (1.07–1.31)	1.53 (1.21–1.95)	1.80 (1.26–2.56)
Acute Respiratory Distress Syndrome Network cohort					
Unadjusted analysis	Not measured	1.16 (1.11–1.20)	1.04 (1.02–1.05)	1.14 (1.10–1.18)	1.47 (1.39–1.56)
Multivariable analysis ^b	Not measured	1.10 (1.05–1.14)	1.03 (1.02–1.05)	1.09 (1.04–1.13)	1.41 (1.30–1.53)

^aOdds ratios for all formulations of dead-space fraction differ significantly from 1.0 with $p < 0.01$ for all analyses.

^bMultivariable model adjusted for Acute Physiology and Chronic Health Evaluation, Berlin Acute Respiratory Distress Syndrome severity, positive end-expiratory pressure, and clinical trial enrolled.

minute ventilation significantly improved mortality risk prediction (3). Sinha et al (42) proposed a similar formulation, the ventilatory ratio, which includes predicted minute ventilation and predicted PaCO_2 . Both approaches may be useful for bedside contemplation of minute ventilation requirements, but neither has been validated as a surrogate for measured dead-space fraction to support use in clinical research.

Important limitations to this study exist. First, estimated dead-space fraction is not intended for use in place of direct measurement in clinical practice or future prospective studies. Rather, this report highlights the need to incorporate measurement of dead-space fraction in future clinical trials and prospectively collected observational data. When measured dead-space fraction is unavailable, the unadjusted Harris-Benedict estimate may be considered for research purposes. The unadjusted Harris-Benedict estimate tends to underestimate dead-space fraction when PaCO_2 is less than 30 mm Hg. Thus, caution should be used when a large proportion of the study population has marked hypocapnia.

Second, the best-performing estimate here relies on the unadjusted Harris-Benedict equation for energy expenditure and assumes a respiratory quotient of 0.8 for all patients. Respiratory quotient fluctuates with feeding, nutritional status, and anaerobiosis (43, 44), factors not considered here. The unadjusted Harris-Benedict equation has been shown previously to be unreliable in predicting energy expenditure in critically ill patients (39), which in part may be due to variation in ventilator settings (15, 45, 46), metabolic stress (47, 48), and fasting status (44) in prior studies. By contrast, a singular focus on early ARDS and protocolized low tidal volume ventilation here likely improved its performance in estimating dead-space fraction. It is unclear how the unadjusted Harris-Benedict dead-space fraction estimate would perform in cohorts managed with different ventilator settings, prone positioning, or experimental interventions.

Finally, differences in mechanical ventilation practices directly affect measured dead-space fraction and may limit its utility as a marker of disease severity absent comparable settings. Tidal volume, PEEP titration, dynamic hyperinflation, and use of a brief end-inspiratory pause all have been shown

to affect measured dead-space fraction irrespective of underlying disease severity (12–16). In this study, these effects likely were mitigated in part by use of protocolized lung-protective ventilation for all included patients, though between-patient variation still occurred (Table 1).

CONCLUSIONS

Dead-space fraction is one of few lung-specific independent predictors of mortality from ARDS. As such, dead-space fraction should be measured whenever possible in clinical trials and prospective epidemiological studies to permit adjustment for its effects in secondary analyses. For analyses of existing data where dead-space fraction was not measured, the unadjusted Harris-Benedict estimate should be considered to estimate the association between dead-space fraction and mortality.

REFERENCES

1. Acute Respiratory Distress Syndrome Network: Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000; 342:1301–1308
2. Seeley E, McAuley DF, Eisner M, et al: Predictors of mortality in acute lung injury during the era of lung protective ventilation. *Thorax* 2008; 63:994–998
3. Ranieri VM, Rubenfeld GD, Thompson BT, et al: ARDS Definition Task Force: Acute respiratory distress syndrome: The Berlin Definition. *JAMA* 2012; 307:2526–2533
4. Nuckton TJ, Alonso JA, Kallet RH, et al: Pulmonary dead-space fraction as a risk factor for death in the acute respiratory distress syndrome. *N Engl J Med* 2002; 346:1281–1286
5. Kallet RH, Alonso JA, Pittet JF, et al: Prognostic value of the pulmonary dead-space fraction during the first 6 days of acute respiratory distress syndrome. *Respir Care* 2004; 49:1008–1014
6. Cepkova M, Kapur V, Ren X, et al: Pulmonary dead space fraction and pulmonary artery systolic pressure as early predictors of clinical outcome in acute lung injury. *Chest* 2007; 132:836–842
7. Hager DN, Krishnan JA, Hayden DL, et al: ARDS Clinical Trials Network: Tidal volume reduction in patients with acute lung injury when plateau pressures are not high. *Am J Respir Crit Care Med* 2005; 172:1241–1245
8. Raurich JM, Vilar M, Colomar A, et al: Prognostic value of the pulmonary dead-space fraction during the early and intermediate phases of acute respiratory distress syndrome. *Respir Care* 2010; 55:282–287

9. Tomashefski JF Jr, Davies P, Boggis C, et al: The pulmonary vascular lesions of the adult respiratory distress syndrome. *Am J Pathol* 1983; 112:112–126
10. Calfee CS, Gallagher D, Abbott J, et al; NHLBI ARDS Network: Plasma angiopoietin-2 in clinical acute lung injury: Prognostic and pathogenetic significance. *Crit Care Med* 2012; 40:1731–1737
11. Günther A, Mosavi P, Heinemann S, et al: Alveolar fibrin formation caused by enhanced procoagulant and depressed fibrinolytic capacities in severe pneumonia. Comparison with the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2000; 161:454–462
12. Suter PM, Fairley B, Isenberg MD: Optimum end-expiratory airway pressure in patients with acute pulmonary failure. *N Engl J Med* 1975; 292:284–289
13. Vieillard-Baron A, Prin S, Augarde R, et al: Increasing respiratory rate to improve CO₂ clearance during mechanical ventilation is not a panacea in acute respiratory failure. *Crit Care Med* 2002; 30:1407–1412
14. Tusman G, Suarez-Sipmann F, Böhm SH, et al: Monitoring dead space during recruitment and PEEP titration in an experimental model. *Intensive Care Med* 2006; 32:1863–1871
15. Kiiski R, Kaitainen S, Karppi R, et al: Physiological effects of reduced tidal volume at constant minute ventilation and inspiratory flow rate in acute respiratory distress syndrome. *Intensive Care Med* 1996; 22:192–198
16. Devaquet J, Jonson B, Niklason L, et al: Effects of inspiratory pause on CO₂ elimination and arterial PCO₂ in acute lung injury. *J Appl Physiol* (1985) 2008; 105:1944–1949
17. Matthay MA, Brower RG, Carson S, et al: Randomized, placebo-controlled clinical trial of an aerosolized β_2 -agonist for treatment of acute lung injury. *Am J Respir Crit Care Med* 2011; 184:561–568
18. Talmor D, Sarge T, Malhotra A, et al: Mechanical ventilation guided by esophageal pressure in acute lung injury. *N Engl J Med* 2008; 359:2095–2104
19. Liu KD, Levitt J, Zhuo H, et al: Randomized clinical trial of activated protein C for the treatment of acute lung injury. *Am J Respir Crit Care Med* 2008; 178:618–623
20. National Heart, Lung, and Blood Institute ARDS Clinical Trials Network: Higher versus lower positive end-expiratory pressures in patients with the acute respiratory distress syndrome. *N Engl J Med* 2004; 351:327–336
21. National Heart, Lung, and Blood Institute ARDS Clinical Trials Network: Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med* 2006; 354:2564–2575
22. Wheeler AP, Bernard GR, Thompson BT, et al: Pulmonary-artery versus central venous catheter to guide treatment of acute lung injury. *N Engl J Med* 2006; 354:2213–2224
23. Rice TW, Wheeler AP, Thompson BT, et al; NIH NHLBI Acute Respiratory Distress Syndrome Network of Investigators: Enteral omega-3 fatty acid, gammalinolenic acid, and antioxidant supplementation in acute lung injury. *JAMA* 2011; 306:1574–1581
24. Rice TW, Wheeler AP, Thompson BT, et al; National Heart, Lung, and Blood Institute ARDS Clinical Trials Network: Initial trophic vs full enteral feeding in patients with acute lung injury: The EDEN randomized trial. *JAMA* 2012; 307:795–803
25. Kallet RH, Daniel BM, Garcia O, et al: Accuracy of physiologic dead space measurements in patients with acute respiratory distress syndrome using volumetric capnography: Comparison with the metabolic monitor method. *Respir Care* 2005; 50:462–467
26. Sinha P, Flower O, Soni N: Dead-space ventilation: A waste of breath! *Intensive Care Med* 2011; 37:735–746
27. Weir JB: New methods for calculating metabolic rate with special reference to protein metabolism. *J Physiol* 1949; 109:1–9
28. Harris JA, Benedict FG: A biometric study of human basal metabolism. *Proc Natl Acad Sci U S A* 1918; 4:370–373
29. Siddiki H, Kojicic M, Li G, et al: Bedside quantification of dead-space fraction using routine clinical data in patients with acute lung injury: Secondary analysis of two prospective trials. *Crit Care* 2010; 14:R141
30. Frankenfield DC, Coleman A, Alam S, et al: Analysis of estimation methods for resting metabolic rate in critically ill adults. *JPEN J Parenter Enteral Nutr* 2009; 33:27–36
31. Frankenfield D: Validation of an equation for resting metabolic rate in older obese, critically ill patients. *JPEN J Parenter Enteral Nutr* 2011; 35:264–269
32. Mifflin MD, St Jeor ST, Hill LA, et al: A new predictive equation for resting energy expenditure in healthy individuals. *Am J Clin Nutr* 1990; 51:241–247
33. Murray JF, Matthay MA, Luce JM, et al: An expanded definition of the adult respiratory distress syndrome. *Am Rev Respir Dis* 1988; 138:720–723
34. Bernard G: The Brussels score. *Sepsis* 1997; 1:43–44
35. Bland JM, Altman DG: Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986; 1:307–310
36. Ferguson ND, Fan E, Camporota L, et al: The Berlin definition of ARDS: An expanded rationale, justification, and supplementary material. *Intensive Care Med* 2012; 38:1573–1582
37. Spragg RG, Bernard GR, Checkley W, et al: Beyond mortality: Future clinical research in acute lung injury. *Am J Respir Crit Care Med* 2010; 181:1121–1127
38. Charron C, Repesse X, Bouferrache K, et al: PaCO₂ and alveolar dead space are more relevant than PaO₂/FiO₂ ratio in monitoring the respiratory response to prone position in ARDS patients: A physiological study. *Crit Care* 2011; 15:R175
39. Frankenfield D, Hise M, Malone A, et al; Evidence Analysis Working Group: Prediction of resting metabolic rate in critically ill adult patients: Results of a systematic review of the evidence. *J Am Diet Assoc* 2007; 107:1552–1561
40. Frankenfield DC, Alam S, Bekteshi E, et al: Predicting dead space ventilation in critically ill patients using clinically available data. *Crit Care Med* 2010; 38:288–291
41. Wexler HR, Lok P: A simple formula for adjusting arterial carbon dioxide tension. *Can Anaesth Soc J* 1981; 28:370–372
42. Sinha P, Sanders RD, Soni N, et al: Acute respiratory distress syndrome: The prognostic value of ventilatory ratio—A simple bedside tool to monitor ventilatory efficiency. *Am J Respir Crit Care Med* 2013; 187:1150–1153
43. Kan MN, Chang HH, Sheu WF, et al: Estimation of energy requirements for mechanically ventilated, critically ill patients using nutritional status. *Crit Care* 2003; 7:R108–R115
44. Zauner C, Schneeweiss B, Kranz A, et al: Resting energy expenditure in short-term starvation is increased as a result of an increase in serum norepinephrine. *Am J Clin Nutr* 2000; 71:1511–1515
45. Kiiski R, Takala J, Kari A, et al: Effect of tidal volume on gas exchange and oxygen transport in the adult respiratory distress syndrome. *Am Rev Respir Dis* 1992; 146:1131–1135
46. Brandi LS, Bertolini R, Santini L, et al: Effects of ventilator resetting on indirect calorimetry measurement in the critically ill surgical patient. *Crit Care Med* 1999; 27:531–539
47. van Lanschot JJ, Feenstra BW, Vermeij CG, et al: Calculation versus measurement of total energy expenditure. *Crit Care Med* 1986; 14:981–985
48. Kiiski R, Takala J: Hypermetabolism and efficiency of CO₂ removal in acute respiratory failure. *Chest* 1994; 105:1198–1203