

UC Irvine

UC Irvine Previously Published Works

Title

Apoptosis in the gastric mucosa: molecular mechanisms, basic and clinical implications.

Permalink

<https://escholarship.org/uc/item/3qn1h0wg>

Journal

Journal of physiology and pharmacology : an official journal of the Polish Physiological Society, 51(1)

ISSN

0867-5910

Authors

Szabó, I
Tarnawski, A S

Publication Date

2000-03-01

Peer reviewed

Review article

I. SZABÓ, A.S. TARNAWSKI

APOPTOSIS IN THE GASTRIC MUCOSA: MOLECULAR MECHANISMS, BASIC AND CLINICAL IMPLICATIONS

Medical Service, Department of Veterans Affairs Medical Center, Long Beach, Department of
Medicine, University of California, Irvine, California, USA,

Apoptosis a programmed cell death, is an essential mechanism of eliminating damaged or aged cells and thus to maintain tissue integrity. There are two central pathways that lead to apoptosis: a) the positive induction by ligands (death factors) binding to plasma membrane receptors (death factor receptors) and b) negative induction by the loss of suppressor activity. The common execution mechanisms of apoptosis consist of the activation of cytosolic aspartate-specific proteases (ICE-proteases) termed caspases, which can be activated via various intracellular pathways. In the stomach, mucosal surface epithelial cells are constantly exfoliating to the gastric lumen and completely replaced within 3–5 days under physiological conditions. Apoptosis has been reported to take place in all regions of the stomach with apoptotic cells occurring predominantly in the superficial parts of the gastric glands, at a rate of 2–3% for all cells. Following mucosal injury (e.g. ulcer development), apoptosis rapidly increases and remains elevated for 2–3 months. In a 3-month old ulcer scar, the apoptosis rate of mucous, parietal, chief and endocrine cells was found to be similar to that of normal gastric mucosa. *Helicobacter pylori* (*H. pylori*) infection induces apoptosis in the gastric mucosa and this action appears to be independent of VacA cytotoxin of *H. pylori* strains. Nonsteroidal anti-inflammatory drugs (NSAIDs), especially cyclooxygenase-2 (COX-2) inhibitors are potent inducers of gastric epithelial cell apoptosis. However, they can abrogate apoptosis or proliferation effects induced by *H. pylori*. Many details of the exact intracellular and molecular mechanisms regulating apoptosis in gastric mucosa remain to be elucidated.

Key words: *apoptosis, gastric mucosa, Helicobacter pylori, NSAIDs, caspases.*

INTRODUCTION

Tissue integrity depends on a balance between the cell renewal and the death of damaged or aged cells. Gastric mucosa is a tissue with a high renewal

This study was supported by the Medical Research Service of the Department of Veterans Affairs. Dr. I. Szabo has been a visiting scientist from the Department of Medicine, University of Pecs, Hungary

rate. For example, it takes approximately 3–5 days to replace all surface epithelial cells (1). Apoptosis — a programmed cell death — is the last stage of a cell destiny (2–4). It is a controlled disassembly of a cell, which does not affect neighbouring cells and does not induce an inflammatory responses (1). Apoptosis can be triggered by a variety of factors e.g. UV or γ -radiation, heat shock, DNA injury, protease inhibitors, and other factors listed in *Fig. 1*.

Factors and injurious stimuli inducing apoptosis

Proinflammatory cytokines: TNF α , IL-1, IL-16, Lymphotoxin β
 Other death factors: Fas Ligand, TRAIL
 Genetic factors: Sek1 null mutation
 UV- and γ -radiation
 Heat shock, hypoxia
 Withdrawal of hormones and growth factors
 DNA damaging agents and anticancer drugs: 5 fluorouracil, cisplatin, doxorubicin, daunorubicine, vincristine, nocodazole, colchicine, arabinofuranosylocytosine,
 Metabolic changes: intracellular acidification
 Reactive oxygen species: H₂O₂, NO donors (S-nitrooglutathione), oxidized dopamin
 Other metabolic drugs: polyphenols, 2-methoxyestradiol, monochloramine, lovastatin, amylin, ceramide, arsenite, curcumin, calphostin C

Fig. 1. Factors and injurious stimuli inducing apoptosis.

There are two central pathways that lead to apoptosis: a) the positive induction by ligands (death factors) binding to plasma membrane receptors (death factor receptors) (*Fig. 3*) and b) negative induction by the loss of suppressor activity. The common execution mechanisms of apoptosis consist of the activation of cytosolic aspartate-specific proteases (ICE-proteases) termed caspases (5, 6), which can be activated via various intracellular pathways. The activated caspases cause endonucleolytical cleavage of DNA (7), and subsequent cleavage of several cellular substrates such as poly(ADP-ribose) polymerase, gelsolin, actin, lamin, and fodrin (8–11). The caspase enzymes (caspase-1–12), the members of the interleukin-1 β converting enzyme (ICE) family, are divided based on their sequence homology into three groups: ICE-, CPP32-, and Ich-1-like proteases. The activation of caspases is accompanied by characteristic morphologic changes of the cell, including membrane blebbing, cell shrinkage, margination of nuclear matrix, chromatin condensation, DNA cleavage and formation of apoptotic bodies (*Fig. 2*).

Positive induction of apoptosis involves ligands such as tumor necrosis factor (TNF), TRAIL (TNF α -related apoptosis inducing ligand), or CD95 ligand/Fas ligand (CD95L/FasL) (*Fig. 3*), which are able to induce programmed cell death by acting on the surface cell death factor receptors [TNF receptor-1 (TNFR1), death receptor 3–6 (DR-3-6) and CD95/Fas receptor]. The ligands are predominantly trimeric and bind to cell surface causing the

aggregation of cell surface receptors (12). Receptor oligomerization orients their cytosolic 80 amino acid death domains (DD) into a configuration that recruits adapter proteins (13). The adapter complex in turn recruits caspase-8, causing its activation and initiating the cascade of caspase-mediated cell disassembly.




Fig. 2. Diagrammatic presentation of cellular changes during apoptosis: (A) normal cell, (B) translocation of chromatin toward the nuclear envelope, (C) condensation of the cytoplasm, followed by (D) nuclear fragmentation and the formation of membrane-bound apoptotic bodies.

The death ligand, $\text{TNF}\alpha$, can lead to the activation of caspases in many cell types through the activation of the two members of mitogen activated protein kinase (MAPK) superfamily; c-Jun N-terminal kinase/stress-activated protein kinases (JNK1/SAPK) (14–16) or p38 MAPK family (17, 18). The upstream mediator of p38 is MAPK kinase 3/6 (MKK3/MKK6), which can be activated by apoptosis signal regulating kinase (ASK1). ASK1 also activates stress signaling kinase (SEK1/MKK4), the inducer of JNK1/SAPK after genotoxic stress (*Fig. 3*). One apoptotic pathway initiated through TNFR1 uses caspase-8 pathway through the interaction of TNFR-associated death domain protein (TRADD) and Fas-associated death domain protein (FADD/MORT1) (19). TRADD additionally recruits receptor interacting protein (RIP), which may trigger a second pathway (*Fig. 3*).

In CD95/Fas-induced signaling pathway (20), the activation of JNK1/SAPK has been described through MKK7 (21) (*Fig. 3*). Binding of the ligand (CD95L/FasL) is thought to induce oligomerization of the receptor with transduction of the death signal by recruitment of FADD/MORT1, which in turn binds to caspase-8, initiating the protease cascade of apoptosis (22). Fas has been shown to be constitutively expressed at a low level in most tissues, including the gastric mucosa and other parts of the gastrointestinal tract (23, 24).




Fig. 3. Diagrammatic presentation of general pathways of positive induction of apoptosis. Abbreviations: ASK1, apoptosis signal regulating kinase-1; DD, death domain; JNK1/SAPK, C-Jun N-terminal kinase or stress-activated protein kinase; CD95L/FasL, CD95 ligand or Fas ligand; FADD/MORT1, Fas-associated death domain protein; MAPK, mitogen activated protein kinase; MKK, mitogen activated protein kinase kinase; NFκB, nuclear factor-κB; RIP, receptor interacting protein; SEK1, stress signal kinase-1; TNFα, tumor necrosis factor receptor-1; TRADD, TNFR-associated death domain; TRAF2, TNF receptor associated factor-2.

The recently identified death receptors, DR-3, -4, -5 and -6 mediate cell death in a wide variety of malignant cell lines and primary tumor cells (25–26), where they exert action similar to TNFR1 but not to Fas (27). They have not been shown to be present in gastric mucosal cells.

Negative induction of apoptosis is generated by the loss of suppressor activity and involves the mitochondrial mechanisms. Release of cytochrome C from mitochondria into the cytosol serves as a trigger to activate caspases (28) (Fig. 4). Proteins of Bcl-2 family regulate the permeability of mitochondrial outer membrane to ions and to cytochrome C. At least 16 genes of Bcl-2 family have been identified (29), which either promote (Bax, Bad, Bak) or antagonise (Bcl-2, Bcl-x) apoptosis. The ratio of apoptosis promoting factors to those inhibiting apoptosis in the cell determines whether the apoptotic cell death will be turned on and the cascade of caspases will be activated. In the cytosol, cytochrome C binds to apoptosis protease activator protein-1 (APAF-1) in the presence of ATP and activates caspase-9, which induces effector caspases such as caspase-3 (30).




Fig. 4. Diagrammatic presentation of negative induction of apoptosis. Abbreviation: APAF-1, apoptosis protease activator protein-1.

The pathway by which apoptosis is induced clearly vary, depending on stimulus and cell types. The sequence of intracellular signaling and events involved in apoptosis of gastric mucosa has been only partly elucidated.

APOPTOSIS IN THE GASTRIC MUCOSA

The surface mucosal cells are constantly exfoliating into the gastric lumen, with a 3–5 day renewal rate under normal physiological conditions. The fasting of the animal prolongs the cell renewal rate, while feeding and injury causes faster renewal. Apoptosis has been reported to take place in all regions of the stomach, occurring predominantly in the upper part of the gastric glands and involving 2–3% of all epithelial cells (31). Exfoliated surface mucosal cells are replaced by cells migrating from the glandular foveolar and neck area (1). During migration these cells differentiate, mature and eventually degenerate. The loss of parietal, chief, and endocrine cells of oxyntic mucosa is much slower than the loss of surface epithelial cells (1).

Apoptosis has been shown to be induced following acute mucosal injury and during gastric ulcer healing (1, 32). At the onset of gastric ulceration, the rate of apoptosis rapidly increases. The 3.9-fold increase in mucosal expression of caspase-3 (effector caspase) activity can be detected as early as 2 hrs after

experimental ulcer induction (32). Caspase-3 activity increases up to 33-fold within 2 days after ulcer induction, and is followed by a subsequent decline throughout the first week (33). The expression of Bax, Bak and p53 was found to be increased at the edge of gastric ulceration, while Bcl-2 expression was similar to control (34). The apoptosis rate of epithelial cells during gastric ulcer healing remained elevated for 2—3 months. However, in a 3-month old ulcer scar, the rate of apoptosis for mucous, parietal, chief and endocrine cells is similar to that present in normal gastric mucosa (1).

EFFECT OF *H. PYLORI* ON GASTRIC EPITHELIAL CELL APOPTOSIS

Several studies showed that incubation of gastric epithelial cells with *H. pylori* or its supernatant causes inhibition of cell proliferation. This action has been demonstrated to occur in a time- and concentration-dependent manner in human gastric AGS (35), MKN 28 (36) and KATO III cells (37).

In addition to inhibition of cell proliferation, growth inhibitory effect of *H. pylori* in gastric epithelial cells is associated with the induction of apoptosis. The bacterial concentration causing DNA fragmentation and reduction of cell number was found to be 100-fold lower than the concentration that causes inhibition of DNA synthesis (31). Using terminal deoxynucleotidyl transferase-mediated deoxyuridine triphosphate nick-end labelling (TUNEL) technique to detect apoptosis in gastric mucosal sections, Moss et al. found that the apoptotic index was increased up to 16% in *H. pylori* infected patients with duodenal ulcers, whereas it is only 2—3% in uninfected healthy mucosa (31). Eradication of *H. pylori* decreased the elevated apoptotic index of the gastric mucosa to the basal level.

Further studies aimed to characterise the anti-proliferative activity of *H. pylori* demonstrated that this effect does not correlate with the presence of vacuolizing cytotoxin (VacA) (32). Knipp et al. (36) found a putative inhibitory protein originated from the cytosol of *H. pylori* (termed proliferation inhibiting protein) responsible for induction of apoptosis in gastric mucosa. Another potential candidate is ammonia, generated by *H. pylori*'s urease. Ammonia has been shown to induce apoptosis *in vitro* in rat gastric epithelial cell lines (38). Wagner et al. (39) suggested that the stimulation of apoptosis is a general feature of *H. pylori* that is induced by all strains, and is specific to *H. pylori*, because *C. jejuni* is not able to affect epithelial cell growth and apoptosis. Clarithromycin has the ability to attenuate *H. pylori*-induced apoptosis of gastric mucosa by altering either the metabolism of the bacteria or the mucosal responses induced by bacteria (40).

In vivo studies using proliferating cell nuclear antigen, BrdU labeling and MIB-1 antibody contradicted the results of the above *in vitro* investigations.

H. pylori infection is accompanied by increased epithelial cell proliferation in patients with *H. pylori*-associated gastritis (41—45). Enhanced apoptosis caused by *H. pylori* occurs predominantly in the superficial compartment of the mucosa, where *H. pylori* is localised. The inflammation of the deeper portion of the mucosal glands, such as the neck area, combined with the high rate of apoptosis may cause a compensatory cell hyperproliferation. Piotrowski *et al.* (46) demonstrated that bacterial lipopolysaccharide (LPS) is the virulence factor responsible for the induction of apoptosis by *H. pylori*. Intra-gastric administration of *H. pylori* LPS induces a concentration dependent apoptosis in rat gastric mucosal cells detected by *in situ* DNA fragmentation assay (46).

In the analysis of other potential mechanisms of *H. pylori*-induced apoptosis, it should be pointed out, that *H. pylori* infection stimulates the release of pro-inflammatory cytokines — interleukin-1, interleukin-2, interleukin-6, interleukin-8, interferon- γ , and TNF α (47—49). These cytokines have been shown to upregulate the expression of CD95/Fas (50). CD95 expression has been shown upregulated and CD95L/FasL mRNA levels increased in gastric surface epithelium of patients infected with *H. pylori*, suggesting that apoptosis involved in *H. pylori*-gastritis includes CD95 and CD95L activation (51, 52). CagA producing *H. pylori* strains are associated with increased cytokine production in the gastric mucosa and in gastric epithelial cells (24, 48, 53—55), which in turn may be responsible for activation of CD95-CD95L system. Incubation of gastric epithelial cells with *H. pylori* supernatant in presence of F(ab')₂ anti-Fas fragments is able to reduce (at least in part) apoptosis. In addition, the number of dead cells detected by cytotoxic assay is higher than the fraction of apoptotic cells measured by FACS analysis, indicating that cytosolic cell death might also occur in addition to apoptosis (51). This additional cytolytic cell death is caused by *H. pylori* VacA cytotoxin, which induces cell degeneration by vacuolar-ATPase (56).

Recent experiments have shown changes in the expression levels of mitochondrial factors (negative inductors) during *H. pylori*-induced apoptosis. Upregulation of pro-apoptotic Bax, Bcl-x_s and downregulation of anti-apoptotic Bcl-2 were found *in vivo* in gastric epithelium and *in vitro* in Kato III cells infected with *H. pylori* (57, 58).

NSAIDS AND APOPTOSIS

NSAIDs have been shown to induce apoptosis in isolated rat gastric mucosal cells (59) and in various cell lines derived from gastrointestinal tract (60—63). Indomethacin and sodium diclofenac inhibit cell growth (64) and induce apoptotic DNA fragmentation in gastric cells in a dose- and time-dependent manner (59, 60). Following indomethacin treatment, a 20-fold

increase in cell apoptosis, and 4-fold increase in caspase-3 activity can be detected in rat gastric mucosa (32). The DNA fragmentation induced by COX inhibitors, particularly COX-2 inhibitors, is not affected by the exogenous addition of 16,16-dimethyl prostaglandin E₂, suggesting that the inhibition of prostaglandin production is not responsible for this process. However, NSAIDs-induced apoptosis was significantly reduced by caspase inhibitors (59, 65).

The mechanisms of NSAIDs-inducing apoptosis are not fully elucidated. NSAIDs-induced TNF α release in the gastric mucosa might be responsible for positive induction of apoptosis (33, 66, 67). The expression of apoptosis-related genes, such as Fas, Bcl-2 and Bax, are not affected by indomethacin or selective COX-2 inhibitors (e.g. NS-398). Intra- and extra-cellular calcium chelators, protein tyrosine kinase inhibitor, protein kinase A (PKA) inhibitor and protein kinase C (PKC) inhibitors do not influence indomethacin-induced apoptosis in colon cancer cells. In contrast, NSAIDs induced p21waf-1 transcription rate, suggesting the possible association of NSAIDs-induced apoptosis with cell-cycle control (68). The up-regulation of c-myc proto-oncogene was associated with NSAID-induced apoptosis in gastric epithelial cell lines. In these cell lines indomethacin increased c-myc mRNA and protein expression. Conversely, down-regulation of c-myc with antisense oligonucleotides significantly reduced indomethacin-induced apoptosis (69).

Nitric oxide plays also a role in indomethacin-induced apoptosis in the gastric mucosa. The activity of nitric oxide synthase-2 (NOS-2) positively correlates with the elevation of caspase-3 activity in indomethacin-induced apoptosis in rat gastric mucosa, suggesting the participation of NOS-2 in the amplification of apoptosis signaling (32). Nitric oxide-releasing NSAIDs (NO-NSAIDs), a new class of NSAID derivatives with markedly reduced gastrointestinal toxicity, are non-peptide caspase inhibitors. Inhibition of caspases by these drugs and their ability to increase gastric blood flow likely contribute to their gastroprotective effect (66).

Increased caspase-3 activity can be observed shortly after aspirin treatment in rat gastric mucosa, while chronic administration of aspirin causes only slight elevation of caspase activity, indicating that the resistance to apoptosis might be one of the mechanisms responsible for gastric mucosal adaptation to chronic aspirin administration (70).

In addition to above described mechanisms of NSAID-induced apoptosis, there are certainly many other factors and intracellular pathways involved in this process, which remain to be elucidated.

It remains controversial whether the harmful effects of *H. pylori* and NSAIDs on gastric mucosa are additive. A significantly higher apoptosis rate can be observed in gastric mucosa of human subjects with *H. pylori* infection or treated with NSAIDs, compared to controls. Unlike NSAIDs treated subjects,

patients with *H. pylori* infection show significantly enhanced gastric mucosal proliferation. Interestingly, a study indicates that NSAIDs do not potentiate but rather abrogate the apoptosis induced by *H. pylori* (71).

Acknowledgements: Supported by the Medical Research Service of Veterans Administration. We thank Dr. Matthew J. Domek for reading the manuscript and a constructive discussion.

ABBREVIATIONS

APAF-1, apoptosis protease activator protein-1; ASK1, apoptosis signal regulating kinase-1; BrdU, bromodeoxyuridine; CagA, cytotoxin associated protein; DD, death domain; JNK1/SAPK, C-Jun N-terminal kinase-1 or stress-activated protein kinase; CD95L/FasL, CD95 ligand or Fas ligand; COX-2, cyclooxygenase-2; FADD/MORT1, Fas-associated death domain protein; FACS, fluorescence-activated cell sorter; ICE, interleukin-1 β converting enzyme; I κ B, inhibitor of NF κ B; LPS, lipopolysaccharide; MAPK, mitogen activated protein kinase; MIB-1, a monoclonal antibody reacting with Ki-67 antigen; MKK, mitogen activated protein kinase kinase; NF κ B, nuclear factor- κ B; NOS-2, nitric oxide synthase-2; NSAIDs, nonsteroidal anti-inflammatory drugs; NO-NSAIDs, nitric oxide-releasing NSAIDs; RIP, receptor interacting protein; PI-3K, phosphatidylinositol-3 kinase; PKA and PKC, protein kinase A and C, respectively; SEK1, stress signal kinase-1; TNF α , tumor necrosis factor-alpha; TNFR1, tumor necrosis factor receptor-1; TNFR1, tumor necrosis factor receptor-1; TRADD, TNFR-associated death domain; TRAF2, TNF receptor associated factor-2; TRAIL, TNF-related apoptosis inducing ligand; TUNEL, terminal deoxynucleotidyl transferase-mediated deoxyuridine triphosphate nick-end labelling; VacA, vacuolating cytotoxin.

REFERENCES

1. Stachura J, Tarnawski A, Dabros W. Apoptosis: genetically programmed physiologic cell loss in normal gastric oxyntic mucosa and in mucosa of grossly healed gastric ulcer. *J Clin Gastroenterol* 1993; 17: S70—7.
2. Bursch W, Obermammer F, Schulte-Herman R. Cell death by apoptosis and its protective role against disease. *Trends Pharmacol Sci* 1992; 13: 245—51.
3. Evans VG. Multiple pathways of apoptosis. *Cell Biol Int* 1993; 17: 461—75.
4. Nagata S. Apoptosis by death factor. *Cell* 1997; 88: 355—65.
5. Alnemri ES, Livingston DJ, Nicholson DW *et al.* Human ICE/CED-3 protease nomenclature. *Cell* 1996; 87: 171.
6. Thornberry NA, Lazebnik Y. Caspases: enemies within. *Science* 1998; 281: 1312—6.
7. Schwartzman RA, Cidlowski JA. Apoptosis the biochemistry and molecular biology of programmed cell death. *Endocr Rev* 1993; 14: 133—51.
8. Lazebnik YA, Kaufmann SH, Desnoyers S, Poirier GG, Earnshaw WC. Cleavage of poly(ADP-ribose) polymerase by a proteinase with properties like ICE. *Nature* 1994; 371: 346—7.
9. Kothakota S, Azuma T, Reinhard C *et al.* Caspase-3-generated fragment of gelsolin: effector of morphological change in apoptosis. *Science* 1997; 278: 294—8.
10. Mashima T, Naito M, Noguchi K, Miller DK, Nicholson DW, Tsuruo T. Actin cleavage by CPP-32/apopain during the development of apoptosis. *Oncogene* 1997; 14: 1007—12.

11. Martin SJ, O'Brien GA, Nishioka WK *et al.* Proteolysis of fodrin (non-erythroid spectrin) during apoptosis. *J Biol Chem* 1995; 270: 6425—28.
12. Ashkenazi A, Dixit VM. Death receptors: signaling and modulation. *Science* 1998; 281: 1305—8.
13. Tartaglia LA, Ayres TM, Wong GH, Goeddel DV. A novel domain within the 55 kd TNF receptor signals cell death. *Cell* 1993; 74: 845—53.
14. Derijard B, Hibi M, Wu IH *et al.* JNK1: a protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain. *Cell* 1994; 76: 1025—37.
15. Kyriakis JM, Avruch J. pp54 microtubule-associated protein 2 kinase. A novel serine/threonine protein kinase regulated by phosphorylation and stimulated by poly-L-lysine. *J Biol Chem* 1990; 265: 17355—63.
16. Kyriakis JM, Banerjee P, Nikolakaki E *et al.* The stress-activated protein kinase subfamily of c-Jun kinases. *Nature* 1994; 369: 156—60.
17. Han J, Lee JD, Jiang Y, Li Z, Feng L, Ulevitch RJ. Characterisation of the structure and function of a novel MAP kinase kinase (MKK6). *J Biol Chem* 1996; 271: 2886—91.
18. Lee JC, Laydon JT, McDonnell PC *et al.* A protein kinase involved in the regulation of inflammatory cytokine biosynthesis. *Nature* 1994; 372: 739—46.
19. Chinnaiyan AM, Tepper CG, Seldin MF *et al.* FADD/MORT1 is a common mediator of CD95 (Fas/APO-1) and tumor necrosis factor receptor-induced apoptosis. *J Biol Chem* 1996; 271: 4961—65.
20. Lynch DH, Ramsdell F, Alderson MR. Fas and FasL in the homeostatic regulation of immune responses. *Immunol Today* 1995; 16: 569—74.
21. Moriguchi T, Toyoshima F, Masuyama N, Hanafusa H, Gotoh Y, Nishida E. A novel SAPK/JNK kinase, MKK7, stimulated by TNF α and cellular stresses. *EMBO J* 1997; 16: 7045—53.
22. Nagata S. Apoptosis mediated by Fas and its related diseases. *Nippon Ika Daigaku Zasshi* 1997; 64: 459—62.
23. French LE, Hahne M, Viard I *et al.* Fas and Fas ligand in embryos and adult mice: ligand expression in several immune-privileged tissues and coexpression in adult tissues characterised by apoptotic cell turnover. *J Cell Biol* 1996; 133: 335—43.
24. Houghton J, Korah RM, Condon MR, Kim KH. Apoptosis in *Helicobacter pylori*-associated gastric and duodenal ulcer disease is mediated via the Fas antigen pathway. *Dig Dis Sci* 1999; 44: 465—78.
25. Chinnaiyan AM, O'Rourke K, Yu GL *et al.* Signal transduction by DR3, a death domain-containing receptor related to TNFR-1 and CD95. *Science* 1996; 274: 990—2.
26. Griffith TS, Lynch DH. TRAIL: a molecule with multiple receptors and control mechanisms. *Curr Opin Immunol* 1998; 10: 559—63.
27. Ashkenazi A, Dixit VM. Apoptosis control by death and decoy receptors. *Curr Opin Cell Biol* 1999; 11: 255—60.
28. Liu X, Kim CN, Yang J, Jemmerson R, Wang X. Induction of apoptotic program in cell-free extracts: requirement for dATP and cytochrome C. *Cell* 1996; 86: 147—157.
29. Korsmeyer SJ. Regulators of cell death. *Trends Genet* 1995; 11: 101—5.
30. Zou H, Henzel WJ, Liu X, Lutschg A, Wang X. Apaf-1, a human protein homologous to *C. elegans* CED-4, participates in cytochrome C-dependent activation of caspase-3. *Cell* 1997; 90: 405—13.
31. Moss SF, Calam J, Agarwal B, Wang S, Holt PR. Induction of gastric epithelial apoptosis by *Helicobacter pylori*. *Gut* 1996; 38: 498—501.
32. Piotrowski J, Slomiany A, Slomiany BL. Activation of apoptotic caspase-3 and nitric oxide synthase-2 in gastric mucosal injury induced by indomethacin. *Scand J Gastroenterol* 1999; 34: 129—34.

33. Slomiany BL, Piotrowski J, Slomiany A. Role of basic fibroblast growth factor in the suppression of apoptotic caspase-3 during chronic gastric ulcer healing. *J Physiol Pharmacol* 1998; 49: 489—500.
34. M Shimada, K Ina, T Ando *et al.* Fas/Fas Ligand pathway is associated with epithelial apoptosis in gastric ulceration. *Gastroenterology* 1999; 116: A310.
35. Knipp U, Birkholz S, Kaup W, Opferkuch W. Partial characterisation of cell proliferation-inhibiting protein produced by *Helicobacter pylori*. *Infect Immun* 1996; 64: 3491—96.
36. Ricci V, Ciacci C, Zarrilli R *et al.* Effect of *Helicobacter pylori* on gastric epithelial cell migration and proliferation in vitro: role of VacA and CagA. *Infect Immun* 1996; 64: 2829—33.
37. Chang K, Fujiwara Y, Wyle F, Tarnawski A. *Helicobacter pylori* toxin inhibits growth and proliferation of cultured gastric cells- KATO III. *J Physiol Pharmacol* 1993; 44: 17—22.
38. Tsuji S, Kawano S, Takei Y, Tsuji M, Kobayashi I, Nagano K. Ammonia induces gastric cell apoptosis: possible implication to *Helicobacter*-related gastric mucosal atrophy. *Gastroenterology* 1995; 108: A244.
39. Wagner S, Beil W, Westermann J *et al.* Regulation of gastric epithelial cell growth by *Helicobacter pylori*: evidence for a major role of apoptosis. *Gastroenterology* 1997; 113: 1836—47.
40. Minohara Y, Fan X, Ernst PB, Crowe SE, Patel JA. The effects of clarithromycin on *Helicobacter pylori* adherence and induction of apoptosis in gastric epithelial cells. *Gastroenterology* 1999; 116: A254.
41. Brenes F, Ruiz B, Correa P, Hunter F, Rhamakrishnan T, Fontham E, Shi TY. *Helicobacter pylori* cause hyperproliferation of the gastric epithelium: pre- and post-eradication indices of proliferating cell nuclear antigen. *Am J Gastroenterol* 1993; 88: 1870—75.
42. Fraser AG, Sim R, Sankey EA, Dhillon AP, Pounder RE. Effect of eradication of *Helicobacter pylori* on gastric epithelial cell proliferation. *Aliment Pharmacol Ther* 1994; 8: 167—73.
43. Lynch DA, Mapstone NP, Clarke AM *et al.* Cell proliferation in *Helicobacter pylori* associated gastritis and the effect of eradication therapy. *Gut* 1995; 36: 346—350.
44. Cahill RJ, Xia H, Kilgallen C, Beattie S, Hamilton H, O'Morain C. Effect of eradication of *Helicobacter pylori* infection on gastric epithelial cell proliferation. *Dig Dis Sci* 1995; 40: 1627—1631.
45. Cahill RJ, O'Morain CA. Gastric epithelial cell proliferation. *Eur J Cancer Prev* 1994; 3 (Suppl 2): 55—60.
46. Piotrowski J, Skrodzka D, Slomiany A, Slomiany BL. *Helicobacter pylori* lipopolysaccharide induces gastric epithelial apoptosis. *Biochem Mol Biol Int* 1996; 40: 597—602.
47. Peek RM Jr, Blaser MJ. Pathophysiology of *Helicobacter pylori*-induced gastritis and peptic ulcer disease. *Am J Med* 1997; 102: 200—7.
48. Sharma SA, Tummuru MK, Miller GG, Blaser MJ. Interleukin-8 response of gastric epithelial cell lines to *Helicobacter pylori* stimulation in vitro. *Infect Immun* 1995; 63: 1681—87.
49. Yamaoka Y, Kita M, Kodama T, Sawai N, Kashima K, Imanishi J. Expression of cytokine mRNA in gastric mucosa with *Helicobacter pylori* infection. *Scand J Gastroenterol* 1995; 30: 1153—59.
50. Estaquier J, Tanaka M, Suda T, Nagata S, Golstein P, Ameisen JC. Fas-mediated apoptosis of CD4⁺ and CD8⁺ T cells from human immunodeficiency virus-infected persons: differential in vitro preventive effect of cytokines and protease antagonists. *Blood* 1996; 87: 4959—66.
51. Rudi J, Kuck D, Strand S *et al.* Involvement of the CD95 (APO-1/Fas) receptor and ligand system in *Helicobacter pylori*-induced gastric epithelial apoptosis. *J Clin Invest* 1998; 102: 1506—14.
52. Censini S, Lange C, Xiang Z *et al.* CagA pathogenicity island of *Helicobacter pylori*, encodes type I-specific and disease-associated virulence factors. *Proc Natl Acad Sci USA* 1996; 93: 14648—53.

53. Tummuru MK, Sharma SA, Blaser MJ. *Helicobacter pylori* *picB*, a homologue of the *Bordetella pertussis* toxin secretion protein, is required for induction of IL-8 in gastric epithelial cells. *Mol Microbiol* 1995; 18: 867—76.
54. Peek RM Jr, Miller GG, Tham KT *et al.* Heightened inflammatory response and cytokine expression in vivo to CagA+ *Helicobacter pylori* strains. *Lab Invest* 1995; 73: 760—70.
55. Yamaoka Y, Kita M, Kodama T, Sawai N, Imanishi J. *Helicobacter pylori* *cagA* gene and expression of cytokine messenger RNA in gastric mucosa. *Gastroenterology* 1996; 110: 1744—52.
56. Papini E, Gottardi E, Satin B *et al.* The vacuolar ATPase proton pump is present on intracellular vacuoles induced by *Helicobacter pylori*. *J Med Microbiol* 1996; 45: 84—89.
57. Konturek P, Faller G, Marlicz K *et al.* Antigastric autoantibodies in *Helicobacter pylori* infected patients with Gastric carcinoma: implication of plasma gastrin release and gastric secretory functions. *Gastroenterology* 1999; 116: A218.
58. Takagi A, Watanabe S, Igarashi M, Koike J, Miwa T. *Helicobacter pylori* induced cell Death is associated with expression of Bcl-xs in gastric epithelial cells. *Gastroenterology* 1999; 116: A326.
59. Kusuvara H, Matsuyuki H, Matsuura M, Imayoshi T, Okumoto T, Matsui H. Induction of apoptotic DNA fragmentation by nonsteroidal anti-inflammatory drugs in cultured rat gastric mucosal cells. *Eur J Pharmacol* 1998; 360: 273—80.
60. Slomiany BL, Piotrowski J, Slomiany A. Role of caspase-3 and nitric oxide synthase-2 in gastric mucosal injury induced by indomethacin: effect of sucralfate. *J Physiol Pharmacol* 1999; 50: 3—16.
61. Zhu GH, Wong BC, Ching CK, Lai KC, Lam SK. Differential apoptosis by indomethacin in gastric epithelial cells through the constitutive expression of wild-type p53 and/or up-regulation of c-myc. *Biochem Pharmacol* 1999; 58: 193—200.
62. Bortuzzo C, Hanif R, Kashfi K, Staiano-Coico L, Shiff SJ, Rigas B. The effect of leukotrienes B and selected HETEs on the proliferation of colon cancer cells. *Biochim Biophys Acta* 1996; 1300: 240—46.
63. Shiff SJ, Koutsos MI, Qiao L, Rigas B. Nonsteroidal anti-inflammatory drugs inhibit the proliferation of colon adenocarcinoma cells: effects on cell cycle and apoptosis. *Exp Cell Res* 1996; 222: 179—88.
64. Kuwayama H, Nakajima N, Matsuo Y, Eastwood GL. Effects of single parenteral indomethacin injection in rat fundic and antral epithelial proliferation. *J Clin Gastroenterol* 1990; 12 (Suppl 1): S72—75.
65. Ahnen DJ. Colon cancer prevention by NSAIDs: what is the mechanism of action? *Eur J Surg Suppl* 1998; 582: 111—14.
66. Fiorucci S, Santucci L, Federici B *et al.* Nitric oxide-releasing NSAIDs inhibit interleukin-1 beta converting enzyme-like cysteine proteases and protect endothelial cells from apoptosis induced by TNFalpha. *Aliment Pharmacol Ther* 1999; 13: 421—35.
67. Fiorucci S, Antonelli E, Migliorati G *et al.* TNFalpha processing enzyme inhibitors prevents aspirin-induced TNFalpha release and protection against gastric mucosal injury in rats. *Aliment Pharmacol Ther* 1998; 12: 1139—53.
68. Hong SP, Ha SH, Park IS, Kim WH. Induction of apoptosis in colon cancer cells by nonsteroidal anti-inflammatory drugs. *Yonsei Med J* 1998; 39: 287—95.
69. Zhu GH, Wong BC, Eggo MC *et al.* Nonsteroidal anti-inflammatory drug-induced apoptosis in gastric cancer cells is blocked by protein kinase C activation through inhibition of c-myc. *Br J Cancer* 1999; 79: 393—400.
70. Alderman BM, Cook GA, Yeomans ND. Resistance to apoptosis is a likely mechanism of gastric mucosal adaptation to aspirin. *Gastroenterology* 1999; 116: A110.

71. Zhu GH, Yang XL, Lai KC *et al.* Nonsteroidal anti-inflammatory drugs could reverse *Helicobacter pylori*-induced apoptosis and proliferation in gastric epithelial cells. *Dig Dis Sci* 1998; 43: 1957—63.

Received: September 8, 1999

Accepted: November 24, 1999

Author's address: Andrzej S. Tarnawski, M.D., D.Sc., Gastroenterology Section (111G, DVA Medical Center Long Beach (CA), 5901 East Seventh Street, Long Beach, CA 90822, Tel: (562) 494—5804, Fax: (562) 494—5675.
E-mail: astarnaw@uci.edu