Lawrence Berkeley National Laboratory

Recent Work

Title

BIS(PENTAMETHYLCYCLOPENTADIENYL)CARBOXYLATO AND DITHIOCARBAMATO DERIVATIVES OF NEODYMIUM(III) AND YTTERBIUM(III) . CRYSTAL STRUCTURE OF BIS (PENTAMETHYL-CYCLOPENTADIENYL) DIETHYLDITHIOCARBAMATOYTTERBIUM(III)

Permalink

<https://escholarship.org/uc/item/3qn6c42n>

Author

Tilley, T. D.

Publication Date

1981-07-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Molecular Research Division

Submitted to Inorganic Chemistry

RECEIVED
LAWRENCE
BERKELEY LABORATORY

AUG 24 1981

LIBRARY
DOCUMENTS SECTION

BIS(PENTAMETHYLCYCLOPENTADIENYL)CARBOXYLATO
AND DITHIOCARBAMATO DERIVATIVES OF NEODYMIUM(III)
AND YTTERBIUM(III). CRYSTAL STRUCTURE OF
BIS(PENTAMETHYLCYCLOPENTADIENYL)DIETHYLDITHIOCARBAMATO-
YTTERBIUM(III)

T. Don Tilley, Richard A. Andersen, Allan Zalkin,
and David H. Templeton

July 1981

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782*

LBL-13010
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

BIS(PENTAMETHYLCYCLOPENTADIENYL) CARBOXYLATO
AND DITHIOCARBAMATO DERIVATIVES OF NEODYMIUM(III)
AND YTTERBIUM(III). CRYSTAL STRUCTURE OF
BIS(PENTAMETHYLCYCLOPENTADIENYL)DIETHYLDITHIOCARBAMATO-
YTTERBIUM(III)

T. Don Tilley, Richard A. Andersen, Allan Zalkin*
and David H. Templeton

Materials and Molecular Research Division
Lawrence Berkeley Laboratory

and

Department of Chemistry
University of California
Berkeley, California 94720

JULY 1981

ABSTRACT

The pentamethylcyclopentadienyl derivatives, $(\text{Me}_5\text{C}_5)_2\text{Yb}(\text{O}_2\text{CR})$ where R is Me_3C or CF_3 , $(\text{Me}_5\text{C}_5)_2\text{Yb}(\text{S}_2\text{CNEt}_2)$ and $(\text{Me}_5\text{C}_5)_2\text{Nd}(\text{S}_2\text{CNEt}_2)$ have been prepared by metathetical exchange reactions on the anionic species, $[(\text{Me}_5\text{C}_5)_2\text{MCl}_2]^-$. The crystal structure of $(\text{Me}_5\text{C}_5)_2\text{Yb}(\text{S}_2\text{CNEt}_2)$ has been determined by single crystal X-ray diffraction methods. The crystals are monoclinic, C2/c, with cell dimensions $a = 12.268(4) \text{ \AA}$, $b = 15.536(6) \text{ \AA}$, $c = 14.269(5) \text{ \AA}$, $\beta = 105.23(3)^\circ$ and $V = 2624 \text{ \AA}^3$. For $Z = 4$ the calculated density is 1.14 g/cm^3 . The structure was refined to a conventional R factor of 0.039 using 2179 data where $F^2 > 3\sigma(F^2)$. The Yb(III) atom is bonded to two sulfur atoms and to two pentamethylcyclopentadienyl

This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division of the U.S. Department of Energy under Contract No. W-7405-ENG-48.

groups. The Yb atom lies on a crystallographic two-fold axis. The dithiocarbamate ligand is disordered in the structure. The Yb-S distance is 2.70(1) Å, and the average of 5 Yb-C distances is 2.63(3) Å.

INTRODUCTION

In attempting to develop the chemistry of reactive organometallic lanthanide systems, we have had an interest in preparing neutral, base-free pentamethylcyclopentadienyl derivatives of the di- and trivalent metals. The strong Lewis-acidic nature of these metals has led to isolation of trivalent species such as $(C_5Me_5)_2YbCl_2Na(OEt_2)_2$ and $(C_5Me_5)_2YbCl(thf)$,^{1,2} as well as the divalent species $Yb(C_5Me_5)_2(thf)$ and $Eu(C_5Me_5)_2(thf)$.³ The neutral base-free systems are of interest since these coordinatively unsaturated species are expected to exhibit a rich chemistry and, perhaps, to lead to compounds of unusual structural types.

Isolation of $(C_5Me_5)_2YbCl_2Na(OEt_2)_2$ and $(C_5Me_5)_2YbCl(thf)$ suggests that a chelating ligand that can donate four electrons to the metal could yield neutral, base-free trivalent compounds. Such compounds might then be useful for preparation of the base-free ytterbium(II) metallocenes, if reduction in a non-coordinating solvent can be effected. Since carboxylate (RCO_2^-) ligands have been effectively utilized as leaving groups in transition metal chemistry, they seemed to be ideally suited for this task. Another chelating ligand, diethyldithiocarbamate ($Et_2NCS_2^-$), is electronically similar, but contains the softer sulfur atoms as donors. Because little is known about the properties of sulfur-based ligands in molecular lanthanide systems, we have conducted a low-temperature magnetic susceptibility study, as well as an X-ray crystal structure determination of $(C_5Me_5)_2Yb(S_2CNEt_2)$.

RESULTS AND DISCUSSION

Preparation of the carboxylate and dithiocarbamate derivatives of ytterbium is straightforward. The analogous neodymium dithio-

carbamate, $(\text{Me}_5\text{C}_5)_2\text{Nd}(\text{S}_2\text{CNEt}_2)$, was prepared similarly. The physical properties are described in the experimental section. Because two donor atoms are present, R_2CO_2^- and the softer $\text{Et}_2\text{NCS}_2^-$ allow isolation of ether-free complexes. Unfortunately the ytterbium(III) species could not be reduced with sodium amalgam in refluxing toluene. Further, $(\text{Me}_5\text{C}_5)_2\text{YbN}(\text{SiMe}_3)_2^1$ could not be reduced under similar conditions.

The similarity in the magnetic behavior of $(\text{C}_5\text{Me}_5)_2\text{Yb}(\text{O}_2\text{CCMe}_3)$ and $(\text{C}_5\text{Me}_5)_2\text{Yb}(\text{S}_2\text{CNEt}_2)$ at low temperature (see experimental section) suggests that the magnetic properties of the ytterbium ions in these complexes are not effected by the change from carboxylate to dithiocarbamate ligand. The magnetic moments (5-45K), 3.29 and 3.39 B.M., respectively, are close in value to those observed for other ytterbium(III) complexes over this temperature range.^{1,4,5}

The ytterbium dithiocarbamate complex crystallizes as well-formed, purple prisms that are monoclinic, space group C2/c, with cell dimensions $a = 12.268(4) \text{ \AA}$, $b = 15.536(6) \text{ \AA}$, $c = 14.269(5) \text{ \AA}$, and $\beta = 105.23(3)^\circ$. For $Z = 4$ and a molecular weight of 652.98, the calculated density is 1.14 g/cm^3 . A few structures containing lanthanide to sulfur bonds have been determined,^{6,7,8} but the present structure is the first example of an Yb-S bond.

Positional parameters, with labelling as shown in Fig. 1, are given in Table I. A list of selected distances and angles are given in Tables II and III.

The molecule consists of the Yb atom bonded to two pentamethylcyclopentadienyl rings and to two sulfur atoms of the diethyldithiocarbamate ligand, Fig. 1. The average of five Yb-C distances is $2.63 \pm 0.03 \text{ \AA}$ and the Yb-S distance is $2.70 \text{ \AA}$. The pentamethylcyclopentadienyl rings are in a staggered configuration with respect to each other, see Fig. 2. The diethyldithiocarbamate ligand is disordered; atoms N, C(12A), C(13A), C(12B) and C(13B) are in general positions with half of the molecules in the given conformation (Fig. 1) and half in the conformation that would result from a 180° rotation about the two-fold axis. The sulfur atoms are most probably disordered as well, as seen by their elliptical shape in Fig. 1, but the separation of the two positions was too small to be treated in the least-squares as separate half-atoms and instead was refined as a large thermally anisotropic atom. Because of this disorder the molecule as a whole has low symmetry C_1 , however the

$(\text{Me}_5\text{C}_5)_2\text{Yb}$ part of the molecule has perfect C_2 symmetry. Attempts to define the structure in a lower symmetry space group were unsuccessful.

The Yb-C distances (to the C_5 ring) average $2.63 \pm 0.03 \text{ \AA}$. The Yb atom is $2.33 \text{ \AA}$ from the mean plane of atoms C(1) through C(5). These ring atoms are all within $0.006 \text{ \AA}$ of this plane. The five methyl carbon atoms of the pnetamethylcyclopentadienyl group are all bent out of the mean plane away from the Yb atom; the distances from the plane are 0.18, 0.19, 0.12, 0.07 and $0.34 \text{ \AA}$ for atoms C(6) through C(10), respectively.

The ytterbium-carbon bond length of $2.63 \pm 0.03 \text{ \AA}$ in the trivalent, dithiocarbamate complex is identical within experimental error to that found for this bond length ($2.66 \pm 0.01 \text{ \AA}$) in the divalent species, $(\text{Me}_5\text{C}_5)_2\text{Yb}(\text{thf})$.³ Since the oxidation state and coordination number of the latter are one less than the former, both quantities are important in determining average metal to ligand bond lengths.⁹

EXPERIMENTAL

All operations were carried out under nitrogen. Microanalyses were performed by the microanalytical laboratory of this department. Infrared spectra were recorded on a Perkin-Elmer instrument as Nujol mulls between cesium iodide windows. The ^1H NMR spectra were recorded on a JEOL-FX90Q instrument on toluene- d_8 solutions and are reported in δ -units (positive values to high field of Me_4Si). Solid state magnetic susceptibilities were obtained as previously described.¹⁰

Bis(pentamethylcyclopentadienyl)(2,2-dimethylpropionato)ytterbium (III). Sodium bis(pentamethylcyclopentadienyl)dichloroytterbium bis(diethylether)⁵ (1.7 g, 0.0025 mol) in diethyl ether (30 mL) was added to a suspension of sodium 2,2-dimethylpropionate [prepared by stirring sodium hydride with pivalic acid in diethyl ether for 12 h] (0.31 g, 0.0025 mol) in diethyl ether (20 mL). After stirring for 6 h the red solution was evaporated to dryness and the residue was extracted with pentane (35 mL). The extract was filtered and cooled (-70°C). The red prisms (0.65 g, 48%) were collected and dried under reduced pressure, mp $154-155^\circ\text{C}$. Anal. Calcd for $\text{C}_{25}\text{H}_{39}\text{O}_2\text{Yb}$: C, 55.1; H, 7.22. Found: C, 55.2; H, 7.19. The mass spectrum gave a monomeric molecular ion (545) as well as peaks due to higher oligomers at 924, 889, 820, 802, 749, 612, and 571. ^1H NMR ($+35^\circ\text{C}$): δ 10.95 ($\nu_{1/2}$ = 107 Hz), 30 H; δ - 26.45 ($\nu_{1/2}$ = 21 Hz), 9 H. IR: 2713 w, 1503 m, 1489 m, 1430 m, 1225 s, 1169 w, 1064 w, 1026 m, 940 w, 896 s, 810 m, 793 m, 617 m,

474 w, 408 m, 388 m, 319 s, cm^{-1} . The magnetic susceptibility followed Curie behavior from 5-35 K, with $\mu_B = 3.29$ B.M.

Bis(pentamethylcyclopentadienyl)(trifluoroacetato)ytterbium(III).

The salt, $\text{Na}(\text{OEt})_2(\text{Me}_5\text{C}_5)_2\text{YbCl}_2$ (1.1 g, 0.0016 mol) in tetrahydrofuran (30 mL) was added to NaO_2CCF_3 (0.22 g, 0.0016 mol) in tetrahydrofuran (10 mL) and the red solution was stirred for 12 h. The tetrahydrofuran was removed under reduced pressure and the residue was extracted with diethyl ether (2 x 40 mL). The extract was filtered, and cooling (-10°C) yielded violet prisms in 57% (0.50 g) yield, mp $262-263^\circ\text{C}$. Anal. Calcd for $\text{C}_{22}\text{H}_{30}\text{F}_3\text{O}_2\text{Yb}$: C, 47.5; H, 5.43. Found: C, 47.7; H, 5.49. The mass spectrum afforded a monomeric molecular ion at 557. IR: 3140 w, 3092 w, 2726 w, 1680 s, 1204 s, 1158 s, 1023 w, 847 m, 785 m, 718 m, 614 w, 592 w, 523 w, 468 w, 385 m, 312 s, cm^{-1} .

Bis(pentamethylcyclopentadienyl)(diethyldithiocarbamato)ytterbium(III).

The anionic complex, $\text{Na}(\text{OEt})_2(\text{Me}_5\text{C}_5)_2\text{YbCl}_2$ (1.4 g, 0.0020 mol) in diethyl ether (40 mL) was added to sodium diethyldithiocarbamate (0.35 g, 0.0020 mol) in diethyl ether (20 mL). After stirring for 8 h the diethyl ether was removed under reduced pressure. The residue was extracted with pentane (40 mL), and filtered. Concentration of the filtrate to ca. 10 mL and cooling (-10°C) yielded purple prisms in 81% (0.97 g) yield, mp $226-227^\circ\text{C}$. Anal. Calcd for $\text{C}_{25}\text{H}_{40}\text{NS}_2\text{Yb}$: C, 50.7; H, 6.81; N, 2.37; S, 10.8. Found: C, 50.7; H, 6.76; N, 2.29; S, 10.7. The mass spectrum

gave a monomeric molecular ion at 592. ^1H NMR (+35°C): δ 7.45 ($\nu_{1/2}$ = 61 Hz), 30 H; δ - 5.76 ($\nu_{1/2}$ = 18 Hz), 4 H; δ - 9.00 ($\nu_{1/2}$ = 13 Hz), 6 H. All of the resonances were singlets. IR: 2723 w, 1487 s, 1424 s, 1360 m, 1306 w, 1277 s, 1209 m, 1241 m, 1089 m, 1023 w, 987 m, 911 m, 841 m, 800 w, 779 w, 608 w, 593 w, 564 m, 473 w, 392 m, 360 m, 311 s, cm^{-1} . The magnetic susceptibility followed Curie behavior (5-55 K), giving μ_B = 3.39 B.M.

Bis(pentamethylcyclopentadienyl)(diethyldithiocarbamato)neodymium (III). The anionic complex, $\text{Li}(\text{OEt})_2(\text{Me}_5\text{C}_5)_2\text{NdCl}_2$, (2.2 g, 0.034 mol) in diethyl ether (40 mL) was added to $\text{NaS}_2\text{CNEt}_2$ (0.58 g, 0.0034 mol) in diethyl ether (20 mL). After stirring for 8 h the green solution was filtered and the filtrate was concentrated to ca. 30 mL. Cooling (-10°C) yielded green prisms in 72% (1.4 g) yield, mp 229-231°C. Anal. Calcd for $\text{C}_{25}\text{H}_{40}\text{NNdS}_2$: C, 53.3; H, 7.16; N, 2.49. Found: C, 53.5; H, 7.03; N, 2.47. The mass spectrum gave a monomeric molecular ion at 562. ^1H NMR (+70°C): δ 7.00 ($\nu_{1/2}$ = 23 Hz), 30 H; δ - 0.10 ($\nu_{1/2}$ = 17 Hz), 4 H; δ - 0.94 ($\nu_{1/2}$ = 16 Hz), 6 H. The resonances were singlets. IR: 2722 w, 1482 s, 1420 s, 1357 m, 1402 w, 1273 s, 1203 s, 1138 m, 1087 m, 1063 w, 1021 w, 979 m, 905 m, 836 m, 798 w, 773 w, 607 w, 560 m, 470 w, 430 w, 382 m, 348 m, 310 s, cm^{-1} . The magnetic susceptibility followed Curie behavior (5-50 K) giving μ_B = 2.75 B.M.

Some of the air sensitive purple crystals were inserted into thin walled quartz capillaries in an argon filled dry box. A crystal was examined with a Picker FACS-I automated diffractometer equipped with a graphite monochromator and a Mo X-ray tube ($\lambda(K\alpha_1)$ 0.70930 Å). Omega scans of several low angle reflections showed peaks with half-widths of 0.13° to 0.20°. The space group is C2/c. The setting angles of twelve manually centered reflections ($4 < 2\theta < 49^\circ$) were used to determine cell parameters by least-squares.

Intensity data were collected using a θ -2 θ scan technique with a scan speed of 2°/min on 2 θ . Each peak was scanned 0.9° before the $K\alpha_1$ peak to 0.9° after the $K\alpha_2$ peak, and backgrounds were counted for 10 s at each end of the scan range. The temperature during the data collection was $24 \pm 1^\circ\text{C}$. Three standard reflections were measured every 250th scan. A total of 4748 scans ($4^\circ < 2\theta < 45^\circ$) yielded 2327 unique data of which 2175 had $F^2 > 3\sigma$. Absorption correction ($\mu = 37.0 \text{ cm}^{-1}$) was applied¹¹ which ranged from 1.53 to 1.98. The intensities of the three standard reflections decayed about 4% during the data collection period and the data were corrected for this effect.

A three dimensional Patterson calculation showed the Yb and S atom positions, and a subsequent least-squares refinement and Fourier calculation revealed all the non-hydrogen atoms in the structure. After a few cycles of least-squares refinements of the structure in which only the Yb and S atoms were assigned anisotropic thermal parameters the R factor, $R = \sum ||F_o| - |F_c|| / \sum |F_o|$, was 0.055.

The ethyl carbon atoms of the diethyldithiocarbamate ligand were ill behaved in the refinements as indicated by large thermal parameters and chemically unsatisfactory bond distances to adjacent atoms. A difference Fourier map showed additional electron density in the region of the ethyl carbon atoms as well as the nitrogen atom, indicating disorder. The two ethyl carbon atoms were subsequently described as four half-atoms and the nitrogen atom was allowed to wander from the special position on the two-fold axis. With further refinements the structural parameters converged to their final and satisfactory results. Several attempts were made to refine the structure in the non-centric space group Cc, but the resulting structure would not converge to a reasonable chemical model, and the resulting R factors, despite the increased number of parameters, were not significantly different from the centric refinements. The least-squares function used minimizes the function $\sum w(|F_o| - |F_c|)^2 / \sum w F_o^2$. The expressions used in processing the data and estimating weights are given in the supplementary material; the "ignorance factor" was set to 0.05. Scattering factors from International Tables¹² and anomalous dispersion corrections¹³ were applied. In the final refinement all of the atoms with the exception of the carbon atoms in the dithiocarbamate ligand were assigned anisotropic thermal parameters. Hydrogen atoms were not included.

The discrepancy indices for 2175 data were $F^2 > 3\sigma$ are

$$R = \sum ||R_o| - |F_c|| / \sum |F_o| = 0.039$$

$$R_w = [\sum w(|F_o| - |F_c|)^2 / \sum w|F_o|^2]^{1/2} = 0.052$$

R for all 2327 data is 0.041. The error in an observation of unit weight is 1.84. In the last cycle, no parameter changed more than 0.1 σ . In the final difference Fourier the largest electron density peak was $\sim 0.5 \text{ e/\AA}^3$ and was near the Yb atom.

ACKNOWLEDGMENT

This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division of the U.S. Department of Energy under contract No. W-7405-Eng-48.

SUPPLEMENTARY MATERIAL AVAILABLE

Data processing formulas, the table of thermal parameters and the listing of structure factor amplitudes (10 pages). Ordering information is given on any current masthead page.

REFERENCES

1. Tilley, T.D.; Andersen, R.A. Inorg. Chem. in press.
2. Watson, P.L.; Whitney, J.F.; Harlow, R.L. Inorg. Chem. in press.
3. Tilley, T.D.; Andersen, R.A.; Spencer, B.; Ruben, H.; Zalkin, A.; Templeton, D.H. Inorg. Chem. 1980, 19, 2999.
4. Baker, E.C.; Raymond, K.N. Inorg. Chem. 1977, 16, 2710.
5. Tilley, T.D.; Zalkin, A.; Andersen, R.A.; Templeton, D.H. Inorg. Chem. 1981, 20, 551.
6. Ciampolini, M.; Nardi, N.; Colamarino, P.; Orioli, P. J. Chem. Soc. Dalton Trans. 1977, 379.
7. Meseri, Y.; Pinkerton, A.A.; Chapuis, G. J. Chem. Soc. Dalton Trans. 1977, 725.
8. Pinkerton, A.A.; Schwarzenbach, D. J. Chem. Soc. Dalton Trans. 1980, 1300.
9. Raymond, K.N.; Eigenbrot, C.W. Acc. Chem. Res. 1980, 13, 276.
10. Andersen, R.A.; Templeton, D.H.; Zalkin, A. Inorg. Chem. 1978, 17, 2317.
11. Templeton, L.K.; Templeton, D.H., American Crystallographic Association Proceedings 1973, Series 2, Vol. 1, p.143.
12. "International Tables for X-ray Crystallography"; Kynoch Press: Birmingham, England, 1974; Vol. IV: p.90.
13. Cromer, D.T.; Liberman, D. J. Chem. Phys. 1970, 53, 1891.

Table I. Positional Parameters^a

Atom	x	y	z
YB	0	.20917(2)	.250
S	.0541(2)	.3539(1)	.3565(1)
N	.0376(8)	.4957(6)	.2433(9)
C(1)	.1806(5)	.1109(4)	.3012(4)
C(2)	.2238(5)	.1953(4)	.2979(5)
C(3)	.1918(6)	.2255(4)	.2007(5)
C(4)	.1295(5)	.1593(4)	.1426(4)
C(5)	.1228(5)	.0883(4)	.2039(4)
C(6)	.2073(6)	.0528(6)	.3897(6)
C(7)	.3044(7)	.2424(7)	.3808(6)
C(8)	.2298(8)	.3101(6)	.1665(8)
C(9)	.0840(8)	.1608(6)	.0334(5)
C(10)	.0880(7)	-.0017(5)	.1665(7)
C(11)	0	.4078(8)	.250
C(12A)	.108(2)	.544(1)	.330(2)
C(13A)	.031(3)	.580(2)	.384(3)
C(12B)	.005(1)	.541(1)	.151(1)
C(13B)	-.116(2)	.577(2)	.125(2)

^aHere and in the following tables the number in parentheses is the estimated standard deviation in the least significant digit.

Table II. Interatomic Distances (Å)

Yb	- 2 C(1)	2.63(1)	C(1) - C(2)	1.42(1)
Yb	- 2 C(2)	2.66(1)	C(2) - C(3)	1.42(1)
Yb	- 2 C(3)	2.64(1)	C(3) - C(4)	1.41(1)
Yb	- 2 C(4)	2.60(1)	C(4) - C(5)	1.42(1)
Yb	- 2 C(5)	2.60(1)	C(5) - C(1)	1.43(1)
Yb	- 2 S	2.70(1)	C(1) - C(6)	1.52(1)
S	- C(11)	1.71(1)	C(2) - C(7)	1.52(1)
C(11)	- N	1.45(2)	C(3) - C(8)	1.52(1)
N	- C(12A)	1.50(2)	C(4) - C(9)	1.51(1)
N	- C(12B)	1.46(2)	C(5) - C(10)	1.52(1)
C(12A)	- C(13A)	1.49(4)		
C(12B)	- C(13B)	1.53(3)		

Table III. Selected angles (deg.)^a

S	- Yb	- S'	67.1(3)	C(5) - C(1) - C(2)	107(1)
C(1)	- Yb	- S	105.2(3)	C(1) - C(2) - C(3)	109(1)
C(2)	- Yb	- S	80.7(2)	C(2) - C(3) - C(4)	108(1)
C(3)	- Yb	- S	88.0(2)	C(3) - C(4) - C(5)	108(1)
C(4)	- Yb	- S	118.9(2)	C(4) - C(5) - C(1)	108(6)
C(5)	- Yb	- S	132.2(2)	C(6) - C(1) - C(2)	125(1)
C(1)	- Yb	- S'	134.8(2)	C(6) - C(1) - C(5)	127(1)
C(2)	- Yb	- S'	107.3(2)	C(7) - C(2) - C(1)	126(1)
C(3)	- Yb	- S'	82.8(2)	C(7) - C(2) - C(3)	124(1)
C(4)	- Yb	- S'	90.8(3)	C(8) - C(3) - C(2)	125(1)
C(5)	- Yb	- S'	122.2(3)	C(8) - C(3) - C(4)	127(1)
S	- C(11)	- S'	121.4(8)	C(9) - C(4) - C(3)	126(1)
S	- C(11)	- N	117.6(6)	C(9) - C(4) - C(5)	125(1)
S'	- C(11)	- N	117.2(6)	C(10) - C(5) - C(1)	126(1)
C(11) - N	- C(12A)		122(1)	C(10) - C(5) - C(4)	124(1)
C(11) - N	- C(12B)		120(1)		
N	- C(12A)- C(13A)		108(2)		
N	- C(12B)- C(13B)		114(2)		

^aPrimed atoms are at positions -x, y, 1/2-z.

FIGURE CAPTIONS

Fig. 1. ORTEP drawing of the molecule.

Fig. 2. ORTEP view of the molecule parallel to a line through the ring centers.

Fig. 1

XBL 812-8216

XBL 816-10260

Fig. 2

SUPPLEMENTARY MATERIALS FOR

BIS(PENTAMETHYLCYCLOPENTADIENYL)CARBOXYLATO
AND DITHIOCARBAMATO DERIVATIVES OF NEODYMIUM(III)
AND YTTERBIUM(III). CRYSTAL STRUCTURE OF
BIS(PENTAMETHYLCYCLOPENTADIENYL)DIETHYLDITHIOCARBAMATO-
YTTERBIUM (III)

T. Don Tilley, Richard A. Andersen, Allan Zalkin*,
and David H. Templeton

Thermal Parameters for $(C_5Me_5)_2Yb(S_2CNEt_2)_2$.^a

ATOM	B11	B22	B33	B12	B13	B23
YB	3.89(2)	2.55(2)	2.58(2)	0	-.00(1)	0
S	7.49(9)	3.33(6)	3.68(6)	.17(6)	-.85(6)	-.73(5)
N	3.9(4)	3.4(4)	5.6(5)	.1(3)	.5(4)	-.4(4)
C(1)	4.0(2)	4.1(3)	3.8(2)	1.2(2)	.4(2)	.6(2)
C(2)	3.3(2)	4.9(3)	3.7(3)	-.3(2)	.1(2)	-.6(2)
C(3)	4.0(2)	5.0(3)	4.0(3)	-.3(2)	1.0(2)	-.0(2)
C(4)	4.6(3)	4.7(3)	2.9(2)	.3(2)	.9(2)	-.8(2)
C(5)	4.0(2)	3.4(2)	4.4(3)	.5(2)	.2(2)	-1.0(2)
C(6)	5.9(3)	7.8(4)	5.6(4)	1.2(3)	.3(3)	2.6(3)
C(7)	5.4(3)	7.8(4)	4.8(4)	-.3(3)	-.9(3)	-2.3(4)
C(8)	7.0(5)	5.9(4)	7.5(5)	-1.7(3)	2.8(4)	.6(4)
C(9)	7.9(4)	7.9(5)	3.1(3)	.3(4)	.2(3)	-.5(3)
C(10)	6.6(4)	3.5(3)	8.7(5)	-.1(3)	.9(3)	-2.4(3)
C(11)	5.8(2)					
C(12A)	6.8(4)					
C(13A)	11.6(8)					
C(12B)	5.5(3)					
C(13B)	9.3(6)					

^aThe anisotropic temperature factor has the form $\exp[-0.25(B_{11}h^2a^{*2} + 2B_{12}hka^*b^* + \dots)]$. The isotropic thermal parameter has the form $\exp(-B\sin^2\theta/\lambda^2)$.

OBSERVED STRUCTURE FACTORS, STANDARD DEVIATIONS, AND DIFFERENCES (ALL X 5.0)
YTTERBIUM BIS(C5ME5)DITHIODIETHYLVARBAMATE F(0,0,0) = 2585

FOB AND FCA ARE THE OBSERVED AND CALCULATED STRUCTURE FACTORS.
SG = ESTIMATED STANDARD DEVIATION OF FOB. DEL = /FOB/ - /FCA/.

* INDICATES ZERO WEIGHTED DATA.

L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL
H,K=	0,	0		6	156	4	2	10	169	4	-1	5	580	15	73	-10	324	8	27
21226	31	-6		7	430	11	5	11	0	14	-1*	6	85	2	-10	-9	18	5	-9*
4	173	5	-44	8	41	2	-0	12	161	4	-2	7	377	10	-0	-8	383	10	-2
6	232	6	-19	9	369	10	18	H,K=	0,	14		8	90	3	9	-7	126	3	3
8	503	13	9	10	123	3	-5	0	296	8	-9	9	417	11	15	-6	358	9	-17
10	396	10	12	11	257	7	-1	1	117	3	4	10	16	17	7*	-5	236	6	-4
12	190	5	15	12	3	19	-4*	2	235	6	-11	11	289	8	29	-4	431	11	-11
14	152	4	-1	13	171	4	-1	3	110	3	4	12	74	3	-0	-3	287	7	16
16	139	4	-2	14	38	3	3	4	158	4	-4	13	185	5	3	-2	552	14	17
H,K=	0,	2		15	129	4	-2	5	97	3	2	14	81	5	-2	-1	14	6	-12*
0	579	15	-44	H,K=	8,	8		6	162	4	1	15	143	4	-3	0	630	16	16
1	526	13	12	0	146	4	-15	7	106	3	-2	16	22	8	-4*	1	205	5	53
2	495	13	-14	1	562	14	-3	8	195	5	-1	H,K=	1,	3		2	456	12	2
3	295	7	-21	2	158	4	-2	9	83	3	4	-16	118	3	3	3	378	10	4
4	519	14	49	3	425	11	6	10	181	5	-5	-15	134	4	-1	4	381	10	-4
5	420	11	26	4	220	6	2	H,K=	0,	16	-14	183	5	6		5	473	12	29
6	356	9	-8	5	280	7	6	0	142	4	1	-13	176	5	-2	6	376	10	-9
7	141	4	-20	6	202	5	-3	1	97	3	-7	-12	133	4	5	7	191	5	5
8	348	9	3	7	287	8	2	2	119	3	6	-11	218	6	22	8	370	10	8
9	51	2	-2	8	112	3	7	3	172	4	-5	-10	113	3	3	9	68	3	-9
10	284	7	20	9	231	6	31	4	90	3	3	-9	243	6	-3	10	322	8	38
11	19	6	2*	10	68	3	-1	5	195	5	-1	-8	298	8	18	11	41	6	6
12	178	5	9	11	173	4	-4	6	82	3	-3	-7	279	7	-11	12	230	6	1
13	121	3	-1	12	91	3	4	7	137	4	-1	-6	534	14	15	13	115	3	-1
14	150	4	0	13	111	3	-3	8	99	3	-2	-5	313	8	-11	14	172	5	1
15	111	3	1	14	93	3	3	H,K=	0,	18	-4	770	20	39		15	54	4	-6
16	128	4	-2	H,K=	0,	10		0	0	18	-2*	-3	340	9	5	H,K=	1,	7	
H,K=	0,	4		0	264	7	2	1	146	4	1	-2	340	9	-26	-15	18	9	-6*
0	205	5	34	1	196	5	-2	2	16	8	7*	-1	342	9	19	-14	127	3	-6
1	297	8	-35	2	392	10	-14	3	164	4	-3	0	387	10	-15	-13	23	5	-1*
2	320	8	-8	3	224	6	-4	H,K=	1,	1		1	349	9	19	-12	220	6	0
3	656	17	36	4	458	12	-8	-16	46	4	-3	2	365	9	-46	-11	43	3	4
4	299	8	-53	5	189	5	7	-15	115	3	-9	3	414	10	13	-10	311	8	19
5	670	17	27	6	312	8	24	-14	61	3	-3	4	502	13	17	-9	151	4	8
6	357	9	24	7	152	4	10	-13	172	4	-1	5	394	10	-8	-8	303	8	0
7	343	9	6	8	129	3	14	-12	0	19	-9*	6	309	8	-6	-7	150	4	-9
8	182	5	10	9	101	3	-1	-11	314	8	40	7	309	8	-4	-6	274	7	-13
9	277	7	6	10	104	3	0	-10	17	6	1*	8	180	5	6	-5	86	2	1
10	82	3	9	11	99	3	3	-9	515	13	3	9	205	6	1	-4	376	10	-17
11	215	6	21	12	167	4	-1	-8	77	3	6	10	222	6	20	-3	28	2	9
12	150	4	-2	13	88	3	-3	-7	389	10	6	11	186	5	15	-2	667	17	-9
13	200	5	-2	H,K=	0,	12		-6	118	3	-13	12	209	5	4	-1	32	2	-8
14	152	4	1	0	258	7	33	-5	332	8	-15	13	180	5	1	0	553	14	-20
15	153	5	-0	1	28	4	-5	-4	221	6	4	14	188	5	6	1	112	3	-9
16	89	3	-5	2	295	8	32	-3	609	15	-4	15	127	4	-2	2	405	10	-24
H,K=	0,	6		3	35	2	4	-2	25	2	3	H,K=	1,	5		3	34	2	-1
0	137	4	12	4	280	7	19	-1	944	24	-38	-16	129	4	-1	4	389	10	9
1	681	17	-38	5	91	3	1	0	47	2	14	-15	107	4	6	5	59	2	-10
2	32	2	0	6	240	6	-5	1	986	28	23	-14	176	5	-2	6	425	11	1
3	406	10	-21	7	34	3	2	2	138	4	-8	-13	91	3	3	7	61	2	-6
4	131	3	23	8	189	5	-5	3	585	15	42	-12	241	6	-3	8	423	11	31
5	340	9	7	9	28	4	3	4	360	9	61	-11	46	3	6	9	97	3	8

STRUCTURE FACTORS CONTINUED FOR
 YTTERBIUM BIS(C5H5)DITHIODIETHYLVARBAMATE

PAGE 2

L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL
10	292	7	15	8	57	2	4	0	114	3	-4	15	88	3	-0	6	28	3	-1
11	54	3	-0	9	145	4	-12	1	66	3	-1	H,K= 2,		4	7	420	11	8	
12	168	4	-6	10	23	5	0*	2	170	4	-7	-16	96	3	3	8	4	13	-5*
13	42	4	4	11	158	4	-8	3	63	2	1	-15	163	4	-1	9	391	10	26
14	94	3	-9	12	68	3	-4	4	199	5	-9	-14	132	4	1	10	5	16	-4*
H,K= 1,		9		H,K= 1,		13		5	67	3	2	-13	227	6	-4	11	266	7	-5
-14	88	3	-2	-11	169	5	-5	6	156	4	-6	-12	139	4	0	12	47	4	2
-13	138	4	3	-10	62	3	-0	H,K= 2,		0	-11	202	5	21	13	160	5	6	
-12	153	4	1	-9	198	5	-2	-16	120	3	-7	-10	115	3	4	14	13	18	9*
-11	97	3	2	-8	74	3	5	-14	172	5	-6	-9	304	8	4	H,K= 2,		8	
-10	189	5	-7	-7	196	5	-1	-12	336	9	28	-8	141	4	-5	-15	98	3	-4
-9	101	3	10	-6	74	3	6	-10	496	13	-4	-7	480	12	7	-14	81	3	3
-8	169	4	21	-5	177	5	-15	-8	551	14	-5	-6	336	9	-3	-13	163	4	-2
-7	257	7	14	-4	88	3	2	-6	309	8	-35	-5	551	14	-18	-12	58	2	1
-6	165	4	-2	-3	237	6	-4	-4	533	14	8	-4	300	8	3	-11	293	8	-2
-5	353	9	-15	-2	138	4	-5	-2	494	13	-59	-3	343	9	-20	-10	78	4	1
-4	204	5	3	-1	294	8	6	0	635	16	-30	-2	129	3	-10	-9	283	7	21
-3	274	7	-13	0	87	3	1	2	592	15	42	-1	227	6	-0	-8	157	4	11
-2	349	9	-11	1	257	7	9	4	566	14	52	0	92	3	-11	-7	217	6	1
-1	186	5	6	2	13	21	-8*	6	567	15	15	1	637	16	46	-6	186	5	-4
0	359	9	-12	3	206	5	-1	8	554	14	36	2	493	13	29	-5	181	5	-10
1	275	7	-1	4	48	3	-9	10	505	13	50	3	539	14	-2	-4	214	6	3
2	324	8	-11	5	188	5	-10	12	259	7	6	4	532	14	29	-3	310	8	-21
3	402	10	-14	6	30	3	-2	14	158	4	-0	5	445	11	20	-2	156	4	-3
4	213	6	-1	7	211	5	-7	H,K= 2,		2	6	235	6	23	-1	445	11	-32	
5	305	8	-9	8	90	3	-3	-16	116	3	-4	7	278	7	11	0	75	2	-17
6	193	5	15	9	200	5	-8	-15	117	3	0	8	103	3	-1	1	320	8	-3
7	153	4	24	10	38	4	-4	-14	173	5	-7	9	273	7	29	2	131	3	3
8	214	6	18	11	153	4	-8	-13	128	4	1	10	110	3	7	3	256	7	1
9	94	3	4	H,K= 1,		15		-12	239	6	9	11	273	7	7	4	228	6	3
10	149	4	-2	-9	141	4	-2	-11	65	4	15	12	173	5	-3	5	263	7	-10
11	126	3	0	-8	109	3	-2	-10	275	7	1	13	225	6	5	6	195	5	2
12	114	3	-1	-7	111	3	-1	-9	180	5	-2	14	116	4	0	7	399	10	20
13	148	4	5	-6	146	4	-2	-8	381	10	0	15	154	4	3	8	105	3	13
H,K= 1,		11		-5	91	3	2	-7	295	8	5	H,K= 2,		6	9	337	9	12	
-13	151	4	-1	-4	148	4	2	-6	488	12	-9	-15	129	4	-3	10	87	3	-4
-12	44	3	1	-3	136	4	-1	-5	289	7	-35	-14	41	3	0	11	162	4	-5
-11	155	4	-9	-2	143	4	6	-4	538	14	26	-13	214	6	-1	12	101	3	4
-10	45	2	2	-1	229	5	3	-3	189	5	23	-12	20	6	-2*	13	98	3	-5
-9	165	4	-0	0	119	3	2	-2	138	4	-39	-11	345	9	7	14	89	3	4
-8	96	3	-1	1	242	6	5	-1	82	3	-26	-10	36	4	4	H,K= 2,		10	
-7	298	8	14	2	129	3	2	0	185	5	-42	-9	357	9	15	-14	143	4	-1
-6	159	4	9	3	157	4	5	1	226	6	29	-8	74	2	-7	-13	83	3	3
-5	389	10	31	4	127	3	-3	2	421	11	25	-7	255	7	-7	-12	144	4	-4
-4	150	4	14	5	121	4	2	3	305	8	-24	-6	48	2	2	-11	123	4	3
-3	355	9	10	6	110	3	-5	4	396	10	-1	-5	268	7	-25	-10	164	4	1
-2	68	2	7	7	157	4	-4	5	350	9	4	-4	43	2	-5	-9	136	4	-4
-1	267	7	-5	8	92	3	2	6	387	10	-11	-3	520	13	-26	-8	265	7	17
0	14	6	2*	9	175	5	-0	7	69	2	-8	-2	35	2	11	-7	183	5	7
1	276	7	1	H,K= 1,		17		8	388	10	17	-1	631	16	-32	-6	374	10	18
2	106	3	8	-6	179	5	2	9	105	3	2	0	156	4	-1	-5	220	6	3
3	350	9	21	-5	54	3	6	10	329	9	51	1	465	12	-21	-4	293	8	-0
4	234	6	13	-4	184	5	-6	11	212	6	5	2	5	12	0*	-3	177	5	-1
5	290	7	39	-3	40	3	-6	12	194	5	2	3	346	9	-16	-2	206	5	-5
6	192	5	7	-2	127	3	-5	13	188	5	10	4	147	4	10	-1	140	4	-6
7	221	6	3	-1	50	3	-5	14	144	4	-6	5	343	9	0	0	212	6	-10

STRUCTURE FACTORS CONTINUED FOR
YTTERBIUM BIS(C5ME5)DITHIODIETHYLVARBAMATE

PAGE 3

L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL
1	146	4	0	6	190	5	0	11	274	7	5	2	537	14	1	-2	252	6	-16
2	334	9	-12	7	89	3	-7	12	96	3	-3	3	324	8	-7	-1	140	4	-2
3	184	5	-7	8	198	5	-6	13	152	4	0	4	461	12	8	0	264	7	-3
4	377	10	8	9	79	3	2	14	58	3	-4	5	82	3	-8	1	229	6	-1
5	161	4	11	H,K=		2,	16	15	161	4	-2	6	384	10	9	2	146	4	-2
6	289	7	26	-8	78	3	-1	H,K=		3,	3	7	6	15	-6*	3	339	9	-5
7	149	4	-1	-7	151	4	3	-16	134	4	1	8	270	7	27	4	167	4	0
8	195	5	3	-6	58	3	-8	-15	126	3	-2	9	76	2	12	5	278	7	14
9	124	3	5	-5	157	4	-5	-14	174	5	-1	10	234	6	2	6	239	6	16
10	142	4	-2	-4	77	2	2	-13	127	4	-3	11	103	3	2	7	170	4	6
11	106	3	4	-3	120	3	-8	-12	154	4	6	12	198	5	-0	8	258	7	-11
12	173	5	-8	-2	118	3	-0	-11	134	4	24	13	102	3	1	9	152	4	2
13	93	3	-3	-1	113	3	-4	-10	145	4	8	14	165	4	5	10	154	4	1
H,K=		2,	12	0	140	4	-1	-9	224	6	-12	H,K=		3,	7	11	164	4	1
-12	155	4	-6	1	142	4	-8	-8	369	9	-2	-15	26	6	2*	12	112	3	-0
-11	39	3	7	2	102	3	3	-7	363	9	-12	-14	142	4	-6	13	136	4	-5
-10	194	5	-3	3	184	5	-5	-6	412	10	-13	-13	32	4	-6	H,K=		3,	11
-9	18	19	0*	4	76	3	1	-5	336	9	-18	-12	264	8	-3	-13	141	4	-1
-8	196	5	-6	5	165	4	-2	-4	333	8	-22	-11	23	5	0*-12	24	7	4*	
-7	45	3	2	6	107	3	2	-3	329	8	-24	-10	338	9	17	-11	164	4	-6
-6	252	6	-1	7	115	3	3	-2	85	3	-16	-9	17	12	9*-10	23	6	0*	
-5	85	3	8	H,K=		2,	18	-1	513	13	12	-8	248	6	10	-9	200	5	-2
-4	295	8	14	-3	129	4	-3	0	246	6	-0	-7	30	3	2	-8	102	3	5
-3	7	13	2*	-2	0	16	-11*	1	400	10	6	-6	246	6	-7	-7	260	7	6
-2	320	8	27	-1	119	3	0	2	497	13	-13	-5	121	3	2	-6	196	5	8
-1	80	2	-9	0	0	21	-2*	3	242	6	-19	-4	350	9	-9	-5	283	7	20
0	245	6	29	1	146	4	-3	4	481	12	15	-3	198	5	-4	-4	174	5	3
1	25	3	10	2	31	11	-2*	5	258	7	-8	-2	503	13	-24	-3	249	6	14
2	196	5	24	H,K=		3,	1	6	371	10	24	-1	87	2	6	-2	84	3	1
3	157	4	9	-16	55	3	-8	7	251	6	3	0	461	12	-24	-1	259	7	16
4	201	5	5	-15	156	4	-4	8	153	4	-1	1	103	3	-0	0	79	2	8
5	115	3	4	-14	76	3	1	9	196	5	30	2	290	7	-16	1	249	6	19
6	214	6	-7	-13	269	7	-7	10	228	6	8	3	131	3	7	2	138	4	13
7	29	4	1	-12	68	3	2	11	171	4	-1	4	256	7	-5	3	308	8	24
8	215	6	-8	-11	332	9	25	12	242	6	8	5	118	3	-9	4	134	4	13
9	47	3	5	-10	40	4	-6	13	162	4	1	6	365	9	6	5	288	7	11
10	174	5	-6	-9	385	10	-5	14	164	4	4	7	42	2	11	6	58	2	3
11	22	6	3*	-8	140	4	0	H,K=		3,	5	8	448	12	18	7	214	6	-6
H,K=		2,	14	-7	407	10	-15	-16	121	4	1	9	17	14	4*	8	24	4	-1
-11	92	3	6	-6	165	4	-14	-15	70	3	-3	10	281	7	-6	9	179	5	-11
-10	176	5	-0	-5	415	10	-31	-14	166	5	-5	11	37	10	1*	10	53	3	2
-9	103	3	2	-4	12	13	-4*	-13	49	6	3	12	133	4	-12	11	180	5	-6
-8	160	4	1	-3	495	13	-22	-12	230	7	-1	13	16	20	-14*	H,K=		3,	13
-7	86	3	-2	-2	501	13	27	-11	43	3	6	H,K=		3,	9	-12	61	3	7
-6	110	3	1	-1	352	9	-38	-10	278	7	27	-14	90	3	-3	-11	169	5	-2
-5	93	3	2	0	267	7	15	-9	72	2	1	-13	121	3	4	-10	78	3	-1
-4	144	4	-7	1	538	14	39	-8	369	9	-1	-12	150	4	0	-9	163	4	-5
-3	148	4	4	2	383	10	40	-7	212	5	-2	-11	123	3	-2	-8	24	4	6
-2	254	6	-13	3	293	7	-28	-6	394	10	-0	-10	237	6	-5	-7	146	4	-3
-1	168	4	4	4	332	8	23	-5	244	6	-9	-9	146	4	1	-6	17	14	-3*
0	280	7	-19	5	478	12	21	-4	381	10	-28	-8	232	6	16	-5	202	5	-3
1	93	3	6	6	98	3	12	-3	30	2	-2	-7	215	6	10	-4	33	3	2
2	210	5	-11	7	578	15	22	-2	419	11	-34	-6	193	5	-1	-3	268	7	-7
3	58	3	0	8	16	18	3*	-1	16	4	2*	-5	248	6	-5	-2	120	3	-4
4	165	4	0	9	478	13	58	0	531	14	-9	-4	145	4	-12	-1	280	7	-11
5	67	3	-1	10	90	3	8	1	197	5	13	-3	184	5	6	0	88	3	-3

STRUCTURE FACTORS CONTINUED FOR
YTTERBIUM BIS(C5ME5)DITHIODIETHYLVARBAMATE

PAGE 4

L	FOB	SG	DEL	L	FOB	SG	DEL	L	FCB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL
1	226	6	-8	12	154	4	-1	5	443	11	8	-1	417	11	-20	2	274	7	-4
2	7	18	-4*	14	171	5	1	6	83	3	-3	0	212	6	0	3	59	4	4
3	188	5	-13	H,K= 4, 2				7	234	6	16	1	233	6	-6	4	217	6	-8
4	36	2	11	-16	114	3	-1	8	66	2	10	2	244	6	-0	5	15	10	4*
5	183	5	-8	-15	105	3	-1	9	156	4	19	3	191	5	-6	6	168	4	-4
6	80	2	4	-14	137	4	-6	10	124	3	-1	4	135	4	3	7	53	2	0
7	199	5	-7	-13	111	3	0	11	207	5	-3	5	286	7	15	8	190	5	-4
8	90	3	-6	-12	198	5	6	12	157	4	-1	6	68	2	12	9	27	4	4
9	174	5	-8	-11	61	2	11	13	169	5	3	7	316	8	5	10	164	4	-5
10	31	6	-8*	-10	235	6	4	14	103	3	-2	8	77	3	-5	H,K= 4, 14			
H,K= 3, 15				-9	127	3	-1	H,K= 4, 6				9	243	6	-3	-10	136	4	0
-9	128	4	4	-8	245	6	-9	-16	18	8	1*	10	111	3	1	-9	67	3	-1
-8	106	3	-2	-7	301	8	-8	-15	116	3	-1	11	126	3	-7	-8	136	4	1
-7	100	3	-2	-6	216	6	-25	-14	19	6	12*	12	97	3	1	-7	58	2	1
-6	116	3	3	-5	224	6	-13	-13	169	4	-3	H,K= 4, 10				-6	159	4	2
-5	112	3	0	-4	479	12	-31	-12	26	4	4	-14	116	3	-2	-5	73	3	7
-4	104	3	3	-3	53	2	-9	-11	234	6	6	-13	49	3	-2	-4	213	6	-2
-3	183	5	2	-2	771	20	13	-10	49	3	3	-12	107	3	-4	-3	86	3	5
-2	113	3	1	-1	130	3	27	-9	262	7	14	-11	79	3	2	-2	236	6	-4
-1	208	5	-2	0	544	14	-2	-8	76	3	-1	-10	166	4	-2	-1	135	4	3
0	140	4	-1	1	280	7	-8	-7	319	8	-3	-9	135	4	-7	0	200	5	2
1	154	4	-3	2	306	8	-18	-6	81	3	-6	-8	238	6	8	1	142	4	8
2	167	4	0	3	406	10	10	-5	420	11	-10	-7	157	4	5	2	143	4	-1
3	98	3	-1	4	319	8	7	-4	53	2	5	-6	272	7	27	3	113	3	2
4	150	4	-3	5	285	7	6	-3	515	13	-16	-5	159	4	8	4	148	4	-4
5	126	3	-1	6	361	9	2	-2	80	2	2	-4	191	5	11	5	94	3	-3
6	100	3	-3	7	121	3	-1	-1	562	14	-12	-3	166	4	8	6	180	5	-7
7	153	4	-3	8	239	6	22	0	16	5	14*	-2	163	4	2	7	101	3	3
8	82	3	4	9	85	3	15	1	439	11	0	-1	214	6	0	8	162	4	-5
H,K= 3, 17				10	190	5	10	2	25	2	6	0	275	7	7	H,K= 4, 16			
-6	151	4	-4	11	163	4	-3	3	317	8	2	1	197	5	1	-8	80	3	-0
-5	59	3	2	12	170	5	1	4	9	12	-1*	2	383	10	16	-7	141	4	3
-4	142	4	-4	13	121	3	0	5	318	8	-4	3	180	5	12	-6	81	3	0
-3	46	3	2	14	148	4	-1	6	52	2	1	4	354	9	20	-5	140	4	-2
-2	114	3	-3	H,K= 4, 4				7	347	9	26	5	136	4	10	-4	112	3	2
-1	31	6	0*	-16	71	3	-3	8	21	6	17*	6	188	5	1	-3	105	4	-4
0	124	3	-6	-15	138	4	-2	9	280	7	-6	7	99	3	2	-2	122	3	3
1	43	4	-2	-14	55	3	1	10	38	3	2	8	143	4	-6	-1	109	3	-3
2	167	4	-8	-13	141	4	-9	11	168	4	-5	9	107	3	6	0	82	3	2
3	60	3	1	-12	26	5	3*	12	65	3	5	10	175	5	-9	1	150	4	-3
4	175	5	-4	-11	146	4	18	13	132	4	-7	11	94	3	-3	2	54	3	1
H,K= 4, 0				-10	125	3	5	H,K= 4, 8				H,K= 4, 12				3	168	5	-1
-16	119	3	-4	-9	229	6	-11	-15	88	3	-5	-12	125	3	-0	4	71	3	6
-14	193	5	-3	-8	261	7	-9	-14	78	3	2	-11	20	7	-7*	5	123	3	1
-12	324	8	7	-7	376	10	-19	-13	137	4	-5	-10	145	4	-5	H,K= 5, 1			
-10	311	8	-11	-6	347	9	2	-12	81	3	-2	-9	17	11	2*-16	47	3	-2	
-8	322	8	-12	-5	424	11	-16	-11	238	6	-3	-8	167	4	-4	-15	112	4	-7
-6	298	8	-14	-4	199	5	-19	-10	60	4	-3	-7	88	3	1	-14	38	3	-6
-4	510	13	-64	-3	494	13	-11	-9	253	6	13	-6	213	6	-5	-13	208	5	-8
-2	698	18	-65	-2	171	4	-26	-8	106	3	17	-5	75	3	-0	-12	18	22	-5*
0	561	14	-22	-1	361	9	-21	-7	182	5	9	-4	229	6	5	-11	265	7	17
2	421	11	1	0	331	8	2	-6	161	4	-4	-3	10	17	8*-10	16	14	5*	
4	418	11	16	1	477	12	-16	-5	227	6	-10	-2	270	7	18	-9	211	6	-12
6	634	16	15	2	346	9	0	-4	153	4	-9	-1	21	4	1*	-8	127	4	4
8	475	13	45	3	560	14	15	-3	350	9	-7	0	310	8	11	-7	249	6	-16
10	263	7	19	4	277	7	-1	-2	142	4	-1	1	51	4	3	-6	151	4	-16

STRUCTURE FACTORS CONTINUED FOR
YTTERBIUM BIS(C5H5)DITHIODIETHYLVARBAMATE

PAGE 5

L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL
-5	400	10	-30	-11	0	16	-8*	-13	64	3	-1	-2	105	3	-3	-12	202	5	-4
-4	81	2	3	-10	200	5	21	-12	101	3	-2	-1	244	6	-5	-11	86	3	2
-3	701	18	-40	-9	124	3	12	-11	104	4	-3	0	86	3	1	-10	195	5	17
-2	49	2	3	-8	264	7	-9	-10	150	4	-9	1	201	6	-5	-9	154	4	4
-1	599	15	-20	-7	195	5	1	-9	206	5	-6	2	9	15	-7*	-8	203	5	-14
0	70	2	-5	-6	366	9	-27	-8	140	4	3	3	166	5	-3	-7	222	6	-4
1	447	11	-5	-5	160	4	-8	-7	263	7	14	4	28	4	-1	-6	280	7	-21
2	135	3	-7	-4	489	13	-25	-6	175	5	20	5	192	5	2	-5	146	4	7
3	400	10	4	-3	14	7	-12*	-5	199	5	15	6	65	3	1	-4	322	8	-5
4	67	2	-6	-2	481	12	-22	-4	241	6	7	7	195	5	0	-3	204	5	-3
5	416	11	9	-1	85	2	-5	-3	118	3	8	8	54	3	-3	-2	330	9	-3
6	55	2	-5	0	364	9	-20	-2	245	6	-2	H,K=	5,	15	-1	267	7	-8	
7	372	10	27	1	224	6	1	-1	202	5	3	-9	92	4	-3	0	332	9	-21
8	23	15	10*	2	385	10	6	0	247	6	2	-8	119	3	4	1	264	7	-3
9	203	5	12	3	208	5	2	1	298	8	9	-7	92	3	-1	2	384	10	1
10	44	6	3	4	362	9	2	2	231	6	12	-6	122	3	-0	3	176	5	-1
11	153	5	-4	5	59	2	1	3	246	6	14	-5	135	4	-2	4	297	8	9
12	66	3	0	6	329	9	29	4	236	6	12	-4	98	3	1	5	41	3	5
13	141	4	-2	7	43	3	-6	5	133	4	8	-3	184	5	-1	6	275	7	23
H,K=	5,	3	3	8	257	7	15	6	173	4	-2	-2	94	3	1	7	52	2	11
-16	103	3	1	9	68	2	-2	7	85	3	0	-1	153	4	4	8	256	7	2
-15	84	3	2	10	196	5	-2	8	148	4	-1	0	125	4	3	9	92	3	-7
-14	95	3	-0	11	119	3	2	9	126	3	-3	1	89	3	3	10	177	5	-6
-13	96	3	1	12	151	4	1	10	117	3	1	2	118	3	-5	11	145	4	-2
-12	117	3	-1	13	89	3	3	11	151	4	-1	3	83	3	1	12	121	3	-7
-11	149	4	14	H,K=	5,	7	7	H,K=	5,	11	4	4	111	3	-1	13	83	3	-5
-10	110	3	4	-15	29	4	8	-13	91	3	1	5	128	3	-0	H,K=	6,	4	4
-9	191	5	-13	-14	134	4	-4	-8	77	3	-2	6	89	3	4	-16	79	3	0
-8	220	6	-8	-13	38	3	-2	-7	227	6	-6	H,K=	5,	17	-15	117	3	-0	
-7	234	6	-7	-12	187	5	-3	-6	70	3	-0	-4	114	3	1	-14	80	3	-2
-6	306	8	-4	-11	54	3	0	-5	236	6	9	-3	51	3	0	-13	153	4	-6
-5	299	8	-23	-10	219	6	-8	-4	45	4	8	-2	105	3	3	-12	54	2	-2
-4	348	9	-19	-9	53	2	7	-3	197	5	19	-1	50	3	-1	-11	196	5	-2
-3	315	8	-14	-8	197	5	13	-2	67	2	6	0	128	4	-2	-10	109	3	7
-2	358	9	-23	-7	67	2	2	-1	262	7	17	1	52	3	3	-9	195	5	11
-1	267	7	-18	-6	264	7	-5	0	151	4	1	H,K=	6,	0	0	-8	197	5	-7
0	347	9	3	-5	35	2	4	1	342	9	17	-16	99	3	0	-7	258	7	-13
1	362	9	-7	-4	407	10	-6	2	181	5	1	-14	139	4	-3	-6	199	5	2
2	479	12	13	-3	93	3	-3	3	314	8	5	-12	228	6	-1	-5	267	7	-26
3	428	11	4	-2	502	13	-6	4	114	3	3	-10	234	6	2	-4	62	2	-1
4	298	8	5	-1	108	3	-18	5	197	5	-0	-8	277	7	-27	-3	303	8	-20
5	254	7	0	0	432	11	-5	6	33	3	3	-6	376	10	-29	-2	23	3	2
6	164	4	1	1	73	2	-8	7	126	3	-6	-4	491	13	-31	-1	329	8	-12
7	114	3	15	2	237	6	-1	8	27	4	-1	-2	573	15	-46	0	227	6	1
8	119	3	18	3	26	3	2	9	158	4	-6	0	372	9	2	1	375	10	-13
9	157	4	-0	4	238	6	-2	10	60	3	-2	2	291	7	-8	2	314	8	4
10	180	5	-8	5	59	2	13	H,K=	5,	13	4	4	323	8	-14	3	335	9	5
11	178	5	-3	6	300	8	26	-11	119	3	-0	6	431	11	31	4	215	6	-10
12	173	5	-2	7	93	3	4	-10	28	7	2*	8	317	8	21	5	220	6	21
13	122	3	-3	8	275	7	-6	-9	140	4	1	10	155	4	-1	6	109	3	16
H,K=	5,	5	5	9	23	5	-5*	-8	15	9	9*	12	133	4	-4	7	245	6	10
-16	115	3	2	10	168	4	-8	-7	153	4	-5	H,K=	6,	2	8	129	3	-4	
-15	54	3	-3	11	25	5	7*	-6	15	11	10*	-16	99	3	-2	9	234	6	-4
-14	136	4	-3	12	118	3	-7	-5	183	5	-9	-15	55	3	5	10	162	4	-1
-13	15	9	-0*	H,K=	5,	9	9	-4	59	2	-1	-14	145	4	-3	11	191	5	-2
-12	173	5	-8	-14	78	3	1	-3	221	6	-6	-13	36	5	-7	12	104	3	-2

STRUCTURE FACTORS CONTINUED FOR
YTTERBIUM BIS(C5ME5)DITHIODIETHYLVARBAMATE

PAGE 6

L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL
H,K=	6,	6		11	90	3	-5	-3	89	3	3	-11	147	4	-2	-12	171	4	2
-15	128	4	-1	H,K=	6,	10		-2	185	5	-1	-10	209	5	5	-11	63	2	4
-14	17	17	-2*	-13	64	3	1	-1	89	3	3	-9	200	5	14	-10	174	5	-3
-13	184	7	-1	-12	71	2	0	0	123	3	1	-8	247	6	2	-9	26	7	3*
-12	41	3	3	-11	66	2	-3	1	59	3	-3	-7	199	5	-2	-8	165	4	5
-11	220	6	-10	-10	135	4	1	2	104	3	-1	-6	162	4	-8	-7	29	4	14
-10	15	18	0*	-9	91	3	-3	3	59	3	-2	-5	97	3	-2	-6	212	6	20
-9	232	6	14	-8	230	6	-1	4	157	4	-2	-4	38	2	3	-5	53	3	12
-8	51	3	-3	-7	102	3	-0	5	83	3	2	-3	68	2	-4	-4	301	8	7
-7	237	6	5	-6	277	7	6	6	180	5	-4	-2	194	5	2	-3	66	3	6
-6	44	3	-5	-5	115	4	12	H,K=	6,	16		-1	205	5	-3	-2	270	7	7
-5	291	8	-9	-4	227	6	18	-6	47	3	-0	0	332	9	-6	-1	18	24	1*
-4	19	5	-1*	-3	99	3	13	-5	110	3	1	1	278	7	5	0	210	5	11
-3	357	9	-17	-2	163	4	21	-4	88	4	0	2	316	8	-13	1	17	8	8*
-2	38	2	3	-1	149	4	13	-3	98	3	2	3	224	6	3	2	224	6	30
-1	366	9	-11	0	230	6	17	-2	116	3	-1	4	155	4	18	3	73	2	4
0	131	3	-12	1	215	6	14	-1	100	3	-1	5	223	6	22	4	306	8	15
1	291	7	-2	2	270	7	16	0	94	3	0	6	105	3	8	5	81	3	-3
2	136	4	4	3	183	5	-8	1	133	4	3	7	227	6	-5	6	286	7	-0
3	276	7	-1	4	201	5	-3	2	73	3	2	8	178	5	-7	7	71	3	-3
4	23	5	7*	5	131	4	-3	H,K=	7,	1		9	181	5	-6	8	170	4	-9
5	345	9	37	6	108	3	-3	-16	15	17	-11*	10	183	5	-4	9	45	3	-3
6	95	3	-8	7	96	3	3	-15	125	4	-2	11	127	3	-6	18	113	3	-10
7	312	8	0	8	118	3	-2	-14	0	15	-8*	12	136	4	-3	H,K=	7,	9	
8	46	3	1	9	90	3	3	-13	169	4	-5	H,K=	7,	5	-14	93	3	1	
9	211	5	-5	10	156	4	-3	-12	24	5	-5*	-15	35	4	-11	-13	66	3	3
10	45	3	1	H,K=	6,	12		-11	229	6	-2	-14	157	4	-1	-12	96	3	-3
11	142	4	-4	-12	110	3	3	-10	73	2	5	-13	10	18	-24*	-11	95	4	5
12	26	5	3*	-11	4	15	1*	-9	216	6	1	-12	195	5	-7	-10	101	3	-4
H,K=	6,	8		-10	139	4	0	-8	87	3	1	-11	32	3	-0	-9	151	4	-4
-15	91	3	-6	-9	28	5	-4	-7	256	7	-16	-10	220	6	-3	-8	145	4	-3
-14	37	4	-1	-8	165	4	-2	-6	100	3	-2	-9	89	3	-4	-7	180	5	-5
-13	140	4	1	-7	39	5	1	-5	309	8	-23	-8	229	6	12	-6	179	5	-1
-12	72	3	1	-6	171	5	-8	-4	116	3	-1	-7	79	2	2	-5	165	4	7
-11	141	4	-1	-5	29	4	2	-3	397	10	-35	-6	228	6	-9	-4	202	5	7
-10	145	4	-3	-4	192	5	-5	-2	138	4	-3	-5	65	3	5	-3	130	3	13
-9	149	4	1	-3	60	2	-1	-1	351	9	-15	-4	208	5	-13	-2	169	4	18
-8	175	5	-2	-2	200	5	-10	0	78	2	6	-3	45	2	-1	-1	164	4	15
-7	175	4	17	-1	0	15	-8*	1	201	5	-3	-2	249	6	-9	0	162	4	17
-6	129	3	6	0	226	6	-10	2	67	2	-5	-1	82	3	-7	1	221	6	8
-5	289	7	-1	1	72	3	3	3	248	6	-10	0	257	7	-15	2	131	4	3
-4	85	2	7	2	210	5	-7	4	28	4	-0	1	159	4	1	3	187	5	3
-3	317	8	-1	3	69	3	-1	5	334	9	22	2	285	7	6	4	158	4	-7
-2	121	3	6	4	200	5	-3	6	11	16	2*	3	118	3	6	5	99	3	1
-1	284	7	-0	5	11	14	-1*	7	350	9	7	4	310	8	29	6	174	5	-3
0	126	3	3	6	171	4	-8	8	44	4	-2	5	68	2	5	7	82	3	1
1	234	6	14	7	8	15	6*	9	203	5	-4	6	314	8	-2	8	127	3	3
2	129	3	9	8	140	4	-5	10	75	3	1	7	6	19	-12*	9	131	4	-0
3	213	6	26	H,K=	6,	14		11	139	4	-2	8	260	7	-5	H,K=	7,	11	
4	119	3	9	-10	97	3	-2	12	58	4	0	9	68	3	-2	-12	29	4	-7
5	246	6	8	-9	75	3	1	H,K=	7,	3		10	180	5	-4	-11	101	3	-1
6	88	3	0	-8	94	3	1	-16	76	3	4	11	80	3	3	-10	58	3	1
7	199	5	-5	-7	62	3	-3	-15	110	3	2	H,K=	7,	7	-9	150	4	2	
8	64	2	-1	-6	127	4	-3	-14	68	3	-1	-15	23	6	-4*	-8	66	3	-4
9	146	4	-7	-5	71	3	-2	-13	127	3	-2	-14	135	4	-5	-7	194	5	-1
10	94	3	1	-4	188	5	-2	-12	125	3	-1	-13	39	7	-0	-6	39	3	-2

STRUCTURE FACTORS CONTINUED FOR
YTTERBIUM BIS(C5ME5)DITHIODIETHYLVARBAMATE

PAGE 7

L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL
-5	202	5	-5	6	274	7	-1	8	94	3	-1	-12	79	3	-1	-13	117	3	-1
-4	25	7	6*	8	220	6	-1	9	176	5	-5	-11	83	3	-7	-12	29	3	6
-3	167	4	-4	10	149	4	-4	10	119	4	-5	-10	103	3	2	-11	133	3	-5
-2	34	3	-2	H,K=		8,	2	H,K=		8,	6	-9	100	3	-3	-10	34	3	0
-1	175	5	-5	-16	90	3	-8	-15	105	4	1	-8	141	4	-0	-9	139	4	-7
0	83	3	2	-15	43	3	7	-14	21	10	15*	-7	95	3	-3	-8	48	2	7
1	214	6	-4	-14	121	3	0	-13	141	4	-3	-6	137	4	-2	-7	210	5	8
2	100	3	2	-13	35	3	7	-12	0	15	-8*	-5	101	3	0	-6	69	2	2
3	186	5	-6	-12	156	4	-1	-11	160	4	-3	-4	147	4	-0	-5	274	7	0
4	75	2	-0	-11	107	3	0	-10	30	3	-2	-3	106	4	-2	-4	45	2	8
5	149	4	-9	-10	189	5	5	-9	166	4	-5	-2	178	5	-3	-3	240	6	5
6	42	3	-4	-9	153	4	2	-8	51	4	-6	-1	86	3	-2	-2	56	2	4
7	129	3	-3	-8	187	5	13	-7	193	5	13	0	215	6	-3	-1	232	6	2
8	49	3	-1	-7	147	4	3	-6	22	6	-3*	1	79	3	-2	0	109	3	8
H,K=		7,	13	-6	205	5	-9	-5	255	7	18	2	179	5	4	1	253	6	14
-10	26	6	6*	-5	56	2	-2	-4	10	15	-12*	3	77	3	-8	2	94	3	9
-9	111	3	1	-4	235	6	-11	-3	260	7	12	4	117	3	-6	3	288	7	12
-8	13	14	4*	-3	45	2	-1	-2	19	5	12*	5	93	3	-2	4	35	5	-3
-7	125	3	-0	-2	267	7	-5	-1	220	6	20	6	121	3	-6	5	224	6	0
-6	34	3	4	-1	181	5	-1	0	29	4	-4	7	76	3	-4	6	24	8	-10*
-5	166	4	-1	0	212	6	-6	1	237	6	27	8	135	4	-1	7	161	4	-5
-4	68	2	6	1	220	6	-4	2	49	2	8	H,K=		8,	12	8	76	3	0
-3	173	5	-2	2	214	6	6	3	286	7	10	-11	13	15	-5*	9	132	3	-1
-2	41	3	6	3	161	4	12	4	0	21	-3*	-10	128	4	5	10	55	3	1
-1	134	4	-5	4	265	7	20	5	283	7	2	-9	28	10	-6*	H,K=		9,	3
0	15	7	10*	5	39	8	7*	6	8	14	-12*	-8	148	5	5	-15	52	2	4
1	129	4	-3	6	240	6	-4	7	206	5	-2	-7	17	11	-5*	-14	47	2	-2
2	15	10	8*	7	79	3	-3	8	14	15	2*	-6	149	4	3	-13	69	2	2
3	163	4	0	8	175	5	-3	9	136	4	-11	-5	0	16	-6*	-12	60	3	-1
4	47	3	0	9	138	4	0	10	25	5	-3*	-4	146	4	-1	-11	109	3	4
5	174	5	-7	10	130	4	-7	H,K=		8,	8	-3	14	10	8*	-10	139	4	3
6	48	4	-1	11	102	3	-4	-14	49	4	5	-2	134	4	-2	-9	149	4	2
H,K=		7,	15	H,K=		8,	4	-13	114	3	-3	-1	32	3	2	-8	158	4	6
-7	63	3	-5	-15	97	3	-2	-12	54	3	1	0	159	4	2	-7	159	4	13
-6	85	3	1	-14	57	3	0	-11	107	3	-1	1	39	5	-3	-6	139	4	13
-5	100	3	-1	-13	96	3	-3	-10	72	2	-1	2	175	5	5	-5	172	4	17
-4	87	3	4	-12	80	3	-1	-9	116	4	-3	3	8	16	4*	-4	89	2	5
-3	127	4	-5	-11	154	4	-3	-8	78	3	2	4	160	4	1	-3	200	5	10
-2	77	4	3	-10	103	3	-2	-7	157	4	-5	5	45	4	5	-2	159	4	5
-1	121	3	-2	-9	207	5	2	-6	71	3	5	6	134	4	-9	-1	210	5	11
0	89	3	1	-8	109	3	9	-5	236	6	3	H,K=		8,	14	0	233	6	20
1	96	3	-4	-7	219	6	3	-4	65	3	10	-8	84	3	3	1	186	5	14
2	104	3	4	-6	84	3	3	-3	249	6	11	-7	46	3	1	2	202	5	14
3	102	3	-3	-5	168	4	-5	-2	116	3	8	-6	117	4	-0	3	140	4	8
H,K=		8,	0	-4	107	3	-9	-1	151	4	7	-5	66	3	3	4	133	3	-1
-16	93	3	-0	-3	168	4	-8	0	187	5	2	-4	148	4	1	5	91	3	1
-14	133	4	2	-2	130	3	7	1	124	4	4	-3	64	3	6	6	86	2	-0
-12	184	5	-7	-1	248	6	-3	2	180	5	-1	-2	133	4	-1	7	84	3	-3
-10	196	5	-9	0	156	4	8	3	177	5	4	-1	41	5	2	8	129	3	1
-8	171	4	4	1	255	7	16	4	93	3	-1	0	109	3	5	9	110	3	-5
-6	254	7	-20	2	143	4	8	5	230	6	-4	1	55	3	3	H,K=		9,	5
-4	321	8	-25	3	242	6	21	6	58	4	4	2	105	3	0	-14	89	3	-2
-2	343	9	-18	4	110	3	7	7	167	5	-4	3	64	3	-3	-13	35	4	3
0	234	6	-16	5	195	5	-5	8	82	3	3	H,K=		9,	1	-12	96	3	-3
2	245	6	-11	6	88	3	1	9	97	3	-4	-15	101	3	2	-11	37	5	2
4	312	8	29	7	195	5	-7	H,K=		8,	10	-14	30	3	3	-10	134	4	-1

STRUCTURE FACTORS CONTINUED FOR
YTTERBIUM BIS(CSME5)DITHIODIETHYLVARBAMATE

PAGE 8

L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL
-9	46	2	-5	0	116	3	-5	-11	59	2	6	-2	34	3	-7	0	130	4	-1
-8	184	5	-4	1	191	5	4	-10	118	3	2	-1	212	5	-3	1	27	4	1
-7	69	2	-5	2	113	3	-2	-9	81	2	4	0	55	2	-3	2	125	3	3
-6	253	6	9	3	123	3	3	-8	171	4	-2	1	177	5	-3	H,K= 11,			
-5	69	2	-2	4	128	3	-2	-7	66	3	5	2	22	4	2	-14	24	6	-5*
-4	281	7	12	5	90	3	-8	-6	227	6	5	3	152	4	-1	-13	88	3	-4
-3	66	2	3	6	129	4	0	-5	52	2	10	4	20	9	14*	-12	32	3	3
-2	275	7	15	7	93	3	0	-4	217	6	7	5	149	4	-1	-11	89	3	0
-1	69	2	2	H,K= 9,				-3	73	2	9	6	39	3	-0	-10	39	3	4
0	242	6	17	-11	109	3	6	-2	177	5	6	7	125	3	-2	-9	113	3	1
1	107	3	3	-10	48	3	-6	-1	162	4	8	H,K= 10,				-8	31	3	4
2	205	5	4	-9	132	4	6	0	176	5	6	-12	45	2	3	-7	142	4	-6
3	33	3	-0	-8	68	3	2	1	145	4	6	-11	80	3	-3	-6	22	5	4*
4	181	5	-2	-7	118	3	1	2	179	5	-6	-10	73	2	3	-5	180	5	-10
5	11	14	2*	-6	55	2	2	3	84	3	-4	-9	81	2	2	-4	35	3	5
6	151	4	-4	-5	107	3	-4	4	171	4	-3	-8	88	3	-0	-3	159	4	-10
7	49	2	-2	-4	30	3	-3	5	64	2	-1	-7	104	3	1	-2	60	2	2
8	138	4	-5	-3	152	4	5	6	129	3	-2	-6	80	2	-6	-1	137	4	-4
9	80	3	2	-2	35	4	-6	7	81	3	0	-5	140	4	1	0	68	2	4
H,K= 9,				-1	168	4	4	8	104	3	-1	-4	72	3	-5	1	139	4	-6
-14	93	3	2	0	61	2	-2	H,K= 10,				-3	164	4	-1	2	38	5	-3
-13	13	16	2*	1	160	4	-8	-14	25	4	4	-2	87	3	3	3	189	5	-6
-12	109	3	1	2	50	3	0	-13	69	3	-2	-1	133	4	-3	4	45	3	0
-11	20	5	6*	3	133	4	-2	-12	52	3	5	0	89	3	2	5	188	5	-4
-10	88	3	-1	4	10	13	-3*	-11	91	3	-3	1	134	4	-4	6	42	6	-2
-9	26	3	4	5	131	3	-1	-10	74	3	-1	2	69	2	-0	7	113	4	-5
-8	125	3	-2	H,K= 9,				-9	127	3	-3	3	154	4	-3	H,K= 11,			
-7	36	6	0	-8	16	6	9*	-8	96	3	2	4	62	3	3	-13	61	2	2
-6	208	5	-3	-7	127	3	6	-7	167	4	-7	5	158	4	-2	-12	69	2	1
-5	54	4	6	-6	32	3	9	-6	74	3	2	6	53	3	-1	-11	84	3	1
-4	251	6	-2	-5	147	4	3	-5	216	6	-1	H,K= 10,				-10	89	3	5
-3	59	2	10	-4	47	3	9	-4	93	3	0	-11	56	3	-3	-9	91	3	-1
-2	194	5	2	-3	146	4	3	-3	198	5	-2	-10	96	3	1	-8	81	2	-2
-1	72	3	3	-2	14	15	2*	-2	151	4	-1	-9	59	2	-5	-7	96	3	1
0	164	4	1	-1	123	4	1	-1	191	5	-3	-8	88	3	-2	-6	91	3	0
1	51	2	-2	0	22	5	11*	0	157	4	2	-7	60	3	-3	-5	113	3	1
2	198	5	4	1	121	4	6	1	168	4	-7	-6	85	2	4	-4	102	3	0
3	44	3	-0	2	13	14	-3*	2	71	2	2	-5	64	2	-0	-3	137	4	-0
4	213	5	1	3	117	3	-0	3	143	4	-1	-4	110	3	2	-2	111	3	-2
5	29	5	1	H,K= 10,				0	18	5	6*	-3	88	3	0	-1	138	4	2
6	191	5	-2	-14	104	3	0	5	99	3	-4	-2	136	4	1	0	128	3	-0
7	22	6	1*	-12	85	3	-2	6	49	3	0	-1	104	3	1	1	115	3	1
8	114	3	-4	-10	110	3	-5	7	97	3	-2	0	138	4	2	2	102	3	1
H,K= 9,				-8	150	4	-14	8	89	3	1	1	102	3	-0	3	93	3	-1
-12	100	3	-0	-6	202	6	5	H,K= 10,				2	125	3	3	4	89	3	-0
-11	83	3	2	-4	177	5	3	-13	91	3	3	3	73	2	-1	5	89	3	-2
-10	92	3	-5	-2	208	5	4	-12	0	13	-13*	4	97	3	1	6	84	3	-1
-9	101	3	2	0	243	6	9	-11	102	3	2	H,K= 10,				7	95	3	3
-8	89	3	-5	2	228	6	-1	-10	17	7	4*	-8	110	3	1	H,K= 11,			
-7	81	2	-2	4	225	6	-7	-9	132	3	-0	-7	16	7	1*	-13	17	9	-3*
-6	103	3	-5	6	172	5	-7	-8	14	10	5*	-6	120	3	1	-12	99	3	-2
-5	77	3	-3	8	116	3	-5	-7	172	4	-8	-5	16	7	13*	-11	42	3	-3
-4	161	4	-2	H,K= 10,				-6	40	3	6	-4	128	4	-2	-10	123	3	-2
-3	128	3	-4	-14	71	3	0	-5	201	5	-12	-3	12	13	6*	-9	41	3	-4
-2	160	4	0	-13	43	3	2	-4	36	7	13*	-2	134	4	-0	-8	132	4	-6
-1	192	5	-0	-12	79	2	-0	-3	219	6	-10	-1	34	3	-3	-7	22	5	10*

STRUCTURE FACTORS CONTINUED FOR
YTTERBIUM BIS(C5ME5)DITHIODIETHYLVARBAMATE

PAGE 9

L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL	L	FOB	SG	DEL
-6	142	4	-7	0	39	4	-7	-6	9	14	5*	-2	92	3	3				
-5	11	15	1*	1	118	3	3	-5	99	3	-1	-1	97	3	4				
-4	157	4	-7		H, K=	12,	0	-4	0	17	-4*	0	74	3	1				
-3	58	2	-0	-12	94	3	-1	-3	98	3	1	1	91	3	-2				
-2	162	4	-6	-10	93	3	-1	-2	13	15	-1*	2	47	4	1				
-1	74	3	-8	-8	90	3	-4	-1	99	4	0	3	78	3	-2				
0	161	4	-2	-6	119	3	-5	0	8	15	-0*		H, K=	13,	5				
1	42	2	-6	-4	149	4	1	1	128	4	0	-10	69	3	3				
2	147	4	-1	-2	110	3	2	2	14	15	5*	-9	21	8	-9*				
3	15	7	-6*	0	100	3	-6	3	148	4	-2	-8	80	3	5				
4	136	4	-3	2	144	4	-2	4	8	17	3*	-7	36	5	2				
5	33	3	-6	4	162	5	-2		H, K=	12,	8	-6	73	3	-9				
6	118	3	-4		H, K=	12,	2	-9	78	3	3	-5	18	11	-5*				
	H, K=	11,	7	-12	80	3	-2	-8	45	3	9	-4	96	3	1				
-12	84	3	3	-11	58	3	3	-7	99	3	7	-3	17	17	-3*				
-11	27	4	9	-10	77	3	-0	-6	40	6	6	-2	107	3	-0				
-10	95	3	2	-9	61	4	6	-5	92	3	6	-1	29	4	1				
-9	37	3	2	-8	92	3	3	-4	58	3	6	0	117	4	-4				
-8	117	3	6	-7	36	3	-1	-3	75	3	-0	1	29	5	0				
-7	55	3	2	-6	118	3	2	-2	61	3	2	2	121	3	-4				
-6	129	3	11	-5	62	2	5	-1	71	3	-4		H, K=	13,	7				
-5	58	3	2	-4	123	3	2	0	49	3	0	-8	86	3	5				
-4	124	3	5	-3	79	3	6	1	97	4	-0	-7	18	9	5*				
-3	23	4	1*	-2	100	3	-3	2	38	6	-5	-6	105	3	9				
-2	112	3	0	-1	74	3	6		H, K=	12,	10	-5	17	12	6*				
-1	0	15	-5*	0	113	3	-3	-6	60	3	2	-4	86	3	4				
0	102	3	-5	1	52	3	3	-5	45	18	-10*	-3	31	4	3				
1	11	14	6*	2	129	4	-5	-4	59	4	-4	-2	74	3	3				
2	137	4	-3	3	44	4	-1	-3	60	3	-2	-1	25	5	-4*				
3	24	7	9*	4	133	4	-4	-2	96	3	3	0	83	3	1				
4	152	4	-3	5	55	3	-6	-1	62	3	-3		H, K=	14,	0				
5	30	4	3		H, K=	12,	4		H, K=	13,	1	-8	101	3	1				
	H, K=	11,	9	-12	55	3	4	-11	76	3	-1	-6	94	3	-5				
-10	61	3	-1	-11	75	3	-5	-10	34	4	5	-4	89	3	-5				
-9	79	3	-1	-10	58	3	1	-9	91	3	1	-2	87	3	-2				
-8	65	2	-2	-9	87	3	-1	-8	32	5	6	0	91	3	-1				
-7	78	3	2	-8	41	3	1	-7	103	3	1		H, K=	14,	2				
-6	73	3	-1	-7	88	3	-2	-6	36	5	3	-8	66	3	-3				
-5	72	3	4	-6	33	3	10	-5	112	3	0	-7	42	3	3				
-4	92	3	2	-5	93	3	-2	-4	44	3	2	-6	74	3	0				
-3	80	3	3	-4	36	5	6	-3	98	3	-5	-5	52	3	3				
-2	111	3	1	-3	92	5	-5	-2	31	4	-2	-4	90	3	-2				
-1	90	3	-1	-2	68	3	5	-1	100	3	-4	-3	64	3	4				
0	116	3	-2	-1	113	4	-3	0	22	6	-3*	-2	98	3	-1				
1	87	3	-1	0	78	3	2	1	110	3	-4	-1	54	4	3				
2	122	3	2	1	122	3	-1	2	23	8	8*	0	87	3	2				
3	65	3	-1	2	72	3	-2	3	100	4	-3		H, K=	14,	4				
	H, K=	11,	11	3	111	3	-3		H, K=	13,	3	-7	51	6	-2				
-8	28	4	2	4	57	3	1	-10	57	3	3	-6	38	3	-1				
-7	90	3	-0	5	114	3	-4	-9	56	3	1	-5	74	3	1				
-6	25	4	-1		H, K=	12,	6	-8	59	3	5	-4	44	3	2				
-5	89	3	-0	-11	90	3	-1	-7	58	3	5	-3	93	3	-6				
-4	36	3	-3	-10	19	7	-2*	-6	58	3	-0	-2	51	4	1				
-3	106	3	-2	-9	99	3	-2	-5	53	3	3								
-2	56	3	-0	-8	8	15	-9*	-4	88	3	4								
-1	125	3	3	-7	100	3	-2	-3	72	3	3								

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720