Lawrence Berkeley National Laboratory

Recent Work

Title

GIANT RESONANCE STUDIES WITH HIGH ENERGY HEAVY IONS

Permalink

<https://escholarship.org/uc/item/3qp497t5>

Author

Doll, P.

Publication Date

1978-11-01

Submitted to Physical Review Letters

RECEIVED
LAWRENCE
BERKELEY LABORATORY

LBL-8289
Preprint c.2

JAN 11 1979

LIBRARY AND
DOCUMENTS SECTION

GIANT RESONANCE STUDIES WITH HIGH ENERGY HEAVY IONS

P. Doll, D. L. Hendrie, J. Mahoney, A. Menchaca-Rocha,
D. K. Scott, T. J. M. Symons, K. Van Bibber,
Y. P. Viyogi, and H. Wieman

November 1978

Prepared for the U. S. Department of Energy
under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782

LBL-8289
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

GIANT RESONANCE STUDIES WITH HIGH ENERGY HEAVY IONS*

P. Doll^a, D. L. Hendrie^b, J. Mahoney, A. Menchaca-Rocha^c,
D. K. Scott, T. J. M. Symons, K. Van Bibber, Y. P. Viyogi^d,
and H. Wieman

Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

October 1978

Abstract

Evidence is presented for the excitation of giant multipole resonances in inelastic scattering of 315 MeV ^{16}O ions on ^{12}C , ^{58}Ni , and ^{208}Pb targets. Highly excited modes around 20 MeV of excitation energy in ^{208}Pb exhaust a large fraction of the energy weighted isoscalar E3 and E5 sum rule strengths.

For the study of energy dissipation processes in heavy-ion reactions it is important to understand the response function¹ of the nuclei involved in the collisions. It is natural to assume that in the initial reaction step the excitation of simple one-particle one-hole states absorbs part of the kinetic energy of relative motion. Therefore, giant multipole resonances may also be involved² if, in the adiabatic limit, the collision times are of the same order of magnitude as the corresponding quantum transition times.³ In addition, angular momentum matching conditions in heavy-ion reactions allow large angular momenta transfers. While several low-energy, heavy-ion scattering experiments (≤ 10 MeV/A)^{4,5,6} have reported the observation of giant quadrupole structures in nuclei, it was the goal

of the present experiment to improve these studies with a high energy ^{16}O beam at 315 MeV on ^{12}C , ^{58}Ni and ^{208}Pb , and to search for excitations of higher multipolarity, which have been reported so far only in electron scattering on ^{208}Pb .^{7,8}

The experiments were performed at the Lawrence Berkeley Laboratory 88-Inch Cyclotron. A beam of 315 MeV $^{16}\text{O}^{6+}$ ions was scattered from self-supporting ^{208}Pb (1.03 mg/cm^2), ^{58}Ni (1.67 mg/cm^2) and ^{12}C (0.36 mg/cm^2) foils. The reaction products were identified in the focal plane of a magnetic QSD spectrometer, using a counter which determines the position ($B\rho$), energy loss (ΔE) and time-of-flight (TOF) of the reaction products. Position spectra of inelastically scattered ^{16}O particles on ^{208}Pb , ^{58}Ni and ^{12}C targets are shown in Fig. 1 a, b, c. Single nucleon transfer reactions were also investigated to assess the competition between transfer and inelastic scattering as the energy loss mechanism in heavy-ion collisions. A more detailed presentation of these data will be given elsewhere.

All spectra exhibit pronounced structures in the continuum. The spectrum for ^{208}Pb , taken close to the grazing angle, shows strong excitation of giant resonance structures, which become pronounced for ^{58}Ni only at angles more forward than the grazing angle ($\theta_g^{\text{lab}} \approx 9^\circ$). For the $^{16}\text{O} + ^{12}\text{C}$ system the grazing angle is $\sim 2^\circ$ in the laboratory system and at such forward angles hydrogen contamination in the ^{12}C target obscured the ^{16}O energy spectrum in the giant resonance region.

Compared to hadron ($A \leq 4$) scattering data, the spectra reveal an improved peak to continuum ratio, possibly equalled only by a recent high-energy ^6Li scattering experiment.⁹ The contributions from sequential decay of ejectiles following α , proton or neutron pick-up lie outside the region of the giant resonance structures in all three nuclei at 315 MeV

incident energy. These processes are known¹⁰ to be a severe problem in inclusive light-ion scattering spectra. At high excitation energies there is some contribution from ^{15}O ground state groups as indicated in Fig. 1. The broad peak at a Q-value of approximately 6 MeV in all spectra is attributed to the 3^- , 6.13 MeV projectile excitation, Doppler-broadened by the subsequent γ -decay and by underlying target states. The kinematically expected positions of projectile excitations built up on the strongest low-lying target states are indicated by arrows on the horizontal axis of each spectrum (see Fig. 1).

We interpret the 10.8 MeV ($\Gamma \approx 2.5$ MeV) structure in ^{208}Pb ,¹⁴ the 16 MeV ($\Gamma \approx 3.2$ MeV) side group in ^{58}Ni ¹⁴ (the 13.1 MeV group is found in electron scattering¹² to be of E3 nature), and the 25.3 to 26.7 MeV ($\Gamma \approx 4$ MeV) groups in ^{12}C ¹⁰ as the known isoscaler giant quadrupole resonances in these nuclei. The structure at 13.6 MeV in ^{208}Pb (see Fig. 1a) which is more pronounced in the present experiment than in light-ion scattering experiments, has been identified¹³ previously as the giant monopole resonance. The distinctive structure of the spectra in the giant resonance region invites a detailed comparison with light-ion coincidence experiments especially for the ^{12}C case.¹⁰ At a scattering angle of 3° the excitation of giant resonance structures seems to be even stronger than at 4° (Fig. 1c), but is superimposed on a 10 to 15 MeV wide bump from a hydrogen contamination in the ^{12}C target.

One of the most striking and novel features of the spectra for ^{208}Pb is the observation of a pronounced gross structure peak at an excitation energy of approximately 20 MeV. No structure around this

excitation energy was found in the inelastic scattering of ^{16}O at 140 MeV,⁶ nor has any hadron scattering experiment revealed gross structure in this region. However, two electron scattering^{7,8} experiments and several theoretical calculations^{1, 14} infer a strong concentration of the $L = 3$ and $L = 5$ strength function at similar excitation energies. In particular the structures found at 18.1 and 19.7 MeV in the present experiment are well reproduced by continuum RPA calculations.^{1,14} An additional hint that these highly excited structures in ^{208}Pb represent higher multipole modes comes from the simple liquid drop model,³ which predicts that, for increasing target mass (A), the response times $\tau \propto A^{2/3}/L^{3/2}$ of higher multipolarities (L) become better matched for the projectile collision times typical of this experiment.

Angular distributions were measured (see Fig. 2) for low-lying discrete states and the highly excited broad structures. Only typical examples are discussed here. The 25.3 MeV and 26.7 MeV components in ^{12}C were analyzed separately (after subtracting a smooth background through the minima in the spectra as indicated in Fig. 1 by dashed curves) and revealed no significant differences in the diffraction patterns. The same result holds for the 18.1 MeV and 19.7 MeV groups in ^{208}Pb . DWBA calculations have been performed using optical model parameters obtained for ^{208}Pb at an incident energy of 315 MeV (set Q in Ref. 15); the same parameters were used for ^{58}Ni . For ^{12}C the E18 potential from Ref. 16 was used. We included 250 partial waves for ^{208}Pb and 150 partial waves for ^{58}Ni and ^{12}C . A complex coupling plus Coulomb excitation form factor was used. The resulting curves are shown in Fig. 2. No attempt was made to improve the fit to the data by adjusting the phase angle between the nuclear and

Coulomb interaction amplitudes. The DWBA cross-sections are normalized to the data at the grazing angle to give the Coulomb excitation strengths β_{CE} ($R_{CE} = 1.2 A^{1/3}$). We obtained, for example, $\beta_{CE} = 0.41, 0.09$ and 0.10 for the low-lying $4.44 \text{ MeV } (2^+)$, $2.46 \text{ MeV } (4^+)$ and $2.61 \text{ MeV } (3^-)$ transitions in ^{12}C , ^{58}Ni and ^{208}Pb , respectively, in good agreement with known $B(EL)$ values.^{10,17,11} The energy-weighted sum rule strengths were deduced on the assumption of a uniform mass distribution¹⁸ ($r_0 = 1.2 \text{ fm}$) and result in $25 \pm 15\%$ E2 strength for the 25.3 to 26.7 MeV groups in ^{12}C , $40 \pm 15\%$ E2 strength for the 16 MeV group in ^{58}Ni , and $87 \pm 20\%$ E2 strength for the 10.8 MeV group in ^{208}Pb . If we assume that the pronounced structures around 20 MeV in ^{208}Pb belong to the $3\hbar\omega$ components of the isoscalar 3^- and 5^- giant resonances, and first exhaust 100% of the E3 sum rule strength (using about 30% of the observed cross section at the grazing angle) we are left with 10 to 20% of the E5 sum rule strength. Approximately the same total strength is predicted by the RPA calculations,^{1,14} but with more equal division between the modes. Although the quoted E2 strengths for ^{58}Ni and ^{208}Pb agree reasonably well with hadron scattering data,¹¹ the E2 strength in ^{12}C is almost 3 times larger than is found in a recent light-ion coincidence experiment.¹⁰ These results, combined with the suggestive l -dependent angular distributions observed for ^{12}C (see Fig. 2), make high energy heavy-ion scattering a promising tool for the study of giant resonances.

In this letter, we have presented the most striking evidence yet for the excitation of giant resonances with heavy-ion beams. Whereas it is clear that improvements in the data will be achieved using coincidence

techniques, as in the case of light-ion scattering,¹⁰ we emphasize that we have already observed pronounced structures of high multipole states in a straightforward singles experiment. This suggests that there are exciting prospects for mapping the nuclear response function of high multipole resonances in nuclei with even higher energy beams, which are required to match the corresponding quantum transition times.

The authors would like to thank G. R. Satchler and A. J. Baltz for help in performing the DWBA calculations, and H. Faraggi, C. K. Gelbke, K. T. Knöpfle and A. Shotter for valuable discussions.

* The work was supported by the Nuclear Physics Division of the Department of Energy

- a. Nato Fellow, on leave from MPI, Heidelberg, West Germany.
 - b. Present address: University of Maryland, College Park, Maryland.
 - c. Present address: Universidad Nacional Autonoma de Mexico, Mexico.
 - d. IAEA Fellow, on deputation from Bhabha Atomic Research Centre, Calcutta, 700064, India.
1. G. Bertsch and S. F. Tsai, Phys. Reports 18C, 126 (1975).
 2. R. A. Broglia, C. H. Dasso, G. Pollarolo, and A. Winther, Phys. Rev. Lett. 41, 25 (1978).
 3. J. P. Bondorf, Proc. Int. School of Physics (Enrico Fermi), eds. H. Faraggi and R. A. Ricci, (North Holland, 1976) p. 221.
 4. R. R. Betts, S. B. DiCenzo, M. H. Mortensen, and R. L. White, Phys. Rev. Lett., 39, 1183 (1977).
 5. M. Buenerd, D. Lebrun, J. Chauvin, Y. Gaillard, J. M. Loiseaux, P. Martin, G. Perrin, and P. de Saintignon, Phys. Rev. Lett. 40, 1482, (1978).
 6. A. Guterman, D. Ashery, J. Alster, D. K. Scott, M. S. Zisman, C. K. Gelbke, H. Wieman, and D. L. Hendrie, submitted to Phys. Rev. Lett.
 7. M. Sasao and Y. Torizuka, Phys. Rev. C15, 217 (1977).
 8. R. Pitthan, F. R. Buskirk, E. B. Dally, J. N. Dyer, and X. K. Maruyama, Phys. Rev. Lett. 33, 849 (1974).
 9. K. T. Knöpfle, G. J. Wagner, P. Doll, D. L. Hendrie and H. Wieman. Contr. Int. Conf. Nuclear Interactions, 1978, Canberra, Australia.

10. H. Riedesel, K. T. Knöpfle, H. Breuer, P. Doll, G. Mairle, and G. J. Wagner, Phys. Rev. Lett. 41 377 (1978), and references therein.
11. D. H. Youngblood, J. M. Moss, C. M. Rozsa, J. D. Bronson, A. D. Bacher, and D. R. Brown, Phys. Rev. 13C, 994 (1976).
12. R. Pitthan, G. M. Bates, J. S. Beachy, F. R. Buskirk, E. B. Dally, D. H. Dubois, J. N. Dyer and S. J. Dowalick, to be published.
13. D. H. Youngblood, C. M. Rozsa, J. M. Moss, D. R. Brown and J. D. Bronson, Phys. Rev. Lett. 39, 1188 (1977).
14. J. Speth, E. Werner and W. Wild, Phys. Reports 33C, 127 (1977).
15. C. Olmer, M. Mermaz, M. Buenerd, C. K. Gelbke, D. L. Hendrie, J. Mahoney, D. K. Scott, M. H. Macfarlane, and S. C. Pieper, Phys. Rev. C18, 205 (1978).
16. J. G. Cramer, R. M. DeVries, D. A. Goldberg, M. S. Zisman and C. F. Maguire, Phys. Rev. C14, 2158 (1976).
17. D. C. Kocher, R. L. Auble, Nucl. Data Sheets 19, 494 (1976).
18. G. R. Satchler, Nucl. Phys. A195 1, (1972), and private communication.

FIGURE CAPTIONS

Fig. 1 Energy spectra for inelastic scattering of ^{16}O at 315 MeV on (a) ^{208}Pb , (b) ^{58}Ni and (c) ^{12}C . The arrows on the horizontal axis indicate expected positions of projectile excitations built up on low-lying target states. The label ^{15}O denotes the position of low-lying transitions in the $(^{16}\text{O}, ^{15}\text{O})$ reaction.

Fig. 2 Angular distributions of low-lying and highly-excited groups for inelastic ^{16}O scattering on ^{208}Pb and ^{12}C . The solid curves represent DWBA calculations with the angular momentum transfers shown.

XBL 7810-11613

Fig. 1

XBL 7810-11525

Fig. 2

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720