UC San Diego

UC San Diego Previously Published Works

Title

Evolutionary links between intra- and extracellular acid-base regulation in fish and other aquatic animals.

Permalink

<https://escholarship.org/uc/item/3rx9d9x0>

Journal

Journal of experimental zoology. Part A, Ecological and integrative physiology, 333(6)

ISSN

2471-5638

Authors

Tresguerres, Martin
Clifford, Alexander M
Harter, Till S
et al.

Publication Date

2020-07-01

DOI

10.1002/jez.2367

Peer reviewed

Evolutionary links between intra- and extracellular acid-base regulation in fish and other aquatic animals

Running title: Acid-base regulatory mechanisms

Martin Tresguerres^{1*}, Alexander M. Clifford¹, Till S. Harter¹, Jinae N. Roa¹, Angus B. Thies¹, Daniel P. Yee¹, Colin J. Brauner²

¹Marine Biology Research Division, Scripps Institution of Oceanography, University of California San Diego, 9500 Gilman Drive, La Jolla, CA, USA

²Department of Zoology, University of British Columbia, 6270 University Boulevard, Vancouver, BC, V6T 1Z4, Canada

*E-mail: mtresguerres@ucsd.edu

Abstract

The acid-base relevant molecules carbon dioxide (CO₂), protons (H⁺), and bicarbonate (HCO₃⁻) are substrates and end products of some of the most essential physiological functions including aerobic and anaerobic respiration, ATP hydrolysis, photosynthesis, and calcification. The structure and function of many enzymes and other macromolecules are highly sensitive to changes in pH, and thus maintaining acid-base homeostasis in the face of metabolic and environmental disturbances is essential for proper cellular function. On the other hand, CO₂, H⁺ and HCO₃⁻ have regulatory effects on various proteins and processes, both directly through allosteric modulation and indirectly through signal transduction pathways. Life in aquatic environments

presents organisms with distinct acid-base challenges that are not found in terrestrial environments. These include a relatively high CO_2 relative to O_2 solubility that prevents internal $\text{CO}_2/\text{HCO}_3^-$ accumulation to buffer pH, a lower O_2 content that may favor anaerobic metabolism, and variable environmental CO_2 , pH and O_2 levels that require dynamic adjustments in acid-base homeostatic mechanisms. Additionally, some aquatic animals purposely create acidic or alkaline microenvironments that drive specialized physiological functions. For example, acidifying mechanisms can enhance O_2 delivery by red blood cells, lead to ammonia trapping for excretion or buoyancy purposes, or lead to CO_2 accumulation to promote photosynthesis by endosymbiotic algae. On the other hand, alkalinizing mechanisms can serve to promote calcium carbonate skeletal formation. This non-exhaustive review summarizes some of the distinct acid-base homeostatic mechanisms that have evolved in aquatic organisms to meet the particular challenges of this environment.

Keywords: ammonia, coral reef, hemoglobin, hypoxia, ocean acidification, oxygen transport, preferential pH_i regulation, proton pump, soluble adenylyl cyclase, symbiosome

Introduction

Proper cellular function depends on a series of tightly regulated, enzymatically catalyzed biochemical reactions, and therefore all organisms have evolved mechanisms to maintain appropriate conditions in their extra- and intracellular compartments. In biological systems, hydrogen ions (H^+) are produced and consumed through various chemical reactions. Due to its extremely large charge to size ratio, H^+ immediately react with the electron cloud of adjacent molecules including the carboxy and amino functional groups of proteins thereby altering their ionization status, conformation, and function. Thus, the ability to regulate intracellular pH (pHi) is essential for life.

The origin of pHi regulatory mechanisms is most likely rooted in the first proto-cells. It has been theorized that an important step in the origin of the first cell was the trapping of ribozymes and substrates within a membrane vesicle bilayer (Koch & Silver, 2005). An ability to maintain protoplasm pH within a range compatible with ribozyme activity would have enabled faster reaction rates and some degree of independence from the surrounding environment, providing a strong selective advantage. Regardless of its evolutionary origin, pHi regulation is essential for all extant organisms because intracellular acid-base homeostasis is continuously challenged by net H^+ production from ATP-consuming reactions. During aerobic conditions, the synthesis of ATP through mitochondrial oxidative phosphorylation acts as a major sink for H^+ that closely matches the H^+ production from ATP hydrolysis (Hochachka & Mommsen, 1983). However, O_2 limitation typically leads to increased H^+ production through anaerobic pathways, which, together with a lag in the mitochondrial H^+ sink, can acidify pHi. Moreover, intracellular acidification can occur even during aerobic conditions if the outwardly directed CO_2

partial pressure (PCO_2) gradient decreases due to an increase in PCO_2 in the extracellular fluids or the environment. In this case, some of the CO_2 that is produced as an end product of cellular respiration will accumulate inside the cell and combine with water to produce H^+ according to the reactions shown in Figure 1A.

These pH-dependent reactions have important implications for acid-base physiology: (1) in the presence of carbonic anhydrase (CA), an enzyme that catalyzes the left most reaction, equilibria are reached virtually instantaneously; (2) an increase in CO_2 leads to increased $[\text{H}^+]$ and thus has an acidifying effect; (3) most biological fluids and aquatic environments have a pH between 6-8, resulting in HCO_3^- as the dominant carbon species (Fig. 1B); and (4) active acidification and alkalinization of a compartment can be used to modulate the ratio between the different carbon species, such as to accumulate CO_2 , HCO_3^- , or CO_3^{2-} . Also important for acid-base regulation, H^+ , HCO_3^- , and CO_3^{2-} (as well as NH_4^+) are charged molecules and therefore require carrier proteins to cross lipid membranes. On the other hand, CO_2 (as well as O_2 and NH_3) as a gas can rapidly diffuse through membranes and the process can be further facilitated by aquaporin and Rhesus (Rh) channel proteins (Musa-Aziz, Chen, Pelletier, & Boron, 2009).

To avoid the adverse effects of intracellular acidification, cells must be able to buffer excess H^+ or actively extrude them at a rate equal to that of production. While all organisms routinely experience acid-base stress, life in aquatic environments poses particular acid-base challenges that are not found in terrestrial environments. In water, acid-base equivalents may be dissolved as ions, such as H^+ and HCO_3^- , whereas in gaseous air, the only acid-base equivalent is CO_2 . In addition, the solubility of CO_2 in

water is about 30 times higher than that of O_2 , and the diffusion ratio between the two gases is higher in water compared to air. Also, the O_2 content in water is much lower than in air. Thus, the ventilatory volume of water required to meet the O_2 demands of water breathers is far in excess of that required to ensure efficient excretion of respiratory CO_2 from blood to the water. Therefore, water breathers cannot elevate blood PCO_2 and $[HCO_3^-]$ to values as high as air-breathers, and thus they have a correspondingly lower internal fluid HCO_3^- -buffering capacity (Dejours, 1994). In addition, ventilatory regulation of CO_2 excretion is not an effective strategy for acid-base regulation in water breathers because decreases in ventilation rate to elevate blood PCO_2 would limit O_2 uptake, and blood PCO_2 is already so low that hyperventilation is largely ineffective in reducing blood PCO_2 (Gilmour, 2001). As a result, systemic acid-base regulation in aquatic animals is largely dependent upon active transport of acid-base equivalents in exchange for counter ions between the animal and the environment. The first part of this article will discuss the most common acid-base disturbances in aquatic environments and some intra- and extra-cellular acid-base regulatory strategies that are uniquely used by aquatic animals.

In addition to challenging acid-base homeostasis, variations in CO_2 , pH and HCO_3^- can be important modulators of physiological processes. The second part of this article will discuss three such examples: (1) the frequent pH_i changes experienced by red blood cells (RBCs) as they circulate between the respiratory surfaces and the tissues, and how these pH_i changes plays an essential role in the delivery of O_2 throughout the body; (2) the effect of pH on ammonia metabolism, and how active acidification of intra- and extracellular compartments can be used to promote ammonia

excretion or accumulation, and (3) the extreme pH microenvironments that are generated by coral cells, and their roles in mediating metabolic communication with their photosynthetic symbionts and in promoting skeletal calcification. Some of the potential evolutionary links between pHi regulation and other physiological processes are emphasized throughout this article.

Common acid-base disturbances in aquatic environments

In many water bodies, photosynthetic activity during the day may outpace respiration and result in elevated environmental O₂ (hyperoxia), reduced PCO₂ (hypocapnia), and elevated pH. However at night, respiration in the absence of photosynthesis can result in hypoxia, elevated PCO₂ (hypercapnia), and decreased pH. Hypercapnia is particularly evident in environments with high densities of organisms and slow water flow such as lakes, swamps, kelp forests, reefs, mangroves, and tide pools (Duarte, Ferreira, Wood, & Val, 2013; Hofmann et al., 2011; Kline et al., 2012; Truchot & Duhamel-Jouve, 1980). Another example of environmental hypercapnia is the gradual elevation of PCO₂ in water bodies due to increased absorption of anthropogenic CO₂ emissions (“ocean acidification” and “freshwater acidification”; Caldeira & Wickett, 2003; Van deWaal, Verschoor, Verspagen, van Donk, E. & Huisman, 2009).

In the majority of naturally occurring cases of environmental hypercapnia, PCO₂ remains lower than that of the internal fluids of the organism. However, the reduced PCO₂ gradient between the internal fluids and the environment limits excretion of endogenously produced CO₂, leading to an elevation in PCO₂ and a higher [H⁺] (and [HCO₃⁻]) by law of mass action (Fig. 1A, Melzner et al., 2009). But in environments with

a very high density of respiring biomass at night (Furch & Junk, 1997), in the proximity of geological CO₂ seeps (Hall-Spencer et al., 2008), and in high-density aquaculture systems (Ellis, Urbina, & Wilson, 2017), environmental PCO₂ can be elevated above the internal PCO₂ of an organism. In these cases, CO₂ will diffuse into the animal down its partial pressure gradient, and induce a much more pronounced acidosis.

A metabolic acidosis in fish is routinely observed after exhaustive exercise (e.g. Milligan & Wood, 1986) and during exposure to environmental hypoxia (e.g. Thomas & Hughes, 1982) due to increased reliance on anaerobic metabolism and the higher H⁺ production associated with ATP depletion relative to ATP production. Sessile aquatic invertebrates living in the intertidal zone may likewise experience hypoxia or anoxia during aerial emersion at low tide, as their gas exchange surfaces are ineffective in air, or the animal needs to minimize gas exchange to avoid desiccation, or both (Bayne, Bayne, Carefoot, & Thompson, 1976). A metabolic alkalosis typically develops in the blood of fish upon the consumption of a large meal (“alkaline tide”), which is due to the secretion of H⁺ into the stomach and the absorption of HCO₃⁻ into the blood (e.g. Wood, Kajimura, Mommsen, & Walsh, 2005). Conversely, a blood metabolic acidosis develops after feeding in agastric fish, which is due to the secretion of HCO₃⁻ into the gastrointestinal tract and the absorption of H⁺ into the blood (“acidic tide”) (Wood, Bucking, & Grosell, 2010).

Basic concepts of pH regulation

The logarithmic pH scale, where $\text{pH} = -\log [\text{H}^+]$ (Sørensen, 1909) easily masks relative changes in the actual [H⁺] within a fluid, which is the ion that directly interacts

with molecules. As such, at any point of the pH scale, a 1 unit pH change represents a 10-fold change in $[H^+]$, a 0.3 pH unit change is a ~2-fold change in $[H^+]$, and a 0.1 pH unit change is a ~25% change in $[H^+]$. Consequently, the range of the pH scale over which changes are observed matters greatly when assessing the magnitude of an acid-base disturbance. For example, a decrease from pH 7.4 to 7.3 reflects a 10 nM increase in $[H^+]$, but a decrease from pH 8.0 to 7.9 reflects only a 2.5 nM increase. Thus, the H^+ load associated with a pH change from 7.4 to 7.3 is four-fold larger than the H^+ load that changes pH from 8.0 to 7.9, and will require proportionally more resources in terms of buffering capacity or energy to remove the excess H^+ from a given compartment and prevent adverse effects on cellular function.

Intracellular buffering is a first line of defense against fluctuations in pH_i . The total intracellular buffering capacity is determined by the sum of the HCO_3^- and non- HCO_3^- buffering systems, which can bind and release H^+ to lessen an acidosis or an alkalosis, respectively. The main component of the non- HCO_3^- system is imidazole of the histidine groups in side chains of amino acids. Their pK_a is in the range of ~6.0-7.0, which is close to the pH_i set point of most cells and therefore histidine groups are particularly effective at buffering excess H^+ during physiologically relevant decreases in pH_i . For instance, muscle contraction generates a large H^+ load that can induce muscle fatigue and contractile failure (Jarvis, Woodward, Debold, & Walcott, 2018) and accordingly, fish muscle cells contain large amounts of the histidine-rich dipeptides carnosine, anserine, and balenine resulting in a non- HCO_3^- buffering capacity that is ~2-3 times greater than that of other tissues (Walsh & Milligan, 1989). Also, different types of muscle fibers create different acidic conditions that determine the cellular strategy of

homeostasis. Fast-twitch white muscle relies largely on anaerobic metabolism that produces H^+ (Kieffer, 2000), whereas slow-twitch, red muscle relies on aerobic metabolism and predominantly produces CO_2 . Thus, to compensate for a more rapid and pronounced metabolic acidosis, white muscle contains more histidine-rich dipeptides that elevate the intracellular buffering capacity over that of red muscle (Dolan et al., 2019).

As a rapid, reversible, and passive strategy buffers are critical to maintaining pH_i homeostasis; however, their capacity is finite and when overwhelmed, pH_i regulation hinges on active mechanisms (Fig. 2). In most animal cells, Na^+/K^+ -ATPase (NKA) activity drives H^+ excretion by secondarily active transporters such as the ubiquitous Na^+/H^+ exchanger isoform 1 (NHE1). NKA activity can also drive HCO_3^- uptake *via* Na^+/HCO_3^- cotransporters (NBCs) and Na^+ -dependent Cl^-/HCO_3^- exchangers (NDCBEs); an increase in intracellular $[HCO_3^-]$ reacts with and decreases $[H^+]$, thus is equivalent to active H^+ extrusion (reviewed in Casey, Grinstein, & Orlowski, 2010). In cancer cells growing in acidic microenvironments, the V-type H^+ -ATPase (VHA) is another active H^+ extruding mechanism that helps counteract intracellular acidification (reviewed in Torigoe et al, 2002). Notably, H^+ excretion by VHA does not depend on NKA activity; however, VHA-dependent H^+ excretion must occur in concert with the net transport of a counter ion (typically Cl^- excretion or Na^+ absorption) (Tresguerres, 2016).

Cells that produce lactate as the end product of fermentation typically excrete H^+ together with lactate through monocarboxylate- H^+ cotransporters (MCTs). However, several important differences exist between aquatic animals and mammals. For example, white muscle in fish expresses MCTs at a relatively low abundance and

203 retains a significant proportion of the lactate that is produced during exhaustive
204 exercise, which allows for the localized replenishment of glycogen stores *in situ* from
205 lactate (reviewed in Weber, Choi, Gonzalez, & Omlin, 2016). Likewise, aquatic
206 invertebrates that produce imino acids (“opines”) in the final step of fermentation may
207 retain these end products intracellularly for later oxidization when aerobic conditions
208 return, or reconvert them into the original pyruvate and amino acid substrates (Ellington,
209 1983). In some cases, the accumulation of fermentative metabolites may be associated
210 with a pronounced intracellular acidification that can inhibit glycogenic and
211 gluconeogenic metabolism through pH effects on enzyme activity and substrate or end-
212 product inhibition (Walsh & Milligan, 1989).

213 Cases of intracellular alkalinization are less common than those of acidification.
214 Nonetheless, when an alkaline load is experienced, some cells use anion exchangers
215 (AEs) that excrete HCO_3^- in exchange for Cl^- , thereby acting as “acid-loading”
216 transporters that help counteract an alkaline load. And some cells experiencing elevated
217 intracellular Ca^{2+} levels use Plasma Membrane Ca^{2+} -ATPases (PMCAs) to extrude Ca^{2+}
218 in exchange for extracellular H^+ thereby acidifying the cytosol (reviewed in Casey et al.,
219 2010).

220 Active pH_i regulation requires the ability to sense disturbances from a set point
221 and to trigger compensatory responses. One such mechanism relies on pH-dependent
222 amino acid conformational changes that render acid-secreting proteins such as NHE1
223 inactive when pH increases, and acid-loading proteins such as AE3 inactive when pH
224 decreases (reviewed in Casey et al., 2010). Other molecular acid-base sensors are
225 coupled to signal transduction pathways (Tresguerres, Buck, & Levin, 2010), of which

the soluble adenylyl cyclase (sAC) is arguably the best characterized in aquatic animals (Tresguerres, Barott, Barron, & Roa, 2014). This evolutionarily conserved enzyme is directly stimulated by HCO_3^- to produce the ubiquitous second messenger cyclic adenosine monophosphate (cAMP) (Chen et al., 2000) that regulates the activity of effector proteins *via* PKA-dependent phosphorylation, Exchange Protein Activated by cAMP (EPAC), and cAMP gating of membrane channels (Fig. 3). In many systems, sAC activity is directly stimulated by HCO_3^- ; however, in the presence of CA, changes in $[\text{HCO}_3^-]$ almost instantaneously reflect changes in $[\text{CO}_2]$ and $[\text{H}^+]$, enabling sAC to indirectly sense extracellular and intracellular acid-base disturbances of any origin (Tresguerres, Levin, & Buck, 2011). Indeed, sensing cytosolic $[\text{HCO}_3^-]$ may be more rapid and reliable for pHi regulation than sensing $[\text{H}^+]$ because the repeated association and dissociation with cytosolic macromolecules slows down H^+ diffusion, which may confound the detection of a H^+ load (Chang & Oude Elferink, 2014). To our knowledge, a role of sAC in pHi regulation has only been established in corals (Barott, Barron, & Tresguerres, 2017). However, given that corals are phylogenetically deeply rooted metazoans, the role of sAC in pHi regulation most likely extends to most other animal Phyla. In addition, the presence of sAC in the nucleus of mammalian (Zippin et al., 2004) and shark (Roa & Tresguerres, 2017) cells suggests a conserved role in regulating gene expression in response to changing acid-base conditions (Fig. 3).

Aquatic animals have specialized cells (“acid-base ionocytes”) on the gills and skin epithelia that actively maintain blood pH by exchanging acid-base equivalents with the environment; this centralized strategy of pHe homeostasis lessens the need for pHi regulation by every individual cell (reviewed in Larsen et al. 2014). The identity and

kinetics of the ion transporting proteins involved in pHe regulation varies greatly between species and environments; however, these proteins are all derived from those involved in pHi regulation (i.e. CAs, NKA, NHEs, NBCs, NDBCEs, VHA, AEs, sAC). The differential placement of transport proteins in the ionocyte's apical or basolateral membrane allows for the vectorial transport of H^+ and HCO_3^- between the internal fluids and the external environment for the purposes of pHe regulation. Similarly, many of the ion-transporting proteins and regulatory pathways involved in pHi regulation take on novel physiological functions when regulating the pH of other internal compartments to promote systemic O_2 transport, ammonia excretion, biomineralization, and CO_2 delivery to photosymbionts.

Coupled pH regulation and preferential pHi regulation

A severe acid-base challenge that overwhelms the capacity for pHe regulation will result in a disturbance to both pHe and pHi. However, some animals are able to tightly regulate pHi even when the pHe defenses have been breached. This capacity gives them an unusual resilience to environmental hypercapnia, and possibly during other acid-base challenges as discussed below.

Environmental hypercapnia results in a sustained elevation in blood PCO_2 and potentially a large acid-base disturbance, and permits investigating the relative contributions of pHi and pHe regulation. Exposure to severe hypercapnia induces a rapid and large reduction in pHe that is often followed by a less pronounced reduction in pHi. During continuous exposure to CO_2 , complete pHi recovery is associated with significant (>50-100%) pHe recovery, and therefore this acid-base regulatory pattern

has been termed “coupled pH regulation” (Shartau, Baker, Crossley & Brauner, 2016). Coupled pH regulation during exposure to a respiratory acidosis has been observed in most amphibians, reptiles, and mammals investigated to date, and thus coupled pH regulation appears to be widespread amongst vertebrates (Shartau, Baker, et al., 2016). This pattern has also been observed in the few invertebrates where simultaneous measurements of tissue pHe and pHi have been made during exposure to hypercapnia, and include the land snail (*Otala lactea*; Barnhart & McMahon, 1988), a deep-sea bivalve (*Acesta excavate*; Hammer, Kristiansen, & Zachariassen, 2011), and the peanut worm (*Sipunculus nudus*; Pörtner, Reipschläger, & Heisler, 1998).

However, some aquatic vertebrates display a different pattern of acid-base regulation, where tissue pHi is completely regulated despite large reductions in pHe during the first few hours of exposure to environmental hypercapnia, and in some cases pHi even increases relative to control values despite a large reduction in pHe (Baker et al., 2009). This pattern of rapid and tight pHi regulation during a transient reduction in pHe during acute CO₂ exposure has been termed “preferential pHi regulation”. Importantly, it does not imply the absence of pHe regulation; just that pHi regulation may be virtually instantaneous and more robust than pHe regulation.

In white sturgeon (*Acipenser transmontanus*) exposed to a PCO₂ of 6 kPa, pHe was reduced by 0.7 pH units within 15 minutes (Baker et al., 2009). Despite the severe blood acidosis, heart pHi increased by 0.05 pH units and was maintained over the subsequent 90 min of hypercapnia (Baker, 2010). Similarly, when sturgeon were exposed to 3 and 6 kPa PCO₂, the pHi of brain, liver and white muscle was tightly regulated. At that time, blood pH was reduced below the blood buffer line indicating a

295 net acid excretion from the cells to the blood, and this reflects the preferential regulation
296 of the intra- over the extracellular compartment (Baker et al., 2009). Therefore in
297 hypercapnic sturgeon pHi regulation occurs more rapidly than pHe regulation, resulting
298 in a H⁺ transfer from the cells to the blood that is faster than their excretion to the
299 environment at the gills.

300 In addition to sturgeon, preferential pHi regulation has been observed in a
301 number of other fishes including the armored catfish (*Pterygoplichthys pardalis*), the
302 marbled swamp eel (*Synbranchus marmoratus*), the striped catfish (*Pangasianodon*
303 *hypophthalmus*), and three species of gar (*Lepisosteus oculatus*, *L. osseus*, and
304 *Atractosteus spatula*) (reviewed in Shartau et al., 2020). Preferential pHi regulation was
305 also observed in the late stage developing embryos of the common snapping turtle
306 (*Chelydra serpentine*; Shartau, Crossley, Kohl, & Brauner, 2016) and American alligator
307 (*Alligator mississippiensis*; Shartau, Crossley, Kohl, Elsey, & Brauner, 2018). Thus, it
308 has been proposed that preferential pHi regulation may be a general trait in vertebrate
309 embryos prior to the complete development of the extracellular compartments and
310 structures for acid-base regulation (Shartau, Baker, et al, 2016). This trait is then either
311 retained or lost during development depending on the animal's life history and/or the
312 environment. For example, the greater siren (*Siren lacertian*) is the only tetrapod known
313 to retain preferential pHi regulation into adulthood (Heisler, Forcht, Ultsch, & Anderson,
314 1982), and this eel-like amphibian inhabits stagnant wetlands where the water is
315 routinely hypercapnic and hypoxic, or even anoxic (Ultsch, 1973, Ultsch & Anthony
316 1973). To our knowledge, there is no evidence for preferential pHi regulation in

invertebrates; however, very few studies have simultaneously measured pHe and pHi in invertebrates during the early stages of exposure to severe hypercapnia.

While exposure to elevated PCO₂ has been used as a tool to induce a large acidosis to investigate the presence or absence of preferential pHi regulation, a more common acid-base disturbance may result from anaerobic metabolism due to an O₂ limitation. Armored catfish can tolerate long periods of severe hypoxia or even anoxia (Armbruster, 1998), where they preferentially regulate pHi of brain, heart, liver and white muscle despite a severe blood acidosis (Harter et al., 2014). Also white sturgeon are considered hypoxia-tolerant (Cech & Crocker, 2002, however not to the extent of the air-breathing armored catfish. During a hypoxic challenge induced by air exposure, sturgeon demonstrated some capacity for preferential pHi regulation in heart and brain; however, the pHi of liver and white muscle decreased during this challenge (Shartau, Baker, & Brauner, 2017). These findings on armored catfish and white sturgeon may point towards a link between preferential pHi regulation and the ability to survive in O₂ limited environments; however, this hypothesis must be tested with further comparative studies. In addition questions remain regarding the molecular and cellular mechanisms underlying preferential pHi regulation that provide exciting avenues to further investigate the evolution and prevalence of preferential pHi regulation.

The role of red blood cell pHi on systemic gas transport

RBCs in the vertebrate circulatory system come in close contact with every other cell type and carry high concentrations of hemoglobin (Hb) and CA that enhance O₂ delivery and CO₂ removal in all tissues. Cardiovascular gas transport is modulated by

the RBC microenvironment and by the fluctuations in pH that occur between arterial and venous blood, with every pass through the circulatory system. The diffusion of CO₂ into the blood at the tissue capillaries causes a decrease in blood pH, whereas the systemic excretion of CO₂ at the gas exchange surfaces causes an increase in pH (Fig. 4A). These cyclical changes in pH between the arterial and venous systems are dampened by the buffer capacity of the blood that, in water breathers, is largely provided by non-bicarbonate buffers. In fishes, these buffers are proteins in the plasma and Hb and organic phosphates within the RBCs and, while most species rely on both intra- and extracellular buffers, their individual contributions may vary greatly among the major fish lineages. On one end of the spectrum are the Antarctic icefishes that have lost RBCs and Hb from the circulation, and where the only non-bicarbonate buffers in the blood are histidine-rich plasma proteins (Feller, Poncin, Aittaleb, Schyns, & Gerday, 1994). On the other hand, lamprey rely almost entirely on buffers within the RBC, which prevent pH_i fluctuations but lead to large arterial-venous changes in blood pH_e (Tufts & Boutilier, 1989). In most vertebrates, Hb is the principal blood buffer and some H⁺ binding sites on Hb are modulated by oxygen. This Haldane effect links the transport of O₂ and CO₂ in the blood and is particularly important in teleost fishes (Harter & Brauner, 2017).

Due to their charge, extracellular H⁺ have no direct access to RBC intracellular buffers, such as Hb. However, the Jacobs-Stewart cycle (Jacobs & Stewart, 1942) links the activities of H⁺ to the transmembrane fluxes of CO₂ and HCO₃⁻ by the reversible hydration and dehydration reactions in the plasma and within the RBC (Fig. 4B). CO₂ crosses the RBC membrane by diffusion (Wagner, 1977), a process that may be facilitated by aquaporin 1 and RhAG (Muza-Asis et al., 2009), while HCO₃⁻ is

transported by the abundant RBC AE, Band 3 (Romano and Passow 1984). Within the RBC the equilibration between CO_2 , HCO_3^- and H^+ is catalyzed by CA (Itada & Forster, 1977), whereas the blood plasma of many vertebrates lacks CA (Henry & Swenson, 2000) and often contains CA inhibitors that ensure an absence of CA activity against a background of constant RBC lysis that releases soluble CA (Henry, Gilmour, Wood, & Perry, 1997). Without CA activity, the uncatalyzed CO_2 hydration and dehydration reactions in the plasma are slow and typically the rate-limiting step in the Jacobs-Stewart cycle (Motaïs, Fievet, Garcia-Romeu, & Thomas, 1989). However, at the tissue capillaries, membrane-bound, plasma-accessible CA (paCA) isoforms that are unaffected by plasma CA inhibitors (Gervais & Tufts, 1998; Heming et al., 1993), will accelerate the Jacobs-Stewart cycle and effectively link pHe and RBC pHi.

In a theoretical steady state, H^+ are passively distributed across the RBC membrane in a Donnan-like equilibrium; however, the negative charge of Hb and organic phosphates favors a higher $[\text{H}^+]$ inside the RBC, resulting in a lower pHi relative to pHe (Jensen, 2004). This pH gradient across the RBC membrane is of physiological significance as it renders the blood an effective sink for CO_2 that removes the gas from the tissues. Due to the higher plasma pHe (typically 7.8-8 in fishes) and the relatively low pK_a of the CO_2 - HCO_3^- equilibrium [~ 6.1 ; (Boutilier et al., 1984)] more than 90% of CO_2 can be transported as HCO_3^- in the plasma. This increases the capacitance of blood for CO_2 far beyond the physical solubility of the gas in plasma and severely reduces the convection requirements for CO_2 excretion (Tufts & Perry, 1998).

The active regulation of RBC pHi is largely driven by the Na^+ and K^+ gradients that are generated by RBC NKA activity (Fig. 4C; Thomas & Egée, 1998). A decrease in

386 RBC pHi is typically due to a net K^+ efflux *via* a K^+ -2 Cl^- -cotransporter (KCC) and the
387 loss of Cl^- displaces H^+ from equilibrium *via* the Jacobs-Stewart cycle (Cossins &
388 Gibson, 1997). Whereas an increase in RBC pHi is driven by a net Na^+ influx, either
389 through Na^+ - K^+ - Cl^- -cotransporters (NKCC) or NHEs (Nikinmaa 2003). The β -
390 adrenergically activated NHEs (β -NHE) of teleosts are particularly effective regulators of
391 RBC pHi and presumably have evolved to protect O_2 transport by pH-sensitive Hb
392 during a systemic acidosis (Berenbrink et al., 2005). The binding of H^+ to Hb decreases
393 its affinity for O_2 and this Bohr effect (Bohr, Hasselbalch, & Krogh, 1904) describes the
394 prominent role of pH in fine-tuning cardiovascular O_2 transport in nearly all vertebrates
395 (Fig. 4A). Teleost fishes have exceptionally pH-sensitive Hbs with large Bohr
396 coefficients and in addition have a Root effect where H^+ binding prevents a complete O_2
397 saturation of Hb even at very high PO_2 (see Berenbrik et al 2005). Based on this high
398 pH-sensitivity of Hb, several physiological mechanisms have evolved in teleosts that
399 actively acidify the blood to increase O_2 unloading to specialized tissues.

400 Perhaps the best-known examples are the teleost *retia mirabilia*, vascular
401 counter-current exchangers that are coupled to acidifying tissues that trigger the Root
402 effect. This mechanism can produce PO_2 values of several hundred atmospheres
403 allowing teleosts with gas-filled bladders to regulate buoyancy at depth (Nielsen &
404 Munk, 1964; Pelster 1997) and to drive O_2 across large diffusion distances to their
405 avascular retinas (Wittenberg & Wittenberg, 1962). Similarly, the intestine of marine
406 teleosts, which secretes large amounts of HCO_3^- into the lumen, may acidify the blood
407 sufficiently to enhance Hb- O_2 unloading and thereby meet its high metabolically
408 demand for O_2 (Cooper, Regan, Brauner, De Bastos, & Wilson, 2014).

More recently, *in vivo* studies have shown that rainbow trout may actively modulate RBC pHi to enable higher tissue PO₂ compared to those in mammals (Rummer, McKenzie, Innocenti, Supuran, & Brauner, 2013) and that can be maintained even in the face of exercise or hypoxia (McKenzie et al., 2004). When the RBC β -NHEs are activated by catecholamines, the extrusion of H⁺ exceeds the rate of re-equilibration *via* the Jacobs-Stewart cycle due to the absence of CA activity in teleost plasma. However, when RBCs reach the tissue capillaries the Jacobs-Stewart cycle accelerates in the presence of paCA and the sudden linkage between pHe and pHi effectively “short-circuits” β -NHE activity. The result is a rapid transfer of H⁺ into the RBC that enhances O₂ unloading to the tissue *via* the Bohr effect. When the RBCs leave the capillaries and the site of CA, β -NHE activity recovers pHi and Hb-O₂ affinity during venous transit, securing the renewed oxygenation of Hb at the gills (Harter, May, Federspiel, Supuran, & Brauner, 2018). This mechanism of β -NHE short-circuiting is not tied to morphological structures, such as the *retes* and therefore, it may be generally available to enhance Hb-O₂ unloading to all tissues in teleosts (Randall, Rummer, Wilson, Wang, & Brauner, 2014). In fact, in swimming Atlantic salmon β -NHE short-circuiting allows for a reduction in cardiac output by nearly a third, which may enable the athletic performance of this migratory species (Harter, Zanuzzo, Supuran, Gamperl, & Brauner, 2019). Many other teleost species, besides salmonids, also have RBC β -NHE that may be short-circuited to enhance O₂ unloading (Berenbrink et al., 2005; Harter & Brauner, 2017). Whether other transporters that create H⁺ gradients across the RBC membrane (i.e. NHE, KCC, NKCC) can also be short-circuited in the presence of CA

remains unexplored, and if substantiated may extend the relevance of this mechanism to species that lack β -NHE, such as other fishes, birds and mammals.

Furthermore, many invertebrate species also have Hbs or hemocyanins, some of which display pH-sensitive O_2 binding characteristics that resemble the Bohr effect of vertebrate Hbs (van Holde & Miller, 1995). Invertebrate respiratory pigments that are dissolved in the plasma lack the cellular mechanism that fine-tune gas transport in vertebrates by modulating the RBC microenvironment. However, as shown in cephalopods, the changes in hemolymph pH during circulatory transit may be sufficient to alter the O_2 binding properties of their hemocyanins, and thus, to modulate cardiovascular O_2 transport and facilitate pH homeostasis, much like in vertebrates (Brix, Lykkeboe, & Johansen, 1981). This remarkable example of convergent evolution illustrates the powerful regulatory effects of pH on physiological systems and its ubiquity across animal taxa.

Links between pH and ammonia metabolism

In solution, NH_3 and NH_4^+ follow the pH-dependent equilibrium shown in Figure 5A. Since the pK_a of this reaction is ~ 9.3 , over 95% of ammonia will be present as NH_4^+ at the pH values found in most biological fluids and the external environment (Fig. 5B). Furthermore, NH_3 is a gas and thus crosses cellular membranes much faster than NH_4^+ ions (Boron, 2010). Thus, small but physiologically relevant pH changes result in relatively large changes in the partial pressure of NH_3 (P_{NH_3}), and the resulting difference in partial pressure can drive the diffusion of the gas across a cellular membrane. Additionally, NH_4^+ has nearly identical hydration shell sizes, ionic

conductance, and water mobility rates compared to those of K^+ , which allows NH_4^+ to “hijack” K^+ carrier proteins such as NKA, NKCC, and K^+ channels (Wiener & Verlander, 2017). In combination, these physico-chemical characteristics permit the unregulated entry of ammonia into cells and subcellular compartments, where it can have toxic effects through disruptions in pH, membrane potential, the inner mitochondrial H^+ gradient, cell volume, and the Krebs cycle (see Ip & Chew 2010 for review).

The main source of endogenous ammonia production (ammoniogenesis) in animals is as a by-product of the transdeamination reactions during amino acid catabolism within the mitochondrial matrix. These reactions result in the equimolar production of NH_4^+ and HCO_3^- , and predominantly take place in the kidney in mammals (Weiner & Verlander, 2017) and in the liver in fish (Ip & Chew 2010). Additionally, the intestine of carnivorous fishes can catabolize amino acids and produce significant ammonia load following a meal (Karlsson, Eliason, Mydland, Farrell, & Kiessling, 2006). The deamination of serine by serine-dehydratase is another important ammoniagenic pathway, especially in mollusks. The purine nucleotide cycle is a third ammoniagenic pathway, and is prominent during pHi acidification induced by anaerobic metabolism in both fish white muscle (Mommsen and Hochachka, 1988) and intertidal invertebrates (Campbell, 1991).

In mammals, the most common causes of ammonia build up are due to diseases that alter ammonia metabolism and excretion (Wiener & Verlander, 2017). However, aquatic animals may be exposed to high environmental ammonia levels resulting from organic matter degradation, during hypoxic conditions that impair nitrification, in overcrowded and confined environments, and from agricultural, sewage, and industrial

run-offs (Alabaster & Lloyd, 1980). These conditions can impair the excretion of endogenous ammonia and in extreme cases result in ammonia influx leading to internal ammonia accumulation in internal fluids. In general, the toxicity of a given environmental ammonia concentration increases as environmental pH increases due to the resulting increase in the proportion of ammonia present as NH_3 , which more readily diffuses into the animal (Randall and Tsui, 2002).

In aquatic animals, waste ammonia is typically excreted to the surrounding water across the gills and the skin (Weihrauch & Allen, 2018). The transport of ammonia across cellular membranes is facilitated by ammonium transporters (AMTs) and Rh glycoproteins (Rhs) (Fig. 6). AMTs are broadly present in bacteria, algae and invertebrates (Huang & Peng, 2005) and can electrogenically excrete NH_4^+ into the external medium (Wacker, Garcia-Celma, Lewe, & Andrade, 2014). In the anal papillae of mosquito larvae, AMT1 was found in the basolateral membrane of epithelial cells (Chasiotis et al., 2016). Although AMTs have also been proposed to be present in the apical membrane of gill epithelial cells of marine polychaetes (Thiel et al., 2017), this putative cellular localization has not been confirmed. In addition to AMTs, invertebrates have the Rh isoform Rhp1, which is expressed in the apical membrane of ammonia excreting epithelial cells (Hu et al., 2014, 2017, Thomsen et al., 2016). On the other hand, vertebrates lack AmtS and express several Rh isoforms. The most comprehensive analysis of Rh localization in fish has been performed in pufferfish (*Takifugu rubripes*), which express Rhcg1 and Rhcg2 in the apical membrane of ionocytes and pavement cells, and Rhbg in the basolateral membrane of pavement cells (Nakada, Westhoff, Kato, & Hirose, 2007). Additionally, Rhag is expressed in

500 RBCs, and in some cases is present in the apical and basolateral membranes of fish
501 epithelial cells (reviewed in Wright & Wood, 2009). Although non-tetrapod vertebrates
502 have an Rhp2 gene, its expression has only been shown in sharks (Nakada et al.,
503 2010). Rhp2 mRNA is highly expressed in shark kidney, and the protein is present in
504 the basolateral membrane of renal tubule cells (Nakada et al., 2010). Lower levels of
505 Rhp2 mRNA were also present in shark blood, gill, brain, intestine, liver, rectal gland
506 and stomach (Nawata, Walsh, & Wood, 2015), however, cellular localization in these
507 tissues has not been explored.

508 The substrate specificity of the various Rh channels has large implications for the
509 reciprocal relationship between pH and ammonia transport. When NH_3 is transported
510 into a compartment its protonation to NH_4^+ consumes H^+ and thus has an alkalinizing
511 effect, a response that is stimulated by a greater H^+ availability in compartments that
512 have a lower pH (the opposite is the case for NH_4^+ transport). Unfortunately, substrate
513 specificity studies are not trivial due to the interrelationship between pH_i , pH_e , $\text{NH}_3/\text{NH}_4^+$
514 ratios, and the greater molecular mass and higher pKa of the radiolabeled NH_4^+
515 analogue, ^{14}C -methyl-ammonium, compared to NH_4^+ (~10.6 vs. ~9.3). Heterologous
516 expression in *Xenopus* oocytes suggests that mammalian Rhag and Rhbg can transport
517 both NH_3 and NH_4^+ , and that Rhcg exclusively transports NH_3 (Caner et al., 2015);
518 however, this remains a highly debated subject (Weiner & Verlander, 2017). Knowledge
519 about the substrates transported by the Rhs from aquatic species is even more limited:
520 the substrate for Rhp1 is unknown, Rhp2 seems to preferentially transport NH_3 (Nakada
521 et al., 2010), and detailed substrate specificity studies for other Rhs from aquatic
522 species are lacking. In addition, some Rhs may facilitate CO_2 transport (Musa-Aziz et

al., 2009), and therefore caution should be used when inferring potential Rh functions in aquatic organisms.

Acid-trapping of ammonia

Acidification of a given compartment favors ammonia speciation into NH_4^+ and reduces PNH_3 , thus facilitating NH_3 diffusion into the compartment and trapping it as NH_4^+ . This mechanism is known as “acid-trapping of ammonia”, and is an effective means for excreting ammonia or for accumulating it into subcellular compartments. The acidification can be generated by the hydration of excreted CO_2 at the external surfaces and by H^+ excretion *via* VHA and NHEs, while the diffusion of NH_3 across cellular membranes is facilitated by Rh channels (Figure 6).

Acid-trapping is a generalized strategy to excrete ammonia by freshwater fishes (Ip & Chew, 2010) and invertebrates (Weihrauch & O'Donnell, 2017). In theory, acid-trapping of ammonia is less advantageous for marine animals due to the challenge of secreting sufficient H^+ to acidify highly buffered seawater (Wilkie, 2002). Indeed, the marine polychaetae (*Eurythoe complanata*) directly excretes NH_4^+ across its gills, possibly through AMTs (Thiel et al., 2017). However, many marine invertebrates use acid-trapping to excrete ammonia into a stagnant or enclosed fluid. For example, the gills of cephalopods have intricate infoldings that create a semi-tubular luminal space that limits ventilation volume, permitting apical excretion of H^+ *via* NHE to acidify the seawater and acid trap NH_3 that diffuses through apical Rhp (Hu et al., 2014). Likewise, excretion of CO_2 and H^+ into the palial cavity of bivalves results in a typical pH of ~7.5 (and as low as pH 6.5 during aerial exposure) (Littlewood & Young, 1994). The

ammonia that is trapped in this acidified fluid can be released into the bulk seawater when the animal opens its valves while submerged, or can potentially be volatilized into air while emerged. Similarly, marine animals may utilize acid-trapping into internal fluids such as that in the renal lumen, a mechanism that has been proposed to contribute to the production of ammonia-rich urine in octopuses (Hu et al., 2017).

A variation of acid-trapping of ammonia occurs in the gills of marine carps and is known as vesicular acid-trapping (Figure 6). Here, acidification of intracellular vesicles by VHA promotes NH_3 diffusion and subsequent trapping as NH_4^+ . The vesicles are thereafter trafficked *via* microtubules to the apical membrane, and NH_4^+ is excreted into the environment *via* exocytosis (Weihrauch, Ziegler, Siebers, & Towle, 2002). Vesicular acid-trapping of ammonia has also been proposed in goldfish kidney (Fehsenfeld, Kolosov, Wood, & O'Donnell, 2019).

Since NH_4^+ is lighter than seawater, acid-trapping may also be used to accumulate ammonia in coelomic cavities or in vacuoles within body tissues for the purpose of buoyancy (Voight, Pörtner, & O'Dor, 1995). Indeed, the tissues of some deep-sea cephalopods can reach $[\text{NH}_4^+]$ upwards of 500 mmol L^{-1} (Seibel, Goffredi, Thuesen, Childress, & Robison, 2004), with ~50-60% of their total body mass being comprised of NH_4^+ -rich fluid (Voight et al., 1995). The protein-rich diet of cephalopods enables this buoyancy strategy by providing the necessary source of ammonia thorough amino acid catabolism. These squids maintain their pHe at ~7.2 while the pH of sequestration sites can reach values as low as 5 (Voight et al., 1995). Together, these pH set-points provide the necessary transmembrane pH and $P\text{NH}_3$ gradients to permit acid-trapping of ammonia, first within the hemolymph and thereafter within the

sequestration sites. Similar NH_4^+ sequestration may provide buoyancy in tunicate embryos (Lambert & Lambert, 1978) and pelagic crustaceans (Sanders & Childress, 1988). Although the molecular mechanisms underlying NH_4^+ sequestration have yet to be elucidated, they likely involve Rhp and VHA.

Extreme pH microenvironments in coral cells

Reef-building corals that host photosymbiotic algae experience some of the most extreme acid-base disturbances found among animals. While a carbon concentrating mechanism (CCM) promotes photosynthesis by acidifying the algal microenvironment to pH values as low as 4, skeleton calcification is promoted by creating an alkaline microenvironment where pH values can be greater than 9 (reviewed in Tresguerres et al, 2017). Remarkably, this 100,000-fold difference in $[\text{H}^+]$ exists over just a few hundred microns that separate the cells that host symbiotic algae from those that build the skeleton (Fig. 7A). Since corals lack specialized organs, acid-base homeostasis relies on regulatory mechanisms within each individual cell.

The enzyme ribulose-1,5-bisphosphate carboxylase/oxygenase (rubisco) catalyzes the first major step in photosynthetic CO_2 fixation (Cooper, Filmer, Wishnick, & Lane, 1969). However, rubisco's relatively low affinity for CO_2 compared to contemporary environmental PCO_2 levels and to its significant affinity for O_2 may lead to photorespiration and diminished carbon fixation efficiency (Tamiya & Huzisige, 1948). In response, many phytoplankton species have developed CCMs that elevate PCO_2 at the site of rubisco (Reinfelder, 2011). Likewise, a CCM is essential for sustaining the photosynthetic activity of coral's symbiotic algae (Yellowlees, Rees, & Leggat, 2008).

However, these algae reside inside an intracellular compartment of coral gastrodermal cells called the symbiosome (Fig. 7), which can be modulated by the coral host cell. Recently, a novel host-controlled CCM has been identified whereby VHA that is abundantly expressed in the symbiosome membrane acidifies the lumen down to pH~4 (Barott, Venn, Perez, Tambutté, & Tresguerres, 2015) (Fig. 7B,C). Together with HCO_3^- transport through yet unidentified mechanisms, this VHA-dependent acidification is thought to drive CO_2 flux into the symbiosome lumen and thereby enhance the delivery of CO_2 to the site of fixation. VHA activity in the coral symbiosome membrane has been proposed to additionally slow down symbiotic alga cell division, as well as to drive the transport of phosphates, amino acids, sugars, and of ammonia by acid-trapping (Tang, 2015; Tresguerres et al., 2017, Fig. 7). The presence of an analogous VHA-driven CCM in giant clams that host symbiotic algae in their gut (Armstrong, Roa, Stillman, & Tresguerres, 2018) suggests that this mechanism has evolved convergently in different species.

While coral photosynthesis requires an acidified microenvironment, it alkalinizes the rest of the coral because it consumes CO_2 and H^+ and it generates OH^- (Allemand, Furla, & Bénazet-Tambutté, 1998). At the onset of photosynthesis, the pH_i of the coral host cells immediately increases from ~7.0 to ~7.4 (Barott et al., 2017; Laurent, Tambutte, Tambutte, Allemand, & Venn, 2013). The rate of photosynthesis increases linearly with light irradiance, and so does the initial alkalinization of the host cell cytoplasm. However, pH_i plateaus after ~20 min despite continuous photosynthetic activity, indicating the activation of pH_i regulatory mechanisms (Laurent et al., 2013). At this time, cytosolic H^+ are being replenished at the same rate as they are being

consumed by photosynthesis and a new steady state is reached. The molecular mechanisms involved in this response are unknown. Although certain AEs are a common mechanism used to counteract an intracellular alkalosis (Fig. 2), they extrude HCO_3^- from the cell and this would conflict with the need for dissolved inorganic carbon transport for photosynthesis. Alternatively, transport of HCO_3^- into the symbiosome as proposed in figure 4B would fulfill the need for both pH_i regulation and CCM.

Intracellular buffering is also important to help cope with the immediate alkalinizing effect of algal photosynthesis, and this is reflected in symbiont-containing cells having ~25% higher buffering capacity compared to symbiont-free cells (Laurent et al., 2014). Indeed, their buffering capacity is higher than that of mussel retractor muscle (Zange, Grieshaber, & Jans, 1990) and squid mantle (Pörtner, Boutilier, Tang, & Toews, 1990), which may imply that the magnitude of the alkaline challenge induced by symbiont photosynthesis is greater than the acidic challenge resulting from muscle contraction.

On the other hand, coral calcification takes place in an actively alkalinized environment and represents a source of acidic stress for the rest of the coral. The cells responsible for coral skeleton formation are called calcicoblastic cells, and form an epithelium that is situated directly above the extracellular calcifying fluid (ECF) that separates it from the skeleton. The calcicoblastic cells express an abundance of SLC4 transporters that presumably help deliver HCO_3^- to the ECF (Barott, Perez, Linsmayer, & Tresguerres, 2015; Tresguerres et al., 2017; Zoccola et al., 2015). These cells also express $\text{Na}^+/\text{Ca}^{2+}$ exchangers (Barron et al, 2018) and plasma membrane Ca^{2+} -ATPases that help deliver the required Ca^{2+} (Zoccola et al. 2004; Barott, Perez, et al 2015); the latter might additionally mediate H^+ removal from the ECF. The combined

638 activities of these transporters generate an elevated aragonite saturation state in the
639 ECF that promotes skeleton calcification and counteracts its dissolution. Some of these
640 transporters are likely under regulatory control by sAC, which is expressed in
641 calciblastic cells and mediates the alkalization of the ECF and the growth of skeletal
642 CaCO_3 crystals (Barott et al., 2020). A similar role of sAC in regulating CaCO_3
643 precipitation has been demonstrated or proposed in the intestine of marine teleosts
644 (Tresguerres, Levin, Buck, & Grosell, 2010) and in the teleost inner ear (Kwan, Smith, &
645 Tresguerres, in review). Thus, an evolutionary pattern is emerging whereby sAC-
646 regulated transepithelial acid-base relevant ion-transport generates alkaline conditions
647 that promote calcification in an extracellular space.

648 Other components of the coral calcification mechanism include acidic proteins
649 that promote Ca^{2+} precipitation (Mass et al., 2013), and abundant vesicles in the
650 calciblastic cells that are formed by macropinocytosis from the ECF (Ganot et al., 2020)
651 and potentially also by transcytosis from the oral tissues (Mass et al., 2017).
652 Interestingly, calcifying foraminifera produce their chambered shells by endocytosis of
653 seawater into vesicles, which they subsequently alkalize to a pH >9.0 thus promoting
654 CaCO_3 precipitation (Bentov, Brownlee, & Erez, 2009; de Nooijer, Toyofuku, & Kitazato,
655 2009). Incubation with the VHA inhibitor bafilomycin A1 significantly decreased H^+ efflux
656 from the newly forming chambers and resulted in weakly calcified chamber walls,
657 indicating the involvement of VHA in calcification (Toyofuku et al., 2017). A similar role
658 for VHA in calcification was proposed in the calcifying vesicle of coccolithophorids, which
659 are another eukaryotic phytoplankton with an external CaCO_3 shell (Corstjens, Araki, &
660 González, 2001; Mackinder et al., 2011). It is worth investigating whether VHA is also

present in the vesicles within coral calicoblastic cells and whether it plays a role in coral skeleton formation.

The H^+ that are removed from the ECF during calcification eventually reach the coral gut cavity, where they combine with HCO_3^- to form CO_2 that may be used by photosynthesis within the symbiotic algae (Fig. 7B). Thus, coral calcification and photosynthesis are linked to each other through the complementary and synergistic production and consumption of CO_2 , H^+ and HCO_3^- . This is one of the mechanisms by which the photosynthetic activity of the symbiotic algae stimulates coral skeletal growth, a phenomenon known as “light enhanced calcification” (Kawaguti & Sakumoto, 1948).

Summary

As substrates and products of many biochemical reactions CO_2 , H^+ , HCO_3^- molecules are intrinsically linked to aerobic and anaerobic metabolism, O_2 transport, ammonia homeostasis, metabolic communication between symbiotic partners, and calcification. Future comparative studies at all levels of organization will undoubtedly continue to reveal novel aspects about the evolutionary links between intra- and extracellular acid-base regulation and their effects on multiple aspects of organismal physiology. From a practical perspective, understanding the effects of metabolic and environmental acid-base disturbances on homeostatic processes may help predict the resilience and vulnerability of species to environmental disturbances, both natural and anthropogenic in origin, as well as to artificial environments such as those experienced in intensive aquaculture.

683

684 **Acknowledgements**

685 Supported by National Science Foundation (NSF) IOS #1754994 to M.T., SIO

686 Postdoctoral Scholar fellowship to A.M.C., NSF Graduate Research Fellowship to

687 A.B.T., NSF Postdoctoral Fellowship to J.N.R., and Natural Sciences and Engineering

688 Research Council (NSERC) Discovery grant (2018-04172) to C.J.B.

689

References

- Alabaster, J.S., & Lloyd, R. (1980). Ammonia. Water Quality criteria for fresh water fish, 2nd edition. London: Butterworth Scientific.
- Allemand, D., Furla, P., & Bénazet-Tambutté, S. (1998). Mechanisms of carbon acquisition for endosymbiont photosynthesis in Anthozoa. *Canadian Journal of Botany*, 76, 925–941.
- Armbruster, J. W. (1998). Modifications of the digestive tract for holding air in Loricariid and Scoloplacid catfishes. *Copeia*, 1998(3), 663-675.
- Armstrong, E. J., Roa, J. N., Stillman, J. H., & Tresguerres, M. (2018). Symbiont photosynthesis in giant clams is promoted by V-type H⁺-ATPase from host cells. *The Journal of Experimental Biology*, 221, jeb177220. <https://doi.org/10.1242/jeb.177220>
- Barnhart, M. C., & McMahon, B. R. (1988). Depression of aerobic metabolism and intracellular pH by hypercapnia in land snails, *Otala Lactea*. *The Journal of Experimental Biology*, 138, 289–299.
- Baker, D. W. (2010). Physiological responses associated with aquatic hypercarbia in the CO₂-tolerant white sturgeon *Acipenser transmontanus*. University of British Columbia, Vancouver, Canada.
- Baker, D. W., Matey, V., Huynh, K. T., Wilson, J. M., Morgan, J. D., & Brauner, C. J. (2009). Complete intracellular pH protection during extracellular pH depression is associated with hypercarbia tolerance in white sturgeon, *Acipenser transmontanus*. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 296, R1868-1880. <https://doi.org/10.1152/ajpregu.90767.2008>

712 Barott, K. L., Barron, M. E., & Tresguerres, M. (2017). Identification of a molecular pH
 713 sensor in coral. *Proceedings of the Royal Society B*, 284, 20171769.
 714 <https://doi.org/10.1098/rspb.2017.1769>

715 Barott, K.L., Perez, S.O., Linsmayer, L.B., & Tresguerres, M. (2015). Differential
 716 localization of ion transporters suggests distinct cellular mechanisms for calcification
 717 and photosynthesis between two coral species. *American Journal of Physiology.
 718 Regulatory, Integrative and Comparative Physiology*, 309:R235-246. doi:
 719 10.1152/ajpregu.00052.2015.

720 Barott, K. L., Venn, A. A., Thies, A. B., Tambutté, S., & Tresguerres, M. (2020).
 721 Regulation of coral calcification by the acid-base sensing enzyme soluble adenylyl
 722 cyclase. *Biochemical & Biophysical Research Communications*, in press
 723 <https://doi.org/10.1016/j.bbrc.2020.02.115>

724 Barott, K. L., Venn, A. A., Perez, S. O., Tambutté, S., & Tresguerres, M. (2015). Coral
 725 host cells acidify symbiotic algal microenvironment to promote photosynthesis.
 726 *Proceedings of the National Academy of Sciences*, 112, 607–612.
 727 <https://doi.org/10.1073/pnas.1413483112>

728 Barron, M.E., Thies, A.B., Espinoza, J.A., Barott, K.L., Hamdoun, A., & Tresguerres, M.
 729 (2018). A vesicular Na⁺/Ca²⁺ exchanger in coral calcifying cells. *PLoS One*, 13,
 730 e0205367. doi: 10.1371/journal.pone.0205367.

731 Bayne, B.L., Bayne, C.J., Carefoot, T.C., & Thompson, R.J. (1976). The physiological
 732 ecology of *Mytilus californianus* Conrad : 2. Adaptations to low oxygen tension and air
 733 exposure. *Oecologia*, 22, 229-250. doi: 10.1007/BF00344794.

734 Bentov, S., Brownlee, C., & Erez, J. (2009). The role of seawater endocytosis in the
 735 biomineralization process in calcareous foraminifera. *Proceedings of the National*
 736 *Academy of Sciences*, 106, 21500–21504.

737 Berenbrink, M., Koldkjaer, P., Kepp, O., & Cossins, A. R. (2005). Evolution of oxygen
 738 secretion in fishes and the emergence of a complex physiological system. *Science*, 307,
 739 1752–1757. <https://doi.org/10.1126/science.1107793>

740 Berenbrink, M., Volkel, S., Koldkjaer, P., Heisler, N., & Nikinmaa, M. (2006). Two
 741 different oxygen sensors regulate oxygen-sensitive K⁺ transport in crucian carp red
 742 blood cells. *The Journal of Physiology*, 575, 37–48.

743 Bohr, C., Hasselbalch, K., & Krogh, A. (1904). About a new biological relation of high
 744 importance that the blood carbonic acid tension exercises on its oxygen binding.
 745 *Skandinavisches Archiv Fuer Physiologie*, 16, 402–412.

746 Boron, W.F. (2010). Sharpey-Schafer lecture: gas channels. *Experimental Physiology*,
 747 95, 1107–1130. doi: 10.1113/expphysiol.2010.055244.

748 Boutilier, R. G., Heming, T. A., & Iwama, G. K. (1984). Appendix: Physicochemical
 749 Parameters for use in Fish Respiratory Physiology. In W. S. H. Randall & D. J (Eds.),
 750 *Gills Anatomy, Gas Transfer, and Acid-Base Regulation* (pp. 403–430).

751 Brix, O., Lykkeboe, G., & Johansen, K. (1981). The significance of the linkage between
 752 the Bohr and Haldane effects in cephalopod bloods. *Respiration Physiology*, 44, 177–
 753 186. [https://doi.org/10.1016/0034-5687\(81\)90036-0](https://doi.org/10.1016/0034-5687(81)90036-0)

754 Caldeira, K., & Wickett, M. (2003). Anthropogenic carbon and ocean pH. *Nature* 425,
 755 365. <https://doi.org/10.1038/425365a>

756 Cameron, J. N., & Kormanik, G. A. (1982). Intracellular and extracellular acid-base
 757 status as a function of temperature in the fresh-water channel catfish, *Ictalurus*
 758 *punctatus*. *The Journal of Experimental Biology*, 99, 127–142.

759 Campbell, J. W. (1991). Excretory nitrogen metabolism. In C. L. Prosser (Ed.),
 760 *Comparative Animal Physiology, Environmental and Metabolic Animal Physiology* (4th
 761 ed., pp. 277–324). Wiley.

762 Caner, T., Abdulnour-Nakhoul, S., Brown, K., Islam, M. T., Hamm, L. L., & Nakhoul, N.
 763 L. (2015). Mechanisms of ammonia and ammonium transport by rhesus-associated
 764 glycoproteins. *American Journal of Physiology-Cell Physiology*, 309, C747–58.
 765 <https://doi.org/10.1152/ajpcell.00085.2015>

766 Casey, J. R., Grinstein, S., & Orlowski, J. (2010). Sensors and regulators of intracellular
 767 pH. *Nature Reviews. Molecular Cell Biology*, 11, 50–61.
 768 <https://doi.org/10.1038/nrm2820>

769 Cech, J.J., & Crocker, C.E. (2002), Physiology of sturgeon: effects of hypoxia and
 770 hypercapnia. *Journal of Applied Ichthyology* 18, 320-324.

771 Chang, J.-C., & Oude Elferink, R. P. (2014). Role of the bicarbonate-responsive soluble
 772 adenylyl cyclase in pH sensing and metabolic regulation. *Frontiers in Physiology*, 5.
 773 <https://doi.org/10.3389/fphys.2014.00042>

774 Chasiotis, H., Ionescu, A., Misyura, L., Bui, P., Fazio, K., Wang, J., Patrick, M.,
 775 Weihrauch, D., & Donini, A. (2016). An animal homolog of plant Mep/Amt transporters
 776 promotes ammonia excretion by the anal papillae of the disease vector mosquito *Aedes*
 777 *aegypti*. *The Journal of Experimental Biology*, 219, 1346–1355.
 778 <https://doi.org/10.1242/jeb.134494>

779 Chen, Y., Cann, M. J., Litvin, T. N., Iourgenko, V., Sinclair, M. L., Levin, L. R., & Buck, J.
780 (2000). Soluble Adenylyl Cyclase as an Evolutionarily Conserved Bicarbonate Sensor.
781 *Science*, 289, 625–628.

782 Cooper, T. G., Filmer, D., Wishnick, M., & Lane, M. D. (1969). The active species of
783 “CO₂” utilized by ribulose diphosphate carboxylase. *Journal of Biological Chemistry*,
784 244, 1081–1083.

785 Cooper, C. A., Regan, M. D., Brauner, C. J., De Bastos, E. S. R., & Wilson, R. W.
786 (2014). Osmoregulatory bicarbonate secretion exploits H⁺-sensitive haemoglobins to
787 autoregulate intestinal O₂ delivery in euryhaline teleosts. *Journal of Comparative*
788 *Physiology B*, 184, 865–876. <https://doi.org/10.1007/s00360-014-0844-x>

789 Corstjens, P. L., Araki, Y., & González, E. L. (2001). A coccolithophorid calcifying
790 vesicle with a vacuolar-type ATPase proton pump: Cloning and immunolocalization of
791 the V₀ subunit c. *Journal of Phycology*, 37, 71–78.

792 Cossins, A. R., & Gibson, J. S. (1997). Volume-sensitive transport systems and volume
793 homeostasis in vertebrate red blood cells. *The Journal of Experimental Biology*, 200,
794 343–352.

795 Dejours, P. (1994). Environmental Factors as Determinants in Bimodal Breathing: An
796 Introductory Overview. *Integrative and Comparative Biology*, 34, 178–183.
797 <https://doi.org/10.1093/icb/34.2.178>

798 de Nooijer, L. J., Toyofuku, T., & Kitazato, H. (2009). Foraminifera promote calcification
799 by elevating their intracellular pH. *Proceedings of the National Academy of Sciences*,
800 106(36), 15374–15378.

801 Dolan, E., Saunders, B., Harris, R.C., Bicudo, J.E.P.W., Bishop, D.J., Sale, C., Gualano,
802 B. (2019). Comparative physiology investigations support a role for histidine-containing
803 dipeptides in intracellular acid-base regulation of skeletal muscle. *Comparative*
804 *Biochemistry and Physiology A*, 234, 77-86. doi: 10.1016/j.cbpa.2019.04.017.

805 Duarte, R. M., Ferreira, M. S., Wood, C. M., & Val, A. L. (2013). Effect of low pH
806 exposure on Na⁺ regulation in two cichlid fish species of the Amazon. *Comparative*
807 *Biochemistry and Physiology A*, 166, 441–448.
808 <https://doi.org/10.1016/j.cbpa.2013.07.022>

809 Ellis, R. P., Urbina, M. A., & Wilson, R. W. (2017). Lessons from two high CO₂ worlds—
810 Future oceans and intensive aquaculture. *Global Change Biology*, 23, 2141–2148.
811 <https://doi.org/10.1111/gcb.13515>

812 Ellington, W. R. (1983). The recovery from anaerobic metabolism in invertebrates. *The*
813 *Journal of Experimental Zoology*, 228, 431-444.

814 Ellory, J. C., & Wolowyk, M. W. (1991). Evidence for bumetanide-sensitive, Na⁺-
815 dependent, partial Na–K–Cl co-transport in red blood cells of a primitive fish. *Canadian*
816 *Journal of Physiology and Pharmacology*, 69, 588–591.

817 Esbaugh, A. J., Gilmour, K. M., & Perry, S. F. (2009). Membrane-associated carbonic
818 anhydrase in the respiratory system of the Pacific hagfish (*Eptatretus stouti*).
819 *Respiratory Physiology & Neurobiology*, 166, 107–116.

820 Fehsenfeld, S., Kolosov, D., Wood, C. M., & O'Donnell, M. J. (2019). Section-specific H⁺
821 flux in renal tubules of fasted and fed goldfish. *The Journal of Experimental Biology*,
822 222, jeb200964. <https://doi.org/10.1242/jeb.200964>

823 Feller, G., Poncin, A., Aittaleb, M., Schyns, R., & Gerday, C. (1994). The blood proteins
824 of the Antarctic icefish *Channichthys rhinoceratus*: Biological significance and
825 purification of the two main components. *Comparative Biochemistry and Physiology, B*,
826 109, 89–97.

827 Furch, J., & Junk, W. J. (1997). Physicochemical conditions in the floodplains. In *The*
828 *Central Amazon Floodplain*, pp. 69-108. Berlin, Heidelberg: Springer, Berlin,
829 Heidelberg.

830 Ganot, P., Tambutté, E., Caminiti-Segonds, N., Toullec, G., Allemand, D., Tambutté, S.
831 (2020). Ubiquitous macropinocytosis in anthozoans. *eLife*, 9, e50022 doi:
832 10.7554/eLife.50022

833 Gervais, M. R., & Tufts, B. L. (1998). Evidence for membrane-bound carbonic
834 anhydrase in the air bladder of bowfin (*Amia calva*), a primitive air-breathing fish. *The*
835 *Journal of Experimental Biology*, 201, 2205–2212.

836 Gilmour, K. M. (2001). The CO₂/pH ventilatory drive in fish. *Comparative Biochemistry*
837 *and Physiology A*, 130, 219-40.

838 Hall-Spencer, J.M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner,
839 S.M. *et al.* (2008). Volcanic carbon dioxide vents show ecosystem effects of ocean
840 acidification. *Nature*, 454, 96–99.

841 Hammer, K. M., Kristiansen, E., & Zachariassen, K. E. (2011). Physiological effects of
842 hypercapnia in the deep-sea bivalve *Acesta excavata* (Fabricius, 1779) (Bivalvia;
843 Limidae). *Marine Environmental Research*, 72, 135–142.
844 <https://doi.org/10.1016/j.marenvres.2011.07.002>

845 Harter, T. S., & Brauner, C. J. (2017). The O₂ and CO₂ transport system in teleosts and
846 the specialized mechanisms that enhance Hb-O₂ unloading to tissues. In A. K. Gamperl,
847 T. E. Gillis, A. P. Farrell, & C. J. Brauner (Eds.), *Fish Physiology: Vol. 36B: The*
848 *Cardiovascular System: Morphology, Control and Function* (pp. 1–107). New York:
849 Academic Press.

850 Harter, T. S., May, A. G., Federspiel, W. J., Supuran, C. T., & Brauner, C. J. (2018). The
851 time-course of red blood cell intracellular pH recovery following short-circuiting in
852 relation to venous transit times in rainbow trout, *Oncorhynchus mykiss*. *American*
853 *Journal of Physiology: Regulatory, Integrative and Comparative Physiology*, 315, R397–
854 R407.

855 Harter, T. S., Shartau, R. B., Baker, D. W., Jackson, D. C., Val, A. L., & Brauner, C. J.
856 (2014). Preferential intracellular pH regulation represents a general pattern of pH
857 homeostasis during acid–base disturbances in the armoured catfish, *Pterygoplichthys*
858 *pardalis*. *Journal of Comparative Physiology B*, 184, 709–718.
859 <https://doi.org/10.1007/s00360-014-0838-8>

860 Harter, T. S., Zanuzzo, F. S., Supuran, C. T., Gamperl, A. K., & Brauner, C. J. (2019).
861 Functional support for a novel mechanism that enhances tissue oxygen extraction in a
862 teleost fish. *Proceedings of the Royal Society B*, 286, 20190339.
863 <https://doi.org/doi:10.1098/rspb.2019.0339>

864 Heisler, N., Forcht, G., Ultsch, G. R., & Anderson, J. F. (1982). Acid-base regulation in
865 response to environmental hypercapnia in two aquatic salamanders, *Siren lacertina* and
866 *Amphiuma means*. *Respiration Physiology*, 49, 141–158. [https://doi.org/10.1016/0034-](https://doi.org/10.1016/0034-5687(82)90070-6)
867 [5687\(82\)90070-6](https://doi.org/10.1016/0034-5687(82)90070-6)

868 Heming, T. A., Vanoye, C. G., Stabenau, E. K., Roush, E. D., Fierke, C. A., & Bidani, A.
 869 (1993). Inhibitor sensitivity of pulmonary vascular carbonic-anhydrase. *Journal of*
 870 *Applied Physiology*, 75, 1642–1649.

871 Henry, R. P., & Swenson, E. R. (2000). The distribution and physiological significance of
 872 carbonic anhydrase in vertebrate gas exchange organs. *Respiration Physiology*, 121,
 873 1–12. [https://doi.org/10.1016/S0034-5687\(00\)00110-9](https://doi.org/10.1016/S0034-5687(00)00110-9)

874 Henry, R. P., Gilmour, K. M., Wood, C. M., & Perry, S. F. (1997). Extracellular carbonic
 875 anhydrase activity and carbonic anhydrase inhibitors in the circulatory system of fish.
 876 *Physiological Zoology*, 70, 650–659

877 Hochachka, P. W., & Mommsen, T. P. (1983). Protons and anaerobiosis. *Science*, 219,
 878 1391–1397. <https://doi.org/10.1126/science.6298937>

879 Hofmann, G. E., Smith, J. E., Johnson, K. S., Send, U., Levin, L. A., Micheli, F., Paytan,
 880 A., Price, N. N., Peterson, B., Takeshita, Y., Matson, P. G., Derse Crook, E., Kroeker, K.
 881 J., Gambi, M. C., Rivest, E. B., Frieder, C. A., Yu, P. C., & Martz, T. R. (2011). High-
 882 frequency dynamics of ocean pH: A multi-ecosystem comparison. *PloS One*, 6, e28983.
 883 <https://doi.org/10.1371/journal.pone.0028983>

884 Hu, M. Y., Sung, P.-H., Guh, Y.-J., Lee, J.-R., Hwang, P.-P., Weihrauch, D., & Tseng,
 885 Y.-C. (2017). Perfused gills reveal fundamental principles of pH regulation and ammonia
 886 homeostasis in the cephalopod *Octopus vulgaris*. *Frontiers in Physiology*, 8, 162.
 887 <https://doi.org/10.3389/fphys.2017.00162>

888 Hu, M. Y., Guh, Y.-J., Stumpp, M., Lee, J.-R., Chen, R.-D., Sung, P.-H., Chen, Y.-C.,
 889 Hwang, P.-P., & Tseng, Y.-C. (2014). Branchial NH₄⁺-dependent acid–base transport
 890 mechanisms and energy metabolism of squid (*Sepioteuthis lessoniana*) affected by

891 seawater acidification. *Frontiers in Zoology*, 11, 55. <https://doi.org/10.1186/s12983-014->
892 0055-z

893 Huang, C.-H., & Peng, J. (2005). Evolutionary conservation and diversification of Rh
894 family genes and proteins. *Proceedings of the National Academy of Sciences*, 102,
895 15512–15517. <https://doi.org/10.1073/pnas.0507886102>

896 Ip, Y.K., & Chew, S.F. (2010). Ammonia production, excretion, toxicity, and defense in
897 fish: a review. *Frontiers in Physiology* 1, 134. doi: 10.3389/fphys.2010.00134.

898 Itada, N., & Forster, R. E. (1977). Carbonic anhydrase activity in intact red blood cells
899 measured with ¹⁸O exchange. *Journal of Biological Chemistry*, 252, 3881–3890.

900 Jacobs, M. H., & Stewart, D. R. (1942). The role of carbonic anhydrase in certain ionic
901 exchanges involving the erythrocyte. *The Journal of General Physiology*, 25, 539–552.

902 Jarvis, K., Woodward, M., Debold, E. P., & Walcott, S. (2018). Acidosis affects muscle
903 contraction by slowing the rates myosin attaches to and detaches from actin. *Journal of*
904 *Muscle Research and Cell Motility*, 39, 135–147. <https://doi.org/10.1007/s10974-018->
905 9499-7

906 Jensen, F. B. (2004). Red blood cell pH, the Bohr effect, and other oxygenation-linked
907 phenomena in blood O₂ and CO₂ transport. *Acta Physiologica Scandinavica*, 182, 215–
908 227.

909 Karlsson, A., Eliason, E.J., Mydland, L.T., Farrell, A.P., & Kiessling, A. (2006).
910 Postprandial changes in plasma free amino acid levels obtained simultaneously from
911 the hepatic portal vein and the dorsal aorta in rainbow trout (*Oncorhynchus mykiss*).
912 *The Journal of Experimental Biology*, 209, 4885-4894.

913 Kawaguti, S., & Sakumoto, D. (1948). The effect of light on the calcium deposition of
 914 corals. *Bulletin of the Oceanographical Institute of Taiwan*, 4, 65–70.

915 Kieffer, J.D. (2000). Limits to exhaustive exercise in fish. *Comparative Biochemistry and*
 916 *Physiology A*, 126, 161–79.

917 Kline, D. I., Teneva, L., Schneider, K., Miard, T., Chai, A., Marker, M., ... Hoegh-
 918 Guldborg, O. (2012). A short-term in situ CO₂ enrichment experiment on Heron Island
 919 (GBR). *Scientific Reports*, 2, 413. <https://doi.org/10.1038/srep00413>

920 Koch, A. L., & Silver, S. (2005). The first cell. *Advances in Microbial Physiology*, 50,
 921 227–259. [https://doi.org/10.1016/S0065-2911\(05\)50006-7](https://doi.org/10.1016/S0065-2911(05)50006-7)

922 Lambert, C. C., & Lambert, G. (1978). Tunicate eggs utilize ammonium ions for flotation.
 923 *Science*, 200, 64–65. <https://doi.org/10.1126/science.200.4337.64>

924 Larsen, E. H., Deaton, L. E., Onken, H., O'Donnell, M., Grosell, M., Dantzler, W. H., &
 925 Weihrauch, D. (2014). Osmoregulation and excretion. *Comprehensive Physiology*, 4,
 926 405–573. <https://doi.org/10.1002/cphy.c130004>

927 Laurent, J., Tambutte, S., Tambutte, E., Allemand, D., & Venn, A. (2013). The influence
 928 of photosynthesis on host intracellular pH in scleractinian corals. *The Journal of*
 929 *Experimental Biology*, 216, 1398–1404. <https://doi.org/10.1242/jeb.082081>

930 Laurent, J., Venn, A., Tambutté, É., Ganot, P., Allemand, D., & Tambutté, S. (2014).
 931 Regulation of intracellular pH in cnidarians: Response to acidosis in *Anemonia viridis*.
 932 *FEBS Journal*, 281, 683–695. <https://doi.org/10.1111/febs.12614>

933 Lawrence, M. J., Wright, P. A., & Wood, C. M. (2015). Physiological and molecular
 934 responses of the goldfish (*Carassius auratus*) kidney to metabolic acidosis, and

935 potential mechanisms of renal ammonia transport. *Journal of Experimental Biology*,
936 218, 2124–2135. <https://doi.org/10.1242/jeb.117689>

937 Mackinder, L., Wheeler, G., Schroeder, D., von Dassow, P., Riebesell, U., & Brownlee,
938 C. (2011). Expression of biomineralization-related ion transport genes in *Emiliana*
939 *huxleyi*. *Environmental Microbiology*, 13, 3250–3265.

940 Mass, T., Drake, J. L., Haramaty, L., Kim, J.D., Zelzion, E., Bhattacharya, D., &
941 Falkowski, P. G. (2013). Cloning and characterization of four novel coral acid-rich
942 proteins that precipitate carbonates *in vitro*. *Current Biology*, 23, 1126-1131.
943 <http://dx.doi.org/10.1016/j.cub.2013.05.007>

944 Mass, T., Giuffre, A. J., Sun, C. Y., Stiffler, C. A., Frazier, M.J., Neder, M., Tamura, N.,
945 Stan, C. V., Marcus, M. A., Gilbert, P. (2017). Amorphous calcium carbonate particles
946 form coral skeletons. *Proceedings of the National Academy of Sciences*, 114:E7670-
947 E7678. doi: 10.1073/pnas.1707890114.

948 McKenzie, D. J., Wong, S., Randall, D. J., Egginton, S., Taylor, E. W., & Farrell, A. P.
949 (2004). The effects of sustained exercise and hypoxia upon oxygen tensions in the red
950 muscle of rainbow trout. *Journal of Experimental Biology*, 207(21), 3629–3637.
951 <https://doi.org/10.1242/jeb.01199>

952 Melzner, F., Gutowska, M. A., Langenbuch, M., Dupont, S., Lucassen, M., Thorndyke,
953 M. C., Bleich, M., & Pörtner, H.-O. (2009). Physiological basis for high CO₂ tolerance in
954 marine ectothermic animals: Pre-adaptation through lifestyle and ontogeny?
955 *Biogeosciences*, 6, 2313–2331. <https://doi.org/10.5194/bg-6-2313-2009>

956 Mommsen, T. P., & Hochachka, P. W. (1988). The purine nucleotide cycle as two
 957 temporally separated metabolic units: A study on trout muscle. *Metabolism*, 37, 552–
 958 556. [https://doi.org/10.1016/0026-0495\(88\)90170-9](https://doi.org/10.1016/0026-0495(88)90170-9)
 959 Morrison, P. R., Gilmour, K. M., & Brauner, C. J. (2015). 3—Oxygen and Carbon
 960 Dioxide Transport in Elasmobranchs. In R. E. Shadwick, A. P. Farrell, & C. J. Brauner
 961 (Eds.), *Fish Physiology* (pp. 127–219). [https://doi.org/10.1016/B978-0-12-801286-](https://doi.org/10.1016/B978-0-12-801286-4.00003-4)
 962 4.00003-4
 963 Motais, R., Fievet, B., Garcia-Romeu, F., & Thomas, S. (1989). Na⁺-H⁺ exchange and
 964 pH regulation in red blood cells: Role of uncatalyzed H₂CO₃⁻ dehydration. *American*
 965 *Journal of Physiology*, 256, C728–C735.
 966 Musa-Aziz, R., Chen, L. M., Pelletier, M. F., Boron, W. F. (2009) Relative CO₂/NH₃
 967 selectivities of AQP1, AQP4, AQP5, AmtB, and RhAG. *Proceedings of the National*
 968 *Academy of Sciences*, 106, 5406-5411. doi: 10.1073/pnas.0813231106.
 969 Nakada, T., Westhoff, C. M., Kato, A., & Hirose, S. (2007). Ammonia secretion from fish
 970 gill depends on a set of Rh glycoproteins. *The FASEB Journal*, 21, 1067–1074.
 971 <https://doi.org/10.1096/fj.06-6834com>
 972 Nakada, T., Westhoff, C. M., Yamaguchi, Y., Hyodo, S., Li, X., Muro, T., Kato, A.,
 973 Nakamura, N., & Hirose, S. (2010). Rhesus glycoprotein p2 (Rhp2) is a novel member
 974 of the Rh family of ammonia transporters highly expressed in shark kidney. *Journal of*
 975 *Biological Chemistry*, 285, 2653–2664. <https://doi.org/10.1074/jbc.M109.052068>
 976 Nawata, C.M., Walsh, P.J., & Wood, C.M. (2015). Physiological and molecular
 977 responses of the spiny dogfish shark (*Squalus acanthias*) to high environmental

978 ammonia: scavenging for nitrogen. *The Journal of Experimental Biology*, 15, 238-48.
 979 doi: 10.1242/jeb.114967.

980 Nielsen, J. G., & Munk, O. (1964). A hadal fish (*Bassogigas profundissimus*) with a
 981 functional swimbladder. *Nature*, 204, 594–595.

982 Nikinmaa, M. (2003). Gas transport. In I. Bernhardt & J. C. Ellory (Eds.), *Red cell*
 983 *membrane transport in health and disease* (pp. 489–509). Berlin: Springer.

984 Pelster, B. (1997). Buoyancy at depth. In *Fish Physiology*, vol. 16: Deep-sea fishes
 985 (eds. D. Randall and A. Farrell), pp. 195-238. New York: Academic Press.

986 Perutz, M. F., Fermi, G., Luisi, B., Shaanan, B., & Liddington, R. C. (1987).
 987 Stereochemistry of cooperative mechanisms in hemoglobin. *Accounts of Chemical*
 988 *Research*, 20, 309–321.

989 Pörtner, H. O., Boutilier, R. G., Tang, Y., & Toews, D. P. (1990). Determination of
 990 intracellular pH and PCO₂ after metabolic inhibition by fluoride and nitrilotriacetic acid.
 991 *Respiration Physiology*, 81, 255–273.

992 Pörtner, H. O., Reipschläger, A., & Heisler, N. (1998). Acid-base regulation, metabolism
 993 and energetics in *Sipunculus nudus* as a function of ambient carbon dioxide level. *The*
 994 *Journal of Experimental Biology*, 201, 43–55.

995 Randall, D. J., Rummer, J. L., Wilson, J. M., Wang, S., & Brauner, C. J. (2014). A
 996 unique mode of tissue oxygenation and the adaptive radiation of teleost fishes. *The*
 997 *Journal of Experimental Biology*, 217, 1205–1214.

998 Randall, D. J., & Tsui, T. K. N. (2002). Ammonia toxicity in fish. *Marine Pollution*
 999 *Bulletin*, 45, 17–23. [https://doi.org/10.1016/S0025-326X\(02\)00227-8](https://doi.org/10.1016/S0025-326X(02)00227-8)

1000 Reinfelder, J. R. (2011). Carbon concentrating mechanisms in eukaryotic marine
 1001 phytoplankton. *Annual Review of Marine Science*, 3, 291–315.
 1002 <https://doi.org/10.1146/annurev-marine-120709-142720>
 1003 Romano, L., & Passow, H. (1984). Characterization of anion transport system in trout
 1004 red blood cell. *American Journal of Physiology*, 246, C330-338.
 1005 Rummer, J. L., McKenzie, D. J., Innocenti, A., Supuran, C. T., & Brauner, C. J. (2013).
 1006 Root effect hemoglobin may have evolved to enhance general tissue oxygen delivery.
 1007 *Science*, 340, 1327–1329. <https://doi.org/10.1126/science.1233692>
 1008 Roa, J. N., & Tresguerres, M. (2017). Bicarbonate-sensing soluble adenylyl cyclase is
 1009 present in the cell cytoplasm and nucleus of multiple shark tissues. *Physiological*
 1010 *Reports*, 5(2). <https://doi.org/10.14814/phy2.13090>
 1011 Sanders, N. K., & Childress, J. J. (1988). Ion replacement as a buoyancy mechanism in
 1012 a pelagic deep-sea crustacean. *The Journal of Experimental Biology*, 138, 333–343.
 1013 Seibel, B. A., Goffredi, S. K., Thuesen, E. V., Childress, J. J., & Robison, B. H. (2004).
 1014 Ammonium content and buoyancy in midwater cephalopods. *Journal of Experimental*
 1015 *Marine Biology and Ecology*, 313, 375–387. <https://doi.org/10.1016/j.jembe.2004.08.015>
 1016 Shartau, R. B., Baker, D. W., & Brauner, C. J. (2017). White sturgeon (*Acipenser*
 1017 *transmontanus*) acid-base regulation differs in response to different types of acidoses.
 1018 *Journal of Comparative Physiology B*, 187, 985–994. [https://doi.org/10.1007/s00360-](https://doi.org/10.1007/s00360-017-1065-x)
 1019 [017-1065-x](https://doi.org/10.1007/s00360-017-1065-x)
 1020 Shartau, R.B., Baker, D.W., Harter, T.S., Aboagye, D.L., Allen, P.J., Val, A.L., Crossley
 1021 II, D.A., Kohl, Z.F., Hedrick, M.S., Damsgaard, C., & Brauner, C.J. (2020). Preferential

1022 intracellular pH regulation is a common trait fish exposed to high environmental CO₂.
 1023 *The Journal of Experimental Biology* 2020, jeb.208868 doi: 10.1242/jeb.208868

1024 Shartau, R. B., Baker, D. W., Crossley, D. A., & Brauner, C. J. (2016). Preferential
 1025 intracellular pH regulation: Hypotheses and perspectives. *The Journal of Experimental*
 1026 *Biology*, 219, 2235–2244. <https://doi.org/10.1242/jeb.126631>

1027 Shartau, R. B., Crossley, D. A., Kohl, Z. F., & Brauner, C. J. (2016). Embryonic common
 1028 snapping turtles (*Chelydra serpentina*) preferentially regulate intracellular tissue pH
 1029 during acid-base challenges. *The Journal of Experimental Biology*, 219, 1994–2002.
 1030 <https://doi.org/10.1242/jeb.136119>

1031 Shartau, R. B., Crossley, D. A., Kohl, Z. F., Elsey, R. M., & Brauner, C. J. (2018).
 1032 American alligator (*Alligator mississippiensis*) embryos tightly regulate intracellular pH
 1033 during a severe acidosis. *Canadian Journal of Zoology*, 96, 723–727.
 1034 <https://doi.org/10.1139/cjz-2017-0249>

1035 Sørensen, S. P. L. (1909). Über die Messung und die Bedeutung der
 1036 Wasserstoffionenkonzentration bei enzymatischen Prozessen. *Biochemische Zeitschrift*,
 1037 21, 131–304.

1038 Szebedinszky, C., & Gilmour, K. (2002). The buffering power of plasma in brown
 1039 bullhead (*Ameiurus nebulosus*). *Comparative Biochemistry and Physiology B*, 131,
 1040 171–183.

1041 Tamiya, H., & Huzisige, H. (1948). Effect of oxygen on the dark reaction of
 1042 photosynthesis. *Acta Phytochimica*, 15, 83–104.

1043 Tang, B. L. (2015). Thoughts on a very acidic symbiosome. *Frontiers in Microbiology*, 6,
 1044 816. <https://doi.org/10.3389/fmicb.2015.00816>

1045 Thiel, D., Hugenschütt, M., Meyer, H., Paululat, A., Quijada-Rodriguez, A. R., Purschke,
 1046 G., & Weihrauch, D. (2017). Ammonia excretion in the marine polychaete *Eurythoe*
 1047 *complanata* (Annelida). *The Journal of Experimental Biology*, 220, 425–436.
 1048 <https://doi.org/10.1242/jeb.145615>

1049 Thomas, S., & Egée, S. (1998). Fish red blood cells: Characteristics and physiological
 1050 role of the membrane ion transporters. *Comparative Biochemistry and Physiology A*,
 1051 119, 79–86.

1052 Thomas, S., & Hughes, G. M. (1982). A study of the effects of hypoxia on acid-base
 1053 status of rainbow trout blood using an extracorporeal blood circulation. *Respiration*
 1054 *Physiology*, 49, 371–382. [https://doi.org/10.1016/0034-5687\(82\)90123-2](https://doi.org/10.1016/0034-5687(82)90123-2)

1055 Thomsen, J., Himmerkus, N., Holland, N., Sartoris, F.J., Bleich, M., & Tresguerres, M.
 1056 (2016). Ammonia excretion in mytilid mussels is facilitated by ciliary beating. *The*
 1057 *Journal of Experimental Biology*, 219, 2300-2310. doi: 10.1242/jeb.139550.

1058 Torigoe, T., Izumi, H., Ise, T., Murakami, T., Uramoto, H., Ishiguchi, H., Yoshida, Y.,
 1059 Tanabe, M., Nomoto, M., & Kohno, K. (2002). Vacuolar H⁺-ATPase: functional
 1060 mechanisms and potential as a target for cancer chemotherapy. *Anticancer Drugs*, 13,
 1061 237-243.

1062 Toyofuku, T., Matsuo, M. Y., De Nooijer, L. J., Nagai, Y., Kawada, S., Fujita, K.,
 1063 Reichart, G.J., Nomaki, H., Tsuchiya, M., Sakaguchi, H., & Kitazato, H. (2017). Proton
 1064 pumping accompanies calcification in foraminifera. *Nature Communications*, 8, 14145.
 1065 <https://doi.org/10.1038/ncomms14145>

1066 Tresguerres, M. (2016). Novel and potential physiological roles of vacuolar-type H⁺-
 1067 ATPase in marine organisms. *The Journal of Experimental Biology*, 219, 2088-2097.
 1068 <https://doi:10.1242/jeb.128389>

1069 Tresguerres, M., Barott, K. L., Barron, M. E., & Roa, J. N. (2014). Established and
 1070 potential physiological roles of bicarbonate-sensing soluble adenylyl cyclase (sAC) in
 1071 aquatic animals. *The Journal of Experimental Biology*, 217, 663–672.
 1072 <https://doi.org/10.1242/jeb.086157>

1073 Tresguerres, M., Barott, K. L., Barron, M. E., Deheyn, D. D., Kline, D. I., & Linsmayer, L.
 1074 B. (2017). Cell biology of reef-building corals: Ion transport, acid/base regulation, and
 1075 energy metabolism. In D. Weihrauch & M. O'Donnell (Eds.), *Acid-Base Balance and*
 1076 *Nitrogen Excretion in Invertebrates* (pp. 193–218). [https://doi.org/10.1007/978-3-319-](https://doi.org/10.1007/978-3-319-39617-0_7)
 1077 [39617-0_7](https://doi.org/10.1007/978-3-319-39617-0_7)

1078 Tresguerres, M., Buck, J., & Levin, L. R. (2010). Physiological carbon dioxide,
 1079 bicarbonate, and pH sensing. *Pflügers Archiv*, 460, 953–964.
 1080 <https://doi.org/10.1007/s00424-010-0865-6>

1081 Tresguerres, M., & Hamilton, T. J. (2017). Acid-base physiology, neurobiology and
 1082 behaviour in relation to CO₂-induced ocean acidification. *The Journal of Experimental*
 1083 *Biology*, 220, 2136–2148. <https://doi.org/10.1242/jeb.144113>

1084 Tresguerres, M., Levin, L. R., & Buck, J. (2011). Intracellular cAMP signaling by soluble
 1085 adenylyl cyclase. *Kidney International*, 79, 1277–1288.
 1086 <https://doi.org/10.1038/ki.2011.95>

1087 Tresguerres, M., Milsom, B., Perry, S.F. (2019) CO₂ and acid-base sensing. In M.
 1088 Grosell, P. L. Munday, A. P. Farrell, & C. J. Brauner (Eds.), *Carbon Dioxide*. San Diego,
 1089 CA: Academic Press.

1090 Tresguerres, M., Levin, L. R., Buck, J., & Grosell, M. (2010). Modulation of NaCl
 1091 absorption by [HCO₃⁻] in the marine teleost intestine is mediated by soluble adenylyl
 1092 cyclase. *American Journal of Physiology-Regulatory, Integrative and Comparative*
 1093 *Physiology*, 299(1), R62–R71. <https://doi.org/10.1152/ajpregu.00761.2009>

1094 Truchot, J.-P., & Duhamel-Jouve, A. (1980). Oxygen and carbon dioxide in the marine
 1095 intertidal environment: Diurnal and tidal changes in rockpools. *Respiration Physiology*,
 1096 39, 241–254. [https://doi.org/10.1016/0034-5687\(80\)90056-0](https://doi.org/10.1016/0034-5687(80)90056-0)

1097 Tufts, B. L., & Boutilier, R. G. (1989). The absence of rapid chloride/bicarbonate
 1098 exchange in lamprey erythrocytes: implications for CO₂ transport and ion distributions
 1099 between plasma and erythrocytes in the blood of *Petromyzon marinus*. *The Journal of*
 1100 *Experimental Biology*, 144, 565–576.

1101 Tufts, B. L., & Perry, S. F. (1998). Carbon dioxide transport and excretion. In S. F. Perry
 1102 & B. L. Tufts (Eds.), *Fish Physiology. Vol. 17: Fish respiration* (pp. 229–282). New York:
 1103 Academic Press.

1104 Ultsch, G. R. (1973). The effects of water hyacinths (*Eichhornia crassipes*) on the
 1105 microenvironment of aquatic communities. *Archiv für Hydrobiologie*. 72, 460-473.

1106 Ultsch, G.R., & D.S. Antony (1973). The role of the aquatic exchange of carbon dioxide
 1107 in the ecology of the water hyacinth (*Eichhornia crassipes*). *Florida Scientist* 36, 16-22.

1108 Van de Waal, D. B., Verschoor, A. M., Verspagen, J. M., van Donk, E. & Huisman, J.
 1109 (2009). Climate-driven changes in the ecological stoichiometry of aquatic ecosystems.

1110 *Frontiers in Ecology and the Environment*, 8, 145, 152.

1111 van Holde, K. E., & Miller, K. I. (1995). Hemocyanins. In C.B. Anfinsen, Frederic M.

1112 Richards, John T. Edsall, & David S. Eisenberg (Eds.), *Advances in Protein Chemistry*

1113 (Vol. 47). New York: Academic Press. [https://doi.org/10.1016/S0065-3233\(08\)60545-8](https://doi.org/10.1016/S0065-3233(08)60545-8)

1114 Voight, J. R., Pörtner, H. O., & O'Dor, R. K. (1995). A review of ammonia-mediated

1115 buoyancy in squids (cephalopoda: Teuthoidea). *Marine and Freshwater Behaviour and*

1116 *Physiology*, 25, 193–203. <https://doi.org/10.1080/10236249409378917>

1117 Wacker, T., Garcia-Celma, J. J., Lewe, P., & Andrade, S. L. A. (2014). Direct

1118 observation of electrogenic NH_4^+ transport in ammonium transport (Amt) proteins.

1119 *Proceedings of the National Academy of Sciences*, 111, 9995–10000.

1120 <https://doi.org/10.1073/pnas.1406409111>

1121 Wagner, P. D. (1977). Diffusion and chemical reaction in pulmonary gas exchange.

1122 *Physiological Reviews*, 57, 257–312.

1123 Walsh, P. J., & Milligan, C. L. (1989). Coordination of metabolism and intracellular acid–

1124 base status: Ionic regulation and metabolic consequences. *Canadian Journal of*

1125 *Zoology*, 67(12), 2994–3004. <https://doi.org/10.1139/z89-422>

1126 Weber, J.-M., Choi, K., Gonzalez, A., & Omlin, T. (2016). Metabolic fuel kinetics in fish:

1127 swimming, hypoxia and muscle membranes. *The Journal of Experimental Biology*, 219,

1128 250–258.

1129 Weihrauch, D., & Allen, G. J. P. (2018). Ammonia excretion in aquatic invertebrates:

1130 New insights and questions. *The Journal of Experimental Biology*, 221, jeb169219.

1131 <https://doi.org/10.1242/jeb.169219>

1132 Weihrauch, D., & O'Donnell, M. (Eds.). (2017). *Acid-Base Balance and Nitrogen*
 1133 *Excretion in Invertebrates: Mechanisms and Strategies in Various Invertebrate Groups*
 1134 *with Considerations of Challenges Caused by Ocean Acidification*.
 1135 <https://doi.org/10.1007/978-3-319-39617-0>
 1136 Weihrauch, D., Ziegler, A., Siebers, D., & Towle, D. W. (2002). Active ammonia
 1137 excretion across the gills of the green shore crab *Carcinus maenas*: Participation of
 1138 Na⁺/K⁺-ATPase, V-type H⁺-ATPase and functional microtubules. *The Journal of*
 1139 *Experimental Biology*, 205, 2765–2775.
 1140 Weiner, I. D., & Verlander, J. W. (2017). Ammonia transporters and their role in acid-
 1141 base balance. *Physiological Reviews*, 97, 465–494.
 1142 <https://doi.org/10.1152/physrev.00011.2016>
 1143 Wilkie, M. P. (2002). Ammonia excretion and urea handling by fish gills: Present
 1144 understanding and future research challenges. *Journal of Experimental Zoology*, 293,
 1145 284–301.
 1146 Wittenberg, J. B., & Wittenberg, B. A. (1962). Active secretion of oxygen into the eye of
 1147 fish. *Nature*, 194(4823), 106–107. <https://doi.org/10.1038/194106a0>
 1148 Wood, C. M., Bucking, C., & Grosell, M. (2010). Acid–base responses to feeding and
 1149 intestinal Cl[−] uptake in freshwater- and seawater-acclimated killifish, *Fundulus*
 1150 *heteroclitus*, an agastric euryhaline teleost. *The Journal of Experimental Biology*, 213,
 1151 2681–2692. <https://doi.org/10.1242/jeb.039164>
 1152 Wood, C. M., Kajimura, M., Mommsen, T. P., & Walsh, P. J. (2005). Alkaline tide and
 1153 nitrogen conservation after feeding in an elasmobranch (*Squalus acanthias*). *The*
 1154 *Journal of Experimental Biology*, 208, 2693–2705. <https://doi.org/10.1242/jeb.01678>

1155 Wright, P. A., & Wood, C. M. (2009). A new paradigm for ammonia excretion in aquatic
 1156 animals: Role of Rhesus (Rh) glycoproteins. *The Journal of Experimental Biology*, 212,
 1157 2303–2312. doi: 10.1242/jeb.023085

1158 Yellowlees, D., Rees, T. A. V., & Leggat, W. (2008). Metabolic interactions between
 1159 algal symbionts and invertebrate hosts. *Plant, Cell & Environment*, 31, 679–694.
 1160 <https://doi.org/10.1111/j.1365-3040.2008.01802.x>

1161 Zange, J., Grieshaber, M. K., & Jans, A. W. H. (1990). The regulation of intracellular pH
 1162 estimated by ³¹P-NMR spectroscopy in the anterior byssus retractor muscle of *Mytilus*
 1163 *edulis* L. *The Journal of Experimental Biology*, 150, 95–109.

1164 Zoccola, D., Ganot, P., Bertucci, A., Caminiti-Segonds, N., Techer, N., Voolstra, C. R.,
 1165 Aranda, M., Tambutté, E., Allemand, D., Casey, J. R., & Tambutté, S. (2015).
 1166 Bicarbonate transporters in corals point towards a key step in the evolution of cnidarian
 1167 calcification. *Scientific Reports*, 5, 9983. <https://doi.org/10.1038/srep09983>

1168 Zoccola, D., Tambutté, E., Kulhanek, E., Puverel, S., Scimeca, J. C., Allemand, D., &
 1169 Tambutté, S. (2004). Molecular cloning and localization of a PMCA P-type calcium
 1170 ATPase from the coral *Stylophora pistillata*. *Biochimica & Biophysica Acta*, 1663, 117-
 1171 126.

1172 Zippin, J. H., Farrell, J., Huron, D., Kamenetsky, M., Hess, K. C., Fischman, D. A.,
 1173 Levin, L. R., Buck, J. (2004). Bicarbonate-responsive “soluble” adenylyl cyclase defines
 1174 a nuclear cAMP microdomain. *The Journal of Cell Biology*, 164, 527–534.
 1175 <https://doi.org/10.1083/jcb.200311119>

1176

1177

Figure Legends

Figure 1. pH-dependent chemical equilibria between CO_2 , HCO_3^- , and CO_3^{2-} . (A)

Equation describing the hydration of CO_2 and its subsequent equilibrium reactions with the other dissolved inorganic carbon (DIC) species. The pKAs shown under each reaction are the negative logarithm of the respective dissociation constant, and indicate the pH at which the relevant DIC species are found in equimolar amounts. The hydration of CO_2 into H_2CO_3 is relatively slow; however, in the presence of carbonic anhydrase (CA) enzymes this reaction takes place virtually instantaneously. **(B)** pH-dependent relative proportion of DIC species. Notice how a more acidic pH favors $\text{CO}_2/\text{H}_2\text{CO}_3$, a relatively neutral pH in the biological range favors HCO_3^- , and a more alkaline pH favors CO_3^{2-} .

Figure 2. Summary of pHi regulatory mechanisms. Schematic of a generic eukaryotic cell depicting the most common pHi regulatory mechanisms. **(1)** The mitochondrial H^+ sink maintains a close balance between H^+ production and consumption during resting aerobic metabolism. However, an increase in **(2)** anaerobic metabolism or **(3)** PCO_2 can result in an intracellular H^+ load that acidifies pHi. **(4)** The activity of carbonic anhydrase (CA) mediates the nearly instantaneous reversible equilibration between CO_2 with HCO_3^- and H^+ . **(5)** Buffering is the first line of defense against pHi fluctuations. **(6)** When an acidic load breaches the buffering capacity, cells actively excrete H^+ through Na^+/H^+ exchanger (NHE), V-type ATPase (VHA) and monocarboxylate transporter (MCTs), or take up HCO_3^- through $\text{Na}^+/\text{HCO}_3^-$

1201 cotransporter (NBCs), and/or Na^+ -dependent $\text{Cl}^-/\text{HCO}_3^-$ exchangers (NDCBEs). **(7)**
1202 Active intracellular acidification can be mediated by anion exchanger (AE) proteins and
1203 by Plasma Membrane Ca^{2+} -ATPase (PMCA) (which links intracellular pH with Ca^{2+}
1204 homeostasis). **(8)** The driving force for most of these transporters is provided by the
1205 inward directed Na^+ electrochemical gradient and the inside negative membrane
1206 potential established by the Na^+/K^+ -ATPase (NKA). However, VHA and PMCA directly
1207 hydrolyze ATP and their activities are not dependent on NKA. The lightning bolts indicate
1208 ATP hydrolysis. **(9)** The end products of anaerobic fermentation can have different
1209 fates. The lactate that is produced by lactate dehydrogenase (LDH) is typically excreted
1210 from cells together with H^+ *via* MCTs; however, some cells can retain the lactate and
1211 use it to replenish their glycogen stores. Similarly, the “opines” produced by opine
1212 dehydrogenases (OpDH) during fermentation in invertebrates are retained intracellularly
1213 and reconverted into the original substrates upon the return of aerobic conditions.

1214

1215 **Figure 3. Acid-base sensing by sAC.** Separate pools of sAC in **(1)** the cytoplasm and
1216 **(2)** the nucleus can be stimulated by HCO_3^- from the following sources: **(a)** Carbonic
1217 anhydrase (CA)-dependent hydration of external CO_2 ; **(b)** CA-dependent hydration of
1218 CO_2 generated through mitochondrial respiration; **(d)** entry through bicarbonate
1219 transporters (BT) such as the ones shown in Figure 2 or through channels such as
1220 cystic transmembrane conductance regulator (CFTR). **(e)** The activities of H^+ extruding
1221 transporters (HE) (Figure 2) remove H^+ from the cell and may prevent slowing down of
1222 the CO_2 hydration reaction. The cyclic adenosine monophosphate (cAMP) that is
1223 generated by sAC can regulate the activities of multiple and diverse target proteins *via*

Protein Kinase A (PKA)-dependent phosphorylation, Exchange Protein Activated by cAMP (EPAC) signaling, and gating of membrane channels. In the nucleus, the sAC-cAMP-PKA pathway can regulate the activity of the gene transcription factor, cAMP responsive element binding (CREB). Modified from Tresguerres et al. (2014).

Figure 4. The role of red blood cell (RBC) pHi on systemic O₂ transport in fish. (A)

At the capillaries CO₂ from the tissues diffuses into the blood and into the RBCs. In the presence of carbonic anhydrase (CA) within the RBC CO₂ is rapidly converted into H⁺ and HCO₃⁻. The binding of H⁺ to Hb decreases its affinity for O₂ (Bohr effect; a right shift in the oxygen equilibrium curve; OEC), which is then released to the tissues. The binding of H⁺ to Hb also buffers arterial-venous pH differences promoting pH homeostasis. Within RBCs, the produced HCO₃⁻ is exported into the plasma by the anion exchanger (AE). At the gills the process is reversed: when CO₂ diffuses out of the blood and into the water, HCO₃⁻ is taken up into the RBC and converted into CO₂, which maintains the diffusion gradient for excretion. At the same time, the binding of O₂ to Hb decreases its affinity for H⁺ (Haldane effect), which are released into the RBC and are used by CA to dehydrate HCO₃⁻. In the absence of H⁺ binding, Hb increases its affinity for O₂, which promotes oxygenation of the blood (left shift of the OEC). **(B)** The Jacobs-Stewart cycle describes the transfer of H⁺ across the RBC membrane. H⁺ are charged ions and do not readily diffuse across lipid membranes. In the plasma H⁺ react with HCO₃⁻ to form CO₂ which rapidly diffuses across the membrane and this is often facilitated by channel proteins such as aquaporin 1 (AQP1) and RhAG. Within the RBC CO₂ will dissociate into H⁺, that bind to intracellular buffers, and HCO₃⁻ that is exported

1247 into the plasma by AE. **(C)** Summary of transporters that regulate RBC pHi in fish. The
1248 Na^+/K^+ -ATPase (NKA) creates trans-membrane gradients for Na^+ and K^+ that are used
1249 by secondary active transporters to drive ion transport. Alkalinizing transporters: Na^+ - H^+
1250 exchangers (NHE) use the Na^+ gradient to extrude H^+ ; β -adrenergically activated NHEs
1251 (β -NHE) are activated by catecholamine binding to a receptor on the RBC membrane;
1252 Na^+ - K^+ - 2Cl^- -cotransporter (NKCC) uses the Na^+ gradient to drive net Cl^- uptake.
1253 Because the activities of Cl^- are linked to those of H^+ (via the Jacobs-Stewart cycle),
1254 NKCC activity will increase RBC pHi. On the other hand the K^+ - 2Cl^- -cotransporter
1255 (KCC) will lead to a decrease in RBC pHi due to a net excretion of Cl^- . See text for
1256 further details and references.

1257

1258 **Figure 5. pH-dependent chemical equilibrium of ammonia. (A)** Equation describing
1259 the hydration of NH_4^+ and the subsequent equilibrium reaction with NH_3 . The pKA
1260 shown under the reaction is the negative logarithm of the dissociation constant, and
1261 indicates the pH at which NH_3 and NH^+ are found in equimolar amounts. **(B)** pH-
1262 dependent relative proportion of NH_3 and NH_4^+ . Notice that at physiological pH, NH_4^+ is
1263 by far the dominant species, that acidification further favors NH_4^+ , and that alkalinization
1264 favors NH_3 .

1265

1266 **Figure 6. Acid-trapping of ammonia. (1)** Intracellular ammonia (the sum of NH_3 and
1267 NH_4^+) is predominantly derived from amino acid catabolism in mitochondria, from
1268 facilitated diffusion through ammonium transporters (AMTs) and Rhesus channel
1269 glycoproteins (Rhs), and through import by K^+ -transporting proteins (K^+T) such as

1270 Na^+/K^+ -ATPase, $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ -cotransporter, $\text{K}^+/\text{2Cl}^-$ -cotransporter, and K^+ -channels. **(2)**
1271 At a typical intracellular pH (pH_i), most ammonia is present as NH_4^+ (see Fig. 5).
1272 However, acidification of **(3)** the external boundary layer or **(4)** intracellular vesicles acts
1273 as a siphon for NH_3 , producing NH_4^+ that gets trapped outside of the cell or in the
1274 vesicle, while sustaining a favorable NH_3 partial pressure gradient ($\Delta P\text{NH}_3$) that
1275 promotes further ammonia transport and trapping. The acidification can take place
1276 through H^+ transport by Na^+/H^+ Exchanger (NHE) or V-type- H^+ -ATPase (VHA), and NH_3
1277 diffusion is facilitated by Rh. In addition, NH_3 may diffuse across cellular membranes or
1278 paracellularly (not shown). **(5)** The vesicles can be trafficked to the apical membrane in
1279 microtubule-dependent manner, and the trapped NH_4^+ excreted *via* exocytosis, as
1280 reported in the gills of some marine crabs (note that the cuticle at the apical side has
1281 been omitted for clarity). In addition, similar acidic and NH_4^+ -rich vesicles can be stored
1282 within the body tissues of deep-sea cephalopods and other marine invertebrates for the
1283 purpose of achieving buoyancy (however, the pathways for NH_3 and H^+ transport remain
1284 unknown).

1285

1286 **Figure 7. Extreme pH microenvironments in corals. (A)** Simplified coral histology
1287 diagram showing the movements of acid-base relevant molecules between seawater,
1288 host cells with algal symbionts, and the site of calcification (ECM: extracellular calcifying
1289 medium). Together with Ca^{2+} transport into the ECM and vesicular transport of
1290 amorphous CaCO_3 (not shown), the alkaline pH in the SCM promotes coral skeleton
1291 formation. DIC: dissolved inorganic carbon ($\text{CO}_2 + \text{HCO}_3^- + \text{CO}_3^{2-}$). The pH of
1292 extracellular and intracellular compartments is noted to the left (sun and moon indicating

1293 day- and nighttime pH for seawater respectively). Photosynthesis and calcification are
1294 linked by translocation of photosynthetic products to the site of calcification (i.e. oxygen
1295 and sugars) and calcification byproducts (H^+) to host cells. **(B)** Schematic of a coral host
1296 cell containing an algal symbiont to illustrate the CCM. The alga is not drawn to scale to
1297 allow for clarity but usually occupies >90% of a host cell's volume. BT: HCO_3^-
1298 transporter; CA: carbonic anhydrase; VHA: V-Type H^+ -ATPase. **(C)** VHA
1299 immunostaining (red signal) of a symbiont-containing coral gastrodermal cell showing
1300 abundant VHA presence in the symbiosome membrane. The other proteins involved in
1301 transport of ions and other molecules are omitted for simplicity, and in many cases their
1302 identities are unknown.

1303

1304

1305

1306

1307

1308

A

B

Figure 1

Figure 2

Figure 4

A

B

Figure 5

Figure 6

Figure 7