

Lawrence Berkeley National Laboratory

Recent Work

Title

Higher-order differencing schemes for modeling intense laser-plasma interactions

Permalink

<https://escholarship.org/uc/item/3sj0v063>

Authors

Tarkenton, G.M.

Shadwick, B.A.

Schroeder, C.B.

Publication Date

2003-10-01



C

Abstract Submitted
for the DPP03 Meeting of
The American Physical Society

Sorting Category: 2.3.1

Higher-Order Differencing Schemes for Modeling Intense Laser-Plasma Interactions G. M. TARKENTON, Institute for Advanced Physics, B. A. SHADWICK, Center for Beam Physics, LBNL and Institute for Advanced Physics, C. B. SCHROEDER, Center for Beam Physics, LBNL Traditionally numerical solutions of PDEs use methods that are second-order in both space and time. This is both for historic reasons and because conservative differencing schemes have important properties especially for hydrodynamic modeling. With the aid of modern symbolic algebraic methods, designing higher-order, stable methods is practical. Higher-order methods are attractive as they offer the promise of increased accuracy compared to second-order methods for a given set of grid parameters. We discuss the stability, dispersion, and computation performance of various higher-order methods. We find that the use of higher-order methods for solving both the full-fluid and envelope models yield and significant performance gains (from reduced grid sizes and a larger stability basin) over the corresponding second-order solvers.

- Prefer Oral Session
 Prefer Poster Session

B. A. Shadwick
shadwick@mailaps.org
LBNL/LAP

Date submitted: 18 Jul 2003

Electronic form version 1.4