UC San Diego

UC San Diego Previously Published Works

Title

Reproductive Stage and Modulation of Stress-Induced Tau Phosphorylation in Female Rats.

Permalink

<https://escholarship.org/uc/item/3t79t28v>

Journal

Journal of neuroendocrinology, 27(11)

ISSN

0953-8194

Authors

Steinmetz, Danielle
Ramos, Eugenia
Campbell, Shannon N
et al.

Publication Date

2015-11-01

DOI

10.1111/jne.12323

Peer reviewed

ORIGINAL ARTICLE

Reproductive Stage and Modulation of Stress-Induced Tau Phosphorylation in Female Rats

Danielle Steinmetz*, Eugenia Ramos†, Shannon N. Campbell*, Teresa Morales† and Robert A. Rissman*

*Department of Neurosciences, University of California, San Diego, La Jolla, CA, USA.

†Instituto de Neurobiología, Universidad Nacional Autónoma de México, Queretaro, México.

Journal of Neuroendocrinology

Chronic stress is implicated as a risk factor for Alzheimer's disease (AD) and other neurodegenerative disorders. Although the specific mechanisms linking stress exposure and AD vulnerability have yet to be fully determined, our laboratory and others have shown that acute and repeated restraint stress in rodents leads to an increase in hippocampal tau phosphorylation (tau-P) and tau insolubility, a critical component of tau pathology in AD. Although tau phosphorylation induced by acute psychological stress is dependent on intact signaling through the type 1 corticotropin-releasing factor receptor, how sex steroids or other modulators contribute to this effect is unknown. A naturally occurring attenuation of the stress response is observed in female rats at the end of pregnancy and throughout lactation. To test the hypothesis that decreased sensitivity to stress during lactation modulates stress-induced tau-P, cohorts of virgin, lactating and weaned female rats were subjected to 30 min of restraint stress or no stress (control) and were killed 20 min or 24 h after the episode. Exposure to restraint stress induced a significant decrease in tau-P in the hippocampus of lactating rats killed 20 min after stress compared to lactating controls and virgins subjected to stress treatment. Lactating rats killed 24 hr after restraint stress exposure showed significant elevation in tau-P compared to lactating cohorts killed 20 min after stress. Levels of tau-P in these latter cohorts did not differ significantly from control animals. Furthermore, glycogen synthase kinase (GSK)3- α levels were significantly decreased in stressed lactating animals at both timepoints. This suggests a steep, yet transient stress-induced dephosphorylation of tau, influenced by GSK3, in the hippocampus of lactating rats.

Key words: Alzheimer's disease, stress, lactation, corticotropin-releasing, tau, hippocampus, GSK3, steroid, Prolactin

Correspondence to: Robert A. Rissman, Department of Neurosciences, UCSD School of Medicine, 9500 Gilman Drive, MTF 314 M/C 0624, La Jolla, CA 92093-0624, USA (e-mail: rrrissman@ucsd.edu)

doi: 10.1111/jne.12323

Stress is implicated as a risk factor for the development of Alzheimer's disease (AD), a neurodegenerative disorder defined pathologically by the accumulation of extracellular beta-amyloid (A β) plaques and intracellular neurofibrillary tangles comprised of hyperphosphorylated tau (tau-P) aggregates. The role of maternal and ovarian hormones in the development or prevention of neurodegenerative diseases such as AD has become a subject of intense study because certain hormones appear to demonstrate neuroprotective and neurogenic qualities. For example, during lactation, remarkable adaptations occur in the female brain, including attenuation of the hypothalamic-pituitary-adrenal (HPA) axis to stress and changes in hippocampal plasticity (1–3). The observed increase in plasticity can be influenced by hormonal fluctuations that occur

during lactation in conjunction with suckling stimulation from the litter (4).

Studies have found that over-activation of the HPA axis results in decreased hippocampal neurogenesis, increased neurodegeneration and increased cognitive impairment (5–7). Stress-inducing environmental factors can play a role in AD development and, more specifically, can induce tau-P (7–14). Studies show increased tau-P in rodents subjected to cold water stress, as well as cognitive deficits as a result of excess glucocorticoid (stress hormone) exposure (9,11). Moreover, exposure to a single episode of an emotional stressor, restraint, rapidly induces tau-P in the hippocampus. Furthermore, with repeated restraint exposure or overexpression of corticotropin-releasing factor (CRF), cumulative increases in

insoluble tau-P is observed (13–15). These studies implicate the CRF pathway as being mechanistically involved in stress-induced tau-P, as this phenomenon was not observed in the presence of pharmacological blockade or genetic knockout of CRF receptor 1 (7,13,14).

The physiological changes that occur during pregnancy and lactation may confer neuroprotection against excitotoxins, such as kainic acid, and lead to a decreased sensitivity to stress (16,17). The morphological and functional changes in the maternal brain occur not only in areas that support lactation, but also in areas of learning and memory, such as the CA1 region of the hippocampus (2,3), as well as areas related to neurogenesis, such as the subventricular zone and the dentate gyrus (18,19). Reproduction also facilitates learning and memory and decreases the prevalence of neuronal markers of ageing (20). The number of studies aiming to determine the relationship between maternal hormones (i.e. prolactin, progesterone and oestrogen), neurodegeneration and AD pathology is rapidly growing, potentially as a result of fluctuations in these hormone levels during pregnancy and lactation, and increasing evidence of their effect on the hippocampus, learning and memory. For example, in rodents, prolactin has been shown to decrease anxiety, prevent stress-induced decreases in neurogenesis (21, 22) and to diminish excitotoxic cell damage in the hippocampus (23). In the hippocampus of AD mice, progesterone significantly reduces tau-P and oestrogen prevents A β accumulation (24). The mechanisms by which these hormones regulate tau and A β are not fully understood; however, these studies suggest that maternal hormones work to attenuate the stress response and could potentially play a role in preventing AD pathology.

Lactation is a reproductive condition in which the circadian fluctuation of corticosterone is abolished but basal levels are chronically elevated (1). Thus, a more in-depth understanding of the mechanisms by which lactation reduced sensitivity to stress may further clarify the specific circuitry and cellular mechanisms that influence AD vulnerability. In the present study, we address the impact of the reproductive condition on stress-induced tau-P and major tau kinases [glycogen synthase kinase (GSK), extracellular signal-regulated kinase (ERK) and *c-Jun*-N-terminal kinase (JNK)] in female rats.

Materials and methods

Animals

Adult (250–300 g), virgin or pregnant female Wistar rats were housed individually under a 12 : 12 h light/dark cycle (lights on 06:00 h) and controlled temperature, with food and water available *ad libitum*. One day after parturition, litter sizes were culled to eight to 10 pups. Mothers and their litters were undisturbed until postpartum days 14–19 (lactating rats) or 1 month after weaning at day 22 (weaned rats). Vaginal smears of virgin and weaned female rats were followed for at least three oestrous cycles, and they were stressed or handled in the diestrous phase of the cycle (virgin rats). The Institutional Animal Care and Use Committee of the Institute for Neurobiology at the Universidad Nacional Autonoma de Mexico approved all experimental protocols. Animals were handled in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

Acute restraint stress

Rats were placed in acrylic restrainers with ventilation for 30 min, and were killed either 20 min or 24 h after the restraint episode. Control rats were similarly handled but not exposed to stress. Five rats were included in each group.

Western blot analysis

One set of rats was killed by decapitation, and brains were immediately removed and hippocampus was collected, immediately frozen on dry ice and stored at -70°C until assay. Tissues were homogenised in radio-immunoprecipitation assay buffer as described previously (14,25). Fifteen micrograms of protein was separated on 10% sodium dodecylsulphate-polyacrylamide gel electrophoresis and transferred to nitrocellulose membrane at 100 V for 50 min. Nonspecific binding was blocked by incubating membranes in 5% milk-phosphate buffered saline containing Tween 20 (PBS-T) for 30 min. Membranes were incubated with primary antibodies diluted in 5% milk-PBS-T overnight at 4°C . The primary antibodies were detected by anti-mouse or anti-rabbit horseradish peroxidase-linked secondary antibodies for 1 h (dilution 1 : 2500; EMD Biosciences, La Jolla, CA, USA) and developed using an enhanced chemiluminescence Western blot detection kit (Supersignal West Pico; Pierce Biotechnology, Rockford, IL, USA). Quantitative band intensity readings were obtained using ImageJ (NIH, Bethesda, MD, USA). All readings were normalised to internal standard band intensity readings.

Corticosterone and prolactin determination

Trunk blood was collected to determine serum corticosterone and prolactin levels in groups of rats either from control or stress treatment at 20 min after 30 min of restraint ($n = 5$). Serum was separated by centrifugation (10 min at 9469 g) and stored at -20°C until assay. Corticosterone was determined using immunoassay kits (catalogue number ADI-900-097; Enzo Life Sciences, Farmingdale, NY, USA) that employed a competitive immunoassay for quantitative determinations. The optical density of the samples and controls from the standard curve was read immediately was read at 450nm using a microplate reader (Bio Rad, Hercules, CA, USA). For prolactin determination, microtitre plates were coated overnight with 10 ng of rat prolactin (NHPP-NIH, Torrance, CA, USA). Serial dilutions of standard rat prolactin (rPRL) (0.06–64 ng/ml) in TPBS were incubated for 16 h with 100 μl of primary anti-rPRL polyclonal antiserum (dilution 1 : 40 000; NHPP-NIH) in TPBS containing 1% (w/v) nonfat dry milk (Bio-Rad, Hercules, CA, USA). Samples and standards (100 μl) were then added to the coated wells and incubated for 2 h at room temperature. Secondary goat anti-rabbit immunoglobulin G peroxidase conjugate (Bio-Rad) was added (dilution 1 : 3000 in TPBS with 1% nonfat dry milk) and incubated for 2 h at room temperature. Bound secondary antibodies were detected by reaction with 2,20 azino-di-(3-ethylbenzothiazoline sulphonate) substrate (Roche, Mannheim, Germany). Plates were read at 450nm using a microplate reader (Bio Rad).

Antibodies

Two well-characterised antibodies were used to probe for specific phosphorylated residues on rat hippocampal tau: S202/T205 (AT8, dilution 1 : 500; Pierce Biotechnology) and S396/404 (PHF-1, dilution 1 : 1000; gift from Dr P. Davies, Albert Einstein College of Medicine, Bronx, NY, USA). Kinase expression and activity was assessed using antibodies specific to total GSK-3 β (dilution 1 : 1000; BD Biosciences, San Diego, CA, USA), inactive GSK-3 (pS⁹, dilution 1 : 1000; Cell Signaling Technology, Danvers, MA, USA), active GSK-3 (pY²¹⁶, dilution 1 : 1000; BD Biosciences), mitogen-activated protein

kinases (ERK1/2), dilution 1 : 500; Cell Signaling Technology] and phosphorylated JNK (pJNK, dilution 1 : 1000; Cell Signaling Technology). Actin (dilution 1 : 2000; Sigma-Aldrich, St Louis, MO, USA) was used as a protein-loading control.

Immunohistochemistry

A separate set of rats were anaesthetised with a lethal dose of urethane (1.5–2 g/kg bw, i.p.), perfused with 4% paraformaldehyde and processed for immunohistochemistry as described previously (13). 30 µm-thick coronal sections were cut on a freezing-sliding microtome and stored at –20 °C in antifreeze until use. PHF-1 (dilution 1 : 5000) was used to probe for tau-P on free-floating sections of rat hippocampus. 0.3% hydrogen peroxide was used to quench endogenous peroxidases and 0.1% sodium borohydride was used to reduce reactive free aldehydes. The staining reaction was carried out using a nickel-enhanced DAB Peroxidase Substrate kit (Vector Laboratories, Inc., Burlingame, CA, USA).

Statistical analysis

Integrated intensity readings from western blots were analysed using either a one- or two-way ANOVA using R (The R Project for Statistical Computing, Vienna, Austria) and PRISM (GraphPad Software Inc., San Diego, CA, USA) software; pair comparison was estimated by a *t*-test. Resultant data were plotted on bar graphs, with data expressed as the mean ± SEM percentage of control values.

Results

Basal levels of tau-P and tau kinase expression in the hippocampus of female rats at various reproductive stages

Western blot analysis was used to determine the effects of stress on hippocampal tau-P and tau kinases in female rats at two well-characterised phospho-epitopes, AT8 and PHF-1. Control animals showed no significant baseline differences in tau-P at the AT8 and PHF-1 phosphorylation sites throughout all reproductive stages studied (Fig. 1). However, as shown in Fig. 2, nonstressed lactating control animals express significantly lower levels of total GSK3-α ($F_{7,28} = 42.4$, $P < 0.001$) compared to weaned ($P = 0.016$) and virgin ($P = 0.008$) nonstressed controls. There were no significant differences in ERK or JNK levels across all controls ($F_{7,28} = 32.1$, $P > 0.05$) (Fig. 2).

Effects of stress on tau-P in the hippocampus of lactating rats

Western blot analysis was used to determine the effects of stress on hippocampal tau phosphorylation using AT8 and PHF-1. As shown in Fig. 1, 20-min after restraint stress exposure, lactating rats showed significantly lower levels of tau-P at both the AT8 ($F_{7,28} = 42.4$, $P = 0.0008$) ($P = 0.032$) and PHF-1 ($F_{7,28} = 2.5$, $P = 0.042$) ($P = 0.03$) epitopes compared to lactating controls. This significant change was also observed compared to stressed virgin ($P = 0.008$) and weaned ($P = 0.008$) rats (Fig. 1). 24 hours after stress, tau-P levels in lactating rats at the PHF-1 site significantly increased ($P = 0.008$) compared to the levels observed at 20 min,

Fig. 1. Acute stress-induced hippocampal tau dephosphorylation in lactating rats. Female rats that were virgin, lactating (14–19 postpartum) or weaned (21 days postpartum) ($N = 5$, each) were sacrificed after either no stress (NS), 20 min (20) or 24 h (24) after a single restraint episode. Lactating rats had significantly reduced levels of stress-induced tau phosphorylation (tau-P) at both PHF-1 and AT8 epitopes compared to virgin (*, $P = 0.008$) and weaned rats ($P = 0.008$); however, tau-P returned to control/basal levels after 24 h. Data are presented as mean ± SEM percentage change.

returning to near-control levels. No significant changes in tau-P levels were observed in virgin or weaned rats in response to stress.

To clarify the possible mechanisms of this stress-induced dephosphorylation of tau, changes in several known tau kinases, such as GSK3, were investigated using western blot analysis. As shown in Fig. 3, overall levels of total GSK3-α significantly decreased in lactating rats compared to lactating controls ($F_{7,28} = 42.4$, $P < 0.001$) ($P = 0.008$) 20 min after stress, and these levels remained low 24 h after stress exposure ($P = 0.008$). Interestingly, active GSK3-β significantly increased ($F_{7,28} = 17.61$, $P < 0.0001$) in lactating rats 24 h after stress exposure compared to lactating controls ($P = 0.016$), whereas there were no significant changes in active GSK3-α or inactive GSK3(α or β) in stress-exposed lactating animals compared to lactating controls. This significant drop in GSK3 20 min after stress could play a key role in the stress-induced dephosphorylation of hippocampal tau observed in restraint-stressed lactating rats.

Analysis of corticosterone levels in response to restraint by enzyme-linked immunosorbent assay showed that basal level of corticosterone was higher ($F_{2,22} = 5.43$, $P = 0.01$) in lactating than in virgin ($P < 0.0001$) (Fig. 4) and weaned rats ($P = 0.0377$). Because of the variation of data, increases in the concentration of corticosterone caused by stress were not detected in virgin and weaned rats (three- and two-fold of corresponding control, respectively). However, a tendency to increase after stress was detected in virgin rats ($P = 0.0787$ versus control). Corticosterone concentration in lactating rats was equally high independently of the treatment, control or stress. For prolactin, a higher concentration was detected

Fig. 2. Western blot analysis of baseline kinase levels in unstressed female rats. We assessed differences in major tau kinases in female rat hippocampus under basal conditions (not stressed) in virgin (V), lactating (L) or weaned (W) ($N = 5$, each). Western blot analysis was performed using antibodies specific to glycogen synthase kinase (GSK3), extracellular signal-regulated kinase (ERK) and *c-Jun*-N-terminal kinase (JNK). We found that no stress (NS) lactating rats have significantly less hippocampal GSK3- α compared to both nonstressed virgin and weaned rats (*, $P = 0.016$). There were no significant differences in ERK or JNK. Data are presented as the mean $\pm$ SEM percentage change from the V condition.

Fig. 3. Glycogen synthase kinase (GSK3) levels in control and stressed lactating rats. Lactating rats were sacrificed after no stress (NS), or 20 min (20) or 24 h (24) ($N = 5$, each) after acute restraint stress. Twenty minutes after stress, lactating rats had significantly lower levels of total hippocampal GSK3- α compared to their controls ($P = 0.008$) and these low levels were sustained 24 h later (*, $P = 0.008$). However, 24 h after stress, active GSK3- β levels increased compared to control ($P = 0.016$), although active GSK3- α levels stayed relatively the same. There were no significant changes in inactive GSK3 (both α and β) levels. Data are presented as mean $\pm$ SEM percentage change from NS condition $\pm$ SEM.

Fig. 4. Corticosterone and prolactin plasma concentration in control and stressed rats. Basal corticosterone and prolactin levels were increased in lactating rats compared to virgin and weaned cohorts ($N = 5$, each) and were unchanged in response to restraint stress. Restraint-virgin rats showed a slight increase in corticosterone, although this was not significantly different from control. Values represent the mean $\pm$ SEM; Corticosterone: *** $P < 0.0001$, * $P = 0.0377$ versus lactating control; ++ $P = 0.0092$ versus virgin stress. Prolactin: ** $P < 0.001$ versus virgin and weaned controls. Data are presented as the mean $\pm$ SEM percentage change.

in lactating animals, and no change was induced by stress ($F_{1,24} = 1.57$, $P = 0.22$) (Fig. 4).

Distribution of tau-P in the female rat hippocampus

Immunohistochemical staining of the hippocampi of virgin and lactating rats was used to assess qualitative differences in tau-P (PHF-1 epitope) intensity and localisation in nonstressed controls (not shown) and in animals 20 min after stress. Figure 5 shows representative micrographs of the CA3 region of the hippocampus demonstrating a decrease in somatic tau-P (possibly a shift from soma to dendrites/axons) in lactating rats exposed to restraint

stress compared to virgin cohorts. The overall decrease in tau-P in stressed lactating rats supports our biochemical data.

Discussion

Stress-induced dephosphorylation of tau in the hippocampus of lactating rats

Although previous studies demonstrate significant increases in hippocampal tau-P after both acute and chronic restraint stress in male mice (13,14), the present study reveals a stress-induced dephosphorylation of tau in the hippocampus of lactating rats, a change that is not observed in virgin or one-month old weaned rats. A number of studies have indicated that females respond differently to stress compared to males, that the hormones involved in lactation help to confer a higher tolerance to stress regulated by the HPA axis, and that these hormones grant neuroprotection that extends to many regions of the brain including the hippocampus (4,17). Our current tau kinase data suggest that a significant decrease in both the α and β isoforms of total GSK3 may play a role in the stress-induced decrease in tau-P observed in lactating rats exposed to restraint stress. Because maternal hormones are known to play a role in stress response during pregnancy and lactation, it may be possible that one or more of these hormones are responsible for the alteration in total GSK3 levels.

Maternal hormones involved in lactation and stress

During lactation, the maternal brain shows a significantly reduced response to stress, a phenomenon that is possibly a result of chronically elevated corticosterone levels (1). Basal plasma levels of

Fig. 5. Localisation of tau phosphorylation (tau-P) in the CA3 region of stressed virgin or lactating rat hippocampus. Thirty micrometer thick sections of the hippocampus were stained with PHF-1 to visualise the localisation of tau-P in stressed virgin and lactating rats 20 min after stress ($N = 5$, each). This $\times 20$ magnification of CA3 pyramidal neurons demonstrates a lack of increase in somatic staining in stressed lactating animals (right) compared to stressed virgin animals (left). Unstressed controls (not shown) had no appreciable tau-P signal. Stressed lactating animals did not significantly differ in tau-P levels compared to unstressed controls ($P > 0.05$). Scale bar = 100 μ m.

corticosterone were higher in lactating cohorts than in virgin and weaned rats. No significant increase was detected in response to stress, although virgin rats exhibited a trend toward increase. This lack of statistical change may be related to the timing of plasma sampling, which was performed at sacrifice (50 min post-stress). A number of other hormones, such as prolactin, oxytocin and ovarian hormones, show increased expression during either pregnancy or lactation, and may play crucial roles in the attenuated stress response that is observed.

Prolactin is involved in milk production and is primarily produced in the anterior pituitary during lactation, although prolactin receptors have been previously documented in the brain (26). Prolactin is involved in maternal behaviour, the sleep–wake cycle, grooming, sexual behaviour and stress coping (26). Animals treated with prolactin demonstrate an attenuated HPA-axis response to stress, although the mechanism by which prolactin is involved in the overall anxiolytic effect is not fully understood (27). In the present study, we found a stress-induced dephosphorylation of tau in conjunction with a decrease in both the α and β isoforms of total GSK3 activity. Furthermore, previous studies have documented relationships between a hyperprolactinaemic stage and this tau kinase. One study suggests that GSK3 can work as a prolactin receptor kinase that tags the receptor for degradation, whereas another describes a pathway that involves prolactin acting to increase Akt activity, which in turn inhibits GSK3 activity (28, 29). It may be that increased prolactin levels during lactation result in decreased GSK3 activity (via an Akt pathway) to preserve prolactin receptors during lactation (a critical time period) and could perhaps be one explanation for the decreased baseline levels of GSK3 in lactating rats. However, additional studies are required to determine how stress-induced prolactin activity could cause a sufficiently large decrease in GSK3 activity to cause a significant dephosphorylation of hippocampal tau.

Oxytocin, another hormone present at elevated levels during lactation, facilitates social behaviour, maternal bonding and milk ejection, and is known to decrease the corticosterone stress response, decrease glucocorticoid receptor expression, stimulate hippocampal neurogenesis and modulate behavioural responses to stress (30). In the hippocampus, oxytocin has been shown to influence a number of signaling molecules, to modulate synaptic transmission, and to increase adult hippocampal neurogenesis, even under stressful conditions (31–33). Because oxytocin has such a large influence in the hippocampus during stress and lactation, further work is needed to determine whether it plays a role in the stress-induced decrease in hippocampal GSK3 and tau-P seen in lactating rats.

The ovarian hormone progesterone regulates reproduction, the release of neurotransmitters and neuronal development. Progesterone is present at constant plasma concentrations throughout pregnancy and lactation (until levels drop on postpartum day 18) and has been shown to have a neuroprotective effect in the nervous system (34). Progesterone administration has been shown to decrease tau-P and increase the dephosphorylated form of tau in the hippocampus of ovariectomised rats but does not affect GSK3 phosphorylation (35). Another study found that chronic proges-

terone treatment increased total hippocampal tau in ovariectomised rats, although it did not address tau-P (36). Interestingly, progesterone increases tau-P at the PHF-1 epitope in rat cerebellum associated with an increase in GSK3 activity, and decreases PP2A activity in the hippocampus of ovariectomised rats (34,37). These findings suggest a role for progesterone in regulating tau-P and associated kinases; however, these apparently conflicting studies demonstrate the need for further research on the specific mechanisms by which progesterone regulates these proteins.

The lactation period in the mother has been well established as neuroprotective (16), and previous studies have shown that GSK3 inhibition increases neuroprotection by inhibiting β -catenin and tau-P, modulating apoptotic pathways, promoting neurogenesis and improving cognitive deficits in models of neurodegenerative diseases such as AD and Parkinson's disease (38,39). Our understanding of the role of GSK3 in the cell is continually expanding and broadening, and it has been established that it not only regulates glycogen synthase activity, but also plays a role in a large number of other processes in the cell. It is possible that one of the mechanisms by which lactation confers neuroprotection to the mother is by inhibiting GSK3 activity (via regulation by some maternal/ovarian hormone), thereby inhibiting stress-induced tau-P and apoptosis, and promoting other neuroprotective mechanisms such as neurogenesis.

Hormones play a complex role in the female brain throughout adolescence, motherhood and senescence, including menopause. Although hormones such as prolactin and oxytocin are important for lactation, their actions in the brain and throughout the body have a much larger impact than just milk production and ejection. Maternal and ovarian hormones are known to have significant anxiolytic effects in response to stress and, in some cases, to increase adult neurogenesis. Furthermore, lactation is considered to comprise an enriched environment as a result of exteroceptive stimuli from the pups, such as olfactory, auditory, tactile and visual signals, in addition to the neurogenic stimulation provided by suckling. The neuroplasticity of the female brain also changes as the female goes through different reproductive stages, and tau is considered to play a role in the changes in brain circuitry that occur as part of this plasticity. The dephosphorylation of tau seen in stressed lactating rats may be a neuroprotective effect driven by elevated hormone levels and signalling by pups. Functional adaptations in the female brain during lactation include many forms of stress-attenuation both behaviourally and molecularly, although the extent to which these changes affect the future development of neurological diseases such as AD remains unclear. Some studies suggest that pregnancy is a risk factor for AD because of decreased levels of oestrogen compared to nulliparous women (40, 41). It may be that the neuroprotection conferred by lactation only lasts as long as the lactation period itself, as suggested by the data from our weaned animals. However, further research including studies focused on chronic stress, multiple maternities, and how these variables affect behaviour and neuropathological markers, such as neurofibrillary tangles and A β plaque formation, are needed to better our understanding of the female brain, the stress response and how these specific circuitries

play a role in the development of neurodegenerative diseases such as AD.

Acknowledgements

This work was supported by a grant from UC MEXUS – CONACYT to RAR and TM, as well as NIH AG032755, AG047484, AG010483 and the Alzheimer's Association (RAR) and UNAM-DGAPA-PAPIIT IN202315 (TM). We thank Dr Michael C. Donohue for statistical assistance; Louise Monte for assistance with manuscript preparation; Martha Carranza for assistance with the ELISA; and Dr Martín García and Dr Alejandra Castilla for animal care assistance. Importantly, we thank Dr Peter Davies (Albert Einstein) for donating antibody PHF-1.

Received 24 February 2015,
revised 24 August 2015,
accepted 2 September 2015

References

- Lightman SL, Windle RJ, Wood SA, Kershaw YM, Shanks N, Ingram CD. Peripartum plasticity within the hypothalamo-pituitary-adrenal axis. *Prog Brain Res* 2001; **133**: 111–129.
- Kinsley CH, Trainer R, Stafisso-Sandoz G, Quadros P, Marcus LK, Hearon C, Meyer EA, Hester N, Morgan M, Kozub FJ, Lambert KG. Motherhood and the hormones of pregnancy modify concentrations of hippocampal neuronal dendritic spines. *Horm Behav* 2006; **49**: 131–142.
- Pawluski JL, Galea LA. Hippocampal morphology is differentially affected by reproductive experience in the mother. *J Neurobiol* 2006; **66**: 71–81.
- Cabrera V, Ramos E, González-Arenas A, Cerbón M, Camacho-Arroyo I, Morales T. Lactation reduces glial activation induced by excitotoxicity in the rat hippocampus. *J Neuroendocrinol* 2013; **25**: 519–527.
- Swaab DF, Bao AM, Lucassen PJ. The stress system in the human brain in depression and neurodegeneration. *Ageing Res Rev* 2005; **4**: 141–194.
- McEwen BS. Sex, stress and the hippocampus: allostasis, allostatic load and the aging process. *Neurobiol Aging* 2002; **23**: 921–939.
- Carroll JC, Iba M, Bangasser DA, Valentino RJ, James MJ, Brunden KR, Lee VM, Trojanowski JQ. Chronic stress exacerbates tau pathology, neurodegeneration, and cognitive performance through a corticotropin-releasing factor receptor-dependent mechanism in a transgenic mouse model of tauopathy. *J Neurosci* 2011; **31**: 14436–14449.
- Korneyev AY. Stress-induced tau phosphorylation in mouse strains with different brain ERK 1 + 2 immunoreactivity. *Neurochem Res* 1998; **23**: 1539–1543.
- Feng Q, Cheng B, Yang R, Sun FY, Zhu CQ. Dynamic changes of phosphorylated tau in mouse hippocampus after cold water stress. *Neurosci Lett* 2005; **388**: 13–16.
- Wilson RS, Arnold SE, Schneider JA, Kelly JF, Tang Y, Bennett DA. Chronic psychological distress and risk of Alzheimer's disease in old age. *Neuroepidemiology* 2006; **27**: 143–153.
- Sotiropoulos I, Catania C, Pinto LG, Silva R, Pollerberg GE, Takashima A, Sousa N, Almeida OFX. Stress acts cumulatively to precipitate Alzheimer's Disease-like tau pathology and cognitive deficits. *J Neurosci* 2011; **31**: 7840–7847.
- Johansson L, Guo X, Hällström T, Norton MC, Waern M, Ostling S, Bengtsson C, Skoog I. Common psychosocial stressors in middle aged women related to longstanding distress and increased risk of Alzheimer's disease: a 38-year longitudinal population study. *BMJ Open* 2013; **2013**: e003142.
- Rissman RA, Lee K, Vale W, Sawchenko PE. Corticotropin-releasing factor receptors differentially regulate stress-induced tau phosphorylation. *J Neurosci* 2007; **27**: 6552–6562.
- Rissman RA, Staup MA, Lee AR, Justice NJ, Rice KC, Vale W, Sawchenko PE. Corticotropin-releasing factor receptor-dependent effects of repeated stress on tau phosphorylation, solubility, and aggregation. *Proc Natl Acad Sci USA* 2012; **109**: 6277–6282.
- Campbell SN, Zhang C, Monte L, Roe AD, Rice KC, Tache Y, Masliah E, Rissman R. Increased tau phosphorylation and aggregation in the hippocampus of mice overexpressing corticotropin-releasing factor. *J Alzheimers Dis* 2015; **43**: 967–976.
- Morales T. Recent findings on neuroprotection against excitotoxicity in the hippocampus of female rats. *J Neuroendocrinol* 2011; **23**: 994–1001.
- Slattery D, Neumann I. No stress please! Mechanisms of stress hyporesponsiveness of the maternal brain. *J Physiol* 2008; **586**: 377–385.
- Furuta M, Bridges RS. Gestation-induced cell proliferation in the rat brain. *Brain Res Dev Brain Res* 2005; **156**: 61–66.
- Leuner B, Mirescu C, Noiman L, Gould E. Maternal experience inhibits the production of immature neurons in the hippocampus during the postpartum period through elevations in adrenal steroids. *Hippocampus* 2007; **17**: 434–442.
- Gatewood JD, Morgan MD, Eaton M, McNamara IM, Stevens LF, Macbeth AH, Meyer EA, Lomas LM, Kozub FJ, Lambert KG, Kinsley CH. Motherhood mitigates aging-related decrements in learning and memory and positively affects brain aging in the rat. *Brain Res Bull* 2005; **66**: 91–98.
- Torner L, Toschi N, Pohlner A, Landgraf R, Neumann ID. Anxiolytic and anti-stress effects of brain prolactin: improved efficacy of anti-sense targeting of the prolactin receptor by molecular modeling. *J Neurosci* 2001; **21**: 3207–3214.
- Blume A, Torner L, Liu Y, Subburaju S, Aguilera G, Neumann ID. Prolactin activates mitogen-activated protein kinase signaling and corticotropin releasing hormone transcription in rat hypothalamic neurons. *Endocrinology* 2008; **150**: 1841–1849.
- Morales T, Lorenson M, Walker AM, Ramos E. Both prolactin (PRL) and a molecular mimic of phosphorylated PRL, S179D-PRL, protect the hippocampus of female rats against excitotoxicity. *Neuroscience* 2014; **258**: 211–217.
- Carroll JC, Rosario ER, Chang L, Stanczyk FZ, Oddo S, LaFerla FM, Pike CJ. Progesterone and estrogen regulate Alzheimer-like neuropathology in Female 3xTg-AD mice. *J Neurosci* 2007; **27**: 13357–13365.
- Roe AD, Staup MA, Serrats J, Sawchenko PE, Rissman RA. Lipopolysaccharide-induced tau phosphorylation and kinase activity-modulation, but not mediation, by corticotropin-releasing factor receptors. *Eur J Neurosci* 2011; **34**: 448–456.
- Ben-Jonathan N, Mershon JL, Allen DL, Steinmetz RW. Extrahypothalamic prolactin: distribution, regulation, functions, and clinical aspects. *Endocr Rev* 1996; **17**: 639–669.
- Torner L, Neumann ID. The Brain Prolactin System: involvement in stress response adaptations in lactation. *Stress* 2002; **5**: 249–257.
- Dominguez-Caceres MA, Garcia-Martinez JM, Calcabrini A, Gonzalez L, Porque PG, Leon J, Martin-Perez J. Prolactin induces c-Myc expression and cell survival through activation of Src/Akt pathway in lymphoid cells. *Oncogene* 2004; **23**: 7378–7390.
- Plotnikov A, Li Y, Tran TH, Tang W, Palazzo JP, Rui H, Fuchs SY. Oncogene-mediated inhibition of glycogen synthase kinase 3-B impairs degradation of prolactin receptor. *Cancer Res* 2008; **68**: 1354–1361.
- Cohen H, Kaplan Z, Kozlovsky N, Gidron Y, Matar MA, Zohar J. Hippocampal Microinfusion of oxytocin attenuates the behavioral response to stress by means of dynamic interplay with the glucocorticoid-catecholamine responses. *J Neuroendol* 2010; **22**: 889–904.
- Petersson M, Uvnäs-Moberg K. Systemic oxytocin treatment modulate glucocorticoid and mineralocorticoid receptor mRNA in the rat hippocampus. *Neurosci Lett* 2003; **343**: 97–100.
- Zaninetti M, Raggenbass M. Oxytocin receptor agonists enhance inhibitory synaptic transmission in the rat hippocampus by activating

- interneurons in stratum pyramidale. *Eur J Neurosci* 2002; **12**: 3975–3984.
- 33 Leuner B, Caponiti JM, Gould E. Oxytocin stimulates adult neurogenesis even under conditions of stress and elevated glucocorticoids. *Hippocampus* 2012; **22**: 861–868.
  - 34 Amorim MA, Guerra-Araiza C, Pernia O, da Cruz e Silva EF, Garcia-Segura ML. Progesterone regulates the phosphorylation of protein phosphatases in the brain. *J Neuro Res* 2010; **88**: 2826–2832.
  - 35 Pinto-Almazan R, Calzada-Mendoza CC, Campos-Lara MG, Guerra-Araiza C. Effect of chronic administration of estradiol, progesterone, and tibolone on the expression and phosphorylation of glycogen synthase kinase-3 β and the microtubule-associated protein tau in the hippocampus and cerebellum of the female rat. *J Neuro Res* 2012; **90**: 878–886.
  - 36 Camacho-Arroyo I, Gonzalez-Arena A, Espinosa-Raya J, Pina-Medina AG, Picazo O. Short- and long-term treatment with estradiol or progesterone modifies the expression of GFAP, MAP2 and Tau in prefrontal cortex and hippocampus. *Life Sci* 2011; **89**: 123–128.
  - 37 Guerra-Araiza C, Amorim MA, Camacho-Arroyo I, Garcia-Segura LM. Effects of progesterone and its reduced metabolites, dihydroprogesterone and tetrahydroprogesterone, on the expression and phosphorylation of glycogen synthase kinase-3 and the microtubule-associated protein tau in the rat cerebellum. *Dev Neurobiol* 2007; **67**: 510–520.
  - 38 Culbert AA, Brown MJ, Frame S, Hagen T, Cross DAE, Bax B, Reith AD. GSK-3 inhibition by adenoviral FRAT1 overexpression is neuroprotective and induces tau dephosphorylation and B-catenin stabilisation without elevation of glycogen synthase activity. *FEBS Lett* 2001; **507**: 288–294.
  - 39 King MK, Pardo M, Cheng Y, Downey K, Jope RS, Beurel E. Glycogen synthase kinase-3 inhibitors: rescuers of cognitive impairments. *Pharmacol Ther* 2014; **141**: 1–12.
  - 40 Colucci M, Cammarata S, Assini A, Croce R, Clerici F, Novello C, Mazzella L, Dagnino N, Mariani C, Tanganelli P. The number of pregnancies is a risk factor for Alzheimer's disease. *Eur J Neurol* 2006; **13**: 1374–1377.
  - 41 Ptak U, Barkow K, Heun R. Fertility and number of children in patients with Alzheimer's disease. *Arch Womens Ment Health* 2002; **5**: 83–86.