

UC San Diego

UC San Diego Previously Published Works

Title

Metagenome Sequencing of the Microbial Community of a Solar Saltern Crystallizer Pond at Cahuil Lagoon, Chile

Permalink

<https://escholarship.org/uc/item/3tf394qm>

Journal

Microbiology Resource Announcements, 2(6)

ISSN

2576-098X

Authors

Plominsky, Alvaro M
Delherbe, Nathalie
Ugalde, Juan A
et al.

Publication Date

2014-12-24

DOI

10.1128/genomea.01172-14

Peer reviewed

Metagenome Sequencing of the Microbial Community of a Solar Saltern Crystallizer Pond at C ahuil Lagoon, Chile

Alvaro M. Plominsky,^a Nathalie Delherbe,^a Juan A. Ugalde,^b Eric E. Allen,^{c,d} Marine Blanchet,^e Priscila Ikeda,^f Francisco Santib a ez,^g Kurt Hanselmann,^h Osvaldo Ulloa,^{g,i} Rodrigo De la Iglesia,^a Peter von Dassow,^a Marcia Astorga,^g Mar a Jes us G alvez,^g Mar a Lorena Gonz alez,^g Carlos Henr iquez-Castillo,^a Daniel Vaulot,^{j,k} Adriana Lopes do Santos,^l Gerrit van den Engh,^m Carla Gimpel,ⁿ Florencia Bertoglio,^o Yolaine Delgado,^p Felipe Docmac,^q Claudia Elizondo-Patrone,^r Silvia Narv ez,^s Fernando Sorroche,^t Marcelo Rojas-Herrera,^b Nicole Trefault^b

Facultad de Ciencias Biol gicas, Pontificia Universidad Cat lica de Chile, Santiago, Chile^a; Facultad de Ciencias, Universidad Mayor, Santiago, Chile^b; Marine Biology Research Division, Scripps Institution of Oceanography, University of California San Diego, San Diego, California, USA^c; Division of Biological Sciences, University of California San Diego, San Diego, California, USA^d; Biogeochemistry and Microbiology, Universit  Pierre et Marie Curie, Paris, France^e; Oceanographic Institute, University of Sao Paulo, S o Paulo, Brazil^f; Departamento de Oceanograf a, Universidad de Concepci n, Concepci n, Chile^g; Department of Earth Sciences, Climate Geology, Geomicrobiology, ETH Z rich, Z rich, Switzerland^h; Instituto Milenio de Oceanograf a, Universidad de Concepci n, Concepci n, Chileⁱ; Sorbonne Universit s, UPMC Universit  Paris 06, UMR 7144, Station Biologique, Roscoff, France^j; CNRS, UMR 7144, Station Biologique, Roscoff, France^k; Federal University of Rio de Janeiro, Rio de Janeiro, Brazil^l; BD Advanced Cytometry Group, Seattle, Washington, USA^m; Departamento Cient fico, Instituto Ant rtico Chileno, Punta Arenas, Chileⁿ; Centro Universitario Regional Este, Universidad de la Rep blica, Rocha, Uruguay^o; Departamento de Microbiolog a, Universidad de La Habana, La Habana, Cuba^p; Facultad de Ciencias del Mar y Recursos Biol gicos, Universidad de Antofagasta, Antofagasta, Chile^q; Departamento de Ecolog a y Recursos Naturales, Universidad Andr s Bello, Santiago, Chile^r; Instituto de Investigaciones Marinas, INVEMAR, Santa Marta, Colombia^s; Facultad de Ciencias Exactas, F sico-Qu micas y Naturales, Universidad Nacional de R o Cuarto, C rdoba, Argentina^t

C ahuil Lagoon in central Chile harbors distinct microbial communities in various solar salterns that are arranged as interconnected ponds with increasing salt concentrations. Here, we report the metagenome of the 3.0- to 0.2- m fraction of the microbial community present in a crystallizer pond with 34% salinity.

Received 1 October 2014 Accepted 6 October 2014 Published 13 November 2014

Citation Plominsky AM, Delherbe N, Ugalde JA, Allen EE, Blanchet M, Ikeda P, Santib a ez F, Hanselmann K, Ulloa O, De la Iglesia R, von Dassow P, Astorga M, G alvez MJ, Gonz alez ML, Henr iquez-Castillo C, Vaulot D, Lopes do Santos A, van den Engh G, Gimpel C, Bertoglio F, Delgado Y, Docmac F, Elizondo-Patrone C, Narv ez S, Sorroche F, Rojas-Herrera M, Trefault N. 2014. Metagenome sequencing of the microbial community of a solar saltern crystallizer pond at C ahuil Lagoon, Chile. *Genome Announc.* 2(6):e01172-14. doi:10.1128/genomeA.01172-14.

Copyright   2014 Plominsky et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Nicole Trefault, nicole.trefault@umayor.cl.

Solar salterns represent unique extreme microbial ecosystems that are best studied by applying different molecular techniques (1). These habitats offer relatively low microbial diversity and are generally dominated by archaea (2). In this study, we generated an additional data set from a solar saltern crystallizer pond in C ahuil Lagoon (Chile). We will continue to use the data for comparative studies on the metabolic potential and diversity of microorganisms inhabiting an environment with 10-fold seawater salt concentration.

C ahuil Lagoon extends several miles into mainland from the coast of the South Pacific, where seawater mixes with various freshwater tributaries. Along the border of the lagoon, local inhabitants build a series of locks to distribute brackish water into a series of interconnected ponds where water evaporates until the ion activity products of the dissolved ions exceed the solubility product of halite.

Samples for this study were collected on 12 January 2012 from a final salt crystallizer pond (34 30'10"S 71 59'09"W), which had a salinity of 34.1%, a temperature of 41.2 C, and a pH of 6.4. A volume of 5 liters was first passed through a 60- m nylon mesh, and then 600 ml was sequentially filtered through 20- m, 3.0- m (polycarbonate; Millipore), and 0.22- m-pore filters (polyethersulfone; Millipore). Community DNA was extracted from the 3.0- to 0.22- m size fraction using the

cetyltrimethylammonium bromide (CTAB) method (3) and sequenced by pyrosequencing (Roche 454 GS-FLX system, Titanium chemistry) at the Center for Genomics and Bioinformatics, Universidad Mayor, Chile.

The raw 222,074 sequencing reads (average length, 450 bp) were quality filtered for ambiguities and homopolymers (≥ 10), and reads with $>75\%$ of the sequence having a Phred quality score of ≤ 20 were removed. Taxonomic assignment (MGTXA [<http://mgtaxa.jcvi.org>] against NCBI RefSeq v01-07-2012) of the remaining 183,299 reads revealed that 61% stem from archaea, 19% from viruses, and 16% from bacteria. At the phylum level, 58% of the reads were assigned to *Euryarchaeota* (all from the family *Halobacteriaceae*), 9% to *Actinobacteria*, and 7% to the virus class *Caudovirales*. Several genera of the family *Halobacteriaceae* share similar levels of representation (expressed as the % of reads assigned to *Archaea*): *Haloarcula* (14%), *Haloquadratum* (11.5%), *Halorubrum* (9.5%), *Halobacterium* (9%), *Natronomonas* (9%), *Halogeometricum* (8.5%), *Halomicrobium* (6.5%), and *Halorhabdus* (6%). Although *Haloquadratum* is still an important part of the microbial community, it is not as abundant in this crystallizer pond as it is in the Santa Pola saltern (1, 4).

The data generated as part of this metagenome add valuable information for ongoing and future studies on the metabolic po-

tential of the microbial communities associated with extreme hypersaline habitats worldwide.

Nucleotide sequence accession number. This metagenome project has been deposited in the NCBI Sequence Read Archive under the accession no. [SRX680116](https://www.ncbi.nlm.nih.gov/sra/SRX680116).

ACKNOWLEDGMENTS

This study was performed during the Ecology and Diversity of Marine microorganisms (ECODIM) VII course as part of the 12th Austral Summer Institute organized by the Universidad de Concepción (UdeC) in 2012. The course was held under the auspices of the UdeC and was supported through access to research facilities at the Marine Station of the Pontificia Universidad Católica de Chile (PUC) in Las Cruces.

Funding for the sampling and sequencing was provided by the Agou-ron Institute, the Gordon and Betty Moore Foundation, the Graduate School of UdeC, the Faculty of Biological Sciences of PUC, and the Center for Genomics and Bioinformatics of the Universidad Mayor. The sequence analyses were performed by A.M.P., N.D., and J.A.U. at E.E.A.'s

Laboratory at the Scripps Institution of Oceanography, UCSD, with the support of a MECESUP-PUC 2012 internship grant to A.M.P.

REFERENCES

1. Ghai R, Pašić L, Fernandez AB, Martin-Cuadrado AB, Mizuno CM, McMahon KD, Papke RT, Stepanauskas R, Rodriguez-Brito B, Rohwer F, Sanchez-Porro C, Ventosa A, Rodriguez-Valera F. 2011. New abundant microbial groups in aquatic hypersaline environments. *Sci. Rep.* 1:135. <http://dx.doi.org/10.1038/srep00135>.
2. Oren A. 2002. Molecular ecology of extremely halophilic *Archaea* and *Bacteria*. *FEMS Microbiol. Ecol.* 39:1–7. <http://dx.doi.org/10.1111/j.1574-6941.2002.tb00900.x>.
3. Wilson K. 1990. Preparation of genomic DNA from bacteria, p 2.4.1–2.4.5. *In* Ausubel F, Brent R, Kingston R, Seidman J, Smith J, Struhl K (ed), *Current protocols in molecular biology*. Wiley Interscience, New York, NY.
4. Fernandez AB, Ghai R, Martin-Cuadrado AB, Sanchez-Porro C, Rodriguez-Valera F, Ventosa A. 2013. Metagenome sequencing of prokaryotic microbiota from two hypersaline ponds of a marine saltern in Santa Pola, Spain. *Genome Announc.* 1(6):e00933-13. <http://dx.doi.org/10.1128/genomeA.00933-13>.