

## **UC Davis**

### **UC Davis Previously Published Works**

#### **Title**

Improving the translation of animal ischemic stroke studies to humans

#### **Permalink**

<https://escholarship.org/uc/item/3th0p3mv>

#### **Journal**

Metabolic Brain Disease, 30(2)

#### **ISSN**

0885-7490

#### **Authors**

Jickling, Glen C

Sharp, Frank R

#### **Publication Date**

2015-04-01

#### **DOI**

10.1007/s11011-014-9499-2

Peer reviewed



# HHS Public Access

Author manuscript

*Metab Brain Dis.* Author manuscript; available in PMC 2016 April 01.

Published in final edited form as:

*Metab Brain Dis.* 2015 April ; 30(2): 461–467. doi:10.1007/s11011-014-9499-2.

## Improving the Translation of Animal Ischemic Stroke Studies to Humans

Glen C Jickling and Frank R Sharp

Department of Neurology and MIND Institute, University of California at Davis

### Abstract

Despite testing more than 1026 therapeutic strategies in models ischemic stroke and 114 therapies in human ischemic stroke, only one agent tissue plasminogen activator has successfully been translated to clinical practice as a treatment for acute stroke. Though disappointing, this immense body of work has led to a rethinking of animal stroke models and how to better translate therapies to patients with ischemic stroke. Several recommendations have been made, including the STAIR recommendations and statements of RIGOR from the NIH / NINDS. In this commentary we discuss additional aspects that may be important to improve the translational success of ischemic stroke therapies. These include use of tissue plasminogen activator in animal studies; modeling ischemic stroke heterogeneity in terms of infarct size and cause of human stroke; addressing the confounding effect of anesthesia; use of comparable therapeutic dosage between humans and animals based on biological effect; modeling the human immune system; and developing outcome measures in animals comparable to those used in human stroke trials. With additional study and improved animal modeling of factors involved in human ischemic stroke, we are optimistic that new therapies for the treatment of acute ischemic stroke will be developed.

### Keywords

Stroke; Animal Models; Clinical Trial; Cerebrovascular disease

### Introduction

The translation of acute ischemic stroke therapies from animals to patients with ischemic stroke has been challenging. With over 1026 agents tested in models of ischemic stroke and 114 tested in humans, only one medical therapy, tissue plasminogen activator (tPA), has received approval by the FDA for the acute treatment of ischemic stroke (O'Collins et al. 2006)(Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group 1995). The reasons for this failure to translate has been the subject of numerous meetings (STAIR recommendation) and reviews (Stroke Therapy Academic Industry 2001; Fisher and Stroke Therapy Academic Industry 2003; Fisher et al. 2005; Fisher et al. 2007; Fisher et al. 2009; Saver et al. 2009;

---

Correspondence to: Glen C Jickling MD, Department of Neurology, MIND Institute Wet Labs Room 2415, University of California at Davis Medical Center, 2805 50<sup>th</sup> Street, Sacramento, CA 95817. Phone: 916-703-0389; FAX: 916-703-0369; [gcjickling@ucdavis.edu](mailto:gcjickling@ucdavis.edu).

Albers et al. 2011; Lapchak et al. 2013; Feuerstein et al. 2008). In addition, the NIH/NINDS have published criteria regarding the rigor of animal stroke studies (Landis et al. 2012; Lapchak et al. 2013). Proposed recommendations to improve the translation of acute stroke therapies have included evaluation of agents in the therapeutic time window of ischemic stroke, need to assess dose response, need to assess aged animals of both sexes with vascular risk factors, therapy not acting on the intended therapeutic target, importance of randomization and blinding, defining inclusion and exclusion criteria, ensuring studies are adequately powered, replication of studies by independent groups, declaration of conflicts of interests, and need to evaluate therapies in multiple animal models of stroke including gyrencephalic species.

The goal of the current commentary is not to review prior recommendations with which we agree, but to discuss additional aspects that may be important to the translational success of ischemic stroke therapies. Frequently many of the aspects incorporated in the design of a human stroke trial are not incorporated into animal studies of acute ischemic stroke. However, modeling features of a clinical stroke trial in animals may be essential to determine whether a therapy will ultimately translate to human ischemic stroke. Presented below is a discussion of several factors that may advance preclinical stroke models, including use of tissue plasminogen activator, modeling ischemic stroke heterogeneity in terms of infarct size and cause of human stroke, addressing the confounding effect of anesthesia, dosing therapeutics based on biological effect, modeling the human immune system, and developing methods to assess animal stroke outcomes similar to measures used in human stroke trials.

## Implications of tPA for Animal Studies

tPA is standard of care for patients acute ischemic stroke (Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group 1995). Thus, in human clinical trials evaluating new treatments for acute ischemic stroke, patients treated with tPA are likely to be enrolled. As a result, animal stroke models evaluating acute stroke therapies should also include a group of animals treated with tPA. The inclusion of a tPA treated animal group has several advantages. It permits evaluation of the interaction between the new treatment and tPA, including assessing the impact on thrombolytic activity and adverse events. In addition, tPA treated animals can be used as a positive control as discussed below.

Compounds being considered for clinical trials may benefit from having preclinical animal data comparing the compound of interest to tPA. The rationale is that tPA is the only drug where the observed beneficial effect in animal ischemic stroke has translated to a benefit in human ischemic stroke. If a test compound can demonstrate a beneficial effect similar to the effect of tPA in animals, then it may have an increased probability to be effective in humans. For example, a lead compound could be compared to tPA and vehicle control in a blood clot embolic model. To move forward, the test compound should be as effective as tPA in treated animals. The blood clot ischemic stroke model is suggested because of its prior success in translating tPA to stroke patients, and thrombosis/clot embolism is an important feature of all three human stroke subtypes (Hsia et al. 2003; Tissue plasminogen activator for acute

ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group 1995).

## Use of Anesthesia

In patients with ischemic stroke anesthesia is generally not used, and most strokes do not occur while a patient is anesthetized. However, in the majority of animal stroke models cerebral ischemia is induced while animals are under anesthesia. Though some studies have suggested anesthesia may not affect outcomes, this may depend on type of anesthetic and possibly the cause of stroke (Gelb et al. 2002; Head and Patel 2007; Kirsch et al. 1996; Koerner and Brambrink 2006; Wang et al. 2008; Zivin et al. 1985). Stroke models without anesthesia are needed to evaluate lead compounds being considered for clinical stroke trials. Alternatively, one might question whether anesthesia should be used in patients with acute ischemic stroke. Anesthesia may have neuroprotective effects and may be an important factor to help translate certain compounds with identified benefit in animal stroke models to patients (Gakuba et al. 2011). However, the routine use of anesthetics in patients with ischemic stroke has a number of limitations, including loss of neurological exam to monitor response to therapy, effects on blood pressure and cerebral perfusion, association with worse outcomes in endovascular treated patients, and need for intensive care that is not widely available in all centers treating patients with ischemic stroke (Froehler et al. 2012; Flexman et al. 2012; Davis et al. 2012; Abou-Chebl et al. 2010).

## Importance of Stroke Cause

Ischemic stroke in humans is heterogeneous. Clinical stroke trials generally enroll stroke patients of diverse age, gender, ethnicity, and importantly, stroke cause. The value of modeling age and gender in animal stroke studies has been addressed in the STAIR criteria. However, the importance of modeling stroke cause warrants further consideration, as it too may influence the translational success of an acute stroke therapy. Given most stroke trials include multiple causes of stroke, having preclinical data demonstrating effectiveness in multiple animal stroke models that emulate the major causes of human ischemic stroke may be important to translational success.

The three major causes of human ischemic stroke are large vessel atherosclerosis, cardioembolism, and lacunar small vessel disease. Each has a different pathogenesis. Large vessel stroke is due to atherosclerotic plaque rupture and embolization; cardioembolic stroke is due to cardiac clot formation and embolization; and lacunar stroke is due to a disease of the small penetrating cerebral vessels. It is not known if the mechanisms of brain injury differ between these three major causes of human stroke. However, different prognosis, mortality, and stroke prevention treatment highlight that differences among cardioembolic, large vessel, and lacunar stroke exist and potentially may influence acute ischemic stroke treatment. We have found molecular differences in inflammation and gene response among the three major causes of ischemic stroke in humans (Jickling et al. 2011; Jickling et al. 2010). This suggests that aspects of the immune response are different for each major cause of stroke. Though further study is required, it is possible that the molecular response to brain

ischemia and neuroinflammation may differ for each cause of stroke. Thus, certain acute stroke treatments may have greater therapeutic effect for certain causes of stroke.

If cause of stroke is important to the treatment of acute ischemic stroke, then animal studies should be performed in models of the three major causes of human stroke (large vessel atherosclerotic, cardioembolic, and small vessel lacunar). Currently most preclinical animal studies use a filament stroke model, which models cerebral ischemia well but does not replicate any of the three major causes of human stroke. This is emphasized by the fact that very few genes regulated in the blood of rats following filament induced stroke are regulated in the blood of human ischemic stroke (Tang et al. 2006; Tang et al. 2001; Stamova et al. 2010). Different patterns of infarct evolution have also been observed in embolic compared to filament stroke models in rodents (Henninger et al. 2006). Furthermore, Hossmann has argued that the filament model is a poor model of human stroke that should be abandoned to test therapies for human ischemic stroke because the pathophysiology of ischemic brain injury induced by mechanical vascular occlusion is different from the thromboembolic occlusion that occurs in humans (Hossmann 2012).

A therapy shown to be effective in the filament stroke model should have additional evaluation performed in stroke models that have greater similarity to the human causes of cerebral ischemia. The blood clot / embolic stroke model emulates aspects of human cardioembolic and arterial embolism. This model has been applied in rodents and rabbits, and as discussed below has the benefit of producing variable infarct size (Lapchak 2010, 2009; Meyer et al. 2013; Chapman et al. 2001). Large vessel atherosclerotic stroke has been modeled in apolipoprotein E deficient mice and in rabbits fed a high fat diet (Zhang et al. 2010; Daugherty et al. 2009; Verbeuren 2006; Russell and Proctor 2006; Yanni 2004). Small vessel ischemic lacunar stroke has been modeled using the Spontaneous Hypertensive Rat (SHR), though additional models of cerebral small vessel disease are needed (Bailey et al. 2011a; Bailey et al. 2009; Bailey et al. 2011b; Yao and Nabika 2012; Hainsworth and Markus 2008). In both atherosclerotic and SHR stroke models, the occurrence of cerebral ischemia is spontaneous and requires monitoring for ischemic stroke occurrence. There are many other stroke models not mentioned here, including a photocoagulation model, the in situ thrombin administration model (Orset et al. 2007), and the ferric chloride application model (Karatas et al. 2011). The relationship of these models to three major causes of stroke in humans is requires further study.

### **Infarct Size, Is Heterogeneity Desirable in Animal Models**

Infarct size of patients enrolled in clinical stroke trials is usually quite variable, ranging from small infarcts causing minor disability to very large infarcts that result in marked disability and death (Lansberg et al. 2012; Lansberg et al. 2011; Kidwell et al. 2013). In contrast, in animal stroke studies an infarct of consistent size is usually sought as this facilitates determination of infarct size reduction in response to treatment (Takano et al. 1997; Belayev et al. 1996). Given the variability in human infarct size, animal stroke studies may better translate to patients if treatments are assessed across a range of infarct sizes. Such a design would require a larger numbers of animals and thus be more difficult and expensive. However, given the expense of human trials these larger animal studies that mimic clinical

stroke trials may be warranted. This might be considered over modeling but strokes of different volume may have different degrees of blood brain barrier damage, inflammatory response, and potentially therapeutic response. The clot stroke model is one model that does produce variable infarct size. In the rabbit blood clot model cerebral infarcts of different sizes have been used to construct treatment response curves to assess therapeutic effect (Lapchak et al. 2002b; Lapchak et al. 2002a).

## What is the appropriate behavioral outcome in animal stroke studies

In clinical studies of acute ischemic stroke the modified Rankin Scale (mRS) is used to assess functional status at 90 days following the index event (Huybrechts and Caro 2007; Banks and Marotta 2007). Frequently it is the primary outcome measure in clinical stroke trials, thus the features assessed by the mRS should also be important in animal studies of ischemic stroke (Kidwell et al. 2013; Hacke et al. 2008; Ginsberg et al. 2013; Broderick et al. 2013). The mRS is a simple six point scale that does not provide a comprehensive assessment of functional status but does provide an index of a person's disability related to the stroke (Fearon et al. 2012; Weisscher et al. 2008). This simplicity permits reasonable inter-rater reliability, which is essential in multicenter clinical trials (Banks and Marotta 2007; Quinn et al. 2009; Zhao et al. 2010). Furthermore, the mRS is validated and familiar to stroke investigators worldwide. Thus, the mRS is likely to remain an important outcome measure in clinical trials of ischemic stroke.

How the mRS scale relates to functional outcomes in animal stroke models remains unclear. A mRS score  $\leq 2$  is generally considered a good outcome; a score  $>2$  is a poor outcome with a score of 6 indicating death. The mRS evaluates disability, with a focus on ability to walk, attend to bodily needs, and carry out activities of daily life. These functions are a reflection of impairment in various aspects of the motor, sensory, visual and cognitive systems. Using behavioral measures in animal stroke models that are reflective of the mRS in humans may help improve therapeutic translation in stroke. Determining which features to assess in animal stroke models that are most reflective of features assessed by the mRS requires further investigation. In a mouse MCAO stroke model, Rosell et al. found that no specific behavioral test reliably assessed long-term functional outcome (Rosell et al. 2013). The optimal time to assess outcome in animal stroke also warrants consideration. Human trials often assess mRS at 90 days after the onset of ischemic stroke. Thus, treatment effects in animals should also be evaluated at time points beyond 24 hours. Given the shorter lifespan of rodents, a timeframe of 21-28 days has been suggested, though sustained benefit at longer time points may be desirable in some studies. Outcome assessment of these later time points permits identification of treatment effects that may be transient or decline with time and would thus be less relevant to the ultimate stroke outcome.

Another difference between human clinical trials and animal studies is the analysis of mortality as an outcome. Mortality is an important outcome in human studies of ischemic stroke. However, in experimental stroke studies animals that die are often excluded from analysis (Sicard and Fisher 2009). To improve preclinical stroke studies, mortality should be among the outcomes measured. Furthermore, stroke severity could be modulated so that animal mortality is similar to that expected in a human stroke trial.

## Immune Differences between Animal and Human Ischemic Stroke

The immune system is increasingly viewed as important in acute ischemic stroke. Humans and rodents have a number of differences in their immune systems that may be relevant to acute ischemic stroke and the evaluation of certain therapies. In humans, 70% of circulating leukocytes are neutrophils, whereas in rodents only 20-30% of leukocytes are neutrophils (Fox 1985). Thus, the baseline number and types of immune cells are very different. Immune responses to ischemic stroke may also differ between rodents and humans. We have found the circulating leukocytes have a very different gene expression profile in rodent MCAO stroke compared to that in patients with acute ischemic stroke (Stamova et al. 2010; Tang et al. 2001; Tang et al. 2006). Primates and rabbits have a leukocyte composition closer to that of humans. Potentially, these or other stroke models may be preferred in the evaluation of certain stroke therapies in which the immune system is deemed important.

## How to translate Dose from Animal Models to Humans

Knowing how to translate the dose of a therapy used in an animal stroke model to an effective dose in patients with ischemic stroke is challenging. A variety of methods and formulas have been developed to aid in this process (Reagan-Shaw et al. 2008; Singh 2006). Many of these methods do not address biological differences between rodents and humans that may be very important to drug delivery and activity on its therapeutic target.

One possible solution is to develop an assay to measure a drug's effect on its biological target. Dosage could then be based on target response and calibrated to achieve a similar response in humans as achieved in animal stroke models (Liou et al. 2012; Plenge et al. 2013). A surrogate blood biomarker of biological effect may be very useful in this regard. Drug levels may also be useful, though a drug level alone may not necessarily indicate biological activity. In some cases evaluation of cerebral spinal fluid may necessary if entry into the central nervous system is required for biological activity. Importantly, this should be performed in both animal and human studies, as a compound's ability to cross the blood brain barrier in rodent ischemic stroke does not necessarily imply it can cross the blood brain barrier of a patient with ischemic stroke.

## Conclusions

Animal stroke studies are performed for a variety of reasons. Many are proof of concept studies evaluating aspects of ischemic stroke biology or identifying treatment targets for further investigation. This is important work that is essential to better understand and develop novel approaches to treat stroke. However, given the many studies that have not translated to humans, we need to be mindful that promising results in proof of concept stroke studies require additional evaluation prior to human study. For the field to advance and translate more therapies to human stroke we need new ideas, stroke models, and approaches. In this commentary we have emphasized several differences between experimental animal stroke and human clinical stroke trials. Though the suggestions made may be contentious, discussion is essential to determine the factors and experimental



protocols needed to better model human ischemic stroke in animals and improve the translation of therapies to patient with ischemic stroke.

## Acknowledgements

This work was supported by the American Heart Association (GCJ) and the National Institutes of Health (NS075035, NS079153) (FRS). We appreciate the support of the MIND Institute and the UCD Department of Neurology.

## References

- Abou-Chebl A, Lin R, Hussain MS, Jovin TG, Levy EI, Liebeskind DS, Yoo AJ, Hsu DP, Rymer MM, Tayal AH, Zaidat OO, Natarajan SK, Nogueira RG, Nanda A, Tian M, Hao Q, Kalia JS, Nguyen TN, Chen M, Gupta R. Conscious sedation versus general anesthesia during endovascular therapy for acute anterior circulation stroke: preliminary results from a retrospective, multicenter study. *Stroke; a journal of cerebral circulation*. 2010; 41(6):1175–1179. doi:10.1161/STROKEAHA.109.574129.
- Albers GW, Goldstein LB, Hess DC, Wechsler LR, Furie KL, Gorelick PB, Hurn P, Liebeskind DS, Nogueira RG, Saver JL, Consortium SV. Stroke Treatment Academic Industry Roundtable (STAIR) recommendations for maximizing the use of intravenous thrombolytics and expanding treatment options with intra-arterial and neuroprotective therapies. *Stroke; a journal of cerebral circulation*. 2011; 42(9):2645–2650. doi:10.1161/STROKEAHA.111.618850.
- Bailey EL, McCulloch J, Sudlow C, Wardlaw JM. Potential animal models of lacunar stroke: a systematic review. *Stroke; a journal of cerebral circulation*. 2009; 40(6):e451–458. doi:10.1161/STROKEAHA.108.528430.
- Bailey EL, Smith C, Sudlow CL, Wardlaw JM. Is the spontaneously hypertensive stroke prone rat a pertinent model of sub cortical ischemic stroke? A systematic review. *International journal of stroke : official journal of the International Stroke Society*. 2011a; 6(5):434–444. doi:10.1111/j.1747-4949.2011.00659.x. [PubMed: 21951409]
- Bailey EL, Wardlaw JM, Graham D, Dominiczak AF, Sudlow CL, Smith C. Cerebral small vessel endothelial structural changes predate hypertension in stroke-prone spontaneously hypertensive rats: a blinded, controlled immunohistochemical study of 5- to 21-week-old rats. *Neuropathology and applied neurobiology*. 2011b; 37(7):711–726. doi:10.1111/j.1365-2990.2011.01170.x. [PubMed: 21392049]
- Banks JL, Marotta CA. Outcomes validity and reliability of the modified Rankin scale: implications for stroke clinical trials: a literature review and synthesis. *Stroke; a journal of cerebral circulation*. 2007; 38(3):1091–1096. doi:10.1161/01.STR.0000258355.23810.c6.
- Belayev L, Alonso OF, Busto R, Zhao W, Ginsberg MD. Middle cerebral artery occlusion in the rat by intraluminal suture. Neurological and pathological evaluation of an improved model. *Stroke; a journal of cerebral circulation*. 1996; 27(9):1616–1622. discussion 1623.
- Broderick JP, Palesch YY, Demchuk AM, Yeatts SD, Khatri P, Hill MD, Jauch EC, Jovin TG, Yan B, Silver FL, von Kummer R, Molina CA, Demaerschalk BM, Budzik R, Clark WM, Zaidat OO, Malisch TW, Goyal M, Schonewille WJ, Mazighi M, Engelter ST, Anderson C, Spilker J, Carrozzella J, Ryckborst KJ, Janis LS, Martin RH, Foster LD, Tomsick TA. Interventional Management of Stroke III. Endovascular therapy after intravenous t-PA versus t-PA alone for stroke. *The New England journal of medicine*. 2013; 368(10):893–903. doi:10.1056/NEJMoa1214300. [PubMed: 23390923]
- Chapman DF, Lyden P, Lapchak PA, Nunez S, Thibodeaux H, Zivin J. Comparison of TNK with wild-type tissue plasminogen activator in a rabbit embolic stroke model. *Stroke; a journal of cerebral circulation*. 2001; 32(3):748–752.
- Daugherty A, Lu H, Howatt DA, Rateri DL. Modes of defining atherosclerosis in mouse models: relative merits and evolving standards. *Methods in molecular biology*. 2009; 573:1–15. doi: 10.1007/978-1-60761-247-6\_1. [PubMed: 19763919]
- Davis MJ, Menon BK, Baghirzada LB, Campos-Herrera CR, Goyal M, Hill MD, Archer DP, Calgary Stroke P. Anesthetic management and outcome in patients during endovascular therapy for acute



stroke. *Anesthesiology*. 2012; 116(2):396–405. doi:10.1097/ALN.0b013e318242a5d2. [PubMed: 2222475]

- Fearon P, McArthur KS, Garrity K, Graham LJ, McGroarty G, Vincent S, Quinn TJ. Prestroke modified rankin stroke scale has moderate interobserver reliability and validity in an acute stroke setting. *Stroke; a journal of cerebral circulation*. 2012; 43(12):3184–3188. doi:10.1161/STROKEAHA.112.670422.
- Feuerstein GZ, Zaleska MM, Krams M, Wang X, Day M, Rutkowski JL, Finklestein SP, Pangalos MN, Poole M, Stiles GL, Ruffolo RR, Walsh FL. Missing steps in the STAIR case: a Translational Medicine perspective on the development of NXY-059 for treatment of acute ischemic stroke. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2008; 28(1):217–219. doi:10.1038/sj.jcbfm.9600516.
- Fisher M, Albers GW, Donnan GA, Furlan AJ, Grotta JC, Kidwell CS, Sacco RL, Wechsler LR, Stroke Therapy Academic Industry Roundtable IV. Enhancing the development and approval of acute stroke therapies: Stroke Therapy Academic Industry roundtable. *Stroke; a journal of cerebral circulation*. 2005; 36(8):1808–1813. doi:10.1161/01.STR.0000173403.60553.27.
- Fisher M, Feuerstein G, Howells DW, Hurn PD, Kent TA, Savitz SI, Lo EH, Group S. Update of the stroke therapy academic industry roundtable preclinical recommendations. *Stroke; a journal of cerebral circulation*. 2009; 40(6):2244–2250. doi:10.1161/STROKEAHA.108.541128.
- Fisher M, Hanley DF, Howard G, Jauch EC, Warach S, Group S. Recommendations from the STAIR V meeting on acute stroke trials, technology and outcomes. *Stroke; a journal of cerebral circulation*. 2007; 38(2):245–248. doi:10.1161/01.STR.0000255951.37434.aa.
- Fisher M, Stroke Therapy Academic Industry R. Recommendations for advancing development of acute stroke therapies: Stroke Therapy Academic Industry Roundtable 3. *Stroke; a journal of cerebral circulation*. 2003; 34(6):1539–1546. doi:10.1161/01.STR.0000072983.64326.53.
- Flexman AM, Donovan AL, Gelb AW. Anesthetic management of patients with acute stroke. *Anesthesiology clinics*. 2012; 30(2):175–190. doi:10.1016/j.anclin.2012.04.002. [PubMed: 22901605]
- Fox JG. Laboratory-Animal Medicine Changes and Challenges. *Cornell Vet*. 1985; 75(1):159–170. [PubMed: 3884262]
- Froehler MT, Fifi JT, Majid A, Bhatt A, Ouyang M, McDonagh DL. Anesthesia for endovascular treatment of acute ischemic stroke. *Neurology*. 2012; 79(13 Suppl 1):S167–173. doi:10.1212/WNL.0b013e31826959c2. [PubMed: 23008394]
- Gakuba C, Gauberti M, Mazighi M, Defer G, Hanouz JL, Vivien D. Preclinical evidence toward the use of ketamine for recombinant tissue-type plasminogen activator-mediated thrombolysis under anesthesia or sedation. *Stroke; a journal of cerebral circulation*. 2011; 42(10):2947–2949. doi:10.1161/STROKEAHA.111.620468.
- Gelb AW, Bayona NA, Wilson JX, Cechetto DF. Propofol anesthesia compared to awake reduces infarct size in rats. *Anesthesiology*. 2002; 96(5):1183–1190. [PubMed: 11981160]
- Ginsberg MD, Palesch YY, Hill MD, Martin RH, Moy CS, Barsan WG, Waldman BD, Tamariz D, Ryckborst KJ, Alias, Neurological Emergencies Treatment Trials I. High-dose albumin treatment for acute ischaemic stroke (ALIAS) Part 2: a randomised, double-blind, phase 3, placebo-controlled trial. *Lancet neurology*. 2013; 12(11):1049–1058. doi:10.1016/S1474-4422(13)70223-0.
- Hacke W, Kaste M, Bluhmki E, Brozman M, Davalos A, Guidetti D, Larrue V, Lees KR, Medeghri Z, Machnig T, Schneider D, von Kummer R, Wahlgren N, Toni D, Investigators E. Thrombolysis with alteplase 3 to 4.5 hours after acute ischemic stroke. *The New England journal of medicine*. 2008; 359(13):1317–1329. doi:10.1056/NEJMoa0804656. [PubMed: 18815396]
- Hainsworth AH, Markus HS. Do in vivo experimental models reflect human cerebral small vessel disease? A systematic review. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2008; 28(12):1877–1891. doi:10.1038/jcbfm.2008.91.
- Head BP, Patel P. Anesthetics and brain protection. *Curr Opin Anaesthesiol*. 2007; 20(5):395–399. [PubMed: 17873591]
- Henninger N, Sicard KM, Schmidt KF, Bardutzky J, Fisher M. Comparison of ischemic lesion evolution in embolic versus mechanical middle cerebral artery occlusion in Sprague Dawley rats

using diffusion and perfusion imaging. *Stroke; a journal of cerebral circulation*. 2006; 37(5):1283–1287. doi:10.1161/01.STR.0000217223.72193.98.

- Hossmann KA. The two pathophysiologies of focal brain ischemia: implications for translational stroke research. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2012; 32(7):1310–1316. doi: 10.1038/jcbfm.2011.186.
- Hsia AW, Sachdev HS, Tomlinson J, Hamilton SA, Tong DC. Efficacy of IV tissue plasminogen activator in acute stroke: does stroke subtype really matter? *Neurology*. 2003; 61(1):71–75. [PubMed: 12847159]
- Huybrechts KF, Caro JJ. The Barthel Index and modified Rankin Scale as prognostic tools for long-term outcomes after stroke: a qualitative review of the literature. *Current medical research and opinion*. 2007; 23(7):1627–1636. doi:10.1185/030079907X210444. [PubMed: 17559756]
- Jickling GC, Stamova B, Ander BP, Zhan X, Tian Y, Liu D, Xu H, Johnston SC, Verro P, Sharp FR. Profiles of lacunar and nonlacunar stroke. *Annals of neurology*. 2011; 70(3):477–485. doi: 10.1002/ana.22497. [PubMed: 21796664]
- Jickling GC, Xu H, Stamova B, Ander BP, Zhan X, Tian Y, Liu D, Turner RJ, Mesias M, Verro P, Khoury J, Jauch EC, Pancioli A, Broderick JP, Sharp FR. Signatures of cardioembolic and large-vessel ischemic stroke. *Annals of neurology*. 2010; 68(5):681–692. doi:10.1002/ana.22187. [PubMed: 21031583]
- Kiratas H, Erdener SE, Gursoy-Ozdemir Y, Gurer G, Soylemezoglu F, Dunn AK, Dalkara T. Thrombotic distal middle cerebral artery occlusion produced by topical FeCl(3) application: a novel model suitable for intravital microscopy and thrombolysis studies. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2011; 31(6):1452–1460. doi:10.1038/jcbfm.2011.8.
- Kidwell CS, Jahan R, Gornbein J, Alger JR, Nenov V, Ajani Z, Feng L, Meyer BC, Olson S, Schwamm LH, Yoo AJ, Marshall RS, Meyers PM, Yavagal DR, Wintermark M, Guzy J, Starkman S, Saver JL, Investigators MR. A trial of imaging selection and endovascular treatment for ischemic stroke. *The New England journal of medicine*. 2013; 368(10):914–923. doi:10.1056/NEJMoa1212793. [PubMed: 23394476]
- Kirsch JR, Traystman RJ, Hurn PD. Anesthetics and cerebroprotection: experimental aspects. *Int Anesthesiol Clin*. 1996; 34(4):73–93. [PubMed: 8956065]
- Koerner IP, Brambrink AM. Brain protection by anesthetic agents. *Curr Opin Anaesthesiol*. 2006; 19(5):481–486. [PubMed: 16960478]
- Landis SC, Amara SG, Asadullah K, Austin CP, Blumenstein R, Bradley EW, Crystal RG, Darnell RB, Ferrante RJ, Fillit H, Finkelstein R, Fisher M, Gendelman HE, Golub RM, Goudreau JL, Gross RA, Gubitza AK, Hesterlee SE, Howells DW, Huguenard J, Kelner K, Koroshetz W, Krainc D, Latic SE, Levine MS, Macleod MR, McCall JM, Moxley RT 3rd, Narasimhan K, Noble LJ, Perrin S, Porter JD, Steward O, Unger E, Utz U, Silberberg SD. A call for transparent reporting to optimize the predictive value of preclinical research. *Nature*. 2012; 490(7419):187–191. doi: 10.1038/nature11556. [PubMed: 23060188]
- Lansberg MG, Lee J, Christensen S, Straka M, De Silva DA, Mlynash M, Campbell BC, Bammer R, Olivot JM, Desmond P, Davis SM, Donnan GA, Albers GW. RAPID automated patient selection for reperfusion therapy: a pooled analysis of the Echoplanar Imaging Thrombolytic Evaluation Trial (EPITHET) and the Diffusion and Perfusion Imaging Evaluation for Understanding Stroke Evolution (DEFUSE) Study. *Stroke; a journal of cerebral circulation*. 2011; 42(6):1608–1614. doi: 10.1161/STROKEAHA.110.609008.
- Lansberg MG, Straka M, Kemp S, Mlynash M, Wechsler LR, Jovin TG, Wilder MJ, Lutsep HL, Czartoski TJ, Bernstein RA, Chang CW, Warach S, Fazekas F, Inoue M, Tipirneni A, Hamilton SA, Zaharchuk G, Marks MP, Bammer R, Albers GW, investigators Ds. MRI profile and response to endovascular reperfusion after stroke (DEFUSE 2): a prospective cohort study. *Lancet neurology*. 2012; 11(10):860–867. doi:10.1016/S1474-4422(12)70203-X.
- Lapchak PA. Effect of internal carotid artery reperfusion in combination with Tenecteplase on clinical scores and hemorrhage in a rabbit embolic stroke model. *Brain research*. 2009; 1294:211–217. doi: 10.1016/j.brainres.2009.07.058. [PubMed: 19643097]

- Lapchak PA. Translational stroke research using a rabbit embolic stroke model: a correlative analysis hypothesis for novel therapy development. *Translational stroke research*. 2010; 1(2):96–107. doi: 10.1007/s12975-010-0018-4. [PubMed: 20539748]
- Lapchak PA, Araujo DM, Pakola S, Song D, Wei J, Zivin JA. Microplasmin: a novel thrombolytic that improves behavioral outcome after embolic strokes in rabbits. *Stroke; a journal of cerebral circulation*. 2002a; 33(9):2279–2284.
- Lapchak PA, Araujo DM, Song D, Wei J, Zivin JA. Neuroprotective effects of the spin trap agent disodium-[(tert-butylimino)methyl]benzene-1,3-disulfonate N-oxide (generic NXY-059) in a rabbit small clot embolic stroke model: combination studies with the thrombolytic tissue plasminogen activator. *Stroke; a journal of cerebral circulation*. 2002b; 33(5):1411–1415.
- Lapchak PA, Zhang JH, Noble-Haeusslein LJ. RIGOR Guidelines: Escalating STAIR and STEPS for Effective Translational Research. *Translational stroke research*. 2013; 4(3):279–285. doi:10.1007/s12975-012-0209-2. [PubMed: 23658596]
- Liou SY, Stringer F, Hirayama M. The impact of pharmacogenomics research on drug development. *Drug metabolism and pharmacokinetics*. 2012; 27(1):2–8. [PubMed: 22214937]
- Meyer DM, Compton P, Eastwood JA, Gyls K, Zivin JA. Antiplatelet loading improves behavioral outcome in a rabbit model of stroke. *Stroke; a journal of cerebral circulation*. 2013; 44(11):3246–3248. doi:10.1161/STROKEAHA.113.002565.
- O'Collins VE, Macleod MR, Donnan GA, Horky LL, van der Worp BH, Howells DW. 1,026 experimental treatments in acute stroke. *Annals of neurology*. 2006; 59(3):467–477. doi:10.1002/ana.20741. [PubMed: 16453316]
- Orset C, Macrez R, Young AR, Panthou D, Angles-Cano E, Maubert E, Agin V, Vivien D. Mouse model of in situ thromboembolic stroke and reperfusion. *Stroke; a journal of cerebral circulation*. 2007; 38(10):2771–2778. doi:10.1161/STROKEAHA.107.487520.
- Plenge RM, Scolnick EM, Altshuler D. Validating therapeutic targets through human genetics. *Nature reviews Drug discovery*. 2013; 12(8):581–594. doi:10.1038/nrd4051.
- Quinn TJ, Dawson J, Walters MR, Lees KR. Reliability of the modified Rankin Scale: a systematic review. *Stroke; a journal of cerebral circulation*. 2009; 40(10):3393–3395. doi:10.1161/STROKEAHA.109.557256.
- Reagan-Shaw S, Nihal M, Ahmad N. Dose translation from animal to human studies revisited. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2008; 22(3):659–661. doi:10.1096/fj.07-9574LSF. [PubMed: 17942826]
- Rosell A, Agin V, Rahman M, Morancho A, Ali C, Koistinaho J, Wang X, Vivien D, Schwaninger M, Montaner J. Distal occlusion of the middle cerebral artery in mice: are we ready to assess long-term functional outcome? *Translational stroke research*. 2013; 4(3):297–307. doi:10.1007/s12975-012-0234-1. [PubMed: 24323300]
- Russell JC, Proctor SD. Small animal models of cardiovascular disease: tools for the study of the roles of metabolic syndrome, dyslipidemia, and atherosclerosis. *Cardiovascular pathology : the official journal of the Society for Cardiovascular Pathology*. 2006; 15(6):318–330. doi:10.1016/j.carpath.2006.09.001. [PubMed: 17113010]
- Saver JL, Albers GW, Dunn B, Johnston KC, Fisher M, Consortium SV. Stroke Therapy Academic Industry Roundtable (STAIR) recommendations for extended window acute stroke therapy trials. *Stroke; a journal of cerebral circulation*. 2009; 40(7):2594–2600. doi:10.1161/STROKEAHA.109.552554.
- Sicard KM, Fisher M. Animal models of focal brain ischemia. *Exp Transl Stroke Med*. 2009; 1:7–7. [PubMed: 20150985]
- Singh SS. Preclinical pharmacokinetics: an approach towards safer and efficacious drugs. *Current drug metabolism*. 2006; 7(2):165–182. [PubMed: 16472106]
- Stamova B, Xu H, Jickling G, Bushnell C, Tian Y, Ander BP, Zhan X, Liu D, Turner R, Adamczyk P, Khoury JC, Pancioli A, Jauch E, Broderick JP, Sharp FR. Gene expression profiling of blood for the prediction of ischemic stroke. *Stroke; a journal of cerebral circulation*. 2010; 41(10):2171–2177. doi:10.1161/STROKEAHA.110.588335.
- Stroke Therapy Academic Industry R, II. Recommendations for clinical trial evaluation of acute stroke therapies. *Stroke; a journal of cerebral circulation*. 2001; 32(7):1598–1606.

- Takano K, Tatlisumak T, Bergmann AG, Gibson DG 3rd, Fisher M. Reproducibility and reliability of middle cerebral artery occlusion using a silicone-coated suture (Koizumi) in rats. *Journal of the neurological sciences*. 1997; 153(1):8–11. [PubMed: 9455971]
- Tang Y, Lu A, Aronow BJ, Sharp FR. Blood genomic responses differ after stroke, seizures, hypoglycemia, and hypoxia: blood genomic fingerprints of disease. *Annals of neurology*. 2001; 50(6):699–707. [PubMed: 11761467]
- Tang Y, Xu H, Du X, Lit L, Walker W, Lu A, Ran R, Gregg JP, Reilly M, Pancioli A, Khoury JC, Sauerbeck LR, Carrozzella JA, Spilker J, Clark J, Wagner KR, Jauch EC, Chang DJ, Verro P, Broderick JP, Sharp FR. Gene expression in blood changes rapidly in neutrophils and monocytes after ischemic stroke in humans: a microarray study. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2006; 26(8):1089–1102. doi:10.1038/sj.jcbfm.9600264.
- Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. *The New England journal of medicine*. 1995; 333(24):1581–1587. doi:10.1056/NEJM199512143332401. [PubMed: 7477192]
- Verbeuren TJ. Experimental models of thrombosis and atherosclerosis. *Therapie*. 2006; 61(5):379–387. [PubMed: 17243266]
- Wang L, Jing W, Hang YN. Glutamate-induced c-Jun expression in neuronal PC12 cells: the effects of ketamine and propofol. *J Neurosurg Anesthesiol*. 2008; 20(2):124–130. [PubMed: 18362774]
- Weisscher N, Vermeulen M, Roos YB, de Haan RJ. What should be defined as good outcome in stroke trials; a modified Rankin score of 0-1 or 0-2? *Journal of neurology*. 2008; 255(6):867–874. doi: 10.1007/s00415-008-0796-8. [PubMed: 18338195]
- Yanni AE. The laboratory rabbit: an animal model of atherosclerosis research. *Laboratory animals*. 2004; 38(3):246–256. doi:10.3201/eid0905.020321. [PubMed: 15207035]
- Yao H, Nabika T. Standards and pitfalls of focal ischemia models in spontaneously hypertensive rats: with a systematic review of recent articles. *Journal of translational medicine*. 2012; 10:139. doi: 10.1186/1479-5876-10-139. [PubMed: 22770528]
- Zhang GY, Chen ZQ, Ling F, Li YJ, Wang Y, Gu BX. Establishment of a novel hemodynamic cerebral ischemia model of atherosclerotic rabbit. *Neurology India*. 2010; 58(2):191–194. doi: 10.4103/0028-3886.63782. [PubMed: 20508334]
- Zhao H, Collier JM, Quah DM, Purvis T, Bernhardt J. The modified Rankin Scale in acute stroke has good inter-rater-reliability but questionable validity. *Cerebrovasc Dis*. 2010; 29(2):188–193. doi: 10.1159/000267278. [PubMed: 20029188]
- Zivin JA, Fisher M, DeGirolami U, Hemenway CC, Stashak JA. Tissue plasminogen activator reduces neurological damage after cerebral embolism. *Science*. 1985; 230(4731):1289–1292. [PubMed: 3934754]