

# UC Merced

## UC Merced Previously Published Works

### Title

High-quality permanent draft genome sequence of the extremely osmotolerant diphenol degrading bacterium *Halotalea alkalilenta* AW-7T, and emended description of the genus *Halotalea*

### Permalink

<https://escholarship.org/uc/item/3v83n122>

### Journal

Environmental Microbiome, 10(1)

### ISSN

2524-6372

### Authors

Ntougias, Spyridon

Lapidus, Alla

Copeland, Alex

et al.

### Publication Date

2015

### DOI

10.1186/s40793-015-0052-7

Peer reviewed



SHORT GENOME REPORT

Open Access



# High-quality permanent draft genome sequence of the extremely osmotolerant diphenol degrading bacterium *Halotalea alkalilenta* AW-7<sup>T</sup>, and emended description of the genus *Halotalea*

Spyridon Ntougias<sup>1</sup>, Alla Lapidus<sup>2,3</sup>, Alex Copeland<sup>4</sup>, T. B. K. Reddy<sup>4</sup>, Amrita Pati<sup>4</sup>, Natalia N. Ivanova<sup>4</sup>, Victor M. Markowitz<sup>5</sup>, Hans-Peter Klenk<sup>6</sup>, Tanja Woyke<sup>4</sup>, Constantinos Fasseas<sup>7</sup>, Nikos C. Kyrpides<sup>4,8</sup> and Georgios I. Zervakis<sup>9\*</sup>

## Abstract

Members of the genus *Halotalea* (family *Halomonadaceae*) are of high significance since they can tolerate the greatest glucose and maltose concentrations ever reported for known bacteria and are involved in the degradation of industrial effluents. Here, the characteristics and the permanent-draft genome sequence and annotation of *Halotalea alkalilenta* AW-7<sup>T</sup> are described. The microorganism was sequenced as a part of the Genomic Encyclopedia of Type Strains, Phase I: the one thousand microbial genomes (KMG) project at the DOE Joint Genome Institute, and it is the only strain within the genus *Halotalea* having its genome sequenced. The genome is 4,467,826 bp long and consists of 40 scaffolds with 64.62 % average GC content. A total of 4,104 genes were predicted, comprising of 4,028 protein-coding and 76 RNA genes. Most protein-coding genes (87.79 %) were assigned to a putative function. *Halotalea alkalilenta* AW-7<sup>T</sup> encodes the catechol and protocatechuate degradation to  $\beta$ -keto adipate via the  $\beta$ -keto adipate and protocatechuate ortho-cleavage degradation pathway, and it possesses the genetic ability to detoxify fluoroacetate, cyanate and acrylonitrile. An emended description of the genus *Halotalea* Ntougias et al. 2007 is also provided in order to describe the delayed fermentation ability of the type strain.

**Keywords:** Alkaline two-phase olive mill waste, *Halomonadaceae*, Protocatechuate ortho-cleavage, Catechol to  $\beta$ -keto adipate degradation pathway, Cyanate and acrylonitrile detoxification, GEBA-KMG

## Introduction

The genus *Halotalea* includes a single species, i.e., *H. alkalilenta*, which is a motile, rod-shaped, alkalitolerant and halotolerant Gram-negative staining heterotrophic bacterium [1]. Strain AW-7<sup>T</sup> (=DSM 17697<sup>T</sup> =CECT 7134<sup>T</sup> =CIP 109710<sup>T</sup>) is the type species of the genus *Halotalea* and of the type strain of the species *H. alkalilenta* [1]. The strain was isolated from alkaline olive mill waste, which was generated by a two-phase centrifugal olive oil extraction system located in the Toplou Monastery area, Sitia, Crete [1]. The Neo-Latin genus name derived from the Greek and the

Latin nouns *halos* and *talea*, meaning salt-living and rod-shaped cells, respectively. The Neo-Latin species epithet *halotalea* composed of the Arabic term *al qaliy* and the Latin epithet *lentus* (*a*), meaning alkali and slow respectively which refer to slowly-growing cells under alkaline conditions (alkalitolerant) [1].

*Halotalea alkalilenta* belongs to the family *Halomonadaceae* [1–4], which has accommodated in chronological order the genera *Halomonas* [5], *Chromohalobacter* [6], *Zymobacter* [7], *Carnimonas* [8], *Cobieta* [9], *Halotalea* [1], *Modicisalibacter* [4], *Salinicola* [10], *Kushneria* [11], *Aidingimonas* [12] and *Larsenimonas* [13–15]. By employing multilocus sequence analysis, de la Haba et al. [16] found that all genera of the family *Halomonadaceae*, apart from *Halomonas* and *Modicisalibacter*, are

\* Correspondence: zervakis@aua.gr

<sup>9</sup>Laboratory of General and Agricultural Microbiology, Agricultural University of Athens, Athens, Greece

Full list of author information is available at the end of the article



phylogenetically distinct. *Carnimonas nigrificans* and *Zymobacter palmae* are the closest phylogenetic relatives of *H. alkalilenta*, and were isolated from cured meat and palm sap respectively [7, 8]. *H. alkalilenta* differs from *C. nigrificans* in its higher DNA G+C content and salt upper limit for growth, colony color, motility, its ability to grow at 5 °C and 37 °C, to utilize mannitol, in its inability to hydrolyze starch, to deaminize phenylalanine and to produce acids from D-mannitol and sucrose, in the proportion of the major membrane fatty acids and in the presence/absence of C<sub>10:0</sub>, C<sub>12:0</sub>, C<sub>12:02</sub>-OH, C<sub>14:0</sub>, C<sub>16:0</sub> 3-OH, cyclo-C<sub>17:0</sub>, C<sub>18:0</sub> and C<sub>18:1</sub>9 [1, 8]. *H. alkalilenta* can be distinguished from *Z. palmae* in its higher DNA G+C content, colony color, pH and emperature range for growth, optimum growth temperature, its higher D-glucose tolerance, its ability to utilize citrate, its inability to give positive methyl red and Voges-Proskauer reactions, in the proportion of the major membrane fatty acids and in the presence/absence of C<sub>10:0</sub>, C<sub>10:0</sub> 3-OH, C<sub>12:0</sub> 2-OH, C<sub>15:0</sub>, C<sub>17:0</sub>, cyclo-C<sub>17:0</sub>, C<sub>18:1</sub> ω9 and C<sub>18:1</sub> ω7 [1, 7].

Here, a summarized classification and key characteristics are presented for *H. alkalilenta* AW-7<sup>T</sup>, together with the description of the high-quality permanent draft genome sequence and annotation.

## Organism information

### Classification and features

The 16S rRNA gene sequence of *H. alkalilenta* AW-7<sup>T</sup> was compared using NCBI BLAST under default settings (e.g., considering only the high-scoring segment pairs (HSPs) from the best 250 hits) with the most recent release of the Greengenes database [17] and the relative frequencies of taxa and keywords (reduced to their stem [18]) were determined and weighted by BLAST scores. The frequency of genera that belonged to the family *Halomonadaceae* was 95.2 %. The closest match of *H. alkalilenta* AW-7<sup>T</sup> in 16S rRNA gene, submitted in INSDC (=EMBL/NCBI/DDBJ) under the accession number DQ421388 (=NR\_043806), were *Zymobacter palmae* ATCC 51623<sup>T</sup> (NR\_041786) [7] and *Carnimonas nigrificans* CTCBS1<sup>T</sup> (NR\_029342) [8] showing BLAST similarities of 96.2 % and 95.3 % respectively and HSP coverages of 99.7 % and 100 % respectively.

Figure 1 shows the phylogenetic allocation of *H. alkalilenta* AW-7<sup>T</sup> within the family *Halomonadaceae* in a 16S rRNA gene sequence-based tree. The sequence of the only 16S rRNA gene copy in the genome differs by 5 nucleotides from the previously published 16S rRNA sequence (DQ421388= NR\_043806, coverage 95.0 %).

*H. alkalilenta* AW-7<sup>T</sup> is a Gram-negative motile rod-shaped bacterium [1] with a length of 1.4–2.1 μm and a width of 0.6–0.9 μm (Table 1 and Fig. 2). The temperature range for growth is 5–45 °C, with an optimum temperature for growth at 32–37 °C [1]. *H. alkalilenta* AW-7<sup>T</sup> is halotolerant and alkalitolerant, growing at salinity and pH ranges

of 0–150 g L<sup>-1</sup> NaCl and 5–11, respectively [1]. The optimum salt and pH for growth are 0–3 % w/v NaCl and 7, respectively [1].

*H. alkalilenta* AW-7<sup>T</sup> is a non-denitrifying chemoorganotroph; it utilizes mostly L-glutamine and L-proline, followed by D-galactose, D-glucose, glycerol, D-mannitol, protocatechuate, L-serine, succinate and sucrose, while it grows weakly on acetate, citrate, D-fructose, maltose, sorbitol and gallate [1]. *H. alkalilenta* AW-7<sup>T</sup> also produces acid aerobically from D-fructose, D-galactose, D-glucose, maltose, D-mannose and melibiose, and hydrolyses Tween 20 [1]. Despite the fact that urea hydrolysis is encoded in *H. alkalilenta* AW-7<sup>T</sup> genome, no positive reaction was detected by Ntougias et al. [1] and the present study (using the EnteroPluri-Test). *H. alkalilenta* AW-7<sup>T</sup> is susceptible to kanamycin, polymixin B, rifampicin, streptomycin and tetracycline (50 mg L<sup>-1</sup> each) [1].

In the past, *H. alkalilenta* AW-7<sup>T</sup> and *C. nigrificans* CTCBS1<sup>T</sup> were reported as oxidase positive [1, 8]. However, genome comparisons showed that both *H. alkalilenta* AW-7<sup>T</sup> and *C. nigrificans* CTCBS1<sup>T</sup> possessed an identical oxidative phosphorylation pathway that lacks cytochrome c oxidase, which was distinct from that of *Z. palmae* T109<sup>T</sup>. In addition, no fermentation ability was previously detected for *H. alkalilenta* AW-7<sup>T</sup> using standard incubation periods [1], although the pyruvate fermentation to acetate II MetaCyc pathway is encoded in both *H. alkalilenta* AW-7<sup>T</sup> and *Z. palmae* T109<sup>T</sup>. For this reason, the fermentation ability of *H. alkalilenta* AW-7<sup>T</sup> was re-examined under prolonged incubation period using the EnteroPluri-Test (BD, USA). No fermentation reaction was observed for incubations up to 4–days, although, thereafter, a positive reaction was obtained for glucose (at the 5th day of incubation, without gas production) and dulcitol (at 9th day of incubation). *H. alkalilenta* AW-7<sup>T</sup> could not ferment adonitol, lactose, arabinose and sorbitol after a 9–days incubation period. In agreement to what was previously reported by Ntougias et al. [1], no growth of *H. alkalilenta* AW-7<sup>T</sup> was observed in the present study on yeast extract-peptone-glucose agar plates placed for an incubation period of 1 month in an anaerobic jar containing the Anaerocult A system (Merck). However, exposure of culture plates to oxygen led to fastidious growth. In this sense, it is concluded that *H. alkalilenta* AW-7<sup>T</sup> can tolerate anaerobic conditions through a slow fermentation mechanism.

### Chemotaxonomy

The main membrane fatty acids of *H. alkalilenta* AW-7<sup>T</sup> are in the descending order of concentration: C<sub>18:1</sub> ω7c, C<sub>16:0</sub>, C<sub>19:0</sub> cyclo ω8c, C<sub>12:0</sub> 3-OH and C<sub>16:1</sub> ω7c/iso-C<sub>15:0</sub> 2-OH [1]. The only respiratory quinone found in *H. alkalilenta* AW-7<sup>T</sup> is ubiquinone-9 [1].




Fig. 1 (See legend on next page.)

(See figure on previous page.)

**Fig. 1** Phylogenetic tree displaying the position of *H. alkalilenta* AW-7<sup>T</sup> among the type strains of other species within the *Halomonadaceae*. The tree was inferred from 1152 aligned characters [38, 39] of the 16S rRNA gene sequence under the maximum likelihood (ML) criterion [40]. Tree branches are constructed on the basis of the expected number of substitutions per site. Values above branches denote support values from 100 ML bootstrap replicates [41]. Members of different genera within the *Halomonadaceae* are depicted in different fonts color. Lineages with strain genome sequencing projects registered in GOLD [24] are labeled with one asterisk, and those also listed as 'Complete and Published' with two asterisks

## Genome sequencing and annotation

### Genome project history


*H. alkalilenta* AW-7<sup>T</sup> was selected for sequencing on the basis of its phylogenetic position [19–21], and is part of Genomic Encyclopedia of Type Strains, Phase I:

the one thousand microbial genomes project [22] which aims not only to increase the sequencing coverage of key reference microbial genomes [23]. The genome project is accessible in the Genomes On Line Database [24] and the entire genome sequence is deposited in GenBank.

**Table 1** Classification and general features of *Halotalea alkalilenta* strain AW-7<sup>T</sup> according to the MIGS recommendations [42], published by the Genome Standards Consortium [43] and the Names for Life database [44]

MIGS ID	Property	Term	Evidence code <sup>a</sup>
	Classification	Domain <i>Bacteria</i>	TAS [45]
		Phylum <i>Proteobacteria</i>	TAS [46, 47]
		Class <i>Gammaproteobacteria</i>	TAS [47–49]
		Order <i>Oceanospirillales</i>	TAS [47, 50]
		Family <i>Halomonadaceae</i>	TAS [1–4, 51]
		Genus <i>Halotalea</i>	TAS [1]
		Species <i>Halotalea alkalilenta</i>	TAS [1]
		Type strain: AW-7 <sup>T</sup>	TAS [1]
	Gram stain	<i>negative</i>	TAS [1]
	Cell shape	<i>rod</i>	TAS [1]
	Motility	<i>motile</i>	TAS [1]
	Sporulation	<i>non-sporulating</i>	TAS [1]
	Temperature range	<i>5-45 °C</i>	TAS [1]
	Optimum temperature	<i>32-37 °C</i>	TAS [1]
	pH range; Optimum	<i>5-11; 7</i>	TAS [1]
	Carbon source	<i>carbohydrates, amino-acids, organic acid anions and alcohols</i>	TAS [1]
MIGS-6	Habitat	<i>olive mill waste</i>	TAS [1]
MIGS-6.3	Salinity	<i>up to 15 % NaCl w/v</i>	TAS [1]
MIGS-22	Oxygen requirement	<i>facultatively anaerobic</i>	IDA
MIGS-15	Biotic relationship	<i>free-living</i>	TAS [1]
MIGS-14	Pathogenicity	<i>none</i>	NAS
	Biosafety level	<i>1</i>	TAS [52]
MIGS-4	Geographic location	<i>Greece, Crete, Toplou Monastery</i>	TAS [1]
MIGS-5	Sample collection	<i>2003</i>	NAS
MIGS-4.1	Latitude	<i>35.220</i>	TAS [1]
MIGS-4.2	Longitude	<i>26.216</i>	TAS [1]
MIGS-4.3	Depth	<i>surface</i>	NAS
MIGS-4.4	Altitude	<i>161 m</i>	NAS

<sup>a</sup>Evidence codes - IDA: Inferred from Direct Assay; TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [53]



**Fig. 2** Electron micrograph of negatively-stained *H. alkalilenta* AW-7<sup>T</sup> cells. Bar denotes 1  $\mu$ m

Sequencing, finishing and annotation were accomplished by the DOE Joint Genome Institute [25] using state of the art genome sequencing technology [26]. The project information is summarized in Table 2.

#### Growth conditions and genomic DNA preparation

*H. alkalilenta* AW-7<sup>T</sup> was cultivated aerobically in trypticase soy yeast extract medium at 28 °C. Genomic DNA was obtained using the Invitrogen PureLink® Genomic DNA Mini Kit (Life Technologies Inc.) following the standard protocol. In addition, DNA prepared by the DSMZ is available via the DNA Bank Network [27].

#### Genome sequencing and assembly

The draft genome of was generated at the DOE Joint Genome Institute using the Illumina technology [28].

An Illumina std shotgun library was constructed and sequenced using the Illumina HiSeq 2000 platform which generated 13,537,536 reads totaling 2,030.6 Mb. All general aspects of library construction and sequencing performed can be found at JGI website [29]. All raw Illumina sequence data was passed through DUK, a filtering program developed at JGI, which removes known Illumina sequencing and library preparation artifacts (Mingkun L, et al., unpublished). Following steps were then performed for assembly: (1) filtered Illumina reads were assembled using Velvet (version 1.2.07) [30], (2) 1–3 kb simulated paired end reads were created from Velvet contigs using wgsim [31], (3) Illumina reads were assembled with simulated read pairs using Allpaths-LG (version r46652) [32]. Parameters for assembly steps were: 1) Velvet (velveth: 63 –shortPaired and velvetg: –very clean yes –exportFiltered yes –min contig lgth 500 –scaffolding no –cov cutoff 10) 2) wgsim (–e 0 –l 100 –r 0 –R 0 –X 0) 3) Allpaths-LG (PrepareAllpathsInputs:PHRED 64 = 1 PLOIDY = 1 FRAG COVERAGE = 125 JUMP COVERAGE = 25 LONG JUMP COV = 50, RunAllpathsLG: THREADS = 8 RUN = std shredpairs TARGETS = standard VAPI WARN ONLY = True OVERWRITE = True). The final draft assembly contained 56 contigs in 40 scaffolds, totaling 4.5 Kb in size. The final assembly was based on 1,500.0 Mb of Illumina data. Based on a presumed genome size of 5.0 Mb, the average input read coverage used for the assembly was 300.0  $\times$ .

#### Genome annotation

Genes were detected using the Prodigal software [33] at the DOE-JGI Genome Annotation pipeline [34, 35]. The CDSs predicted were translated and searched against the National Center for Biotechnology Information non-redundant database, UniProt, TIGRFam, Pfam, PRIAM, KEGG, COG, and InterPro databases. Additional gene prediction and functional annotation analysis was carried

**Table 2** *H. alkalilenta* AW-7<sup>T</sup> genome sequencing project details

MIGS ID	Property	Term
MIGS-31	Finishing quality	High-Quality Draft
MIGS-29	Sequencing platforms	Illumina HiSeq 2000
MIGS-31.2	Sequencing coverage	300 $\times$
MIGS-30	Assemblers	vpAllpaths v. r46652
MIGS-32	Gene calling method	Prodigal 2.5
	INSDC ID	JHYY00000000
	Genbank Date of Release	May 5, 2014
	GOLD ID	Gp0040002
	NCBI project ID	221047
MIGS-13	Source material identifier	DSM 17697 <sup>T</sup>
	Project relevance	GEBA-KMG, Tree of Life, Biodegradation, Extremophiles

out in the Integrated Microbial Genomes – Expert Review platform [36]. The genome sequence and the annotations described in this paper are available from the Integrated Microbial Genome system [37].

### Genome properties

The genome is 4,467,826 bp long and comprised of 40 scaffolds with 64.62 % average GC content (Table 3). A total of 4,104 genes were predicted, consisting of 4,028 protein-coding and 76 RNA genes. The majority of protein-coding genes (87.79 %) were assigned to a putative function, whereas the remaining ones were annotated as hypothetical proteins. Distribution of genes into COGs functional categories is displayed in Table 4.

### Insights into the genome sequence

The genome size of *H. alkalilenta* AW-7<sup>T</sup> (4.47 Mbp) is 50 % and 60 % greater than those of *Z. palmae* T109<sup>T</sup> and *C. nigrificans* CTCBS1<sup>T</sup> (2.73 and 2.98 Mbp) respectively. In *H. alkalilenta* AW-7<sup>T</sup>, protein coding genes involved in the major functional categories (i.e., amino acid, carbohydrate and lipid metabolism, membrane transport, energy metabolism) are 50 % and 30 % greater in number than those detected in *Z. palmae* T109<sup>T</sup> and *C. nigrificans* CTCBS1<sup>T</sup>, respectively. Moreover, genes encoding xenobiotic metabolic proteins are 69 % and 57 % more in *H. alkalilenta* AW-7<sup>T</sup> than those identified in *Z. palmae* T109<sup>T</sup> and *C. nigrificans* CTCBS1<sup>T</sup> respectively.

Genome data uncovered the genetic ability of *H. alkalilenta* AW-7<sup>T</sup> to degrade several recalcitrant substrates.

**Table 3** Genome statistics

Attribute	Value	% of Total <sup>a</sup>
Genome size (bp)	4,467,826	100.00
DNA coding region (bp)	3,922,088	87.79
DNA G + C content (bp)	2,887,209	64.62
DNA scaffolds	40	
Total genes	4,104	100.00
RNA genes	76	1.85
tRNA genes	52	1.27
Protein-coding genes	4,028	98.15
Pseudo genes	0	0.00
Genes with function prediction (proteins)	3,603	87.79
Genes in paralog clusters	3,380	82.36
Genes assigned to COGs	3,246	79.09
Genes assigned Pfam domains	3,637	88.62
Genes with signal peptides	343	8.36
Genes with transmembrane helices	905	22.05
CRISPR repeats	3	

<sup>a</sup>The total is based on either the size of the genome in base pairs or the total number of protein coding genes in the annotated genome

**Table 4** Number of genes associated with the general COG functional categories

Code	Value	% Age	Description
J	188	5.15	Translation, ribosomal structure and biogenesis
A	1	0.03	RNA processing and modification
K	313	8.58	Transcription
L	124	3.40	Replication, recombination and repair
B	3	0.08	Chromatin structure and dynamics
D	30	0.82	Cell cycle control, cell division, chromosome partitioning
V	29	0.79	Defense mechanisms
T	125	3.42	Signal transduction mechanisms
M	169	4.63	Cell wall/membrane biogenesis
N	58	1.59	Cell motility
U	56	1.53	Intracellular trafficking, secretion and vesicular transport
O	111	3.04	Posttranslational modification, protein turnover, chaperones
C	233	6.38	Energy production and conversion
G	259	7.10	Carbohydrate transport and metabolism
E	525	14.38	Amino acid transport and metabolism
F	86	2.36	Nucleotide transport and metabolism
H	164	4.49	Coenzyme transport and metabolism
I	120	3.29	Lipid transport and metabolism
P	261	7.15	Inorganic ion transport and metabolism
Q	98	2.68	Secondary metabolites biosynthesis, transport and catabolism
R	440	12.05	General function prediction only
S	257	7.04	Function unknown
-	858	20.91	Not in COGs

*H. alkalilenta* AW-7<sup>T</sup> encodes the bioconversion of catechol and protocatechuate to  $\beta$ -keto adipate via the  $\beta$ -keto adipate and protocatechuate degradation II (ortho-cleavage) pathway respectively, as verified by the ability of strain AW-7<sup>T</sup> to catabolize certain phenolic compounds. Aerobic benzoate degradation I is also encoded, permitting its catabolism via the catechol degrading pathway. Genes encoding fluoroacetate dehalogenase were identified in the genome of *H. alkalilenta* AW-7<sup>T</sup>, indicating its ability for fluoroacetate degradation. The detection of genes involved in cyanate and acrylonitrile degradation was also verified. Lastly, *H. alkalilenta* AW-7<sup>T</sup> is genetically able to produce ectoine and glycine betaine, which appear to serve as the main osmolytes for the adaptation of this species under high osmotic conditions.

Based on genome metabolic features, *H. alkalilenta* AW-7<sup>T</sup> is prototrophic for L-arginine, L-histidine, L-isoleucine, L-leucine, L-lysine, L-phenylalanine, L-

tryptophan, L-tyrosine and L-valine auxotroph, and L-aspartate, L-glutamate, L-glutamine and glycine. Strain AW-7<sup>T</sup> can synthesize selenocysteine but not biotin.

## Conclusions

Genome sequence and biochemical data of the highly osmotolerant species *Halotalea alkalilenta* AW-7<sup>T</sup> revealed the presence of an oxidative phosphorylation pathway that lacks cytochrome c oxidase, and the encoding of the pyruvate fermentation to acetate II (MetaCyc pathway). *H. alkalilenta* AW-7<sup>T</sup> could ferment glucose and ducitol after a prolonged incubation period, which is indicative of the induction of a slow fermentation mechanism, and results in the emendation of the genus *Halotalea* Ntougias et al. 2007. Comparisons to its closest phylogenetic relatives *Zymobacter palmae* T109<sup>T</sup> and *Carnimonas nigrificans* CTCBS1<sup>T</sup>, confirm the distinct taxonomic position of *H. alkalilenta* AW-7 on the basis of its larger genome size and number of protein coding genes involved in the major functional categories and in xenobiotics metabolism. Furthermore, *H. alkalilenta* AW-7<sup>T</sup> encodes the biotransformation of catechol and protocatechuate to  $\beta$ -keto adipate via the  $\beta$ -keto adipate and protocatechuate degradation II (ortho-cleavage) pathway respectively, verifying at the genome level the ability of strain AW-7<sup>T</sup> to degrade phenolic compounds.

## Emended description of the genus *Halotalea* Ntougias et al. 2007

The description of the genus *Halotalea* is the one given by Ntougias et al. 2007 [1], with the following modification: Facultative anaerobe, which exhibits delayed glucose and dulcitol fermentation ability, and lacks cytochrome c oxidase activity.

### Abbreviations

KMG: One thousand microbial genomes; GEBA: Genomic encyclopedia of Bacteria and Archaea; MIGS: Minimum information about a genome sequence; TAS: Traceable; NAS: Non-traceable.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

SN and GIZ performed the fermentation tests, conducted the phylogenetic analyses and drafted the paper, CF prepared the electron micrographs, and all other authors were involved in either sequencing/annotation of the genome and/or editing the paper. All authors read and approved the final manuscript.

### Acknowledgements

This work was performed under the auspices of the US Department of Energy Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley National Laboratory under contract No. DE-AC02-05CH11231, A.L. was supported in part by Russian Ministry of Science Mega-grant no.11.G34.31.0068 (PI. Dr Stephen J O'Brien).

### Author details

<sup>1</sup>Laboratory of Wastewater Management and Treatment Technologies, Department of Environmental Engineering, Democritus University of Thrace, Xanthi, Greece. <sup>2</sup>Theodosius Dobzhansky Center for Genome Bioinformatics, St. Petersburg State University, St. Petersburg, Russia. <sup>3</sup>Algorithmic Biology Lab, St. Petersburg Academic University, St. Petersburg, Russia. <sup>4</sup>Department of Energy Joint Genome Institute, Genome Biology Program, Walnut Creek, CA, USA. <sup>5</sup>Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, CA, USA. <sup>6</sup>Leibniz Institute DSMZ – German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany. <sup>7</sup>Electron Microscopy Laboratory, Agricultural University of Athens, Athens, Greece. <sup>8</sup>Department of Biological Sciences, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia. <sup>9</sup>Laboratory of General and Agricultural Microbiology, Agricultural University of Athens, Athens, Greece.

Received: 23 December 2014 Accepted: 28 July 2015

Published online: 13 August 2015

### References

- Ntougias S, Zervakis GI, Fasseas C. *Halotalea alkalilenta* gen. nov., sp. nov., a novel osmotolerant and alkalitolerant bacterium from alkaline olive mill wastes, and emended description of the family *Halomonadaceae* Franzmann et al. 1989, emend. Dobson and Franzmann 1996. *Int J Syst Evol Microbiol.* 2007;57:1975–83.
- Franzmann PD, Wehmeyer U, Stackebrandt E. *Halomonadaceae* fam. nov., a new family of the class *Proteobacteria* to accommodate the genera *Halomonas* and *Deleya*. *Syst Appl Microbiol.* 1988;11:16–9.
- Dobson SJ, Franzmann PD. Unification of the genera *Deleya* (Baumann et al. 1983), *Halomonas* (Vreeland et al. 1980), and *Halovibrio* (Fendrich 1988) and the species *Paracoccus halodenitrificans* (Robinson and Gibbons 1952) into a single genus, *Halomonas*, and placement of the genus *Zymobacter* in the family *Halomonadaceae*. *Int J Syst Bacteriol.* 1996;46:550–8.
- Ben Ali Gam Z, Abdelkafi S, Casalot L, Tholozan JL, Oueslati R, Labat M. *Modicisalibacter tunisiensis* gen. nov., sp. nov., an aerobic, moderately halophilic bacterium isolated from an oilfield-water injection sample, and emended description of the family *Halomonadaceae* Franzmann et al. 1989 emend Dobson and Franzmann 1996 emend. Ntougias et al. 2007. *Int J Syst Evol Microbiol.* 2007;57:2307–13.
- Vreeland RH, Litchfield CD, Martin EL, Elliot E. *Halomonas elongata*, a new genus and species of extremely salt-tolerant bacteria. *Int J Syst Bacteriol.* 1980;30:485–95.
- Ventosa A, Gutierrez MC, Garcia MT, Ruiz-Berraquero F. Classification of '*Chromobacterium marismortui*' in a new genus, *Chromohalobacter* gen. nov., as *Chromohalobacter marismortui* comb. nov., nom. rev. *Int J Syst Bacteriol.* 1989;39:382–6.
- Okamoto T, Taguchi H, Nakamura K, Ikenaga H, Kuraishi H, Yamasato K. *Zymobacter palmae* gen. nov., sp. nov., a new ethanol-fermenting peritrichous bacterium isolated from palm sap. *Arch Microbiol.* 1993;160:333–7.
- Garriga M, Ehrmann MA, Arnau J, Hugas M, Vogel RF. *Carnimonas nigrificans* gen. nov., sp. nov., a bacterial causative agent for black spot formation on cured meat products. *Int J Syst Bacteriol.* 1998;48:677–86.
- Arahal DR, Castillo AM, Ludwig W, Schleifer KH, Ventosa A. Proposal of *Cobetia marina* gen. nov., comb. nov., within the family *Halomonadaceae*, to include the species *Halomonas marina*. *Syst Appl Microbiol.* 2002;25:207–11.
- Anan'ina LN, Plotnikova EG, Gavrish EY, Demakov VA, Evtushenko LI. *Salinicola socius* gen. nov., sp. nov., a moderately halophilic bacterium from a naphthalene-utilizing microbial association. *Mikrobiologiya.* 2007;76:324–30.
- Sanchez-Porro C, de la Haba RR, Soto-Ramirez N, Marquez MC, Montalvo-Rodriguez R, Ventosa A. Description of *Kushneria aurantia* gen. nov., sp. nov., a novel member of the family *Halomonadaceae*, and a proposal for reclassification of *Halomonas marisflavi* as *Kushneria marisflavi* comb. nov., of *Halomonas indaliniina* as *Kushneria indaliniina* comb. nov. and of *Halomonas avicenniae* as *Kushneria avicenniae* comb. nov. *Int J Syst Evol Microbiol.* 2009;59:397–405.
- Wang Y, Tang S-K, Lou K, Lee J-C, Jeon CO, Xu L-H, et al. *Aidingimonas halophila* gen. nov., sp. nov., a moderately halophilic bacterium isolated from a salt lake. *Int J Syst Evol Microbiol.* 2009;59:3088–94.



13. León MJ, Sánchez-Porro C, de la Haba RR, Llamas I, Ventosa A. *Larsenia salina* gen. nov., sp. nov., a new member of the family *Halomonadaceae* based on multilocus sequence analysis. *Syst Appl Microbiol*. 2014;37:480–7.
14. León MJ, Sánchez-Porro C, de la Haba RR, Llamas I, Ventosa A. Corrigendum to *Larsenia salina* gen. nov., sp. nov., a new member of the family *Halomonadaceae* based on multilocus sequence analysis [*Syst. Appl. Microbiol*, 37 (October (7)) (2014) 480–487]. *Syst Appl Microbiol*. 2015;38:77.
15. Oren A, Garrity GM. List of new names and new combinations previously effectively, but not validly, published. *Int J Syst Evol Microbiol*. 2015;65:1105–11.
16. de la Haba RR, Carmen Márquez M, Thane Papke R, Ventosa A. Multilocus sequence analysis (MLSA) of the family *Halomonadaceae*. *Int J Syst Evol Microbiol*. 2012;62:520–38.
17. DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, et al. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol*. 2006;72:5069–72.
18. Porter MF. An algorithm for suffix stripping. *Program*. 1980;14:130–137.
19. Klenk HP, Göker M. En route to a genome-based classification of *Archaea* and *Bacteria*? *Syst Appl Microbiol*. 2010;33:175–82.
20. Göker M, Klenk HP. Phylogeny-driven target selection for large-scale genome-sequencing (and other) projects. *Stand Genomic Sci*. 2013;8:360–74.
21. Wu D, Hugenholtz P, Mavromatis K, Pukall R, Dalin E, Ivanova NN, et al. A phylogeny-driven Genomic Encyclopaedia of *Bacteria* and *Archaea*. *Nature*. 2009;462:1056–60.
22. Kyrpides NC, Woyske T, Eisen JA, Garrity G, Lilburn TG, Beck BJ, et al. Genomic Encyclopedia of Type Strains, Phase I: the one thousand microbial genomes (KMG-I) project. *Stand Genomic Sci*. 2013;9:628–34.
23. Kyrpides NC, Hugenholtz P, Eisen JA, Woyske T, Göker M, Parker CT, et al. Genomic encyclopedia of *Bacteria* and *Archaea*: sequencing a myriad of type strains. *PLoS Biol*. 2014;8, e1001920.
24. Reddy TB, Thomas AD, Stamatis D, Bertsch J, Isbandi M, Jansson J, et al. The Genomes OnLine Database (GOLD) v.5: a metadata management system based on a four level (meta)genome project classification. *Nucleic Acids Res*. 2015;43(Database issue):D1099–106.
25. DOE Joint Genome Institute. [<http://www.jgi.doe.gov>].
26. Mavromatis K, Land ML, Brettin TS, Quest DJ, Copeland A, Clum A, et al. The fast changing landscape of sequencing technologies and their impact on microbial genome assemblies and annotation. *PLoS One*. 2012;7, e48837.
27. Gemeinholzer B, Dröge G, Zetzsche H, Haszprunar G, Klenk HP, Güntsch A, et al. The DNA Bank Network: the start from a German initiative. *Biopreserv Biobank*. 2011;9:51–5.
28. Bennett S. Solexa Ltd. *Pharmacogenomics*. 2004;5:433–8.
29. JGI Website. [<http://www.jgi.doe.gov>].
30. Zerbino D, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res*. 2008;18:821–9.
31. Wgsim. [<https://github.com/lh3/wgsim>].
32. Gnerre S, MacCallum I. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci U S A*. 2011;108:1513–8.
33. Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: Prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*. 2010;11:119.
34. Mavromatis K, Ivanova NN, Chen IM, Szeto E, Markowitz VM, Kyrpides NC. The DOE-JGI Standard operating procedure for the annotations of microbial genomes. *Stand Genomic Sci*. 2009;1:63–7.
35. Chen IM, Markowitz VM, Chu K, Anderson I, Mavromatis K, Kyrpides NC, et al. Improving microbial genome annotations in an integrated database context. *PLoS One*. 2013;8, e54859.
36. Markowitz VM, Mavromatis K, Ivanova NN, Chen IM, Chu K, Kyrpides NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics*. 2009;25:2271–8.
37. Markowitz VM, Chen I-M A, Palaniappan K, Chu K, Szeto E, Grechkin Y, et al. IMG: the integrated microbial genomes database and comparative analysis system. *Nucleic Acids Res*. 2012;40:D115–22.
38. Lee C, Grasso C, Sharlow MF. Multiple sequence alignment using partial order graphs. *Bioinformatics*. 2002;18:452–64.
39. Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol*. 2000;17:540–52.
40. Stamatakis A, Hoover P, Rougemont J. A rapid bootstrap algorithm for the RAxML web servers. *Syst Biol*. 2008;57:758–71.
41. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol*. 2013;30:2725–9.
42. Field D, Garrity GM, Gray T, Morrison N, Selengut J, Sterk P, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol*. 2008;26:541–7.
43. Field D, Amaral-Zettler L, Cochrane G, Cole JR, Dawyndt P, Garrity GM, et al. The Genomic Standards Consortium. *PLoS Biol*. 2011;9, e1001088.
44. Garrity GM. Names for Life. Browser Tool takes expertise out of the database and puts it right in the browser. *Microbiol Today*. 2010;37:9.
45. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains *Archaea*, *Bacteria*, and *Eucarya*. *Proc Natl Acad Sci U S A*. 1990;87:4576–9.
46. Garrity GM, Bell JA, Lilburn T. Phylum XIV. *Proteobacteria* phyl. nov. In: Brenner DJ, Krieg NR, Staley JT, Garrity GM, editors. *Bergey's Manual of Systematic Bacteriology*. Volume 2, part B. 2nd ed. New York: Springer; 2005. p. 1.
47. Validation List no. 106. Validation of publication of new names and new combinations previously effectively published outside the IJSEM. *Int J Syst Evol Microbiol*. 2005;55:2235–8.
48. Garrity GM, Bell JA, Lilburn T. Class III. *Gammaproteobacteria* class. nov. In: Brenner DJ, Krieg NR, Staley JT, Garrity GM, editors. *Bergey's Manual of Systematic Bacteriology*. Volume 2, part B. 2nd ed. New York: Springer; 2005. p. 1.
49. Williams KP, Kelly DP. Proposal for a new class within the phylum *Proteobacteria*, *Acidithiobacillia* classis nov., with the type order *Acidithiobacillales*, and emended description of the class *Gammaproteobacteria*. *Int J Syst Evol Microbiol*. 2013;63:2901–6.
50. Garrity GM, Bell JA, Lilburn T. Order VIII. *Oceanospirillales* ord. nov. In: Brenner DJ, Krieg NR, Staley JT, Garrity GM, editors. *Bergey's Manual of Systematic Bacteriology*. Volume 2, part B. 2nd ed. New York: Springer; 2005. p. 270.
51. Validation List no. 29. Validation of the publication of new names and new combinations previously effectively published outside the IJSEM. *Int J Syst Evol Microbiol*. 1989;39:205–6.
52. Bundesanstalt für Arbeitsschutz und Arbeitsmedizin (BAuA). Classification of prokaryotes (bacteria and archaea) into risk groups. Germany: Technical Rule for Biological Agents 466 (TRBA 466); 2010. p. 157.
53. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet*. 2000;25:25–9.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

