

UC Irvine

UC Irvine Previously Published Works

Title

Constraining remote oxidation capacity with ATom observations

Permalink

<https://escholarship.org/uc/item/3vd5w0d8>

Journal

Atmospheric Chemistry and Physics, 20(13)

ISSN

1680-7316

Authors

Travis, Katherine R
Heald, Colette L
Allen, Hannah M
[et al.](#)

Publication Date

2020-07-01

DOI

10.5194/acp-20-7753-2020

Peer reviewed



Published in final edited form as:

Atmos Chem Phys. 2020 July ; 20(13): 7753–7781. doi:10.5194/acp-20-7753-2020.

Constraining remote oxidation capacity with ATom observations

Katherine R. Travis^{1,a}, Colette L. Heald^{1,2}, Hannah M. Allen³, Eric C. Apel⁴, Stephen R. Arnold⁵, Donald R. Blake⁶, William H. Brune⁷, Xin Chen⁸, Róisín Commane⁹, John D. Crounse¹⁰, Bruce C. Daube¹¹, Glenn S. Diskin¹², James W. Elkins¹³, Mathew J. Evans^{14,15}, Samuel R. Hall⁴, Eric J. Hints^{13,16}, Rebecca S. Hornbrook⁴, Prasad S. Kasibhatla¹⁷, Michelle J. Kim^{10,18}, Gan Luo¹⁹, Kathryn McKain^{13,16}, Dylan B. Millet⁸, Fred L. Moore^{13,16}, Jeffrey Peischl^{16,20}, Thomas B. Ryerson²⁰, Tomás Sherwen^{14,15}, Alexander B. Thames⁷, Kirk Ullmann⁴, Xuan Wang^{11,21}, Paul O. Wennberg^{3,18}, Glenn M. Wolfe²², Fangqun Yu¹⁹

¹Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA

²Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA, USA

³Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA, USA

⁴Atmospheric Chemistry Observations & Modeling Laboratory, National Center for Atmospheric Research, Boulder, CO, USA

⁵Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, UK

⁶Department of Chemistry, University of California Irvine, Irvine, CA, USA

⁷Department of Meteorology, Pennsylvania State University, University Park, PA, USA

⁸University of Minnesota, Department of Soil, Water and Climate, St. Paul, MN, USA

⁹Dept. of Earth & Environmental Sciences of Lamont-Doherty Earth Observatory and Columbia University, Palisades, NY, USA

¹⁰Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA

This work is distributed under the Creative Commons Attribution 4.0 License.

Correspondence: Katherine R. Travis (katherine.travis@nasa.gov) and Colette L. Heald (heald@mit.edu).

^anow at: NASA Langley Research Center, Hampton, VA, USA

Author contributions.

CLH and KRT designed the study and wrote the paper with input from the co-authors. KRT modified the code, performed the simulations, and led the analysis. HMA, ECA, DRB, WHB, RC, JDC, BCD, GSD, JWE, SRH, EJH, SRH, MJK, KM, FLM, JP, TBR, ABT, KU, POW, and GMW provided ATom measurements used in the analysis. XW provided the model code for the sensitivity runs including acid displacement of chloride on coarse-mode sea-salt aerosols. TS, ME, and PSK provided the model code for the photolysis of particulate nitrate. GL and FY were responsible for the code for the revised treatment of wet scavenging in the model. DBM and XC provided the methanol seawater concentration and assisted in the ocean budget analysis. SRA provided the biogenic ocean isoprene emissions.

Data availability. The ATom-1 and ATom-2 data (Wofsy et al., 2018) are available here: <https://doi.org/10.3334/ORNDAAC/1581>.

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-20-7753-2020-supplement>.

Competing interests. The authors declare that they have no conflict of interest.

¹¹Harvard John A. Paulson School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, USA

¹²NASA Langley Research Center, Hampton, VA, USA

¹³Global Monitoring Division, NOAA Earth System Research Laboratory, Boulder, CO, USA

¹⁴Wolfson Atmospheric Chemistry Laboratories (WACL), Department of Chemistry, University of York, York, UK

¹⁵National Centre for Atmospheric Science (NCAS), University of York, York, UK

¹⁶Cooperative Institute for Research in Environmental Science, University of Colorado, CO, USA

¹⁷Nicholas School of the Environment, Duke University, Durham, NC, USA

¹⁸Division of Engineering and Applied Science, California Institute of Technology, Pasadena, CA, USA

¹⁹Atmospheric Sciences Research Center, University of Albany, Albany, NY, USA

²⁰Chemical Sciences Division, NOAA Earth System Research Laboratory, Boulder, CO, USA

²¹School of Energy and Environment, City University of Hong Kong, Hong Kong, China

²²Atmospheric Chemistry and Dynamics Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD, USA

Abstract

The global oxidation capacity, defined as the tropospheric mean concentration of the hydroxyl radical (OH), controls the lifetime of reactive trace gases in the atmosphere such as methane and carbon monoxide (CO). Models tend to underestimate the methane lifetime and CO concentrations throughout the troposphere, which is consistent with excessive OH. Approximately half of the oxidation of methane and non-methane volatile organic compounds (VOCs) is thought to occur over the oceans where oxidant chemistry has received little validation due to a lack of observational constraints. We use observations from the first two deployments of the NASA ATom aircraft campaign during July–August 2016 and January–February 2017 to evaluate the oxidation capacity over the remote oceans and its representation by the GEOS-Chem chemical transport model. The model successfully simulates the magnitude and vertical profile of remote OH within the measurement uncertainties. Comparisons against the drivers of OH production (water vapor, ozone, and NO_y concentrations, ozone photolysis frequencies) also show minimal bias, with the exception of wintertime NO_y. The severe model overestimate of NO_y during this period may indicate insufficient wet scavenging and/or missing loss on sea-salt aerosols. Large uncertainties in these processes require further study to improve simulated NO_y partitioning and removal in the troposphere, but preliminary tests suggest that their overall impact could marginally reduce the model bias in tropospheric OH. During the ATom-1 deployment, OH reactivity (OHR) below 3 km is significantly enhanced, and this is not captured by the sum of its measured components (cOHR_{obs}) or by the model (cOHR_{mod}). This enhancement could suggest missing reactive VOCs but cannot be explained by a comprehensive simulation of both biotic and abiotic ocean sources of VOCs. Additional sources of VOC reactivity in this region are difficult to reconcile with the full suite of ATom measurement constraints. The model generally reproduces the magnitude and

seasonality of $c\text{OHR}_{\text{obs}}$ but underestimates the contribution of oxygenated VOCs, mainly acetaldehyde, which is severely underestimated throughout the troposphere despite its calculated lifetime of less than a day. Missing model acetaldehyde in previous studies was attributed to measurement uncertainties that have been largely resolved. Observations of peroxyacetic acid (PAA) provide new support for remote levels of acetaldehyde. The underestimate in both model acetaldehyde and PAA is present throughout the year in both hemispheres and peaks during Northern Hemisphere summer. The addition of ocean sources of VOCs in the model increases $c\text{OHR}_{\text{mod}}$ by 3% to 9% and improves model–measurement agreement for acetaldehyde, particularly in winter, but cannot resolve the model summertime bias. Doing so would require 100 Tg yr^{-1} of a long-lived unknown precursor throughout the year with significant additional emissions in the Northern Hemisphere summer. Improving the model bias for remote acetaldehyde and PAA is unlikely to fully resolve previously reported model global biases in OH and methane lifetime, suggesting that future work should examine the sources and sinks of OH over land.

1 Introduction

The hydroxyl radical (OH) is the main oxidant responsible for removing trace gases from the atmosphere, and its concentration defines the tropospheric oxidation capacity. OH is primarily produced by the photolysis of ozone in the presence of water vapor. The lifetimes of key atmospheric trace gases are governed by how quickly they are removed by reaction with OH. Oxidation of volatile organic compounds (VOCs) by OH produces tropospheric ozone and fine particulate matter which are detrimental to human health and vegetation and impact climate. The oxidation of VOCs, carbon monoxide (CO), and methane provides the main sink of OH in the troposphere. Oxidation of methane and VOCs accounts for over half of the global CO production (Duncan et al., 2007; Safieddine et al., 2017), resulting in a tight coupling of these compounds.

Models generally overestimate global mean tropospheric OH and its ratio in the Northern Hemisphere to Southern Hemisphere (Naik et al., 2013; Patra et al., 2014). These biases may be linked to the persistent CO underestimate in models (Shindell et al., 2006), as prescribing OH from observations improves simulated CO (Müller et al., 2018). However, constraining models with observations of ozone and water vapor cannot resolve biases in model OH (Strode et al., 2015), which is impacted by additional complex factors such as the chemical mechanism and the ozone photolysis frequency (Nicely et al., 2017). Constraining the performance of model chemical mechanisms has largely focused on regions of strong biogenic and anthropogenic activity (Emmerson and Evans, 2009; Yu et al., 2010; Marvin et al., 2017), but at least half of the oxidation of methane occurs over the ocean, where models have received little evaluation due to a lack of observational constraints.

The introduction of airborne measurements of OH reactivity (OHR) provides a method to evaluate the sink of OH across a range of altitudes and a variety of locations and chemical environments (Mao et al., 2009; Thames et al., 2020). Previous work compared surface observations of OHR at a single site to the sum of individually calculated OHR components from measurements (Di Carlo, 2004; Yoshino et al., 2006; Sinha et al., 2008, 2010; Mao et al., 2010; Dolgorouky et al., 2012; Hansen et al., 2014; Nakashima et al., 2014; Nölscher et

al., 2012, 2016; Ramasamy et al., 2016; Zannoni et al., 2016, 2017) or from simple models (Ren et al., 2006; Lee et al., 2009; Lou et al., 2010; Mogensen et al., 2011; Mao et al., 2012; Edwards et al., 2013; Kaiser et al., 2016; Whalley et al., 2016). Thames et al. (2020) found evidence of missing OHR between measurements and an observationally constrained box model during the first three ATom deployments. Chen et al. (2019) compared calculated OHR from a global model to OHR determined from a suite of VOCs but did not have measurements of OHR itself. Ferracci et al. (2018) found that missing OHR estimated from surface observations could result in a small increase in the methane lifetime in a global model. Safieddine et al. (2017) and Lelieveld et al. (2016) presented the first global model simulations of OHR but with only qualitative comparison to observations. No study has quantitatively compared simulated and observed OHR in a global model in an effort to constrain the OH sink.

The ATom campaign (Wofsy et al., 2018) provides an unprecedented opportunity to test models in the remote atmosphere with a detailed suite of chemical observations. We simulate the first two deployments (ATom-1: July–August 2016, ATom-2: January–February 2017) using the GEOS-Chem chemical transport model (CTM) as our tool to explore potential sources of systematic errors that could explain the community-wide model overestimate in global mean OH and underestimate of the methane lifetime. We include model evaluation with measurements of OHR, a relatively new constraint available for assessing atmospheric oxidation capacity. To our knowledge, this is the first quantitative use of this measurement to evaluate a CTM.

2 Description of model and observations

2.1 The GEOS-Chem model

We use the GEOS-Chem global 3-D CTM in v12.3.0 (<http://www.geos-chem.org>, last access: 2 July 2020) driven by assimilated meteorological data from the Goddard Earth Observing System Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2; Gelaro et al., 2017). The native MERRA-2 model has a horizontal resolution of $0.5^\circ \times 0.625^\circ$ and 72 vertical levels which we degrade to $2^\circ \times 2.5^\circ$ and 47 vertical levels for use in GEOS-Chem. The midpoint of the first model layer is 58 m. We use time steps of 20 min for chemistry and 10 min for transport as recommended by Philip et al. (2016). GEOS-Chem includes detailed treatment of HO_x – NO_x –VOC–halogen–aerosol chemistry, with recent improvements for isoprene (Chan Miller et al., 2017; Fisher et al., 2016; Marais et al., 2016; Travis et al., 2016), peroxyacetyl nitrate (PAN) (Fischer et al., 2014), and halogen chemistry (Sherwen et al., 2016). The production of organic aerosols is calculated using fixed yields from isoprene, monoterpenes, biomass burning, and anthropogenic fuel combustion (Pai et al., 2020). Aerosol uptake of HO_2 is parameterized with a reactive uptake coefficient (γ) of 0.2 (Jacob, 2000) to produce H_2O (Mao et al., 2013). Aerosol thermodynamic equilibrium is calculated by ISORROPIA II v2.0 (Pye et al., 2009). Surface methane concentrations are prescribed monthly using spatially interpolated observations from the NOAA GMD flask network. We simulate the 2016–2017 period with an 18-month initialization.

Global fire emissions at 3-hourly resolution (Mu et al., 2011) for 2016 and 2017 are from the Global Fire Emissions Database (GFED4s; van der Werf et al., 2017). The GFED4s burned area (Giglio et al., 2013) includes a parameterization of small fires (Randerson et al., 2012). Biogenic VOC emissions are from MEGANv2.1 (Guenther et al., 2012; Hu et al., 2015). Global anthropogenic emissions are from the Community Emissions Data System (CEDS) inventory (Hoesly et al., 2018), overwritten by ethanol from the POET inventory (Olivier et al., 2003; Granier et al., 2005), ethane from Tzompa-Sosa et al. (2017), and regional inventories for the United States (NEI11v1, Travis et al., 2016), Canada (CAC, <https://www.canada.ca/en/services/environment/pollution-waste-management/national-pollutant-release-inventory.html>, last access: 31 July 2013), Mexico (BRAVO, Kuhns et al., 2003), Europe (EMEP, <http://www.emep.int/index.html>, last access: 31 March 2015), Asia (MIX, Li et al., 2017), and Africa (DICE, Marais and Wiedinmyer, 2016). Lightning emissions are constrained with satellite data according to Murray et al. (2012) with a global flash rate of 280 mol NO flash⁻¹ (Marais et al., 2018). Air–sea exchange is calculated for acetaldehyde (Millet et al., 2010), acetone (Fischer et al., 2012), and dimethyl sulfide (Breider et al., 2017). All emissions are processed by the Harvard Emissions Component (HEMCO, Keller et al., 2014). Table 1 gives the 2016 emission budget for CO and NO_x.

The standard simulation includes prescribed methanol concentrations. We expand this simulation to include emissions and chemistry for methanol as well as unsaturated C₂ compounds. Air–sea exchange of methanol is specified using the methodology of Millet et al. (2008) with a constant seawater concentration of 142 nM. Terrestrial biogenic methanol emissions are from MEGANv2.1, and anthropogenic and biomass burning emissions are from the inventories described above. We likewise include biomass burning and anthropogenic emissions of ethyne (C₂H₂) and ethene (C₂H₄) along with terrestrial biogenic emissions of C₂H₄. Oxidation of C₂H₂ by OH proceeds according to the Master Chemical Mechanism (MCM) v3.3.1 (Jenkin et al., 1997, 2015; Saunders et al., 2003), via <http://mcm.leeds.ac.uk/MCM> (last access: 8 February 2018). Simplified C₂H₄ chemistry is included based on Lamarque et al. (2012) with an updated OH rate constant from the MCM v3.3.1. Table S1 in the Supplement shows the reactions and species included for unsaturated C₂ compounds. The standard model does not consider the OH reactivity of a subset of organic acids (RCOOH) from the oxidation of VOCs. We implement oxidation of RCOOH and evaluate the impact of excluding this species, which is minor, in Table S2 and Fig. S1 in the Supplement. The model concentration of H₂ is fixed at 500 ppt, consistent with observed H₂ from ATom-1 and ATom-2 (520 ppt).

The GEOS-Chem global mean tropospheric OH ([OH]_{GM}) is calculated as an air-mass-weighted quantity below the model tropopause (see http://wiki.seas.harvard.edu/geos-chem/index.php/Mean_OH_concentration, last access: 4 May 2020, for the calculation methodology). The [OH]_{GM} for 2016 is 11.9 × 10⁵ molecules cm⁻³ and the corresponding methane lifetime (τ_{CH₄}) is 9.0 years. This result is comparable to the multi-model [OH]_{GM} of 11.1 × 10⁵ molecules cm⁻³ and τ_{CH₄} of 9.7 years from Naik et al. (2013). The best observationally derived estimate of τ_{CH₄} is 11.2 ± 1.3 years (Prather et al., 2012), suggesting a model bias here of 20%. We calculate the ratio of total 2016 air-mass-weighted OH in the Northern (> 0° N) to Southern Hemisphere (< 0° S) to be 1.14. This exceeds observationally

derived ratios of 0.85 to 0.97 (Montzka et al., 2000; Patra et al., 2014) but is at the low end of previous model estimates ranging from 1.13 to 1.42 (Naik et al., 2013).

2.2 Calculated OH reactivity

The atmosphere contains thousands of reactive organic compounds (Goldstein and Galbally, 2007). Transforming the concentrations of these compounds and reactive inorganics to calculated OH reactivity (cOHR) ranks them in order of their importance as OH sinks. The cOHR from a model (cOHR_{mod}) can then be compared to cOHR from a suite of measurements (cOHR_{obs}) where cOHR is defined by Eq. (1).

$$\text{cOHR}(\text{s}^{-1}) = k_{\text{OH,CH}_4}[\text{CH}_4] + k_{\text{OH,CO}}[\text{CO}] + k_{\text{OH,NO}_2}[\text{NO}_2] + \sum k_{\text{OH,VOC}}[\text{VOC}] + \dots \quad (1)$$

Figure 1a shows annual surface cOHR_{mod} for the year 2016 based on the 90 components listed in Table S3. Figure 1b shows the zonal mean profile below 12 km. Approximately 80% of air-mass-weighted cOHR_{mod} resides below 3 km. The average annual surface cOHR_{mod} is 1.8 s⁻¹, with 40% present over the ocean (average of 1.0 s⁻¹). Higher cOHR_{mod} occurs in coastal outflow regions and the lowest cOHR_{mod} is present over the Southern Ocean. The maximum cOHR_{mod} (48 s⁻¹) over northern China is due to high concentrations of SO₂, NO_x, and CO. In the tropics, elevated cOHR_{mod} is mainly from isoprene, other biogenic species, and CO.

2.3 ATom observations

The NASA ATom field campaign (Wofsy et al., 2018) sampled the remote troposphere with the DC-8 aircraft over the Atlantic and Pacific oceans from approximately 200 m to 12 km altitude in four seasons from 2016 to 2018 with a goal of improving the representation of trace gases and short-lived greenhouse gases in models of atmospheric chemistry and climate. We use data here from the first two deployments (ATom-1 and ATom-2), which sampled winter and summer conditions in each hemisphere. We consider only observations over the ocean (73% of measurements). Flight tracks for ATom-1 with land crossings removed are shown in Fig. 2; ATom-2 flight tracks are nearly identical. We sample the model along the flight tracks, and both the model and observations are averaged to the model grid and time step for all the following comparisons. The aircraft carried an extensive chemical payload including observations of water vapor, methane, CO, OH, NO_x, VOCs, photolysis frequencies, and OHR. Table 2 describes the observations used in this work.

3 Comparison of simulated and measured OH

We compare observed and simulated OH concentrations to evaluate whether differences are consistent with the bias in τ_{CH_4} discussed in Sect. 2.1. Figure 3 shows modeled OH sampled along the flight tracks and compared to observed OH (Table 2) for ATom-1 (boreal summer 2016) and ATom-2 (boreal winter 2017) in each hemisphere from the lowest sampled altitude (~ 200 m) to 10 km. There is no evidence of a systematic overestimate in modeled OH throughout the troposphere. Figure S2 shows similarly good agreement across the observed frequency distributions of OH concentration. A model OH overestimate is apparent in the lowest 2 km in the Northern Hemisphere summer that could indicate excessive OH

production or an underestimated sink from emissions of ocean VOCs. Global models tend to overestimate OH against constraints from methyl chloroform observations (Shindell et al., 2006; Naik et al., 2013; Nicely et al., 2017), but we find here that tropospheric OH is successfully simulated within observational uncertainty (74% to 135%, 2σ confidence level). This result from a global CTM is consistent with good agreement between OH measurements and a box model during NASA's Pacific Exploratory Mission – Tropics (PEM-Tropic B) campaign in the clean remote Pacific (Tan et al., 2001) and a similar analysis by Brune et al. (2020) for ATom 1 through 4.

We calculate the median air-mass-weighted column average OH (OH_{col}) from the median OH concentrations in Fig. 3 and the total tropospheric air mass over the ocean. During ATom-1, the modeled OH_{col} in the Northern (Southern) Hemisphere is $4.5(1.4) \times 10^6$ molecules cm^{-3} compared against the observations of $4.4(1.1) \times 10^6$ molecules cm^{-3} during ATom-1. Similarly, during ATom-2, OH_{col} is $0.8(2.8) \times 10^6$ molecules cm^{-3} in the model and $0.9(2.6) \times 10^6$ molecules cm^{-3} in the observations. Median model OH_{col} is within 30% of observations during both deployments, with the smallest bias in the total column during Northern Hemisphere summer when OH is at a maximum. As discussed above, model OH is overestimated in the lowest 2 km during this period, but this bias is minimized in the column average. The observed air-mass-weighted ratio of Northern to Southern Hemisphere OH, calculated in the same manner as described in Sect. 2, is 2.8 during ATom-1 and 0.2 during ATom-2, indicating a strong seasonality that the model largely reproduces (ratios of 2.3 and 0.2). This ratio is less than the ratio of OH_{col} because there is approximately 30% less air mass over the ocean in the Northern Hemisphere ocean than over the Southern Hemisphere. This seasonality is masked by calculations performed on an annual mean basis. The seasonality in this ratio reported by Wolfe et al. (2019) for satellite-derived OH during ATom-1 and ATom-2 is more modest because they calculate a daily average OH that extends to the tropopause, while here, we use largely daytime aircraft observations below 10 km.

The model is in good agreement with OH measurements during ATom, but the uncertainty in the observations is similar to a recent estimate of the GEOS-Chem model uncertainty for OH of 25% to 40% (Christian et al., 2018). In addition, the lifetime of OH is short (seconds) and atmospheric concentrations are highly variable; thus, direct model comparison to measured OH is insufficient to demonstrate model skill in capturing the broader remote oxidation capacity. Agreement between the model and observations could also result from compensating errors in the OH source and sink. We support the model comparison in Fig. 3 with an evaluation of the key factors governing OH production and loss measured by ATom and investigate potential missing sources of VOCs from the ocean during summertime.

4 Constraints on the remote source of OH

In the remote troposphere, OH is primarily produced from the photolysis of ozone in the presence of water vapor (Monks, 2005) and is enhanced by nitrogen oxides (NO_x) from lightning and transport from continental sources. Methane, CO, and VOCs provide the main OH sinks (Murray et al., 2014). We compare the model to ATom-1 and ATom-2 observations of the drivers of the tropospheric OH source (water vapor, ozone, ozone photolysis frequency, NO_x) to determine possible broader sources of model bias.

Figure 4 compares observations of water vapor mixing ratios to the NASA MERRA-2 reanalysis product used by the model. MERRA-2 is generally successful at reproducing observed tropospheric water vapor (Gelaro et al., 2017), and we also find good agreement compared with ATom-1 and ATom-2 observations throughout the troposphere. We evaluate the model treatment of the incoming actinic flux and the resulting ozone photolysis frequency ($j(\text{O}^1\text{D})$) in Fig. 5. Hall et al. (2018) showed that GEOS-Chem actinic fluxes in both cloudy and clear skies were well simulated during the ATom-1 deployment. Figure 5 confirms the minimal model bias in $j(\text{O}^1\text{D})$ and successful representation of the observed seasonality with median summertime values below 3 km ($\sim 4 \times 10^{-5} \text{ s}^{-1}$) approximately 4 times higher than in winter ($\sim 1 \times 10^{-5} \text{ s}^{-1}$).

The GEOS-Chem ozone simulation has been extensively tested against ozonesondes, aircraft, and satellite observations and shows no systematic overestimates (Hu et al., 2017), with the exception of continental surface concentrations (Fiore et al., 2009; Travis et al., 2016). Figure 6 shows that the highest (54–63 ppb) and lowest (14 ppb) tropospheric ozone observed during ATom-1 and ATom-2 occurs during summer in the mid to upper troposphere and marine boundary layer, respectively. Ozone is less variable in wintertime, with values between 30 and 50 ppb. The model generally reproduces the magnitude and shape of the tropospheric ozone profiles as well as the seasonality observed during both deployments. There is no evidence of the systematic Northern Hemisphere ozone bias previously seen in global model evaluations (Young et al., 2013) that was suggested as a cause of excessive OH (Naik et al., 2013). This may be reflected in the improved model interhemispheric OH ratio (Sect. 2.1) seen here over previous studies. Upper tropospheric ozone is overestimated in all cases but Northern Hemisphere summer, but this would not have a large influence on primary OH production (or the methane lifetime) at these altitudes (Brune et al., 2020).

OH is enhanced in the presence of NO_x ($\equiv \text{NO} + \text{NO}_2$). We use NO_y here (Fig. 7a) as our constraint as observed NO_2 was generally near the detection limit in both deployments. We also show NO (Fig. 7b) given its role in secondary OH production. The model reproduces the maximum in NO_y that occurs in the Northern Hemisphere upper troposphere in summertime due to lightning (Marais et al., 2018). Observations show little variability between summer and winter NO_y in the lower troposphere. Southern Hemisphere NO_y is underestimated in the lowest few kilometers in both seasons, which could be due to missing ocean production of methyl nitrate (Fisher et al., 2018). The largest model discrepancy is an overestimate of approximately 70% in the Northern Hemisphere wintertime. Observations of NO reflect the structure of NO_y , with the exception of Northern Hemisphere winter.

Causes of the remote model bias in NO_y

Figure 8 shows that the model NO_y overestimate in winter is primarily caused by nitric acid (HNO_3). Excessive remote HNO_3 is a long-standing model deficiency (Bey et al., 2001; Staudt et al., 2003; Brunner et al., 2003, 2005). The model bias identified here is unlikely to result from overestimated continental emissions due to the short lifetime of NO_y against deposition (~ 3 d in the Northern Hemisphere winter). Models suggest that less than 40% of emitted NO_x in the US is exported downwind (Dentener et al., 2006; Zhang et al., 2012). However, the standard model configuration here does not address the large possible bias in

the US anthropogenic NO_x inventory of ~ 40% (Anderson et al., 2014; Travis et al., 2016) or the downward trend in NO_x emissions from Asia of ~ 30% since 2011 (Krotkov et al., 2016). As expected, scaling Asian and US NO_x emissions by these percentages improves the model bias in winter by only 15% below 3 km (Fig. 8). Recent improvements to the simulation of continental wintertime HNO_3 (Jaeglé et al., 2018) would similarly be expected to have a marginal effect in our study region.

Kasibhatla et al. (2018) showed that acid displacement of chloride (Cl^-) by HNO_3 on sea-salt aerosols (SSA) could resolve model overestimates of gas-phase HNO_3 in the marine boundary layer using the GEOS-Chem model. A more comprehensive simulation of this process was developed by X. Wang et al. (2019). Figure 8 shows sensitivity tests with the mechanism from X. Wang et al. (2019) incorporated into our simulation in the Northern Hemisphere winter. Model HNO_3 decreases by approximately 100 ppt below 3 km, which would significantly improve the wintertime NO_y bias in this region, but the free tropospheric bias remains. The displacement of Cl^- described above generates particulate nitrate on coarse-mode SSA (NITs). Photolysis of nitrate has been proposed as a source of NO_x to the marine boundary layer (Ye et al., 2016; Romer et al., 2018), which might increase HNO_3 . We include NITs photolysis at a frequency of 50 times that of HNO_3 (Kasibhatla et al., 2018). Figure 8 shows that this mechanism is consistent with observations of NO and ozone below 3 km and does not increase HNO_3 but increases the free tropospheric NO_y bias due to PAN formation and exacerbates the overestimate in upper tropospheric ozone during this season.

The difficulty in resolving the bias in wintertime may be due to an overestimate of the NO_y lifetime as demonstrated by our sensitivities discussed above. Luo et al. (2019) proposed a new treatment of model wet scavenging using spatially and temporally varying cloud condensation water content and an empirical description of HNO_3 wet removal. This scheme drastically reduced persistent model biases in nitric acid and nitrate at the surface in the United States (Zhang et al., 2012; Heald et al., 2012). As shown in Fig. 8, the revised wet scavenging scheme could fully resolve the remote bias in HNO_3 throughout the troposphere. However, this parameterization has only received testing over the surface of the continental US, and more evaluation is needed before it can be adopted widely in models.

We find that scaling NO_x , implementing chlorine chemistry, and revised wet scavenging (except in Northern Hemisphere winter) have negative impacts on the modeled OH_{col} along the flight tracks of -1%, -7%, and -4%, respectively. The addition of NITs photolysis to the chlorine chemistry simulation increases OH_{col} by 11% over the base model. In Northern Hemisphere winter only, revised wet scavenging increases OH_{col} by 16%, possibly due to the effect of reduced heterogeneous chemistry. Overall, the annual mean impact of revised wet scavenging from our preliminary sensitivity tests is a -3% reduction in global mean air-mass-weighted OH and a +2% increase in the model methane lifetime. These preliminary sensitivities suggest that resolving the model wintertime NO_y bias in the Northern Hemisphere could marginally reduce the overestimates of global mean OH on an annual basis if the photolysis frequency of NITs is smaller than 50 times the rate of HNO_3 photolysis. Recent work from the NASA KORUS-AQ field campaign found that a rate of 1 to 30 might be more consistent with observational constraints (Romer et al., 2018).

Overall, the main drivers of remote tropospheric OH production in our base-case simulation are in good agreement with observations from the first two ATom deployments, with the exception of an NO_y overestimate in the Northern Hemisphere wintertime. Acid displacement of Cl^- by HNO_3 on SSA (Kasibhatla et al., 2018; X. Wang et al., 2019) may somewhat improve remote HNO_3 below 3 km, but if the resulting NITs undergoes photolysis at a rate of 50 times that of HNO_3 (Kasibhatla et al., 2018), the impact on remote NO_y may be lessened due to the formation of PAN. Both mechanisms require significant further study as tropospheric halogen sources and chemistry and the photolysis frequency of NITs are highly uncertain. A new parameterization of wet scavenging (Luo et al., 2019) would greatly improve modeled remote HNO_3 and NO_y but requires further testing and evaluation of its broader impacts on atmospheric chemistry.

5 Constraints on the remote sink of OH

The primary sinks of tropospheric OH are CO, methane, and VOCs; OHR measurements represent the sum effect of these species. Previous aircraft measurements of OHR provided evidence of missing reactivity in the remote atmosphere linked to unknown highly reactive VOCs (Mao et al., 2009). During ATom, Thames et al. (2020) measured OHR over the Atlantic and Pacific oceans and determined that missing OHR also correlated with oxygenated VOCs, suggesting the presence of unknown ocean emissions. We compare directly measured OHR during the ATom-1 and ATom-2 deployments to calculated OHR (cOHR_{obs}) according to Eq. (1) from the full ATom measurement suite and from the model (cOHR_{mod}) sampled along the flight path. Table 2 describes the observations used to calculate cOHR.

Figure 9 shows the comparison of OHR and cOHR from the model and observations. The observed cOHR is typically less than observed OHR. Along the flight tracks, cOHR_{obs} and cOHR_{mod} show good agreement and strong correlation ($r^2 = 0.95$ for ATom-1 and ATom-2). The model underestimates cOHR_{obs} by 10% to 12% in the lowest 3 km; we discuss this difference below. The measured relationship between OHR and cOHR_{obs} is weaker ($r^2 = 0.53$ for ATom-1, $r^2 = 0.56$ for ATom-2) and cOHR_{obs} is less than OHR below 3 km by 0.2 to 0.4 s^{-1} . Thames et al. (2020) showed that median missing reactivity (between OHR and an observationally constrained box model) below 4 km during the ATom-1, ATom-2, and ATom-3 deployments was between 0.2 and 0.8 s^{-1} . They provided statistical evidence that while near the level of the instrument accuracy, missing OHR in the marine boundary layer was statistically significant. We find that missing OHR is not associated with acetonitrile or CO ($r^2 < 0.06$), indicating that biomass burning is not the cause. Acetaldehyde in Northern Hemisphere summer has the strongest relationship with missing OHR ($r^2 = 0.19$, p -value $\ll 0.01$, Fig. S3), which suggests a potential role for unmeasured reactive VOCs or their oxidation products from the ocean, as also suggested by Read et al. (2012) and Thames et al. (2020).

Ocean emissions of VOCs have been suggested as a source of remote secondary organic aerosols (Gantt et al., 2010; Kim et al., 2017; Mungall et al., 2017), but their impact on remote reactivity has not been quantified. Our base simulation, described in Sect. 2.1, only includes air–sea exchange of acetone, acetaldehyde, methanol, and dimethyl sulfide. We

determine whether additional compounds emitted from the ocean, but not generally included in models, could increase cOHR_{mod} and reconcile the observed discrepancy described above. We follow the standard methodology for air–sea exchange described in Millet et al. (2008) to include emission of the species listed in Table 3 using available measured seawater concentrations, with the addition of isoprene implemented as a direct emission according to Arnold et al. (2009). As shown in Table 3, air–sea exchange represents a net sink of VOCs on an annual basis (-75 Tg C yr^{-1}), but this is mainly due to ocean uptake of acetone, which is a negligible component of cOHR .

Interfacial photochemistry may provide an additional abiotic source of VOCs from the ocean. We model abiotic ocean emissions of VOCs according to Brüggemann et al. (2018) by applying species-specific emission factors to the monthly ocean photochemical potential derived in their study. We use the emission factor appropriate for the upper bound of this source according to Brüggemann et al. (2017) (Table S4). Table 4 provides a breakdown of these additional VOCs with a total annual emission of 28 Tg C yr^{-1} .

Figure 10 shows the annual mean impact of all ocean emissions described in Tables 3 and 4 (including an adjustment to the acetaldehyde seawater concentration described below in Sect. 5.1) on cOHR_{mod} by turning off those ocean sources in a 1-year simulation. Average annual surface cOHR_{mod} over the ocean increases by 6% over the base simulation and 12% over the simulation with no ocean emissions. The largest increases occur in regions of higher biogenic activity along coastlines and in the Southern Ocean due to the adjustment to acetaldehyde emissions discussed in Sect. 5.1. The incremental impact of the additional ocean emissions over the base simulation is shown in Fig. S4. Without any ocean emissions, global mean OH would be 2% greater than in the case with comprehensive treatment of ocean VOCs. Figure 9 shows that along the flight tracks, cOHR_{mod} increases below 3 km by 3% to 9%, which reduces the model bias against cOHR_{obs} . However, the majority of the added species (Tables 3 and 4) were measured during ATom, would therefore contribute to cOHR_{obs} , and cannot explain the gap in OHR.

We evaluate the impact of further expanding the oceanic source of reactive VOCs to reconcile the discrepancy between cOHR_{obs} and OHR in a similar manner to Mao et al. (2009). Here, we test a source of alkanes as previously suggested by Read et al. (2012), using the model species ALK4 ($> \text{C}_4$ alkanes) that has a calculated lifetime of less than 2 d in the Northern Hemisphere summer ($k_{\text{OH}} 2.3 \times 10^{-12} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$ at 298 K). Known alkanes have been measured in seawater (Plass-Dülmer et al., 1993), but the implied source is small. Consequently, we use ALK4 for testing only. Generating the missing OHR in this way requires an implausibly large oceanic ALK4 source of approximately 340 Tg C yr^{-1} compared against all other sources of VOCs in the model (Tables 3 and 4). A sensitivity test with this source, shown in Fig. 9, largely closes the gap between cOHR_{mod} and OHR but would result in a 20% to 50% reduction in OH below 3 km, biasing the model OH simulation (Fig. 3) and degrading model NO_y (Fig. 7) due to increased PAN formation.

Thames et al. (2020) found that a partial recycling of OH would be required to maintain consistency with observed OH and HO_2 during ATom when adding an unknown source of reactivity. If the unknown VOC we suggest includes some OH recycling in its oxidation

mechanism and does not produce PAN, the model bias in OH could be mitigated. We use isoprene as our test of a more reactive VOC that includes OH recycling by scaling the ALK4 emission source by the reaction rate of isoprene with OH to obtain a more reasonable emission source of approximately 9 Tg C yr⁻¹. Figure 9 shows that this source actually has a minimal impact on cOHR_{mod} of no more than 0.1 s⁻¹. Only one-third in summer and two-thirds in winter of the additional cOHR_{mod} from the ocean source of ALK4 are attributable to ALK4; the rest is due to CO, acetaldehyde, and other aldehydes from both increased chemical production and longer lifetimes from suppressed OH. Therefore, a larger source of even a reactive VOC like isoprene is required to close the gap in missing OHR. Reconciling cOHR_{mod} and OHR is therefore difficult using the existing suite of ATom measurement constraints and possible known precursors; further investigation of the accuracy of the OHR measurements in challenging remote conditions may be needed.

We also assess whether the model accurately represents the components of cOHR_{obs} and explore potential additional sources of missing cOHR_{mod}. Figures 11 and 12 show the components of median cOHR in the base simulation below 3 km for each deployment. The composition of cOHR_{mod} is generally consistent with cOHR_{obs}. CO and methane make up half or greater of both cOHR_{obs} and cOHR_{mod}. There is no systematic underestimate in CO reactivity as might be expected from the general model bias described by Shindell et al. (2006), with the exception of a 9% underestimate during Northern Hemisphere winter when the lifetime of CO is longer and biases in continental sources could have a larger impact. During the ATom-1 deployment, cOHR_{obs} is 50% higher in the Northern Hemisphere (summer) than in the Southern Hemisphere (winter) primarily due to the increase in methyl hydroperoxide (MHP) concentrations. During the ATom-2 deployment, cOHR_{obs} is 60% higher in the Northern Hemisphere (winter) than in the Southern Hemisphere (summer) due to the large contribution of CO in Northern Hemisphere wintertime. The model successfully represents the observed seasonality during both deployments but underestimates cOHR_{obs} by 12% in the Northern Hemisphere and 9% in the Southern Hemisphere.

The difference between measured and simulated cOHR is mainly due to differences between measured and simulated OVOCs. These compounds contribute on average 25% to cOHR_{obs} but 17% to cOHR_{mod}. The largest difference in reactivity is from acetaldehyde. Differences between simulated and measured MHP (Fig. S5) are also important and could reflect an error in the MHP lifetime (Müller et al., 2016). However, interferences in the MHP measurement in the boundary layer (Supplement, Sect. S6) have yet to be resolved, and therefore we do not further evaluate causes of underestimated MHP here. We do consider potential missing sources of model acetaldehyde constrained by the ATom measurements over the ocean and assess their impact on simulated OH and CO in Sect. 5.1.

Evaluation of the remote sources of acetaldehyde

Inability to reconcile remote acetaldehyde observations with models is a long-standing problem (Singh et al., 2001., 2003; Millet et al., 2010; Nicely et al., 2016). Singh et al. (2001) proposed that a large, diffuse, and as-yet unknown source of OVOCs such as acetaldehyde must exist in the troposphere to solve this discrepancy. Read et al. (2012) determined that missing cOHR_{mod} from OVOCs (mainly acetaldehyde) in the marine

tropical atmosphere, possibly from terrestrial or ocean sources of alkanes, could cause up to an 8% underestimation of the methane lifetime. Nicely et al. (2016) showed that constraining a box model with observed acetaldehyde reduced tropospheric column OH by 9% and that this acetaldehyde bias was present across eight different CTMs. Therefore, understanding the source of missing acetaldehyde may be part of the cause of the multi-model bias in the methane lifetime and global mean OH.

Figure 13 compares the model simulation of acetaldehyde against observations. Average observed concentrations peak in the Northern Hemisphere during ATom-1 with an average mixing ratio of 230 ppt below 3 km and 100 ppt above 3 km despite a lifetime of only several hours in summer. The maximum model underestimate occurs during this period. Observed concentrations are at a minimum during the ATom-2 deployment, indicating a strong seasonality in the source. In each deployment, concentrations remain as high as 70 to 100 ppt as far south as 60° S (Fig. S6), which the model does not reproduce. There is no apparent difference in model bias between observations over the Atlantic or Pacific Ocean (Fig. S7). The model underestimates acetaldehyde on average by 60 to 90% (50 to 200 ppt) below 3 km and does not capture the observed elevated levels throughout the troposphere.

In earlier studies, measurement uncertainties prevented interpretation of model–measurement disagreements in the remote atmosphere, including difficulties in background subtraction (Apel et al., 2008), with uncertainties as high as 70 ppt (Apel et al., 2003), which hindered analysis of clean conditions. The ATom measurement uncertainty is reduced to 10 ppt/20% (Table 2) and does not have the biases present in previous campaigns (S. Wang et al., 2019). Studies have also disputed whether observed acetaldehyde was compatible with observed PAN due to the significant role of acetaldehyde as a PAN precursor through production of the peroxyacetyl (PA) radical (Singh et al., 2001, 2003; Millet et al., 2010). Global simulations estimate that acetaldehyde is responsible for approximately 40% of PA radical production (Fischer et al., 2014), which would be even larger if acetaldehyde is severely underestimated by models. Reaction of the PA radical with HO₂ is more prevalent in remote environments and produces peroxyacetic acid (PAA) preferentially over PAN, making PAA a more useful constraint for the conditions sampled by ATom. Figure 14 shows the average model underestimate of PAA below 3 km of 70% to 90% (60 to 250 ppt). The model biases for PAA and acetaldehyde both peak with similar magnitude during Northern Hemisphere summer. Figure 15 shows the model comparison with PAN, which is generally well simulated during this period.

S. Wang et al. (2019) used an observationally constrained box model to show that the levels of acetaldehyde observed during ATom are required to explain the observed PAA. The reaction rate of PAA + OH may be 3 times larger (Wu et al., 2017) than the maximum value used by S. Wang et al. (2019), which could result in even better agreement between PAA and acetaldehyde in the marine boundary layer. We evaluate the standard GEOS-Chem acetaldehyde budget, described in detail by Millet et al. (2010), against available ATom observations. The 2016 model budget for the base simulation is provided in Table 5. Acetaldehyde is produced from oxidation of VOCs (ethane, propane, C₄ alkanes, C₃ alkenes, isoprene, ethanol) and is directly emitted from the ocean, terrestrial plant growth, biomass burning, and anthropogenic activities. The model parameterization of acetaldehyde

ocean emissions is dependent on satellite-based observations of colored dissolved organic matter (CDOM) (Millet et al., 2010).

The model free tropospheric bias suggests that long-lived oxidation of VOCs must be underestimated due to the short lifetime of acetaldehyde (< 1 d). The longest-lived precursor VOCs in the model are ethane (2 months) and propane (2 weeks). Ethane has the highest concentration of any measured non-methane VOC during ATom, with an average of 1.5 ppb below 3 km during the Northern Hemisphere winter. The model underestimates average ethane and propane below 10 km by approximately 25% and 60%, respectively (Figs. S8 and S9), which could be due to underestimated natural geologic and fossil fuel emissions (Dalsøren et al., 2018). However, the oxidation of these species is too slow to provide the missing model acetaldehyde and would only marginally increase remote background levels even if it was produced at higher yield at low NO_x (model yields are ~ 50% for ethane and ~ 20% for propane, Millet et al., 2010). The chemical mechanism used for these species is provided in Table S5. One or more precursors able to resolve the model acetaldehyde bias must therefore be present at higher cumulative concentrations than ethane or propane. Modeled ALK4, parameterized as a butane–pentane mixture, maintains a high acetaldehyde yield at low NO_x and has a shorter lifetime (~ 5 d), contributing to a larger perturbation to atmospheric acetaldehyde levels than ethane or propane for a given concentration change. The sensitivity test adding substantial ALK4 emissions from the ocean described in Sect. 4 would not resolve the free tropospheric bias in the Northern Hemisphere but would result in a 40% overestimate below 1 km. Furthermore, ALK4 is too short-lived to substantially perturb the remote atmosphere from a continental source; thus, the potential missing acetaldehyde precursors (from either a marine or terrestrial source) must have a longer lifetime.

As shown in Table 5, primary ocean emissions of acetaldehyde in the base simulation (22 Tg yr^{-1}) are lower than previous work (57 Tg yr^{-1}), likely due to updates to the model parameterization of the water transfer velocity (Johnson, 2010). Additional independent estimates of the ocean source are also much larger (34 to 42 Tg yr^{-1} , Read et al., 2012; S. Wang et al., 2019). However, an increased primary ocean source would not address the bias in the free troposphere or in winter when biogenic activity from CDOM is zero in the model at high latitudes. Ship-borne measurements generally measure non-zero acetaldehyde seawater concentrations of approximately 5 nM (Read et al., 2012), and a recent trans-Atlantic campaign found that acetaldehyde concentrations from 47° S to 50° N did not always correlate with levels of CDOM (Yang et al., 2014). Therefore, we set a minimum seawater concentration of 5 nM in the model parameterization regardless of CDOM level. This change adds 2 Tg C yr^{-1} in emissions and increases concentrations over the remote ocean in winter by up to 50 ppt.

Figure 13 shows the combined effect of adding new ocean VOCs in Sect. 5 and improving the seawater parameterization described above on modeled acetaldehyde (labeled “Improve Ocean VOCs”). Although the direct ocean source in this work is lower than previous estimates as described above, the secondary source from precursor VOCs is enhanced. Of the additional marine VOCs described in Sect. 5, 19 Tg C yr^{-1} produce acetaldehyde as an oxidation product (Tables 3 and 4). This is compared to 12 Tg C yr^{-1} of direct emissions in

the base model. These sources substantially increase average modeled acetaldehyde below 3 km, with the largest improvement during winter (40 to 60 ppt) when atmospheric lifetimes are longer and the influence of the ocean can extend aloft. In summer, the average model increase below 3 km is only 10 to 20 ppt due to higher OH concentrations. Recent work over North America suggested that free tropospheric VOCs may be underestimated due to errors in model vertical mixing (Chen et al. 2019), but in Northern Hemisphere summer slower mixing would not be expected to compensate for the short lifetime of acetaldehyde in this region (~ 5 h). Thus, the pervasive model bias in the free troposphere cannot be explained by an increase in known direct or indirect ocean sources.

Photodegradation of organic aerosols (OA) is another potential source of oxygenated VOCs such as acetaldehyde to the troposphere (Kwan et al., 2006; Epstein et al., 2014; Wong et al., 2015; S. Wang et al., 2019). The source of secondary organic aerosols (SOA) is uncertain and has been suggested to be up to 4 times larger than current estimates given an implied underestimate of the photochemical loss term (Hodzic et al., 2016). We test the potential impact of the maximum possible source of acetaldehyde from photochemical loss of OA by increasing the overall model production of SOA by a factor of 4 to maximize the impact of Reaction (R2) below. We apply a photolysis frequency for OA of $4 \times 10^{-4} J_{\text{NO}_2}$ (Hodzic et al., 2015) to Reactions (R1) and (R2) as an upper limit and describe the formulation of Reactions (R1) and (R2) below.



The model species OCPI and SOAS represent the majority of simulated OA in the remote atmosphere. OCPI is aged (hydrophilic) organic carbon (12 g C mol^{-1}) and SOAS is SOA from all emissions categories (150 g mol^{-1}). Both are assumed for the purposes of the sensitivity tests here to have an OA/OC ratio of 2.1. In Reaction (R1), one molecule of carbon (0.5 ALD2) is produced per reaction. In Reaction (R2), one acetaldehyde molecule (ALD2) is produced per reaction. The resulting impact on acetaldehyde is only appreciable in the Northern Hemisphere winter (Fig. 13), when modeled aerosol amounts are highest and the lifetime of acetaldehyde is long. Given that this test represents an upper limit, we conclude that photolysis of organic aerosols cannot provide a sufficient source of acetaldehyde to reconcile the model with observations.

We consider whether an entirely unknown VOC with moderate lifetime and a high yield of acetaldehyde at low NO_x could resolve the free tropospheric model bias. We emit such a species with a lifetime of approximately 1 month against oxidation by OH, emissions of 100 Tg yr^{-1} from either anthropogenic, biomass burning, or ocean sources, and a yield of one acetaldehyde molecule per reaction with OH. We do not test a terrestrial biogenic source here but expect the results would be similar to the biomass burning case. These simulations result in average tropospheric concentrations of 1 to 5 ppb. The effect of the unknown VOC is compatible with the model simulation of OH (unlike the addition of oceanic ALK4 needed to reconcile OHR observations as described in Sect. 5). The maximum cOHR_{mod} of this species is small ($< 0.03 \text{ s}^{-1}$). The impact on modeled acetaldehyde (Fig. 13) is generally

similar across all three source categories due to the long lifetime of this precursor. As shown in Figs. 13 and 14, the addition of this unknown VOC modestly improves the simulation of acetaldehyde and PAA everywhere, but a large residual underestimate in Northern Hemisphere summer remains. The impact on PAN is minor, with the exception of Northern Hemisphere winter (Fig. 15), but this is likely driven by the model overestimate in NO_y (Fig. 7, Sect. 4.1).

Emission inventories of VOCs are known to be incomplete, for example neglecting emissions from volatile consumer products (McDonald et al., 2018) or failing to identify as much as half of emitted VOCs from biomass burning (Akagi et al. 2011), both of which peak in summer. The average emission factor for unidentified VOCs from biomass burning roughly corresponds to 75 Tg yr^{-1} , similar to our sensitivity tests of 100 Tg yr^{-1} described above. However, recent attempts to quantify these unidentified VOCs (Stockwell et al., 2015; Koss et al., 2018) find that newly identified compounds tend to be too reactive to impact the remote atmosphere, as needed here; however, this work is ongoing and future efforts should investigate potential precursors of acetaldehyde that could be transported to the remote atmosphere. The missing source of precursor VOCs would need to have substantial additional summertime emissions above and beyond the sensitivity tests shown in Fig. 13 to address the Northern Hemisphere summertime bias. The required magnitude of this perturbation is difficult to reconcile within known measurement and emission uncertainty constraints.

6 Conclusions

The detailed set of chemical information available from the ATom field campaign provides the most comprehensive dataset ever collected to evaluate models in the remote atmosphere. The sampling strategy of collecting observations throughout the troposphere in multiple seasons is ideally suited for improving our understanding of tropospheric chemistry in a poorly observed region of the atmosphere. We use the first two deployments of the ATom field campaign during July–August 2016 and January–February 2017 to investigate sources of bias in model simulations of OH. Global models such as the GEOS-Chem CTM used here tend to overestimate the loss of methane by OH and underestimate CO, which provides the main tropospheric sink of OH. Comparisons of the model with observations from the first two ATom deployments do not show systematic bias in the simulation of OH or the drivers of remote OH production (water vapor, photolysis of ozone, ozone, and NO_y), with the exception of wintertime NO_y , which is overestimated by 70%.

The model overestimate of wintertime NO_y is largely attributable to nitric acid. This bias is not due to an anthropogenic inventory overestimate but may reflect insufficient wet scavenging as well as loss to sea-salt aerosols by nitric acid, although the former mechanism may be counteracted by photolysis of the resulting nitrate aerosols. The impact of resolving this wintertime NO_y bias is uncertain but could marginally reduce the model overestimate of OH. Future work should improve constraints on these mechanisms, which have all received only preliminary validation, and carefully examine their impact in the context of broader atmospheric chemistry, particularly NO_y partitioning throughout the troposphere.

We present the first comparison of measured OH reactivity (OHR) from aircraft with a global model to evaluate the tropospheric sink of OH. We calculate OH reactivity (cOHR_{obs}) from relevant species observed during ATom and compare this to cOHR from the model (cOHR_{mod}). Measured OHR is higher than cOHR_{obs} by approximately 0.2 to 0.4 s^{-1} below 3 km . This missing OHR correlates with acetaldehyde during summer, indicating a potential source of missing reactive VOCs, similar to the findings of Mao et al. (2009) and S. Wang et al. (2020). The addition of a comprehensive set of ocean emissions of VOCs increases global mean cOHR by 6% but cannot reproduce the observed OHR enhancement during ATom-1. Adding sufficient alkanes to the model to resolve this bias requires an improbably large ocean source (340 Tg C yr^{-1}) and would degrade the model simulation of OH and NO_y . Only one-third of the increase in cOHR in summer in this test is due to the alkanes; the rest is from oxidation products and changes in OH. Therefore, a more reactive VOC would still need to be emitted in large amounts.

The model successfully simulates the seasonality and hemispheric gradient in cOHR but has a persistent underestimate of up to 12% in the lowest 3 km , primarily due to missing model acetaldehyde. The model does not underestimate CO, with the exception of Northern Hemisphere winter, which has been previously recognized by Kopacz et al. (2010) and attributed to underestimated fossil fuel emissions. The inability to reproduce observations of remote acetaldehyde was first observed during the PEM-Tropics campaign (Singh et al., 2001, 2003; Millet et al., 2010), but the measurement was uncertain. Improvements in measurement precision and the accompanying measurement of PAA during ATom (S. Wang et al., 2019) strengthen the conclusion that there is a large amount of acetaldehyde present in the atmosphere that cannot be explained by current models. We investigate possible underestimates in known sources of acetaldehyde, including emissions of VOCs from anthropogenic, biomass, or oceanic sources or production from the photolysis of organic aerosols. No known source can fully resolve the bias in acetaldehyde throughout the troposphere, and particularly in the Northern Hemisphere summer. We consider the possibility that there is a large, diffuse source of unknown VOCs by implementing 100 Tg yr^{-1} of such a compound from ocean, biomass burning, or anthropogenic sources. This hypothetical source modestly reduces the model acetaldehyde bias and is compatible with the simulation of OH and cOHR; however, an additional source is required to resolve the largest bias in the Northern Hemisphere summer. Errors or omissions in the oxidation mechanism of known VOCs could be another source of bias. For example, significant uncertainties exist in peroxy radical (RO_2) chemistry for large RO_2 molecules (Praske et al., 2017), although the flux of carbon through a minor pathway would have to be large, restricting the possible known sources. Further laboratory and field observations are needed to understand which precursors and sources could lead to the sustained production of acetaldehyde observed during ATom and prior campaigns.

This study demonstrates that long-standing model biases in global mean OH are unlikely to be due to errors in simulating tropospheric chemistry over the ocean. This implies that a large bias must be present in OH production or loss over land and future work should focus on evaluating continental OH sources and sinks. Errors in modeled OH were recently investigated by Strode et al. (2015), and when overestimates related to production terms

were corrected, model OH remained too high in the Northern Hemisphere, suggesting that future studies should focus on errors in OH loss.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements.

We are grateful for helpful conversations and advice from Andrea Molod, Rachel Silvern, Eloïse Marais, Sarah Safieddine, Martin Brüggemann, Christian George, and James Crawford. We acknowledge Tom Hanisco and Jason St. Clair for the use of their formaldehyde observations from ATom and Barbara Barletta and Simone Meinardi for their contribution to the UCI WAS measurements.

Financial support. This research has been supported by the National Science Foundation (grant no. AGS-1564495), the National Center for Atmospheric Research (grant no. 1852977), the National Oceanic and Atmospheric Administration (grant no. NA18OAR4310110), and the National Aeronautics and Space Administration (grant no. NNX14AP89G, grant no. IAT NNH15AB12I, grant no. NNX17AG35G, grant no. NNX15AG61A, grant no. NNX15AG71A).

References

- Akagi SK, Yokelson RJ, Wiedinmyer C, Alvarado MJ, Reid JS, Karl T, Crouse JD, and Wennberg PO: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, 10.5194/acp-11-4039-2011, 2011.
- Anderson DC, Loughner CP, Diskin G, Weinheimer A, Canty TP, Salawitch RJ, Worden HM, Fried A, Mikoviny T, Wisthaler A, and Dickerson RR: Measured and modeled CO and NO_y in DISCOVER-AQ: An evaluation of emissions and chemistry over the eastern US, *Atmos. Environ.*, 96, 78–87, 10.1016/j.atmosenv.2014.07.004, 2014.
- Apel EC, Hills AJ, Lueb R, Zindel S, Eisele S, and Riemer DD: A fast-GC/MS system to measure C₂ to C₄ carbonyls and methanol aboard aircraft, *J. Geophys. Res.-Atmos.*, 108, 8794, 10.1029/2002JD003199, 2003.
- Apel EC, Brauers T, Koppmann R, Bandowe B, Boßmeyer J, Holzke C, Tillmann R, Wahner A, Wegener R, Brunner A, Jocher M, Ruuskanen T, Spirig C, Steigner D, Steinbrecher R, Gomez Alvarez E, Müller K, Burrows JP, Schade G, Solomon SJ, Ladstätter-Weissenmayer A, Simmonds P, Young D, Hopkins JR, Lewis AC, Legreid G, Reimann S, Hansel A, Wisthaler A, Blake RS, Ellis AM, Monks PS, and Wyche KP: Intercomparison of oxygenated volatile organic compound measurements at the SAPHIR atmosphere simulation chamber, *J. Geophys. Res.-Atmos.*, 113, D20307, 10.1029/2008JD009865, 2008.
- Apel EC, Hornbrook RS, Hills AJ, Blake NJ, Barth MC, Weinheimer A, Cantrell C, Rutledge SA, Basarab B, Crawford J, Diskin G, Homeyer CR, Campos T, Flocke F, Fried A, Blake DR, Brune W, Pollack I, Peischl J, Ryerson T, Wennberg PO, Crouse JD, Wisthaler A, Mikoviny T, Huey G, Heikes B, O'Sullivan D, and Riemer DD: Upper tropospheric ozone production from lightning NO_x-impacted convection: Smoke ingestion case study from the DC3 campaign, *J. Geophys. Res.-Atmos.*, 120, 2505–2523, 10.1002/2014JD022121, 2015.
- Arnold SR, Spracklen DV, Williams J, Yassaa N, Sciare J, Bonsang B, Gros V, Peeken I, Lewis AC, Alvaïn S, and Moulin C: Evaluation of the global oceanic isoprene source and its impacts on marine organic carbon aerosol, *Atmos. Chem. Phys.*, 9, 1253–1262, 10.5194/acp-9-1253-2009, 2009.
- Beale R, Liss PS, and Nightingale PD: First oceanic measurements of ethanol and propanol, *Geophys. Res. Lett.*, 37, L24607, 10.1029/2010GL045534, 2010.
- Bey I, Jacob DJ, Yantosca RM, Logan JA, Field BD, Fiore AM, Li Q, Liu HY, Mickley LJ, and Schultz MG: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.*, 106, 23073–23095, 10.1029/2001JD000807, 2001.
- Breider TJ, Mickley LJ, Jacob DJ, Ge C, Wang J, Sulprizio MP, Croft B, Ridley DA, McConnell JR, Sharma S, Husain L, Dutkiewicz VA, Eleftheriadis K, Skov H, and Hopke PK: Multi-decadal trends

- in aerosol radiative forcing over the Arctic: contribution of changes in anthropogenic aerosol to Arctic warming since 1980, *J. Geophys. Res.*, 122, 3573–3594, 10.1002/2016JD025321, 2017.
- Brüggemann M, Hayeck N, Bonnineau C, Pesce S, A. Alpert P, Perrier S, Zuth C, Hoffmann T, Chen J, and George C: Interfacial photochemistry of biogenic surfactants: a major source of abiotic volatile organic compounds, *Faraday Discuss*, 200, 59–74, 10.1039/C7FD00022G, 2017. [PubMed: 28598469]
- Brüggemann M, Hayeck N, and George C: Interfacial photochemistry at the ocean surface is a global source of organic vapors and aerosols, *Nat. Commun*, 9, 2101, 10.1038/s41467-018-04528-7, 2018. [PubMed: 29844311]
- Brune WH, Miller DO, Thames AB, Allen HM, Apel EC, Blake DR, Bui TP, Commane R, Crouse JD, Daube BC, Diskin GS, DiGangi JP, Elkins JW, Hall SR, Hanisco TF, Hannun RA, Hintsä E, Hornbrook RS, Kim MJ, McKain K, Moore FL, Neuman JA, Nicely JM, Peischl J, Ryerson TB, St. Clair JM, Sweeney C, Teng AP, Thompson C, Ullmann K, Veres PR, Wennberg PO, and Wolfe GM: Exploring Oxidation in the Remote Free Troposphere: Insights From Atmospheric Tomography (ATom), *J. Geophys. Res. Atmos*, 125, e2019JD031685, 10.1029/2019JD031685, 2020.
- Brunner D, Staehelin J, Rogers HL, Köhler MO, Pyle JA, Hauglustaine D, Jourdain L, Berntsen TK, Gauss M, Isaksen ISA, Meijer E, van Velthoven P, Pitari G, Mancini E, Grewe G, and Sausen R: An evaluation of the performance of chemistry transport models by comparison with research aircraft observations. Part 1: Concepts and overall model performance, *Atmos. Chem. Phys*, 3, 1609–1631, 10.5194/acp-3-1609-2003, 2003.
- Brunner D, Staehelin J, Rogers HL, Köhler MO, Pyle JA, Hauglustaine DA, Jourdain L, Berntsen TK, Gauss M, Isaksen ISA, Meijer E, van Velthoven P, Pitari G, Mancini E, Grewe V, and Sausen R: An evaluation of the performance of chemistry transport models - Part 2: Detailed comparison with two selected campaigns, *Atmos. Chem. Phys*, 5, 107–129, 10.5194/acp-5-107-2005, 2005.
- Cazorla M, Wolfe GM, Bailey SA, Swanson AK, Arkinson HL, and Hanisco TF: A new airborne laser-induced fluorescence instrument for in situ detection of formaldehyde throughout the troposphere and lower stratosphere, *Atmos. Meas. Tech*, 8, 541–552, 10.5194/amt-8-541-2015, 2015.
- Chan Miller C, Jacob DJ, Marais EA, Yu K, Travis KR, Kim PS, Fisher JA, Zhu L, Wolfe GM, Hanisco TF, Keutsch FN, Kaiser J, Min K-E, Brown SS, Washenfelder RA, González Abad G, and Chance K: Glyoxal yield from isoprene oxidation and relation to formaldehyde: chemical mechanism, constraints from SENEX aircraft observations, and interpretation of OMI satellite data, *Atmos. Chem. Phys*, 17, 8725–8738, 10.5194/acp-17-8725-2017, 2017.
- Chen X, Millet DB, Singh HB, Wisthaler A, Apel EC, Atlas EL, Blake DR, Bourgeois I, Brown SS, Crouse JD, de Gouw JA, Flocke FM, Fried A, Heikes BG, Hornbrook RS, Mikoviny T, Min K-E, Müller M, Neuman JA, O'Sullivan DW, Peischl J, Pfister GG, Richter D, Roberts JM, Ryerson TB, Shertz SR, Thompson CR, Treadaway V, Veres PR, Walega J, Warneke C, Washenfelder RA, Weibring P, and Yuan B: On the sources and sinks of atmospheric VOCs: an integrated analysis of recent aircraft campaigns over North America, *Atmos. Chem. Phys*, 19, 9097–9123, 10.5194/acp-19-9097-2019, 2019.
- Christian KE, Brune WH, Mao J, and Ren X: Global sensitivity analysis of GEOS-Chem modeled ozone and hydrogen oxides during the INTEX campaigns, *Atmos. Chem. Phys*, 18, 2443–2460, 10.5194/acp-18-2443-2018, 2018.
- Colman J, Swanson A, Meinardi S, Sive B, Blake DR, and Rowland FS: Description of the analysis of a wide range of volatile organic compounds in whole air samples collected during PEM-Tropics A and B, *Anal. Chem*, 73, 3723–3731, 10.1021/ac010027g, 2001. [PubMed: 11510840]
- Crouse JD, McKinney KA, Kwan AJ, and Wennberg PO: Measurement of gas-phase hydroperoxides by chemical ionization mass spectrometry, *Anal. Chem*, 78, 6726–6732, 2006. [PubMed: 17007490]
- Dalsøren SB, Myhre G, Hodnebrog Ø, Myhre CL, Stohl A, Pizzo I, Schwietzke S, Höglund-Isaksson L, Helmig D, Reimann S, Sauvage S, Schmidbauer N, Read KA, Carpenter LJ, Lewis AC, Punjabi S, and Wallasch M: Discrepancy between simulated and observed ethane and propane levels explained by underestimated fossil emissions, *Nat. Geosci*, 11, 178–184, 10.1038/s41561-018-0073-0, 2018.

- Dentener F, Drevet J, Lamarque JF, Bey I, Eickhout B, Fiore AM, Hauglustaine D, Horowitz LW, Krol M, Kulshrestha UC, Lawrence M, Galy-Lacaux C, Rast S, Shindell D, Stevenson D, Noije TV, Atherton C, Bell N, Bergman D, Butler T, Cofala J, Collins B, Doherty R, Ellingsen K, Galloway J, Gauss M, Montanaro V, Müller JF, Pitari G, Rodriguez J, Sanderson M, Solomon F, Strahan S, Schultz M, Sudo K, Szopa S, and Wild O: Nitrogen and sulfur deposition on regional and global scales: A multimodel evaluation, *Global Biogeochem. Cy*, 20, GB4003, 10.1029/2005GB002672, 2006.
- Di Carlo P: Missing OH Reactivity in a Forest: Evidence for Unknown Reactive Biogenic VOCs, *Science*, 304, 722–725, 10.1126/science.1094392, 2004. [PubMed: 15118159]
- DiGangi JP, Boyle ES, Karl T, Harley P, Turnipseed A, Kim S, Cantrell C, Maudlin III RL, Zheng W, Flocke F, Hall SR, Ullmann K, Nakashima Y, Paul JB, Wolfe GM, Desai AR, Kajii Y, Guenther A, and Keutsch FN: First direct measurements of formaldehyde flux via eddy covariance: implications for missing in-canopy formaldehyde sources, *Atmos. Chem. Phys*, 11, 10565–10578, 10.5194/acp-11-10565-2011, 2011.
- Diskin GS, Podolske JR, Sachse GW, and Slate TA: Open-Path Airborne Tunable Diode Laser Hygrometer, in: *Diode Lasers and Applications in Atmospheric Sensing*, SPIE Proceedings, 4817, edited by: Fried A, 196–204, 2002.
- Dolgorouky C, Gros V, Sarda-Estevé R, Sinha V, Williams J, Marchand N, Sauvage S, Poulain L, Sciare J, and Bonsang B: Total OH reactivity measurements in Paris during the 2010 MEGAPOLI winter campaign, *Atmos. Chem. Phys*, 12, 9593–9612, 10.5194/acp-12-9593-2012, 2012.
- Duncan BN, Logan JA, Bey I, Megretskaia IA, Yantosca RM, Novelli PC, Jones NB, and Rinsland CP: Global budget of CO, 1988–1997: Source estimates and validation with a global model, *J. Geophys. Res.*, 112, D22301, 10.1029/2007JD008459, 2007.
- Edwards PM, Evans MJ, Furneaux KL, Hopkins J, Ingham T, Jones C, Lee JD, Lewis AC, Moller SJ, Stone D, Whalley LK, and Heard DE: OH reactivity in a South East Asian tropical rainforest during the Oxidant and Particle Photochemical Processes (OP3) project, *Atmos. Chem. Phys*, 13, 9497–9514, 10.5194/acp-13-9497-2013, 2013.
- Elkins JW, Moorem F, and Kline ES: Next Generation Airborne Gas chromatograph for NASA Airborne Platforms, *Earth Science Technology Conference 2001*, 28–30 August 2001, College Park, MD, 2001.
- Emmerson KM and Evans MJ: Comparison of tropospheric gas-phase chemistry schemes for use within global models, *Atmos. Chem. Phys*, 9, 1831–1845, 10.5194/acp-9-1831-2009, 2009.
- Epstein SA, Blair SL, and Nizkorodov SA: Direct Photolysis of α -Pinene Ozonolysis Secondary Organic Aerosol: Effect on Particle Mass and Peroxide Content, *Environ. Sci. Technol*, 48, 11251–11258, 10.1021/es502350u, 2014. [PubMed: 25165890]
- Faloona IC, Tan D, Leshner RL, Hazen NL, Frame CL, Simpkins JB, Harder H, Martinez M, Di Carlo P, Ren X, and Brune WH: A Laser-induced Fluorescence Instrument for Detecting Tropospheric OH and HO₂: Characteristics and Calibration, *J. Atmos. Chem*, 47, 139–167, 10.1023/B:JOCH.0000021036.53185.0e, 2004.
- Ferracci V, Heimann I, Abraham NL, Pyle JA, and Archibald AT: Global modelling of the total OH reactivity: investigations on the “missing” OH sink and its atmospheric implications, *Atmos. Chem. Phys*, 18, 7109–7129, 10.5194/acp-18-7109-2018, 2018.
- Fiore AM, Dentener FJ, Wild O, Cuvelier C, Schultz MG, Hess P, Textor C, Schulz M, Doherty RM, Horowitz LW, MacKenzie IA, Sanderson MG, Shindell DT, Stevenson DS, Szopa S, Van Dingenen R, Zeng G, Atherton C, Bergmann D, Bey I, Carmichael G, Collins WJ, Duncan BN, Faluvegi G, Folberth G, Gauss M, Gong S, Hauglustaine D, Holloway T, Isaksen ISA, Jacob DJ, Jonson JE, Kaminski JW, Keating TJ, Lupu A, Marmer E, Montanaro V, Park RJ, Pitari G, Pringle KJ, Pyle JA, Schroeder S, Vivanco MG, Wind P, Wojcik G, Wu S, and Zuber A: Multimodel estimates of intercontinental source-receptor relationships for ozone pollution, *J. Geophys. Res.*, 114, D04301, 10.1029/2008JD010816, 2009.
- Fischer EV, Jacob DJ, Millet DB, Yantosca RM, and Mao J: The role of the ocean in the global atmospheric budget of acetone, *Geophys. Res. Lett.*, 39, L01807, 10.1029/2011GL050086, 2012.
- Fischer EV, Jacob DJ, Yantosca RM, Sulprizio MP, Millet DB, Mao J, Paulot F, Singh HB, Roiger A, Ries L, Talbot RW, Dzepina K, and Pandey Deolal S: Atmospheric peroxyacetyl nitrate (PAN): a

global budget and source attribution, *Atmos. Chem. Phys.*, 14, 2679–2698, 10.5194/acp-14-2679-2014, 2014.

- Fisher JA, Jacob DJ, Travis KR, Kim PS, Marais EA, Chan Miller C, Yu K, Zhu L, Yantosca RM, Sulprizio MP, Mao J, Wennberg PO, Crounse JD, Teng AP, Nguyen TB, St. Clair JM, Cohen RC, Romer P, Nault BA, Wooldridge PJ, Jimenez JL, Campuzano-Jost P, Day DA, Hu W, Shepson PB, Xiong F, Blake DR, Goldstein AH, Misztal PK, Hanisco TF, Wolfe GM, Ryerson TB, Wisthaler A, and Mikoviny T: Organic nitrate chemistry and its implications for nitrogen budgets in an isoprene- and monoterpene-rich atmosphere: constraints from aircraft (SEAC4RS) and ground-based (SOAS) observations in the Southeast US, *Atmos. Chem. Phys.*, 16, 5969–5991, 10.5194/acp-16-5969-2016, 2016. [PubMed: 29681921]
- Fisher JA, Atlas EL, Barletta B, Meinardi S, Blake DR, Thompson CR, Ryerson TB, Peischl J, Tzompa-Sosa ZA, and Murray LT: Methyl, ethyl, and propyl nitrates: global distribution and impacts on reactive nitrogen in remote marine environments, *J. Geophys. Res.*, 123, 12429–12451, 10.1029/2018JD029046, 2018.
- Gantt B, Meskhidze N, Zhang Y, and Xu J: The effect of marine isoprene emissions on secondary organic aerosol and ozone formation in the coastal United States, *Atmos. Environ.*, 44, 115–121, 10.1016/j.atmosenv.2009.08.027, 2010.
- Gelaro R, McCarty W, Suárez MJ, Todling R, Molod A, Takacs L, Randles CA, Darmenov A, Bosilovich MG, Reichle R, Wargan K, Coy L, Cullather R, Draper C, Akella S, Buchard V, Conaty A, da Silva AM, Gu W, Kim G-K, Koster R, Lucchesi R, Merkova D, Nielsen JE, Partyka G, Pawson S, Putman W, Rienecker M, Schubert SD, Sienkiewicz M, and Zhao B: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454, 10.1175/JCLI-D-16-0758.1, 2017.
- Giglio L, Randerson JT, and van der Werf GR: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4): ANALYSIS OF BURNED AREA, *J. Geophys. Res.-Biogeo.*, 118, 317–328, 10.1002/jgrg.20042, 2013.
- Goldstein AH and Galbally IE: Known and unexplored organic constituents in the Earth's Atmosphere, *Environ. Sci. Technol.*, 41, 1514–1521, 10.1021/es072476p, 2007. [PubMed: 17396635]
- Granier C, Lamarque JF, Mieville A, Muller JF, Olivier J, Orlando J, Peters J, Petron G, Tyndall G, and Wallens S: POET, a database of surface emissions of ozone precursors, available at: <http://www.aero.jussieu.fr/projet/ACCENT/POET.php> (last access: 2 July 2020), 2005.
- Guenther AB, Jiang X, Heald CL, Sakulyanontvittaya T, Duhl T, Emmons LK, and Wang X: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Mod. Dev.*, 5, 1471–1492, 10.5194/gmd-5-1471-2012, 2012.
- Hackenberg SC, Andrews SJ, Airs RL, Arnold SR, Bouman HA, Cummings D, Lewis AC, Minaeian JK, Reifel KM, Small A, Tarran GA, Tilstone GH, and Carpenter LJ: Basin-Scale Observations of Monoterpenes in the Arctic and Atlantic Oceans, *Environ. Sci. Technol.*, 51, 10449–10458, 10.1021/acs.est.7b02240, 2017. [PubMed: 28752764]
- Hall SR, Ullmann K, Prather MJ, Flynn CM, Murray LT, Fiore AM, Correa G, Strode SA, Steenrod SD, Lamarque J-F, Guth J, Josse B, Flemming J, Huijnen V, Abraham NL, and Archibald AT: Cloud impacts on photochemistry: building a climatology of photolysis rates from the Atmospheric Tomography mission, *Atmos. Chem. Phys.*, 18, 16809–16828, 10.5194/acp-18-16809-2018, 2018.
- Hansen RF, Griffith SM, Dusanter S, Rickly PS, Stevens PS, Bertman SB, Carroll MA, Erickson MH, Flynn JH, Grossberg N, Jobson BT, Lefer BL, and Wallace HW: Measurements of total hydroxyl radical reactivity during CABINEX 2009 – Part 1: field measurements, *Atmos. Chem. Phys.*, 14, 2923–2937, 10.5194/acp-14-2923-2014, 2014.
- Heald CL, Collett JL Jr., Lee T, Benedict KB, Schwandner FM, Li Y, Clarisse L, Hurtmans DR, Van Damme M, Clerbaux C, Coheur P-F, Philip S, Martin RV, and Pye HOT: Atmospheric ammonia and particulate inorganic nitrogen over the United States, *Atmos. Chem. Phys.*, 12, 10295–10312, 10.5194/acp-12-10295-2012, 2012.
- Hintsä E, Moore FL, Hurst DF, Dutton GS, Hall BD, Nance JD, Patrick L, McClure-Begley A, Elkins JW, Hall EG, Jordan AF, Gao R-S, Rollins AW, Thornberry TD, Watts LA, Thompson C, Peischl J, Bourgeois I, Ryerson T, Daube BC, Pittman JV, Wofsy SC, Diskin G, Bui TP, Montzka S, and

- Miller B: UAS Chromatograph for Atmospheric Trace Species (UCATS) – a versatile instrument for trace gas measurements on different airborne platforms, *Atmos. Meas. Tech.*, in preparation, 2020.
- Hodzic A, Madronich S, Kasibhatla PS, Tyndall G, Aumont B, Jimenez JL, Lee-Taylor J, and Orlando J: Organic photolysis reactions in tropospheric aerosols: effect on secondary organic aerosol formation and lifetime, *Atmos. Chem. Phys.*, 15, 9253–9269, 10.5194/acp-15-9253-2015, 2015.
- Hodzic A, Kasibhatla PS, Jo DS, Cappa CD, Jimenez JL, Madronich S, and Park RJ: Rethinking the global secondary organic aerosol (SOA) budget: stronger production, faster removal, shorter lifetime, *Atmos. Chem. Phys.*, 16, 7917–7941, 10.5194/acp-16-7917-2016, 2016.
- Hoesly RM, Smith SJ, Feng L, Klimont Z, Janssens-Maenhout G, Pitkanen T, Seibert JJ, Vu L, Andres RJ, Bolt RM, Bond TC, Dawidowski L, Kholod N, Kurokawa J-I, Li M, Liu L, Lu Z, Moura MCP, O'Rourke PR, and Zhang Q: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11, 369–408, 10.5194/gmd-11-369-2018, 2018.
- Hofzumahaus A, Lefer BL, Monks PS, Hall SR, Kylling A, Mayer B, Shetter RE, Junkermann W, Bais A, Calvert JG, Cantrell CA, Madronich S, Edwards GD, and Kraus A: Photolysis frequency of O₃ to O(¹D): Measurements and modeling during the International Photolysis Frequency Measurement and Modeling Intercomparison (IPMMI), *J. Geophys. Res.-Atmos.*, 109, D08S90, 10.1029/2003JD004333, 2004.
- Hottle JR, Huisman AJ, DiGangi JP, Kammrath A, Galloway MM, Coens KL, and Keutsch FN: A Laser Induced Fluorescence-Based Instrument for In-Situ Measurements of Atmospheric Formaldehyde, *Environ. Sci. Technol.*, 43, 790–795, 10.1021/es801621f, 2009. [PubMed: 19245018]
- Hu L, Millet DB, Baasandorj M, Griffis TJ, Turner P, Helmig D, Curtis AJ, and Hueber J: Isoprene emissions and impacts over an ecological transition region in the U.S. Upper Midwest inferred from tall tower measurements: Isoprene emissions in US Upper Midwest, *J. Geophys. Res.-Atmos.*, 120, 3553–3571, 10.1002/2014JD022732, 2015.
- Hu L, Jacob DJ, Liu X, Zhang Y, Zhang L, Kim PS, Sulprizio MP, and Yantosca RM: Global budget of tropospheric ozone: Evaluating recent model advances with satellite (OMI), aircraft (IAGOS), and ozonesonde observations, *Atmos. Environ.*, 167, 323–334, 10.1016/j.atmosenv.2017.08.036, 2017.
- Jacob DJ: Heterogeneous chemistry and tropospheric ozone, *Atmos. Environ.*, 34, 2131–2159, 2000.
- Jaeglé L, Shah V, Thornton JA, Lopez-Hilfiker FD, Lee BH, McDuffie EE, Fibiger D, Brown SS, Veres P, Sparks TL, Ebben CJ, Wooldridge PJ, Kenagy HS, Cohen RC, Weinheimer AJ, Campos TL, Montzka DD, Digangi JP, Wolfe GM, Hanisco T, Schroder JC, Campuzano-Jost P, Day DA, Jimenez JL, Sullivan AP, Guo H, and Weber RJ: Nitrogen Oxides Emissions, Chemistry, Deposition, and Export Over the Northeast United States During the WINTER Aircraft Campaign, *J. Geophys. Res.-Atmos.*, 123, 12368–12393, 10.1029/2018JD029133, 2018.
- Jenkin ME, Saunders SM, and Pilling ME: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmos. Environ.*, 31, 81–104, 1997.
- Jenkin ME, Young JC, and Rickard AR: The MCM v3.3.1 degradation scheme for isoprene, *Atmos. Chem. Phys.*, 15, 11433–11459, 10.5194/acp-15-11433-2015, 2015.
- Johnson MT: A numerical scheme to calculate temperature and salinity dependent air-water transfer velocities for any gas, *Ocean Sci.*, 6, 913–932, 10.5194/os-6-913-2010, 2010.
- Kaiser J, Skog KM, Baumann K, Bertman SB, Brown SB, Brune WH, Crouse JD, de Gouw JA, Edgerton ES, Feiner PA, Goldstein AH, Koss A, Miszta PK, Nguyen TB, Olson KF, St. Clair JM, Teng AP, Toma S, Wennberg PO, Wild RJ, Zhang L, and Keutsch FN: Speciation of OH reactivity above the canopy of an isoprene-dominated forest, *Atmos. Chem. Phys.*, 16, 9349–9359, 10.5194/acp-16-9349-2016, 2016.
- Karion A, Sweeney C, Wolter S, Newberger T, Chen H, Andrews A, Kofler J, Neff D, and Tans P: Long-term greenhouse gas measurements from aircraft, *Atmos. Meas. Tech.*, 6, 511–526, 10.5194/amt-6-511-2013, 2013.
- Kasibhatla P, Sherwen T, Evans MJ, Carpenter LJ, Reed C, Alexander B, Chen Q, Sulprizio MP, Lee JD, Read KA, Bloss W, Crilley LR, Keene WC, Pszenny AAP, and Hodzic A: Global impact of nitrate photolysis in sea-salt aerosol on NO_x, OH, and O₃ in the marine boundary layer, *Atmos. Chem. Phys.*, 18, 11185–11203, 10.5194/acp-18-11185-2018, 2018.

- Keller CA, Long MS, Yantosca RM, Da Silva AM, Pawson S, and Jacob DJ: HEMCO v1.0: a versatile, ESMF-compliant component for calculating emissions in atmospheric models, *Geosci. Model Dev*, 7, 1409–1417, 10.5194/gmd-7-1409-2014, 2014.
- Kim MJ, Novak GA, Zoerb MC, Yang M, Blomquist BW, Huebert BJ, Cappa CD, and Bertram TH: Air-Sea exchange of biogenic volatile organic compounds and the impact on aerosol particle size distributions, *Geophys. Res. Lett*, 44, 3887–3896, 10.1002/2017GL072975, 2017.
- Kopacz M, Jacob DJ, Fisher JA, Logan JA, Zhang L, Megretskaia IA, Yantosca RM, Singh K, Henze DK, Burrows JP, Buchwitz M, Khlystova I, McMillan WW, Gille JC, Edwards DP, Eldering A, Thouret V, and Nedelec P: Global estimates of CO sources with high resolution by adjoint inversion of multiple satellite datasets (MOPITT, AIRS, SCIAMACHY, TES), *Atmos. Chem. Phys*, 10, 855–876, 10.5194/acp-10-855-2010, 2010.
- Koss AR, Sekimoto K, Gilman JB, Selimovic V, Coggon MM, Zarzana KJ, Yuan B, Lerner BM, Brown SS, Jimenez JL, Krechmer J, Roberts JM, Warneke C, Yokelson RJ, and de Gouw J: Non-methane organic gas emissions from biomass burning: identification, quantification, and emission factors from PTR-ToF during the FIREX 2016 laboratory experiment, *Atmos. Chem. Phys*, 18, 3299–3319, 10.5194/acp-18-3299-2018, 2018.
- Krotkov NA, McLinden CA, Li C, Lamsal LN, Celarier EA, Marchenko SV, Swartz WH, Bucsela EJ, Joiner J, Duncan BN, Boersma KF, Veefkind JP, Levelt PF, Fioletov VE, Dickerson RR, He H, Lu Z, and Streets DG: Aura OMI observations of regional SO₂ and NO₂ pollution changes from 2005 to 2015, *Atmos. Chem. Phys*, 16, 4605–4629, 10.5194/acp-16-4605-2016, 2016.
- Kuhns H, Green M, and Etyemezian V: Big Bend Regional Aerosol and Visibility Observational (BRAVO) Study Emissions Inventory, Desert Research Institute, Las Vegas, NV, 2003.
- Kwan AJ, Crounse JD, Clarke AD, Shinzuka Y, Anderson BE, Crawford JH, Avery MA, McNaughton CS, Brune WH, Singh HB, and Wennberg PO: On the flux of oxygenated volatile organic compounds from organic aerosol oxidation, *Geophys. Res. Lett*, 33, 10.1029/2006GL026144, 2006.
- Lamarque J-F, Emmons LK, Hess PG, Kinnison DE, Tilmes S, Vitt F, Heald CL, Holland EA, Lauritzen PH, Neu J, Orlando JJ, Rasch PJ, and Tyndall GK: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev*, 5, 369–411, 10.5194/gmd-5-369-2012, 2012.
- Lee JD, Young JC, Read KA, Hamilton JF, Hopkins JR, Lewis AC, Bandy BJ, Davey J, Edwards P, Ingham T, Self DE, Smith SC, Pilling MJ, and Heard DE: Measurement and calculation of OH reactivity at a United Kingdom coastal site, *J. Atmos. Chem*, 64, 53–76, 10.1007/s10874-010-9171-0, 2009.
- Lelieveld J, Gromov S, Pozzer A, and Taraborrelli D: Global tropospheric hydroxyl distribution, budget and reactivity, *Atmos. Chem. Phys*, 16, 12477–12493, 10.5194/acp-16-12477-2016, 2016.
- Li M, Zhang Q, Kurokawa J-I, Woo J-H, He K, Lu Z, Ohara T, Song Y, Streets DG, Carmichael GR, Cheng Y, Hong C, Huo H, Jiang X, Kang S, Liu F, Su H, and Zheng B: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys*, 17, 935–963, 10.5194/acp-17-935-2017, 2017.
- Li SM, Leithead A, Moussa SG, Liggio J, Moran MD, Wang D, Hayden K, Darlington A, Gordon M, Staebler R, Makar PA, Stroud CA, McLaren R, Liu PSK, O'Brien J, Mittermeier RL, Zhang J, Marson G, Cober SG, Wolde M, and Wentzell JJB: Differences between measured and reported volatile organic compound emissions from oil sands facilities in Alberta, Canada, *P. Natl. Acad. Sci. USA*, 114, E3756–E3765, 10.1073/pnas.1617862114, 2017.
- Lou S, Holland F, Rohrer F, Lu K, Bohn B, Brauers T, Chang CC, Fuchs H, Häseler R, Kita K, Kondo Y, Li X, Shao M, Zeng L, Wahner A, Zhang Y, Wang W, and Hofzumahaus A: Atmospheric OH reactivities in the Pearl River Delta – China in summer 2006: measurement and model results, *Atmos. Chem. Phys*, 10, 11243–11260, 10.5194/acp-10-11243-2010, 2010.
- Luo G, Yu F, and Schwab J: Revised treatment of wet scavenging processes dramatically improves GEOS-Chem 12.0.0 simulations of surface nitric acid, nitrate, and ammonium over the United States, *Geosci. Model Dev*, 12, 3439–3447, 10.5194/gmd-12-3439-2019, 2019.
- Mao J, Ren X, Brune WH, Olson JR, Crawford JH, Fried A, Huey LG, Cohen RC, Heikes B, Singh HB, Blake DR, Sachse GW, Diskin GS, Hall SR, and Shetter RE: Airborne measurement of OH reactivity during INTEX-B, *Atmos. Chem. Phys*, 9, 163–173, 10.5194/acp-9-163-2009, 2009.

- Mao J, Ren X, Chen S, Brune WH, Chen Z, Martinez M, Harder H, Lefer B, Rappenglück B, Flynn J, and Leuchner M: Atmospheric oxidation capacity in the summer of Houston 2006: Comparison with summer measurements in other metropolitan studies, *Atmos. Environ.*, 44, 4107–4115, 10.1016/j.atmosenv.2009.01.013, 2010.
- Mao J, Ren X, Zhang L, Van Duin DM, Cohen RC, Park J-H, Goldstein AH, Paulot F, Beaver MR, Crouse JD, Wennberg PO, DiGangi JP, Henry SB, Keutsch FN, Park C, Schade GW, Wolfe GM, Thornton JA, and Brune WH: Insights into hydroxyl measurements and atmospheric oxidation in a California forest, *Atmos. Chem. Phys.*, 12, 8009–8020, 10.5194/acp-12-8009-2012, 2012.
- Mao J, Fan S, Jacob DJ, and Travis KR: Radical loss in the atmosphere from Cu-Fe redox coupling in aerosols, *Atmos. Chem. Phys.*, 13, 509–519, 10.5194/acp-13-509-2013, 2013.
- Marais EA and Wiedinmyer C: Air Quality Impact of Diffuse and Inefficient Combustion Emissions in Africa (DICE-Africa), *Environ. Sci. Technol.*, 50, 10739–10745, 10.1021/acs.est.6b02602, 2016. [PubMed: 27611340]
- Marais EA, Jacob DJ, Jimenez JL, Campuzano-Jost P, Day DA, Hu W, Krechmer J, Zhu L, Kim PS, Miller CC, Fisher JA, Travis K, Yu K, Hanisco TF, Wolfe GM, Arkinson HL, Pye HOT, Froyd KD, Liao J, and McNeill VF: Aqueous-phase mechanism for secondary organic aerosol formation from isoprene: application to the southeast United States and co-benefit of SO₂ emission controls, *Atmos. Chem. Phys.*, 16, 1603–1618, 10.5194/acp-16-1603-2016, 2016. [PubMed: 32742280]
- Marais EA, Jacob DJ, Choi S, Joiner J, Belmonte-Rivas M, Cohen RC, Beirle S, Murray LT, Schiferl LD, Shah V, and Jaeglé L: Nitrogen oxides in the global upper troposphere: interpreting cloud-sliced NO₂ observations from the OMI satellite instrument, *Atmos. Chem. Phys.*, 18, 17017–17027, 10.5194/acp-18-17017-2018, 2018.
- Marvin MR, Wolfe GM, Salawitch RJ, Canty TP, Roberts SJ, Travis KR, Aikin KC, de Gouw JA, Graus M, Hanisco TF, Holloway JS, Hübler G, Kaiser J, Keutsch FN, Peischl J, Pollack IB, Roberts JM, Ryerson TB, Veres PR, and Warneke C: Impact of evolving isoprene mechanisms on simulated formaldehyde: An inter-comparison supported by in situ observations from SENEX, *Atmos. Environ.*, 164, 325–336, 10.1016/j.atmosenv.2017.05.049, 2017.
- McDonald BC, Gouw J. A. de, Gilman JB, Jathar SH, Akherati A, Cappa CD, Jimenez JL, Lee-Taylor J, Hayes PL, McKeen SA, Cui YY, Kim S-W, Gentner DR, Isaacman-VanWertz G, Goldstein AH, Harley RA, Frost GJ, Roberts JM, Ryerson TB, and Trainer M: Volatile chemical products emerging as largest petrochemical source of urban organic emissions, *Science*, 359, 760–764, 10.1126/science.aag0524, 2018. [PubMed: 29449485]
- McManus JB, Nelson DD, Shorter JH, Jimenez R, Herndon S, Saleska S, and Zahniser M: A high precision pulsed quantum cascade laser spectrometer for measurements of stable isotopes of carbon dioxide, *J. Mod. Opt.*, 52, 2309–2321, 10.1080/09500340500303710, 2005.
- Millet DB, Jacob DJ, Custer TG, de Gouw JA, Goldstein AH, Karl T, Singh HB, Sive BC, Talbot RW, Warneke C, and Williams J: New constraints on terrestrial and oceanic sources of atmospheric methanol, *Atmos. Chem. Phys.*, 8, 6887–6905, 10.5194/acp-8-6887-2008, 2008.
- Millet DB, Guenther A, Siegel DA, Nelson NB, Singh HB, de Gouw JA, Warneke C, Williams J, Eerdeken G, Sinha V, Karl T, Flocke F, Apel E, Riemer DD, Palmer PI, and Barkley M: Global atmospheric budget of acetaldehyde: 3-D model analysis and constraints from in-situ and satellite observations, *Atmos. Chem. Phys.*, 10, 3405–3425, 10.5194/acp-10-3405-2010, 2010.
- Mogensen D, Smolander S, Sogachev A, Zhou L, Sinha V, Guenther A, Williams J, Nieminen T, Kajos MK, Rinne J, Kulmala M, and Boy M: Modelling atmospheric OH-reactivity in a boreal forest ecosystem, *Atmos. Chem. Phys.*, 11, 9709–9719, 10.5194/acp-11-9709-2011, 2011.
- Monks PS: Gas-phase radical chemistry in the troposphere, *Chem. Soc. Rev.*, 34, 376–395, 10.1039/b307982c, 2005. [PubMed: 15852151]
- Montzka SA, Spivakovsky CM, Butler JH, Elkins JW, Lock LT, and Mondeel DJ: New observational constraints for atmospheric hydroxyl on global and hemispheric scales, *Science*, 288, 5465, 10.1126/science.288.5465.500, 2000.
- Mu M, Randerson JT, van der Werf GR, Giglio L, Kasibhatla P, Morton D, Collatz GJ, DeFries RS, Hyer EJ, Prins EM, Griffith DWT, Wunch D, Toon GC, Sherlock V, and Wennberg PO: Daily and 3-hourly variability in global fire emissions and consequences for atmospheric model predictions of carbon monoxide, *J. Geophys. Res.-Atmos.*, 116, D24303, 10.1029/2011JD016245, 2011.

- Müller J-F, Liu Z, Nguyen VS, Stavrakou T, Harvey JN, and Peeters J: The reaction of methyl peroxy and hydroxyl radicals as a major source of atmospheric methanol, *Nat. Commun.*, 7, 13213, 10.1038/ncomms13213, 2016. [PubMed: 27748363]
- Müller J-F, Stavrakou T, Bauwens M, George M, Hurtmans D, Coheur P-F, Clerbaux C, and Sweeney C: Top-Down CO Emissions Based On IASI Observations and Hemispheric Constraints on OH Levels, *Geophys. Res. Lett.*, 45, 2017GL076697, 10.1002/2017GL076697, 2018.
- Mungall EL, Abbatt JPD, Wentzell JJB, Lee AKY, Thomas JL, Blais M, Gosselin M, Miller LA, Papakyriakou T, Willis MD, and Liggio J: Microlayer source of oxygenated volatile organic compounds in the summertime marine Arctic boundary layer, *P. Natl. Acad. Sci. USA*, 114, 6203–6208, 10.1073/pnas.1620571114, 2017.
- Murray LT, Jacob DJ, Logan JA, Hudman RC, and Koshak WJ: Optimized regional and interannual variability of lightning in a global chemical transport model constrained by LIS/OTD satellite data, *J. Geophys. Res.-Atmos.*, 117, D20307, 10.1029/2012JD017934, 2012.
- Murray LT, Mickley LJ, Kaplan JO, Sofen ED, Pfeiffer M, and Alexander B: Factors controlling variability in the oxidative capacity of the troposphere since the Last Glacial Maximum, *Atmos. Chem. Phys.*, 14, 3589–3622, 10.5194/acp-14-3589-2014, 2014.
- Naik V, Voulgarakis A, Fiore AM, Horowitz LW, Lamarque J-F, Lin M, Prather MJ, Young PJ, Bergmann D, Cameron-Smith PJ, Cionni I, Collins WJ, Dalsøren SB, Doherty R, Eyring V, Faluvegi G, Folberth GA, Josse B, Lee YH, MacKenzie IA, Nagashima T, van Noije TPC, Plummer DA, Righi M, Rumbold ST, Skeie R, Shindell DT, Stevenson DS, Strode S, Sudo K, Szopa S, and Zeng G: Preindustrial to present-day changes in tropospheric hydroxyl radical and methane lifetime from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 5277–5298, 10.5194/acp-13-5277-2013, 2013.
- Nakashima Y, Kato S, Greenberg J, Harley P, Karl T, Turnipseed A, Apel E, Guenther A, Smith J, and Kajii Y: Total OH reactivity measurements in ambient air in a southern Rocky mountain ponderosa pine forest during BEACHON-SRM08 summer campaign, *Atmos. Environ.*, 85, 1–8, 10.1016/j.atmosenv.2013.11.042, 2014.
- Nicely JM, Anderson DC, Canty TP, Salawitch RJ, Wolfe GM, Apel EC, Arnold SR, Atlas EL, Blake NJ, Bresch JF, Campos TL, Dickerson RR, Duncan B, Emmons LK, Evans MJ, Fernandez RP, Flemming J, Hall SR, Hanisco TF, Honomichl SB, Hornbrook RS, Huijnen V, Kaser L, Kinnison DE, Lamarque J-F, Mao J, Monks SA, Montzka DD, Pan LL, Riemer DD, Saiz-Lopez A, Steenrod SD, Stell MH, Tilmes S, Turquety S, Ullmann K, and Weinheimer AJ: An observationally constrained evaluation of the oxidative capacity in the tropical western Pacific troposphere: Observationally Constrained OH in TWP, *J. Geophys. Res.-Atmos.*, 121, 7461–7488, 10.1002/2016JD025067, 2016.
- Nicely JM, Salawitch RJ, Canty T, Anderson DC, Arnold SR, Chipperfield MP, Emmons LK, Flemming J, Huijnen V, Kinnison DE, Lamarque J-F, Mao J, Monks SA, Steenrod SD, Tilmes S, and Turquety S: Quantifying the causes of differences in tropospheric OH within global models, *J. Geophys. Res.-Atmos.*, 122, 1983–2007, 10.1002/2016JD026239, 2017.
- Nölscher AC, Williams J, Sinha V, Custer T, Song W, Johnson AM, Axinte R, Bozem H, Fischer H, Pouvesle N, Phillips G, Crowley JN, Rantala P, Rinne J, Kulmala M, Gonzales D, Valverde-Canossa J, Vogel A, Hoffmann T, Ouwersloot HG, Vilà-Guerau de Arellano J, and Lelieveld J: Summertime total OH reactivity measurements from boreal forest during HUMPPA-COPEC 2010, *Atmos. Chem. Phys.*, 12, 8257–8270, 10.5194/acp-12-8257-2012, 2012.
- Nölscher AC, Yañez-Serrano AM, Wolff S, de Araujo AC, Lavri JV, Kesselmeier J, and Williams J: Unexpected seasonality in quantity and composition of Amazon rainforest air reactivity, *Nat. Commun.*, 7, 10383, 10.1038/ncomms10383, 2016. [PubMed: 26797390]
- Olivier J, Peters J, Granier C, Petron G, Muller JF, and Wallens S: Present and future surface emissions of atmospheric compounds, *Eur. Union, Brussels*, 2003.
- Pai SJ, Heald CL, Pierce JR, Farina SC, Marais EA, Jimenez JL, Campuzano-Jost P, Nault BA, Middlebrook AM, Coe H, Shilling JE, Bahreini R, Dingle JH, and Vu K: An evaluation of global organic aerosol schemes using airborne observations, *Atmos. Chem. Phys.*, 20, 2637–2665, 10.5194/acp-20-2637-2020, 2020.
- Patra PK, Krol MC, Montzka SA, Arnold T, Atlas EL, Lintner BR, Stephens BB, Xiang B, Elkins JW, Fraser PJ, Ghosh A, Hintsa EJ, Hurst DF, Ishijima K, Krummel PB, Miller BR, Miyazaki K,

- Moore FL, Mühle J, O'Doherty S, Prinn RG, Steele LP, Takigawa M, Wang HJ, Weiss RF, Wofsy SC, and Young D: Observational evidence for interhemispheric hydroxyl-radical parity, *Nature*, 513, 219–223, 10.1038/nature13721, 2014. [PubMed: 25209800]
- Petropavlovskikh I, Shetter R, Hall S, Ullmann K, and Bhartia PK: Algorithm for the charge-coupled-device scanning actinic flux spectroradiometer ozone retrieval in support of the Aura satellite validation, *J. Appl. Remote Sens.*, 1, 013540, 10.1117/1.2802563, 2007.
- Philip S, Martin RV, and Keller CA: Sensitivity of chemistry-transport model simulations to the duration of chemical and transport operators: a case study with GEOS-Chem v10–01, *Geosci. Model Dev.*, 9, 1683–1695, 10.5194/gmd-9-1683-2016, 2016.
- Plass-Dülmer C, Khedim A, Koppmann R, Johnen FJ, Rudolph J, and Kuosa H: Emissions of light nonmethane hydrocarbons from the Atlantic into the atmosphere, *Global Biogeochem. Cy.*, 7, 211–228, 10.1029/92GB02361, 1993.
- Podolske JR, Sachse GW, and Diskin GS: Calibration and data retrieval algorithms for the NASA Langley/Ames Diode Laser Hygrometer for the NASA Transport and Chemical Evolution Over the Pacific (TRACE-P) mission, *J. Geophys. Res.*, 108, 8792, 10.1029/2002JD003156, 2003.
- Pollack IB, Lerner BM, and Ryerson TB: Evaluation of ultra-violet light-emitting diodes for detection of atmospheric NO₂ by photolysis – chemiluminescence, *J. Atmos. Chem.*, 65, 111–125, 10.1007/s10874-011-9184-3, 2010.
- Praske E, Otkjær RV, Crounse JD, Hethcox JC, Stoltz BM, Kjaergaard HG, and Wennberg PO: Atmospheric autoxidation is increasingly important in urban and suburban North America, *P. Natl. Acad. Sci. USA*, 115, 64–69, 10.1073/pnas.1715540115, 2017.
- Prather MJ, Holmes CD, and Hsu J: Reactive greenhouse gas scenarios: Systematic exploration of uncertainties and the role of atmospheric chemistry, *Geophys. Res. Lett.*, 39, L09803, 10.1029/2012GL051440, 2012.
- Pye HOT, Liao H, Wu S, Mickley LJ, Jacob DJ, Henze DK, and Seinfeld JH: Effect of changes in climate and emissions on future sulfate-nitrate-ammonium aerosol levels in the United States, *J. Geophys. Res.*, 114, D01205, 10.1029/2008JD010701, 2009.
- Ramasamy S, Ida A, Jones C, Kato S, Tsurumaru H, Kishimoto I, Kawasaki S, Sadanaga Y, Nakashima Y, Nakayama T, Matsumi Y, Mochida M, Kagami S, Deng Y, Ogawa S, Kawana K, and Kajii Y: Total OH reactivity measurement in a BVOC dominated temperate forest during a summer campaign, 2014, *Atmos. Environ.*, 131, 41–54, 10.1016/j.atmosenv.2016.01.039, 2016.
- Randerson JT, Chen Y, van der Werf GR, Rogers BM, and Morton DC: Global burned area and biomass burning emissions from small fires, *J. Geophys. Res.-Biogeo.*, 117, G04012, 10.1029/2012JG002128, 2012.
- Read KA, Carpenter LJ, Arnold SR, Beale R, Nightingale PD, Hopkins JR, Lewis AC, Lee JD, Mendes L, and Pickering SJ: Multiannual Observations of Acetone, Methanol, and Acetaldehyde in Remote Tropical Atlantic Air: Implications for Atmospheric OVOC Budgets and Oxidative Capacity, *Environ. Sci. Technol.*, 46, 11028–11039, 10.1021/es302082p, 2012. [PubMed: 22963451]
- Ren X, Brune WH, Oliger A, Metcalf AR, Simpkins JB, Shirley T, Schwab JJ, Bai C, Roychowdhury U, Li Y, Cai C, Demerjian KL, He Y, Zhou X, Gao H, and Hou J: OH, HO₂, and OH reactivity during the PMTACS–NY Whiteface Mountain 2002 campaign: Observations and model comparison, *J. Geophys. Res.-Atmos.*, 111, D10S03, 10.1029/2005JD006126, 2006.
- Romer PS, Wooldridge PJ, Crounse JD, Kim MJ, Wennberg PO, Dibb JE, Scheuer E, Blake DR, Meinardi S, Brosius AL, Thames AB, Miller DO, Brune WH, Hall SR, Ryerson TB, and Cohen RC: Constraints on aerosol nitrate photolysis as a potential source of HONO and NO_x, *Environ. Sci. Technol.*, 52, 13738–13746, 10.1021/acs.est.8b03861, 2018. [PubMed: 30407797]
- Ryerson TB, Buhr MP, Frost GJ, Goldan PD, Holloway JS, Hübler G, Jobson BT, Kuster WC, McKeen SA, Parrish DD, Roberts JM, Sueper DT, Trainer M, Williams J, and Fehsenfeld FC: Emissions lifetimes and ozone formation in power plant plumes, *J. Geophys. Res.-Atmos.*, 103, 22569–22583, 10.1029/98JD01620, 1998.
- Ryerson TB, Williams EJ, and Fehsenfeld FC: An efficient photolysis system for fast-response NO₂ measurements, *J. Geophys. Res.-Atmos.*, 105, 26447–26461, 10.1029/2000JD900389, 2000.

- Safieddine SA, Heald CL, and Henderson BH: The global nonmethane reactive organic carbon budget: A modeling perspective, *Geophys. Res. Lett.*, 44, 3897–3906, 10.1002/2017GL072602, 2017.
- Santoni GW, Daube BC, Kort EA, Jiménez R, Park S, Pittman JV, Gottlieb E, Xiang B, Zahniser MS, Nelson DD, McManus JB, Peischl J, Ryerson TB, Holloway JS, Andrews AE, Sweeney C, Hall B, Hintsä E, Moore FL, Elkins JW, Hurst DF, Stephens BB, Bent J, and Wofsy SC: Evaluation of the airborne quantum cascade laser spectrometer (QCLS) measurements of the carbon and greenhouse gas suite – CO₂, CH₄, N₂O, and CO – during the Cal-Nex and HIPPO campaigns, *Atmos. Meas. Tech.*, 7, 1509–1526, 10.5194/amt-7-1509-2014, 2014.
- Saunders SM, Jenkin ME, Derwent RG, and Pilling MJ: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, 10.5194/acp-3-161-2003, 2003.
- Schlundt C, Tegtmeyer S, Lennartz ST, Bracher A, Cheah W, Krüger K, Quack B, and Marandino CA: Oxygenated volatile organic carbon in the western Pacific convective center: ocean cycling, air–sea gas exchange and atmospheric transport, *Atmospheric Chem. Phys.*, 17, 10837–10854, 10.5194/acp-17-10837-2017, 2017.
- Sherwen T, Schmidt JA, Evans MJ, Carpenter LJ, Groß-mann K, Eastham SD, Jacob DJ, Dix B, Koenig TK, Sinreich R, Ortega I, Volkamer R, Saiz-Lopez A, Prados-Roman C, Mahajan AS, and Ordóñez C: Global impacts of tropospheric halogens (Cl, Br, I) on oxidants and composition in GEOS-Chem, *Atmos. Chem. Phys.*, 16, 12239–12271, 10.5194/acp-16-12239-2016, 2016.
- Shetter RE and Muller M: Photolysis frequency measurements using actinic flux spectroradiometry during PEM-Tropics Mission: Instrumentation description and some results, *J. Geophys. Res.-Atmos.*, 104, 5647–5661, 10.1029/98JD01381, 1999.
- Shindell DT, Faluvegi G, Stevenson DS, Krol MC, Emmons LK, Lamarque J-F, Pétron G, Dentener FJ, Ellingsen K, Schultz MG, Wild O, Amann M, Atherton CS, Bergmann DJ, Bey I, Butler T, Cofala J, Collins WJ, Derwent RG, Doherty RM, Drevet J, Eskes HJ, Fiore AM, Gauss M, Hauglustaine DA, Horowitz LW, Isaksen ISA, Lawrence MG, Montanaro V, Müller J-F, Pitari G, Prather MJ, Pyle JA, Rast S, Rodriguez JM, Sanderson MG, Savage NH, Strahan SE, Sudo K, Szopa S, Unger N, van Noije TPC, and Zeng G: Multimodel simulations of carbon monoxide: Comparison with observations and projected near-future changes, *J. Geophys. Res.*, 111, D19306, 10.1029/2006JD007100, 2006.
- Simpson IJ, Blake NJ, Barletta B, Diskin GS, Fuelberg HE, Gorham K, Huey LG, Meinardi S, Rowland FS, Vay SA, Weinheimer AJ, Yang M, and Blake DR: Characterization of trace gases measured over Alberta oil sands mining operations: 76 speciated C₂–C₁₀ volatile organic compounds (VOCs), CO₂, CH₄, CO, NO, NO₂, NO_y, O₃ and SO₂, *Atmos. Chem. Phys.*, 10, 11931–11954, 10.5194/acp-10-11931-2010, 2010.
- Singh H, Chen Y, Staudt A, Jacob D, Blake D, Heikes B, and Snow J: Evidence from the Pacific troposphere for large global sources of oxygenated organic compounds, *Nature*, 410, 1078, 10.1038/35074067, 2001. [PubMed: 11323667]
- Singh HB, Tabazadeh A, Evans MJ, Field BD, Jacob DJ, Sachse G, Crawford JH, Shetter R, and Brune WH: Oxygenated volatile organic chemicals in the oceans: Inferences and implications based on atmospheric observations and air-sea exchange models, *Geophys. Res. Lett.*, 30, 1862, 10.1029/2003GL017933, 2003.
- Sinha V, Williams J, Crowley JN, and Lelieveld J: The Comparative Reactivity Method – a new tool to measure total OH Reactivity in ambient air, *Atmos. Chem. Phys.*, 8, 2213–2227, 10.5194/acp-8-2213-2008, 2008.
- Sinha V, Williams J, Lelieveld J, Ruuskanen TM, Kajos MK, Patokoski J, Hellen H, Hakola H, Mogensen D, Boy M, Rinne J, and Kulmala M: OH Reactivity Measurements within a Boreal Forest: Evidence for Unknown Reactive Emissions, *Environ. Sci. Technol.*, 44, 6614–6620, 10.1021/es101780b, 2010. [PubMed: 20687598]
- Staudt AC, Jacob DJ, Ravetta F, Logan JA, Bachiochi D, Krishnamurti TN, Sandholm S, Ridley B, and Singh HB: Sources and chemistry of nitrogen oxides over the tropical Pacific, *J. Geophys. Res.*, 108, 8239, 10.1029/2002JD002139, 2003.
- St. Clair JM, McCabe DC, Crouse JD, Steiner U, and Wennberg PO: Chemical ionization tandem mass spectrometer for the in situ measurement of methyl hydrogen peroxide, *Rev. Sci. Instrum.*, 81, 094102, 10.1063/1.3480552, 2010.

- Stockwell CE, Veres PR, Williams J, and Yokelson RJ: Characterization of biomass burning emissions from cooking fires, peat, crop residue, and other fuels with high-resolution proton-transfer-reaction time-of-flight mass spectrometry, *Atmos. Chem. Phys.*, 15, 845–865, 10.5194/acp-15-845-2015, 2015.
- Strode SA, Duncan BN, Yegorova EA, Kouatchou J, Ziemke JR, and Douglass AR: Implications of carbon monoxide bias for methane lifetime and atmospheric composition in chemistry climate models, *Atmos. Chem. Phys.*, 15, 11789–11805, 10.5194/acp-15-11789-2015, 2015.
- Tan D, Faloon I, Simpas JB, Brune W, Olson J, Crawford J, Avery M, Sachse G, Vay S, Sandholm S, Guan H-W, Vaughn T, Mastromarino J, Heikes B, Snow J, Podolske J, and Singh H: OH and HO₂ in the tropical Pacific: Results from PEM-Tropics B, *J. Geophys. Res.-Atmos.*, 106, 32667–32681, 10.1029/2001JD900002, 2001.
- Thames AB, Brune WH, Miller DO, Allen HM, Apel EC, Blake DR, Bui TP, Commane R, Crounse JD, Daube BC, Diskin GS, DiGangi JP, Elkins JW, Hall SR, Hanisco TF, Hannun RA, Hints E, Hornbrook RS, Kim MJ, McKain K, Moore FL, Nicely JM, Peischl J, Ryerson TB, St. Clair JM, Sweeney C, Teng A, Thompson CR, Ullmann K, Wennberg PO, and Wolfe GM: Missing OH reactivity in the global marine boundary layer, *Atmos. Chem. Phys.*, 20, 4013–4029, 10.5194/acp-20-4013-2020, 2020.
- Travis KR, Jacob DJ, Fisher JA, Kim PS, Marais EA, Zhu L, Yu K, Miller CC, Yantosca RM, Sulprizio MP, Thompson AM, Wennberg PO, Crounse JD, St. Clair JM, Cohen RC, Laughner JL, Dibb JE, Hall SR, Ullmann K, Wolfe GM, Pollack IB, Peischl J, Neuman JA, and Zhou X: Why do models overestimate surface ozone in the Southeast United States?, *Atmos. Chem. Phys.*, 16, 13561–13577, 10.5194/acp-16-13561-2016, 2016. [PubMed: 29619045]
- Tzompa-Sosa ZA, Mahieu E, Franco B, Keller CA, Turner AJ, Helmig D, Fried A, Richter D, Weibring P, Walega J, Yacovitch TI, Herndon SC, Blake DR, Hase F, Hannigan JW, Conway S, Strong K, Schneider M, and Fischer EV: Revisiting global fossil fuel and biofuel emissions of ethane, *J. Geophys. Res.-Atmos.*, 122, 2493–2512, 10.1002/2016JD025767, 2017.
- van der Werf GR, Randerson JT, Giglio L, van Leeuwen TT, Chen Y, Rogers BM, Mu M, van Marle MJE, Morton DC, Collatz GJ, Yokelson RJ, and Kasibhatla PS: Global fire emissions estimates during 1997–2016, *Earth Syst. Sci. Data*, 9, 697–720, 10.5194/essd-9-697-2017, 2017.
- Wang S, Hornbrook RS, Hills A, Emmons LK, Tilmes S, Lamarque J-F, Jimenez JL, Campuzano-Jost P, Nault BA, Crounse JD, Wennberg PO, Kim M, Allen H, Ryerson TB, Thompson CR, Peischl J, Moore F, Nance D, Hall B, Elkins J, Tanner D, Huey LG, Hall SR, Ullmann K, Orlando JJ, Tyndall GS, Flocke FM, Ray E, Hanisco TF, Wolfe GM, Clair JS, Commane R, Daube B, Barletta B, Blake DR, Weinzierl B, Dollner M, Conley A, Vitt F, Wofsy SC, Riemer DD, and Apel EC: Atmospheric Acetaldehyde: Importance of Air-Sea Exchange and a Missing Source in the Remote Troposphere, *Geophys. Res. Lett.*, 46, 5601–5613, 10.1029/2019GL082034, 2019. [PubMed: 32606484]
- Wang X, Jacob DJ, Eastham SD, Sulprizio MP, Zhu L, Chen Q, Alexander B, Sherwen T, Evans MJ, Lee BH, Haskins JD, Lopez-Hilfiker FD, Thornton JA, Huey GL, and Liao H: The role of chlorine in global tropospheric chemistry, *Atmos. Chem. Phys.*, 19, 3981–4003, 10.5194/acp-19-3981-2019, 2019.
- Whalley LK, Stone D, Bandy B, Dunmore R, Hamilton JF, Hopkins J, Lee JD, Lewis AC, and Heard DE: Atmospheric OH reactivity in central London: observations, model predictions and estimates of in situ ozone production, *Atmos. Chem. Phys.*, 16, 2109–2122, 10.5194/acp-16-2109-2016, 2016.
- Wofsy SC: HIAPER Pole-to-Pole Observations (HIPPO): fine-grained, global-scale measurements of climatically important atmospheric gases and aerosols, *Philos. T. Roy. Soc. A*, 369, 2073–2086, 10.1098/rsta.2010.0313, 2011.
- Wofsy SC, Afshar S, Allen HM, Apel E, Asher EC, Barletta B, Bent J, Bian H, Biggs BC, Blake DR, Blake N, Bourgeois I, Brock CA, Brune WH, Budney JW, Bui TP, Butler A, Campuzano-Jost P, Chang CS, Chin M, Commane R, Correa G, Crounse JD, Cullis PD, Daube BC, Day DA, Dean-Day JM, Dibb JE, DiGangi JP, Diskin GS, Dollner M, Elkins JW, Erdesz F, Fiore AM, Flynn CM, Froyd K, Gesler DW, Hall SR, Hanisco TF, Hannun RA, Hills AJ, Hints E, Hoffman A, Hornbrook RS, Huey LG, Hughes S, Jimenez JL, Johnson BJ, Katich JM, Keeling RF, Kim MJ, Kupc A, Lait LR, Lamarque J-F, Liu J, McKain K, McLaughlin RJ, Meinardi S, Miller DO,

Montzka SA, Moore FL, Morgan EJ, Murphy DM, Murray LT, Nault BA, Neuman JA, Newman PA, Nicely JM, Pan X, Paplawsky W, Peischl J, Prather MJ, Price DJ, Ray E, Reeves JM, Richardson M, Rollins AW, Rosenlof KH, Ryerson TB, Scheuer E, Schill GP, Schroder JC, Schwarz JP, St. Clair JM, Steenrod SD, Stephens BB, Strode SA, Sweeney C, Tanner D, Teng AP, Thames AB, Thompson CR, Ullmann K, Veres PR, Vieznor N, Wagner NL, Watt A, Weber R, Weinzierl B, Wennberg P, Williamson CJ, Wilson JC, Wolfe GM, Woods CT, and Zeng LH: ATom: Merged Atmospheric Chemistry, Trace Gases, and Aerosols. ORNL DAAC, Oak Ridge, Tennessee, USA, 10.3334/ORNLDAAC/1581, 2018.

- Wolfe GM, Nicely JM, St. Clair JM, Hanisco TF, Liao J, Oman LD, Brune WB, Miller D, Thames A, González Abad G, Ryerson TB, Thompson CR, Peischl J, Mc-Cain K, Sweeney C, Wennberg PO, Kim M, Crouse JD, Hall SR, Ullmann K, Diskin G, Bui P, Chang C, and Dean-Day J: Mapping hydroxyl variability throughout the global remote troposphere via synthesis of airborne and satellite formaldehyde observations, *P. Natl. Acad. Sci. USA*, 116, 11171–11180, 10.1073/pnas.1821661116, 2019.
- Wong JPS, Zhou S, and Abbatt JPD: Changes in Secondary Organic Aerosol Composition and Mass due to Photolysis: Relative Humidity Dependence, *J. Phys. Chem. A*, 119, 4309–4316, 10.1021/jp506898c, 2015. [PubMed: 25196234]
- Wu. H, Wang Y, Li H, Huang L, Huang D, Shen H, Xing Y, and Chen Z: The OH-initiated oxidation of atmospheric peroxyacetic acid: Experimental and model studies, *Atmos. Environ*, 164, 61–70, 10.1016/j.atmosenv.2017.05.038, 2017.
- Yang M, Beale R, Liss P, Johnson M, Blomquist B, and Nightingale P: Air–sea fluxes of oxygenated volatile organic compounds across the Atlantic Ocean, *Atmos. Chem. Phys*, 14, 7499–7517, 10.5194/acp-14-7499-2014, 2014.
- Ye C, Zhou X, Pu D, Stutz J, Festa J, Spolaor M, Tsai C, Cantrell C, Mauldin RL, Campos T, Weinheimer A, Hornbrook RS, Apel EC, Guenther A, Kaser L, Yuan B, Karl T, Haggerty J, Hall S, Ullmann K, Smith JN, Ortega J, and Knote C: Rapid cycling of reactive nitrogen in the marine boundary layer, *Nature*, 532, 489–491, 10.1038/nature17195, 2016. [PubMed: 27064904]
- Young PJ, Archibald AT, Bowman KW, Lamarque J-F, Naik V, Stevenson DS, Tilmes S, Voulgarakis A, Wild O, Bergmann D, Cameron-Smith P, Cionni I, Collins WJ, Dalsøren SB, Doherty RM, Eyring V, Faluvegi G, Horowitz LW, Josse B, Lee YH, MacKenzie IA, Nagashima T, Plummer DA, Righi M, Rumbold ST, Skeie RB, Shindell DT, Strode SA, Sudo K, Szopa S, and Zeng G: Preindustrial to end 21st century projections of tropospheric ozone from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys*, 13, 2063–2090, 10.5194/acp-13-2063-2013, 2013.
- Yoshino A, Sadanaga Y, Watanabe K, Kato S, Miyakawa Y, Matsumoto J, and Kajii Y: Measurement of total OH reactivity by laser-induced pump and probe technique—comprehensive observations in the urban atmosphere of Tokyo, *Atmos. Environ*, 40, 7869–7881, 10.1016/j.atmosenv.2006.07.023, 2006.
- Yu S, Mathur R, Sarwar G, Kang D, Tong D, Pouliot G, and Pleim J: Eta-CMAQ air quality forecasts for O₃ and related species using three different photochemical mechanisms (CB4, CB05, SAPRC-99): comparisons with measurements during the 2004 ICARTT study, *Atmos. Chem. Phys*, 10, 3001–3025, 10.5194/acp-10-3001-2010, 2010.
- Zannoni N, Gros V, Lanza M, Sarda R, Bonsang B, Kalogridis C, Preunkert S, Legrand M, Jambert C, Boissard C, and Lathiere J: OH reactivity and concentrations of biogenic volatile organic compounds in a Mediterranean forest of downy oak trees, *Atmos. Chem. Phys*, 16, 1619–1636, 10.5194/acp-16-1619-2016, 2016.
- Zannoni N, Gros V, Sarda Esteve R, Kalogridis C, Michoud V, Dusanter S, Sauvage S, Locoge N, Colomb A, and Bonsang B: Summertime OH reactivity from a receptor coastal site in the Mediterranean Basin, *Atmos. Chem. Phys*, 17, 12645–12658, 10.5194/acp-17-12645-2017, 2017.
- Zhang L, Jacob DJ, Knipping EM, Kumar N, Munger JW, Carouge CC, van Donkelaar A, Wang YX, and Chen D: Nitrogen deposition to the United States: distribution, sources, and processes, *Atmos. Chem. Phys*, 12, 4539–4554, 10.5194/acp-12-4539-2012, 2012.




Figure 1. Annual mean 2016 (a) surface (log scale) and (b) zonal mean cOHR calculated from individual model species. The GEOS-Chem species included in the calculation of cOHR are listed in Table S3.




Figure 2.
ATom-1 ocean-only flight tracks colored by altitude.




Figure 3. Median OH concentrations for the Northern Hemisphere ($> 0^\circ \text{N}$) and Southern Hemisphere ($< 0^\circ \text{S}$) from the ATHOS instrument described in Table 2 during ATom-1 (July–August 2016) and ATom-2 (January–February 2017) compared against the GEOS-Chem model in 0.5 km altitude bins. The observations have been filtered to remove biomass burning (acetonitrile > 200 ppt) and stratospheric ($\text{O}_3/\text{CO} > 1.25$) influence. The dashed lines show the observed 25th–75th percentiles.




Figure 4. The same as Fig. 3 for median water vapor concentrations. Water vapor mixing ratio was measured by the DLH instrument as described in Table 2.




Figure 5.

The same as Fig. 3 for median photolysis frequencies for ozone ($j(\text{O}^1\text{D})$). The actinic flux measured by the CAFS instrument is used to calculate $j(\text{O}^1\text{D})$ as described in Table 2.




Figure 6. The same as Fig. 3 for median ozone concentrations. Ozone was measured by the NOAA NO_yO_3 instrument as described in Table 2.




Figure 7. The same as Fig. 3 for median NO_y (a) and NO (b) concentrations. NO_y and NO were measured by the NOAA NO_yO_3 instrument as described in Table 2.




Figure 8.

Comparison of modeled and observed HNO₃, ozone, NO_y, and NO with sensitivity studies including scaling emissions from the US and Asia, improved chlorine chemistry (X. Wang et al., 2019), and the photolysis of particulate nitrate on coarse-mode sea-salt aerosols (Kasibhatla et al., 2018) as described in Sect. 4.1. HNO₃ was measured by the Caltech CIMS; ozone, NO_y, and NO were measured by the NOAA NO_yO₃ instrument (Table 2).




Figure 9.

The same as Fig. 3 for median OHR. OHR was measured by the ATHOS instrument as described in Table 2. The calculation of cOHR in the model and observations includes the species described in Table 2. In order to allow for a point-by-point comparison of cOHR in the model and observations, missing values are filled in the observational components of cOHR using linear interpolation. All calculated reactivity values are determined using the temperature and pressure of the ATHOS instrument inlet, which differ from ambient values. The sensitivity tests are described in Sect. 5.




Figure 10.
Impact of all ocean emissions (Tables 3 and 4) on annual simulated 2016 surface cOHR as described in the text.




Figure 11.

Median observed and modeled OHR and cOHR (see text) below 3 km in the Northern Hemisphere ($> 0^\circ \text{N}$) and Southern Hemisphere ($< 0^\circ \text{S}$) during ATom-1. The “Other” category is the following species as described in Table 2: ethanol, propane, ethane, acetone, $> \text{C}_3$ aldehydes, methyl ethyl ketone, methyl vinyl ketone, methacrolein, benzene, toluene, $> \text{C}_4$ alkanes, peroxyacetic acid, peroxyacetic acid, dimethyl sulfide, nitric acid, NO, and NO₂. The diameter of each pie chart is scaled relative to the maximum cOHR for ATom-1.




Figure 12. Same as Fig. 10 but for ATom-2. The diameter of each pie chart is scaled relative to the maximum cOHR for ATom-2.




Figure 13.

The same as Fig. 3 for median acetaldehyde profiles. Acetaldehyde was measured by the TOGA instrument as described in Table 2. The sensitivity studies are described in Sect. 5.1.




Figure 14.

The same as Fig. 3 for median peroxyacetic acid (PAA) profiles. PAA was measured by the Caltech CIMS instrument as described in Table 2. The sensitivity studies are described in Sect. 5.1.




Figure 15.

The same as Fig. 3 for median peroxyacetyl nitrate (PAN) profiles. PAN was measured by the PANTHER instrument as described in Table 2. The sensitivity studies are described in Sect. 5.1.

Table 1.

Annual emissions of CO and NO_x for 2016 used in the GEOS-Chem simulations.

Emissions category	CO, Tg	Emissions category	NO _x , Tg N
Fuel combustion *	590	Fuel combustion *	32.9
Biomass burning	311	Biomass burning	6.0
NM VOC oxidation	698	Soil emissions	7.8
Methane oxidation	936	Lightning emissions	6.0
Total	2535	Total	52.7

* Anthropogenic fossil fuel and biofuel combustion.

Table 2.

Description of ATom measurements used to evaluate the model simulation.

Measurement	Instrument	Accuracy	Detection limit/precision	Reference
OHR	Airborne Tropospheric Hydrogen Oxides Sensor (ATHOS)	0.8 s^{-1}	$\pm 0.4 \text{ s}^{-1}$	Faloona et al. (2004); Mao et al. (2009)
Water vapor	Diode laser hygrometer (DLH)	5%	0.1% or 50 ppb	Diskin et al. (2002); Podolske et al. (2003)
NO_y	NOAA nitrogen oxides and ozone (NO_yO_3)		0.05 ppb^2	Pollack et al. (2010); Ryerson et al. (1998, 2000)
Photolysis frequencies via actinic flux	Charged-coupled device Actinic flux Spectroradiometers (CAFS)	J_{O_3} 20% J_{NO_2} 12%	$J_{\text{O}_3} 10^{-7} \text{ s}^{-1}$ $J_{\text{NO}_2} 10^{-6} \text{ s}^{-1}$	Shetter and Mueller (1999), Petropavloskikh et al. (2007), Hofzumahaus et al. (2004)
Peroxyacetyl nitrate (PAN)	PAN and trace Hydrohalocarbon Experiment (PANTHER)	10%	2 ppt $\pm 10\%$	Elkins et al. (2001); Wofsy et al. (2011)
Components of OH reactivity ³				
CH_4	NOAA Picarro	0.6 ppb	0.3 ppb	Karion et al. (2013)
CO	Harvard Quantum Cascade Laser System (QCLS)	3.5 ppb	0.15 ppb	McManus et al. (2005); Santoni et al. (2014)
H_2 ⁴	UAS Chromatograph for Atmospheric Trace Species (UCATS)/PANTHER		7.5 ppb^5	Hintsa et al. (2020)
$\text{NO}, \text{NO}_2, \text{O}_3$	NOAA NO_yO_3		$0.006, 0.03, 1.7 \text{ ppb}^2$	Pollack et al. (2010); Ryerson et al. (1998, 2000)
Methyl hydroperoxide, nitric acid, hydrogen peroxide, peroxyacetic acid, peroxyacetic acid	Caltech Chemical ionization mass spectrometer (CIMS)	$\pm 30\%, \pm 30\%, \pm 30\%$, $\pm 50\%, \pm 30\%$	25, 50, 50, 30, 100 ppt	St. Clair et al. (2010); Crouse et al. (2006)
Formaldehyde	NASA In Situ Airborne Formaldehyde (ISAF)	10%	10 ppt	Cazorla et al. (2015); DiGangi et al. (2011); Hottle et al. (2009)
Methanol, acetaldehyde, propane, dimethyl sulfide, ethanol, acetone, methyl ethyl ketone, propanal, butanal ⁶ , toluene, methyl vinyl ketone, methacrolein i-Butane + n-butane + i-pentane + n-pentane ⁷	NCAR Trace Organic Gas Analyzer (TOGA)	30%, 20%, 30%, 15%, 15%, 15%, 15%	10, 10, 20, 2, 30, 10, 2, 20, 2, 0.6, 4, 2, 2, 2, 4, 4 ppt	Apel et al. (2015)
OH, HO_2	ATHOS	74% to 135%	0.018, 0.2 ppt	Faloona et al. (2004); Brune et al. (2020)
Ethane, benzene	UCI Whole air sampler (WAS)	5%, 5%	3, 3 ppt	Colman et al. (2001); Simpson et al. (2010)

¹ Model NO_y is defined as $\text{NO} + \text{NO}_2 + \text{HONO} + \text{HONO}_3 + \text{HNO}_4 + 2 \times \text{N}_2\text{O}_5 + \text{ClNO}_2 + \Sigma\text{PNs} + \Sigma\text{ANs}$.

² Average of 2σ uncertainty for each individual 1 Hz measurement for ATom-1 and ATom-2.

³ Included in cOHR are observations of species where at least 20% of the possible available measurements below 3 km are not missing.

⁴ The GEOS-Chem concentration of H_2 is set to a constant value of 500 ppt.

⁵ Average of reported error for each individual measurement for ATom-1 and ATom-2.

⁶ Lumped as > C4 alkanes (ALK4) in GEOS-Chem.

⁷ Lumped as > C3 aldehydes (RCHO) in GEOS-Chem.

Table 3.

Biogenic ocean emissions of VOCs.

GEOS-Chem 2 species ²	No. of lumped species	Produces acetaldehyde?	Annual net emissions (Tg C) ¹	Reference for seawater concentration
ALD2	1	Yes	12.02	Millet et al. (2010)
MOH	1	No	-1.54	Personal communication, Dylan B. Millet, 2018
ACET	1	No	-75.65	Fischer et al. (2012)
LIMO	1	Yes	0.04	Hackenberg et al. (2017)
MTPA	3	Yes	0.05	Hackenberg et al. (2017)
MTPQ	2	Yes	0.06	Hackenberg et al. (2017)
EOH	1	Yes	-5.52	Beale et al. (2010)
C2H6	1	Yes	0.33	Plass-Dülmer et al. (1993)
C2H4	1	No	0.75	Plass-Dülmer C. et al. (1993)
PRPE	2	Yes	0.95	Plass-Dülmer C. et al. (1993)
C3H8	1	Yes	0.16	Plass-Dülmer et al. (1993)
ALK4	2	Yes	0.12	Plass-Dülmer et al. (1993)
C2H2	1	No	0.02	Plass-Dülmer et al. (1993)
ISOP	1	Yes	1.64	Arnold et al. (2009)
RCHO	1	Yes	-1.03	Singh et al. (2003)
MEK	1	Yes	-7.214	Schlundt et al. (2017)
Total net emission			-74.82	
Total net emission producing acetaldehyde			1.60	

¹Net ocean emissions = upward flux out of the ocean–ocean uptake.²More information on the GEOS-Chem species definitions can be found here: http://wiki.seas.harvard.edu/geos-chem/index.php/Species_in_GEOS-Chem (last access: 21 May 2020).

Table 4.

Abiotic ocean emissions of VOCs according to Brüggemann et al. (2018)¹.

GEOS-Chem species ²	No. of lumped species	Produces acetaldehyde?	Annual emission (Tg C)
ACET	1	No	10.07
EOH	1	Yes	5.16
ALD2	1	Yes	2.26
MOH		No	0.79
RCHO	21	Yes	3.88
ISOP	1	Yes	1.04
PRPE	13	Yes	4.44
MACR	1	Yes	0.42
ACTA	1	Yes	0.10
CH2O	1	No	0.03
XYLE	1	No	0.05
TOLU	1	No	0.04
BENZ	1	No	0.02
Total net emission			28.30
Total net emission producing acetaldehyde			17.30

¹Table S2 shows the emission factor assumed for each species and the lumping methodology for Table 4.

²More information on the GEOS-Chem species definitions can be found here: http://wiki.seas.harvard.edu/geos-chem/index.php/Species_in_GEOS-Chem (last access: 21 May 2020).

Table 5.

Model sources of acetaldehyde in 2016.

Sources (Tg yr ⁻¹)*	Millet et al. (2010)	This work
Photochemical production	128	166
Net ocean emission	57	22
Terrestrial plant growth + decay	23	26
Biomass burning	3	3
Anthropogenic emission	2	2
Total source	213	219

* Emissions are given in Tg of acetaldehyde per year for comparison to Millet et al. (2010). These totals are for the baseline model simulation described in Sect. 2.1.