

Lawrence Berkeley National Laboratory

Recent Work

Title

THE CRYSTAL STRUCTURE OF ACETIC ACID

Permalink

<https://escholarship.org/uc/item/3x45b5nd>

Author

Jones, Robert E.

Publication Date

1956-12-14

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

THE CRYSTAL STRUCTURE OF ACETIC ACID

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-3641
Chemistry Distribution

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California
Contract No. W-7405-eng-48

THE CRYSTAL STRUCTURE OF ACETIC ACID

Robert E. Jones

December 14, 1956

Printed for the U. S. Atomic Energy Commission

THE CRYSTAL STRUCTURE OF ACETIC ACID

Contents

Abstract	3
I. Introduction	4
II. Experimental Procedure	4
A. Single Crystal Photographs	4
B. Powder Photographs	5
III. Unit Cell and Space Group	6
IV. Intensity Corrections	6
V. Determination of the Structure	7
VI. Least Square Refinement	8
VII. Discussion of the Structure	12
Acknowledgments	14
References	15

THE CRYSTAL STRUCTURE OF ACETIC ACID

Robert E. Jones

Radiation Laboratory and
Department of Chemistry and Chemical Engineering
University of California, Berkeley 4, California

December 14, 1956

ABSTRACT

The crystal structure of acetic acid was determined using x-ray diffraction techniques and a least squares method of refinement. The molecules are planar within experimental error and, except for the single bond carbon oxygen distance which is shortened to 1.29 A, the bond lengths were found to be in agreement with previous determinations in the gas phase by electron diffraction and infrared spectroscopy. The molecules are linked together by hydrogen bonds in the infinite chains similar to the structure of crystalline formic acid. It appears that van der Waals forces can account for the adhesion between chains.

THE CRYSTAL STRUCTURE OF ACETIC ACID

Robert E. Jones

Radiation Laboratory and
Department of Chemistry and Chemical Engineering
University of California, Berkeley 4, California

December 14, 1956

I. INTRODUCTION

The structure of acetic acid (CH_3COOH) has been investigated in the gas by electron diffraction¹ and by infrared spectroscopy.² This work was undertaken to give further information on the structure of the molecule and on the forces between molecules in the solid state. No x-ray diffraction determination of the structure has been reported before.

The normal fatty acids, of which acetic acid is a member, are thought to form dimers in the gaseous and liquid state. Investigations of high molecular weight members of this series have shown the dimers are also formed in the solid.^{3,4} However, the lowest member of this series, formic acid, was shown to form infinite chains rather than dimers with hydrogen bonds forming the intermolecular linkages.⁵ There was some spectroscopic evidence that acetic acid also formed polymers rather than dimers.⁶ The results of this work confirm this and show the molecules in the solid are linked together as in formic acid.

II. EXPERIMENTAL PROCEDURE

A. Single Crystal Photographs

Single crystals of acetic acid were prepared by freezing C. P. grade glacial acetic acid (99.8% pure) in a jet of cold nitrogen gas. The liquid was sealed in a glass capillary and quickly frozen into a translucent aggregate of small crystals. By then raising the temperature and maintaining it near the melting point for several hours, a transparent single crystal was grown. Its growth was followed by periodically taking oscillation pictures until the diffraction pattern showed that the small crystallites had vanished.

The flow of cold nitrogen was evolved from liquid nitrogen electrically heated by a resistor in a dewar flask. The rate of flow was controlled by adjusting the current in the resistor. Since the crystal had to be maintained for many days, two duplicate nitrogen boiling units were made so that one could be taken out of operation and refilled without interrupting the flow for more than a few seconds.

The single crystal photographs were taken on a Weissenberg camera using Cu K α x-rays ($\lambda = 1.5418$ A). The single crystals showed no tendency to be formed with a unit cell axis parallel to the capillary axis, so they were aligned using oscillation photographs. The Weissenberg photographs were taken using a modified film holder. The geometry of the pictures was that of an ordinary Weissenberg photograph with only the lower half of the film recording reflections. The upper half was absent to permit cooling of the sample.

The single crystal Weissenberg pictures were taken at a temperature within a few degrees of the melting point (16.6°C). While pictures were being run the capillary contained liquid acetic acid, probably either because of impurities reducing the melting point or because of a temperature gradient along the capillary.

B. Powder Photographs

Powder photographs were taken to obtain lattice constants. A modified Norelco camera was used and the temperature of the sample recorded by an iron-constantan thermocouple. The cooling was by a flow of nitrogen produced and controlled in the same way as the single crystal experiments.

The powder photographs used in measuring the lattice constants were taken at $-4 \pm 2^{\circ}\text{C}$ using Cu K α x-rays. Although the single crystal photographs were taken at about 16° , it was impossible to take powder pictures at this temperature because the sample would anneal and form large crystals in less time than necessary to take good powder photographs. To determine the effect of lattice expansion from -4 to 16°C , and possibly extrapolate the values of the cell constants to the higher temperature, pictures were also taken at $-22 \pm 2^{\circ}\text{C}$. However, the difference in lattice constants measured at -4°C and -22°C could not be

detected because of the high error in the determination and the values measured at -4°C are reported below.

III. UNIT CELL AND SPACE GROUP

From oscillation, rotation and three layers of Weissenberg pictures it was deduced that the unit cell was orthorhombic, in agreement with the observations of Steinmetz⁷ on the cleavage and optical properties of acetic acid crystals. From the powder photographs cell constants were measured:

$$a = 13.31 \pm .04 \text{ \AA} \quad b = 4.085 \pm .012 \text{ \AA} \quad c = 5.77 \pm .02 \text{ \AA}$$

The inaccuracy of these determinations is high because high angle θ lines were not observed. With four molecules per unit cell, the calculated density (1.26 g/cm^3) agrees with that observed by deVisser⁸ (1.26585 g/cm^3) for solid acetic acid at its melting point.

Zero, first, and second layer Weissenberg pictures were taken rotating about the b axis. On these photographs reflections were observed for $h0l$ only if $h = 2n$ and for okl only if $k + l = 2n$. These extinction rules correspond to probable space groups $\text{Pna}2_1$ and Pnam . A satisfactory structure was found using the space group $\text{Pna}2_1 - \text{C}_{2v}^9$ with atoms in the positions $4(a):^3$

$$(x, y, z; \bar{x}, \bar{y}, 1/2 + z; 1/2 - x, 1/2 + y, 1/2 + z; 1/2 + x, 1/2 - y, z)$$

IV. INTENSITY CORRECTIONS

Estimates of the intensities of the reflections were made visually from the Weissenberg films and corrections made for the Lorentz and polarization factors and for the velocity effect. No correction was made for absorption by the sample or the capillary, since this correction was estimated to be small.

The corrections were made using an IBM 650 computer. In addition to making the above corrections, the program also calculated for each reflection the magnitude of the observed structure factors (F_o), $\sin^2 \theta$ values, and scattering factors; determined which formula would be used for calculating the structure factor; and determined the use of the

weighting factor used in the program for least squares refinement. The output cards for this program could be used as input cards in the least squares refinement program.

V. DETERMINATION OF THE STRUCTURE

A trial structure was constructed with molecules hydrogen-bonded in chains approximately parallel to the $\{011\}$ and $\{0\bar{1}1\}$ planes similar to the structure of formic acid. Using estimated bond lengths and angles, x and z parameters were calculated. Using these parameters and the observed structure factors, a projection of the electron density along $[010]$ was calculated. The peaks were well resolved and very close to the trial positions; however, the reliability factors (defined in Section VI) showed that the fit of observed structure factors to calculated structure factors was very poor and furthermore the parameters could not be properly refined to give better reliability factors. Since well resolved peaks in a Fourier projection occurring very close to the trial positions were not adequate criteria for telling whether the position parameters for the atoms were correct, no further Fouriers were computed and the parameters were adjusted by trial and error until a set was found that could be refined to give a reliability factor

$$(R_1 = \frac{\sum | | F_o | - | F_c | |}{\sum | F_o |}), \text{ where } F_o \text{ is the observed}$$

structure factor and F_c the calculated structure factor) of approximately 18% and satisfactory bond lengths. Fig. 1 is a $[010]$ projection calculated using observed structure factors from the final refinement using all the data.

Knowing approximate bond lengths and values for the x and z parameters, the differences in the y parameters of the atoms were estimated. It remained, however, to fix the y parameter of one of the atoms. This was done approximately by studying a model of the chain-like structure and adjusting the y parameters so that no intermolecular distances were too close. The final trial y parameters were determined by trial and error attempts to find parameters that would refine properly using the least squares refinement program.

VI. LEAST SQUARES REFINEMENT

A least squares refinement of the final trial parameters was made using an IBM 650 program written by Senko and Templeton.⁹ The results are listed in Table I.

Table I

	x	σ_x	y	σ_y	\bar{z}	σ_z	B
O _I	.1295	.0003	.1032	.0017	.0000	.0011	3.92
O _{II}	.2526	.0004	.3690	.0018	.1780	.0014	4.51
C _I	.1641	.0006	.2761	.0026	.1701	.0023	3.73
C _{II}	.0868	.0006	.3720	.0030	.3570	.0018	4.01

σ_x , σ_y , and σ_z are the standard deviations of the respective coordinates calculated by the program. The values for B were used in the expression for the temperature factor, $\exp \{ -B \sin^2 \theta / \lambda^2 \}$, and correspond to mean square vibrational amplitudes ($\bar{\mu}^2 = B/8\pi^2$) from .047 to .057 Å². In addition to the three coordinates and B for each atom, the scale factor for each layer of reflections was refined.

Three reliability factors were calculated using all the data and with the observed structure factor (F_o) set equal to zero for unobserved reflections. The results were:

$$R_1 = \frac{\sum | | F_o | - | F_c | |}{\sum | F_o |} = 15.8\%$$

$$R_2 = \left[\frac{\sum | | F_o | - | F_c | |^2}{\sum | F_o |^2} \right]^{1/2} = 16.0\%$$

$$R_3 = \left[\frac{\sum w | | F_o | - | F_c | |^2}{\sum w | F_o |^2} \right]^{1/2} = 16.1\%$$

F_c is the magnitude of the calculated structure factor and w is a factor which weights each reflection according to the accuracy of the intensity measurement. In Senko's program $w = 1/4 F_{\min}$ if $F_o < 4 F_{\min}$ and $w = 1/F_o$ if $F_o > F_{\min}$. F_{\min} is the minimum observed structure factor.

Table II

Calculated and Observed Structure Factors of Acetic Acid

Index	Observed F	Calculated F	Index	Observed F	Calculated F
h00			h04		
0	--	280	0	24	26
2	*	79	2	27	28
4	50	45	4	19	19
6	34	33	6	12	10
8	46	42	8	16	15
10	24	22	10	13	12
12	16	15	12	15	14
14	< 11	4	h05		
h01			2	20	23
2	68	90	4	< 12	3
4	44	37	6	13	11
6	37	40	h06		
8	27	25	0	23	25
10	23	23	2	11	7
12	< 12	5	h10		
14	11	11	1	*	6
h02			2	*	96
0	31	38	3	28	28
2	72	74	4	5	9
4	72	65	5	< 5	3
6	19	21	6	25	28
8	22	20	7	25	26
10	23	23	8	10	10
12	22	20	9	8	11
14	10	9	10	< 8	6
h03			11	13	17
2	32	34	12	< 8	1
4	30	24	13	< 8	5
6	22	20	14	< 7	4
8	20	18	15	7	5
10	16	14	h11		
12	< 11	1	0	*	110

*reflection blocked by beam catcher

Table II (cont'd)

Index	Observed	F	Calculated	F	Index	Observed	F	Calculated	F
h11 (cont'd)					5	7		6	
1	60		55		6	8		6	
2	54		54		7	11		12	
3	52		56		8	< 8		3	
4	17		15		9	22		21	
5	45		40		10	14		13	
6	18		19		11	< 8		2	
7	21		23		12	10		10	
8	21		21		h14				
9	< 7		5		1	15		18	
10	17		18		2	9		6	
11	13		12		3	11		12	
12	14		16		4	10		11	
13	11		10		5	11		13	
14	8		5		6	9		7	
h12					7	9		6	
1	34		33		8	10		10	
2	10		11		9	< 8		4	
3	24		25		10	10		10	
4	27		27		h15				
5	21		22		0	22		26	
6	13		14		1	11		8	
7	13		18		2	11		9	
8	17		19		3	10		9	
9	< 8		6		4	8		5	
10	14		14		5	9		11	
11	< 8		6		6	< 8		3	
12	< 8		2		7	8		5	
h13					8	9		6	
0	13		9		h16				
1	31		34		1	< 7		3	
2	33		34		2	9		11	
3	15		16		h17				
4	28		30		0	6		6	

Table II (cont'd)

Index	Observed F	Calculated F	Index	Observed F	Calculated F
h20			6	11	11
1	*	28	7	16	18
2	*	11	8	8	3
3	35	33	9	18	21
4	< 4	1	10	< 7	6
5	41	49	11	9	6
6	9	10	h23		
7	11	12	1	< 6	7
8	7	8	2	15	16
9	7	8	3	16	16
10	< 7	5	4	9	10
11	16	17	5	< 7	4
12	8	9	6	8	5
13	14	16	7	17	19
h21			8	9	8
1	42	37	9	< 7	1
2	13	16	10	< 7	2
3	24	22	11	7	8
4	13	13	h24		
5	24	26	0	16	12
6	< 6	5	1	22	22
7	9	8	2	< 7	5
8	11	9	3	18	19
9	12	10	4	10	6
10	9	8	5	< 7	7
11	10	9	6	< 7	7
h22			7	11	10
0	21	23	8	< 7	2
1	42	44	9	9	8
2	12	10	h25		
3	32	34	1	9	7
4	8	11	2	8	5
5	9	9			

VII. DISCUSSION OF THE STRUCTURE

Molecules of acetic acid are linked together by hydrogen bonds in infinite chains. Forces between chains are presumed to be van der Waals attractions.

The bond lengths and angles in the carboxyl group fall within the range of best values listed in a review paper on the structure of the carboxyl group by Davies and Thomas.¹⁰ The carbon-oxygen bond lengths are $1.29 \pm .02$ A and $1.24 \pm .02$ A; the latter is assumed to be the double bond. The carbon-carbon distance is $1.54 \pm .02$ A. The molecule is planar within experimental error and the bond angles are: $\angle C - C = O = 122 \pm 2^\circ$, $\angle C - C - O = 116 \pm 2^\circ$, and $\angle O = C - O = 122 \pm 2^\circ$. These values are compared with the results of determinations in the gas phase^{1,2} in Table III.

The $C = O$ and $C - C$ bond lengths are within experimental error of those found for the monomer and dimer by electron diffraction,¹ however the $C - O$ distance is considerably shorter. This difference in the $C - O$ bond length has also been found in formic acid gas and solid.^{5,11}

These results for the $C - O$ and $C = O$ bond lengths are also within the limits of error assigned to the Davies and Sutherland results; however their lengths lie outside of the error of our work.

Adjacent molecules in the chain are bound together by hydrogen bonds. Figure 2 shows a section of the chain which is roughly parallel to the $\{011\}$ plane. Adjacent molecules in the chain are not coplanar or parallel. For comparison a section is shown in Figure 3 of the formic acid chain which is likewise roughly parallel to the $\{011\}$ plane in that crystal.

The $O - H \dots O$ bond distance was found to be $2.61 \pm .02$ A. Ubbelohde and Gallagher¹² have compiled a list of oxygen-oxygen hydrogen bond distances in monocarboxylic acids and this value is within the range of values reported in their paper.

The $C = O \dots H - O$ angle is $144 \pm 2^\circ$ and the $C - O - H \dots O$ angle is $122 \pm 2^\circ$. Holtzberg, Post, and Fankuchen⁵ have reported $122 \pm 1^\circ$ and $114 \pm 1^\circ$ for these two angles in formic acid. The larger values in acetic acid are probably due to the size of the methyl group which has made an increase in the carbon-carbon intermolecular distance.

Table III
Bond Lengths and Angles for Acetic Acid

	Electron Diffraction		Infrared Spectroscopy		Author
	J. Karle and L. O. Brockway ¹		M. Davies and G. B. B. M. Sutherland ²		
	Monomer	Dimer	Monomer	Dimer	
C = O	1.24 ± .03 A	1.25 ± .03 A	1.195 A (± 3%)	1.211 (± 3%)	1.24 ± .02 A
C - O	1.43 ± .03	1.36 ± .04	1.298 (± 3%)	1.285 (± 3%)	1.29 ± .02
C - C	1.54 ± .04	1.54 ± .04	-	-	1.54 ± .02
∠C - C = O	113 - 128°	120 ± 5°	-	-	122 ± 2°
∠C - C - O	125 - 94°	110° ± 5°	-	-	116 ± 2°
∠O = C - O	122 - 138°	130 ± 3°	-	-	122 ± 2°

Aside from the oxygen-oxygen hydrogen bond distance, all other intermolecular distances are consistent with van der Waals radii. The shortest of these intermolecular distances is a carbon-oxygen distance of 3.47 Å between adjacent molecules in the chain. This distance is larger than the sum of the methyl and oxygen van der Waals radii listed by Pauling.¹³

Figure 4 is a picture of the structure as a whole.


ACKNOWLEDGMENTS

I am grateful for the guidance and assistance of Professor David H. Templeton throughout this research. I also wish to express thanks to Mrs. Carol H. Dauben and Mrs. Helena W. Ruben for their help in many facets of this work, to Dr. Michael E. Senko and to Dr. Allan Zalkin in particular for the use of their computing programs, and to Mrs. Jane Waite and Miss Pat Hagen for help in preparing this final manuscript.

This work was performed under the auspices of the U.S. Atomic Energy Commission.


REFERENCES

1. J. Karle and L. O. Brockway, J. Am. Chem. Soc. 66, 574 (1944).
2. M. Davies and G. B. B. M. Sutherland, J. Chem. Phys. 6, 755 (1938).
3. V. Vand, W. M. Morley, and T. R. Lomer, Acta Cryst. 4, 324 (1951).
4. E. von Sydow, Arkiv for Kemi 9, 231 (1956).
5. F. Holtzberg, B. Post, and I. Fankuchen, Acta Cryst. 6, 127 (1953).
6. C. Rigaux, Compt. Rend. 238, 783 (1954).
7. H. Steinmetz, Z. Krist. 56, 157 (1921).
8. M. L. E. O. de Visser, Rec. trav. chim. Pays-Bas 12, 139 (1893).
9. M. Senko, Dissertation, University of California (1956).
10. M. Davies and O. Thomas, Disc. Faraday Soc. 9, 335 (1951).
11. I. L. Karle and J. Karle, J. Chem. Phys. 22, 43 (1954).
12. A. R. Ubbelohde and K. J. Gallagher, Acta Cryst. 8, 71 (1955).
13. L. Pauling, The Nature of the Chemical Bond, 2nd ed., Ithaca: Cornell University Press (1942).




MU-12738

Fig. 1 [010] projection of acetic acid. Contours are at intervals of $1 \text{ e}/\text{\AA}^2$. The zero interval is omitted.




MU-12739

Fig. 2 Acetic acid chain. Lengths are in angstrom units.



MU-12740

Fig. 3 Formic acid chain.⁵ Lengths are in angstrom units.



MU-12741

Fig. 4 Schematic drawing of acetic acid structure.