

UC Davis

UC Davis Previously Published Works

Title

Chemical shift assignments of calmodulin bound to a C-terminal site (residues 1120-1147) in the β -subunit of a retinal cyclic nucleotide-gated channel (CNGB1)

Permalink

<https://escholarship.org/uc/item/3x6980ds>

Journal

Biomolecular NMR Assignments, 16(2)

ISSN

1874-2718

Authors

Bej, Aritra
Ames, James B

Publication Date

2022-10-01

DOI

10.1007/s12104-022-10101-7

Peer reviewed



Chemical shift assignments of calmodulin bound to a C-terminal site (residues 1120–1147) in the β -subunit of a retinal cyclic nucleotide-gated channel (CNGB1)

Aritra Bej¹ · James B. Ames¹

Received: 7 June 2022 / Accepted: 3 August 2022 / Published online: 20 August 2022
© The Author(s) 2022

Abstract

Retinal cyclic nucleotide-gated (CNG) channels consist of two protein subunits (CNGA1 and CNGB1). Calmodulin (CaM) binds to two separate sites within the cytosolic region of CNGB1: CaM binding to an N-terminal site (human CNGB1 residues 565–587, called CaM1) decreases the open probability of CNG channels at elevated Ca^{2+} levels in dark-adapted photoreceptors, whereas CaM binding to a separate C-terminal site (CNGB1 residues 1120–1147, called CaM2) may increase channel open probability in light activated photoreceptors. We recently reported NMR chemical shift assignments of Ca^{2+} -saturated CaM bound to the CaM1 site of CNGB1 (BMRB no. 51222). Here, we report complete NMR chemical shift assignments of Ca^{2+} -saturated CaM bound to the C-terminal CaM2 site of CNGB1 (BMRB no. 51447).

Keywords CaM · Calcium · CNGB1 · Retina · Photoreceptor · NMR

Biological context

Retinal CNG channels in rod photoreceptors conduct a cation current in response to light-dependent changes in intracellular levels of cGMP that occur during visual phototransduction (Baylor 1996; Fesenko et al. 1985). CaM binding to retinal CNG channels mediates Ca^{2+} -dependent modulation of channel open probability, which may contribute to light adaptation in retinal rod cells (Bradley et al. 2005; Fain et al. 2001; Hsu and Molday 1993). Retinal CNG channels consist of two protein subunits, CNGA1 and CNGB1 (Bradley et al. 2001). The CNGA1 subunit forms a functional homo-tetrameric channel in the absence of CNGB1, whereas CNGB1 does not form a functional homomeric channel (Finn et al. 1998). In native rod cells, CNG channels form a hetero-tetramer that consists of 3 CNGA1 bound to 1 CNGB1 in a Ca^{2+} -dependent fashion (Shuart et al. 2011). A recent cryoEM structure of the retinal CNG channel (Barret et al. 2021) revealed that a C-terminal site in CNGB1 called CaM2 (residues 1120–1147) is bound to the C-terminal

domain of CaM (residues 80–149, called C-lobe). However, the cryoEM image lacked sufficient resolution to discern atomic-level structural interactions between CaM and CNGB1, and the structure of the CaM N-lobe in the complex was completely missing. CaM was also suggested to bind to a separate N-terminal site in CNGB1 called CaM1 (residues 565–589) (Trudeau and Zagotta 2002) that may regulate CNGB1 binding to CNGA1 (Shuart et al. 2011) and perhaps mediate Ca^{2+} -induced CNG channel inactivation in rod cells (Hsu and Molday 1993; Trudeau and Zagotta 2003). Defects in the Ca^{2+} -dependent regulation of CNG channels are genetically linked to autosomal recessive retinitis pigmentosa and other inherited forms of blindness (Bareil et al. 2001). Elucidating the Ca^{2+} -dependent CNG channel interaction with CaM bound at two separate sites may provide insights for the treatment of retinal diseases. We report here NMR resonance assignments of Ca^{2+} -saturated CaM bound to the CaM2 site of CNGB1 (hereafter called CaM/CaM2). These assignments are an important step toward elucidating the complete structure of CaM bound to CNGB1.

✉ James B. Ames
jbames@ucdavis.edu

¹ Department of Chemistry, University of California, Davis, CA 95616, USA

Methods and experiments

Expression and purification of CaM

Human CaM was overexpressed in *E. coli* strain BL21(DE3) using pET11b (Novagen) and the expressed protein was purified as described previously (Bej and Ames 2022a). The purity of the protein samples was confirmed by sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE). The CaM2 peptide (CNGB1 residues 1122–1143) was purchased from GenScript. The CaM2 peptide was added in threefold excess to Ca²⁺-bound CaM and concentrated to 0.4 mM in a final volume of 0.3 ml.

NMR spectroscopy

All NMR samples of isotopically labeled CaM bound to the unlabeled CaM2 peptide (called CaM/CaM2) were prepared in 20 mM Tris-d₁₁ (pH 7.0) and 1 mM CaCl₂ containing either 8% or 100% (v/v) D₂O and placed in a Shigemi NMR tube (Shigemi Inc.). NMR experiments (at 308 K) were performed on a Bruker Avance III 800 MHz spectrometer equipped with a triple resonance cryogenic (TCI) probe. The ¹⁵N-¹H HSQC spectrum (Fig. 1A, B) contained 200 × 2048 complex points for ¹⁵N(F1) and ¹H(F2). Backbone resonances were assigned by analyzing triple resonance spectra: HNCACB, CBCA(CO)NH, HNCOC, HBHA(CO)NH, and HBHANH. Side chain resonances (aliphatic (Fig. 1C) and aromatic) were assigned by analyzing HCCCONH-TOCSY, HCCH-TOCSY, HBCBCGCDHD and HBCBCGCDCEHE as described previously (Ikura et al. 1991). The NMR data were processed using NMRPipe (Delaglio et al. 1995) and assignment was performed using Sparky (Lee et al. 2015).

Extent of assignments and data deposition

Backbone resonance assignments of CaM/CaM2 are illustrated by labeled peaks in the ¹⁵N-¹H HSQC spectrum of CaM/CaM2 (Fig. 1A, B). Side chain methyl resonance assignments are illustrated by the labeled peaks in the constant-time ¹³C-¹H HSQC spectrum (Fig. 1C). NMR assignments were derived from 3D heteronuclear NMR experiments performed on ¹³C/¹⁵N-labeled CaM bound to unlabeled CaM2 peptide. The high degree of chemical shift dispersion and uniform peak intensities indicate that CaM/CaM2 complex is stably folded. The large downfield chemical shifts of the amide resonances assigned to G26, G62, G99 and G135 indicate that Ca²⁺ is bound to each of the four EF-hands (Fig. 1A). The large upfield chemical shifts of methyl resonances assigned to residues I28, V36, I101 and




Fig. 1 NMR spectroscopy of isotopically labeled CaM bound to unlabeled CaM2 peptide. **A** ¹⁵N-¹H HSQC spectrum of CaM/CaM2 recorded at 800 MHz ¹H frequency. **B** Close-up view of resonance assignments in the spectrally crowded region. **C** Constant-time ¹³C-¹H HSQC spectrum of CaM/CaM2. Complete resonance assignments are available at BMRB 51447

V109 (Fig. 1C) suggest that these residues may be located in the hydrophobic core near aromatic residues. The NMR linewidth of the V109 resonance for CaM/CaM2 is much sharper than it is for CaM/CaM1 (Bej and Ames 2022a), suggesting that the CaM C-lobe binds to CaM2 with higher affinity than it binds to CaM1. At least 92% of the backbone resonances (¹HN, ¹⁵N, ¹³C α , ¹³C β , and ¹³CO) and 85% of side-chain resonances were assigned. Three residues in the

second EF-hand of CaM (A74, R75, and K76) could not be assigned, because their HSQC peaks could not be detected. These same resonances are exchange broadened in CaM bound to the CNGB1 CaM1 peptide (Bej and Ames 2022a) and the α -subunit of the retinal cyclic nucleotide-gated channel (CNGA2) (Contessa et al. 2005), but are not exchange broadened in free CaM (Bej and Ames 2022b; Kainosho et al. 2006). The chemical shift assignments (^1H , ^{15}N , ^{13}C) for CaM/CaM2 have been deposited in the BioMagResBank (<http://www.bmrb.wisc.edu>) under accession number 51447.

The secondary structure of CaM/CaM2 was calculated on the basis of chemical shift index (Wishart et al. 1992) and ANN-Secondary structure prediction using TALOS+ (Shen et al. 2009) (Fig. 2). The secondary structure of CaM/CaM2 is identical to that reported previously for CaM/CaM1 (Bej and Ames 2022a), and is depicted by cylinders and triangles in Fig. 2A. A preliminary analysis of NOESY-derived distances indicate that the eight α -helices and four β -strands combine to form four EF-hands (EF1: residues 7–39, EF2:

residues 45–76, EF3: residues 83–112 and EF4: residues 119–144) as seen in the crystal structure of Ca^{2+} -bound CaM (Babu et al. 1988). The two N-terminal EF-hands (EF1 and EF2) interact with one another to form the CaM N-lobe, and the two C-terminal EF-hands (EF3 and EF4) form the C-lobe. The CaM2 peptide binds to CaM and causes chemical shift perturbations (CSPs) observed for CaM residues in both the N-lobe and C-lobe (Fig. 3), suggesting that the CaM2 peptide is making contact with both lobes of CaM as seen in previous structures of CaM bound to other peptides (Hoefflich and Ikura 2002). The CSP values for the CaM/CaM2 complex in this study are somewhat higher on average than the CSP values observed previously for CaM/CaM1 (Bej and Ames 2022a). The higher CSPs caused by CaM2 binding might be explained by higher affinity CaM binding to CaM2 compared to CaM1. The CSP values for C-lobe residues (I86, A89, F93, H108, M110, L113, M145 and T147) are detectably higher than the CSPs for the corresponding residues in the N-lobe. However, N-lobe residues




Fig. 2 Secondary structure and order parameters of CaM/CaM2. **A** Secondary structure probability and **B** RCI order parameter (RCI-S^2) were predicted using TALOS+ (Shen et al. 2009). Secondary struc-

tural elements are depicted by cylinders (helix) and triangles (strand) derived from the CaM crystal structure (PDB ID—2VAY (Halling et al. 2009))




Fig. 3 Chemical shift perturbation (CSP) versus residue number for CaM/CaM2. CSP was calculated as: $CSP = \sqrt{(\Delta H^N)^2 + (\Delta N)^2}$, where ΔH^N and ΔN are the difference in the $^1H^N$ and ^{15}N chemical

shifts, respectively for CaM/CaM2 versus free CaM in the absence of CaM2 (Bej and Ames 2022b). CSP values are superimposed on the CaM crystal structure (PDB ID: 2VAY (Halling et al. 2009))

F13, A16 and F20 have relatively high CSP values (above 0.5) that suggest these residues may be contacting the CaM2 peptide. On average, the C-lobe has higher CSP values than the N-lobe, which suggests that CaM2 may bind to the CaM C-lobe with higher affinity than that of the N-lobe. This is in stark contrast to CaM binding to the N-terminal CNGB1 peptide (CaM1) in which the CaM N-lobe exhibits higher CSP values (Bej and Ames 2022a). On the basis of our CSP analysis, we suggest that a single CaM may bind to CNGB1 in which the CaM C-lobe preferentially binds to the C-terminal CNGB1 site (CaM2) and the CaM N-lobe preferentially binds to the N-terminal CNGB1 site, CaM1. Future studies are needed to test this possibility by measuring the binding stoichiometry of CaM bound to the full-length CNG channel. The NMR assignments of CaM/CaM2 presented here are an important first step toward determining the full three-dimensional structure of CaM bound to CaM2.

Acknowledgements We thank Derrick Kaseman and Ping Yu for technical support and help with NMR experiments.

Author contributions AB performed all experiments, analyzed data and helped write the manuscript. JBA directed the overall project and wrote the manuscript.

Funding Work supported by NIH Grants to J.B.A (R01 EY012347) and to the UC Davis NMR Facility (RR11973).

Data availability The assignments have been deposited to the BMRB under the accession code: 51447.

Declarations

Conflict of interest The authors declare they have no competing conflict of interest.

Ethical approval The experiments comply with the current laws of the United States.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Babu YS, Bugg CE, Cook WJ (1988) Structure of calmodulin refined at 2.2 Å resolution. *J Mol Biol* 204:191–204
- Bareil C, Hamel CP, Delague V, Arnaud B, Demaille J, Claustres M (2001) Segregation of a mutation in CNGB1 encoding the beta-subunit of the rod cGMP-gated channel in a family with autosomal recessive retinitis pigmentosa. *Hum Genet* 108:328–334
- Barret DCA, Schertler GFX, Kaupp UB, Marino J (2021) Structural basis of the partially open central gate in the human CNGA1/CNGB1 channel explained by additional density for calmodulin in cryo-EM map. *J Struct Biol* 214:107828
- Baylor D (1996) How photons start vision. *Proc Natl Acad Sci USA* 93:560–565

- Bej A, Ames JB (2022a) Chemical shift assignments of calmodulin bound to the beta-subunit of a retinal cyclic nucleotide-gated channel (CNGB1). *Biomol NMR Assign* 16:147–151
- Bej A, Ames JB (2022b) Chemical shift assignments of calmodulin under standard conditions at neutral pH. *Biomol NMR Assign*. <https://doi.org/10.1007/s12104-022-10082-7>
- Bradley J, Frings S, Yau K, Reed R (2001) Nomenclature for ion channel subunits. *Science* 294:2095–2096
- Bradley J, Reisert J, Frings S (2005) Regulation of cyclic nucleotide-gated channels. *Curr Opin Neurobiol* 15:343–349
- Contessa GM, Orsale M, Melino S, Torre V, Paci M, Desideri A, Cicero DO (2005) Structure of calmodulin complexed with an olfactory CNG channel fragment and role of the central linker: residual dipolar couplings to evaluate calmodulin binding modes outside the kinase family. *J Biomol NMR* 31:185–199
- Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeiffer J, Bax A (1995) NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J Biomol NMR* 6:277–293
- Fain GL, Matthews HR, Cornwall MC, Koutalos Y (2001) Adaptation in vertebrate photoreceptors. *Physiol Rev* 81:117–151
- Fesenko EE, Kolesnikov SS, Lyubarsky AL (1985) Induction by cyclic GMP of cationic conductance in plasma membrane of retinal rod outer segment. *Nature* 313:310–313
- Finn JT, Krautwurst D, Schroeder TY, Chen TY, Reed RR, Yau KW (1998) Functional co-assembly among subunits of cyclic-nucleotide-activated, nonselective cation channels, and across species from nematode to human. *Biophys J* 74:1333–1345
- Halling DB, Georgiou DK, Black DJ, Yang G, Fallon JL, Quijcho FA, Pedersen SE, Hamilton SL (2009) Determinants in CaV1 channels that regulate the Ca²⁺ sensitivity of bound calmodulin. *J Biol Chem* 284:20041–20051
- Hoeflich KP, Ikura M (2002) Calmodulin in action: diversity in target recognition and activation mechanisms. *Cell* 108:739–742
- Hsu YT, Molday RS (1993) Modulation of the cGMP-gated channel of rod photoreceptor cells by calmodulin. *Nature* 361:76–79
- Ikura M, Spera S, Barbato G, Kay LE, Krinks M, Bax A (1991) Secondary structure and side-chain ¹H and ¹³C resonance assignments of calmodulin in solution by heteronuclear multidimensional NMR spectroscopy. *Biochemistry* 30:9216–9228
- Kainosho M, Torizawa T, Iwashita Y, Terauchi T, Mei Ono A, Guntert P (2006) Optimal isotope labelling for NMR protein structure determinations. *Nature* 440:52–57
- Lee W, Tonelli M, Markley JL (2015) NMRFAM-SPARKY: enhanced software for biomolecular NMR spectroscopy. *Bioinformatics* 31:1325–1327
- Shen Y, Delaglio F, Cornilescu G, Bax A (2009) TALOS+: a hybrid method for predicting protein backbone torsion angles from NMR chemical shifts. *J Biomol NMR* 44:213–223
- Shuart NG, Haitin Y, Camp SS, Black KD, Zagotta WN (2011) Molecular mechanism for 3:1 subunit stoichiometry of rod cyclic nucleotide-gated ion channels. *Nat Commun* 2:457
- Trudeau MC, Zagotta WN (2002) Mechanism of calcium/calmodulin inhibition of rod cyclic nucleotide-gated channels. *Proc Natl Acad Sci USA* 99:8424–8429
- Trudeau MC, Zagotta WN (2003) Calcium/calmodulin modulation of olfactory and rod cyclic nucleotide-gated ion channels. *J Biol Chem* 278:18705–18708
- Wishart DS, Sykes BD, Richards FM (1992) The chemical shift index: a fast and simple method for the assignment of protein secondary structure through NMR spectroscopy. *Biochemistry* 31:1647–1651

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.