

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

Energy conservation mechanisms for syntrophic growth of desulfovibrio vulgaris and methanococcus maripaludis

### Permalink

<https://escholarship.org/uc/item/3x8390xx>

### Authors

Walker, C.B.

Martens-Habbena, W.



Yang, Z.K.

et al.

### Publication Date

2007-05-14






## ABSTRACT


In the absence of electron acceptors, many *Desulfovibrio* species grow on non-fermentable substrates via syntrophic association with hydrogen consuming methanogens. We examined the physiology of *D. vulgaris* Hildenborough growing syntrophically with *Methanococcus maripaludis* LL using a combination of transcriptional and deletion mutant analyses. Syntrophic cocultures were established in chemostats on minimal media amended with lactate but lacking electron acceptor. Replicated whole genome transcriptional analyses identified 169 and 254 genes that were significantly up- or down-regulated, respectively, relative to sulfate-limited monocultures grown at the same generation time. The majority of up-regulated genes were associated with energy production/conservation, signal transduction mechanisms, and amino acid transport/metabolism. A number of the down-regulated genes were associated with signal transduction mechanisms, inorganic ion transport/metabolism and amino acid transport and metabolism. In order to elucidate possible roles of several highly up-regulated genes associated with electron transfer and energy conservation, we constructed mutants of *Desulfovibrio* deleted in a subset of these genes. Cocultures developed with these mutants displayed a range of growth yields, implicating a putative carbon-monoxide induced hydrogenase (Coo, DVU2286-93) and a high-molecular weight cytochrome (Hmc, DVU0531-6) in energy conservation during syntrophic growth. Mutant monocultures grew to the same density on lactate/sulfate as the wildtype. The *cooL* and *hmc* mutants grew significantly slower and to approximately 25% yields of wildtype cocultures. Together, these data suggest a role of these genes in energy conservation of *D. vulgaris* Hildenborough during syntrophic growth.

## CHEMOSTAT CONFIGURATION



Chemostats are run using a 24 hr retention time at 37 °C and a stirring speed of 250 rpm. The headspace of the chemostat is flushed with a mixture of N<sub>2</sub>:CO<sub>2</sub> (90:10) at a rate of 0.20 - 0.50 ml/min. The headspace gas composition is sampled in 15 min. intervals using a Hiden QIC-20 mass spectrometer.


## BIOMASS & HEADSPACE GAS MEASUREMENTS



Steady-state was assumed when O.D.<sub>600</sub> measurements varied by less than 10% of initial value for 3 retention times. *D. vulgaris*:*M. maripaludis* cell ratio was ~4:1 throughout steady-state as determined by DAPI-stained cell counts.

## TRANSCRIPTIONAL ANALYSIS

Triplicate biological replicates of cocultures and sulfate-limited *D. vulgaris* monocultures were analyzed by the ESPP Functional Genomics Core using custom-designed whole-genome microarrays. Microarray slides were designed with duplicate spots of each open-reading frame (ORF) for both organisms. At least three slides were used for each biological replicate. The ESPP Computational Core calculated RNA/DNA expression ratios for each ORF and the log<sub>2</sub> ratio comparing the coculture versus sulfate-limited monoculture growth conditions was determined. Z-scores for each ORF were calculated to determine statistical significance. Operon-based estimates of local accuracy indicate an absolute Z-score of 1.0 accurately predicts expression changes between the two conditions. Using this value, 169 ORFs were statistically down-regulated. Genes were assigned clusters of orthologous group (COG) functional codes based on previous genome annotations.




Cluster of orthologous groups up- and down-regulated during syntrophic growth. Categories are amino acid transport (E), carbohydrate transport and metabolism (G), cell division and chromosome partitioning (D), cell envelope biogenesis (M), cell motility and secretion (N), chromatin structure and dynamics (B), coenzyme metabolism (H), DNA replication, recombination and repair (I), defense mechanisms (V), energy production and conservation (C), function unknown (S), general function prediction only (R), inorganic ion transport and metabolism (P), intracellular trafficking and secretion (U), lipid metabolism (L), nucleotide transport and metabolism (F), post-translational modification, protein turnover, chaperones (O), secondary metabolites biosynthesis, transport and catabolism (Q), signal transduction mechanisms (T), transcription (K) and translation, ribosomal structures and biogenesis (J)

## TRANSCRIPTIONAL RESPONSE

Gene	TIGR annotation	Log R	ZI score	RNA/DNA expression ratio in coculture	Gene	TIGR annotation	Log R	ZI score	RNA/DNA expression ratio in coculture
DVU2285	lactate permease	1.37	1.92	0.44	DVU2297	glycine/betaine/L-proline ABC transporter	-2.48	3.69	0.45
DVU3025	pyruvate-ferredoxin oxidoreductase ( <i>por</i> )	1.96	2.51	1.82	DVU2298	glycine/betaine/L-proline ABC transporter	-1.99	2.84	0.27
DVU1179	aldehyde-ferredoxin oxidoreductase ( <i>aox</i> )	1.93	1.91	1.64	DVU2299	glycine/betaine/L-proline ABC transporter	-2.22	2.90	0.20
DVU2287	Coo hydrogenase ( <i>cooL</i> )	1.67	2.56	1.88	DVU2300	hypothetical protein	-1.05	1.20	1.73
DVU2289	Coo hydrogenase ( <i>cooL</i> )	1.73	2.77	1.56	DVU2571	ferrous iron transport protein ( <i>feoB</i> )	-4.71	6.36	0.15
DVU2402	heterodisulfide reductase ( <i>hdsA</i> )	1.42	2.03	0.29	DVU2572	ferrous iron transport protein ( <i>feoA</i> )	-4.60	7.29	0.16
DVU2405	alcohol dehydrogenase	2.94	3.51	3.39	DVU2383	tonB dependent receptor domain protein	-4.37	6.14	0.03
DVU1769	periplasmic [Fe] hydrogenase ( <i>hydA</i> )	2.33	1.86	0.37	DVU2384	ABC transporter	-2.44	2.5	0.14
DVU1922	periplasmic [NiFe] hydrogenase ( <i>hynA</i> )	1.00	1.09	0.23	DVU2386	ABC transporter	-1.05	1.46	0.05
DVU0536	high-molecular weight cytochrome ( <i>hmcA</i> )	2.22	2.85	0.54	DVU2390	tonB domain protein	-2.34	3.23	0.09
DVU0531	high-molecular weight cytochrome ( <i>hmcF</i> )	1.93	2.76	0.26	DVU1295	sulfate adenylyltransferase ( <i>sat</i> )	0.75	1.17	4.48
DVU1295	sulfate adenylyltransferase ( <i>sat</i> )	0.75	1.17	4.48	DVU0847	adenylyl-sulfate reductase ( <i>apsrA</i> )	1.21	1.20	4.84
DVU1073	thiosulfate reductase ( <i>ptsA</i> )	2.08	3.41	0.48	DVU0402	dissimilatory sulfite reductase ( <i>dsrA</i> )	0.94	1.56	3.59
DVU0053	sulfate permease, putative	-0.84	1.43	0.07	DVU1636	pyrophosphatase ( <i>ppoC</i> )	1.38	1.82	3.16
DVU1307	cytochrome c <sub>33</sub>	-0.82	1.56	0.15	DVU0694	methylgluteryl oxidoreductase	0.98	1.73	0.56
DVU2680	flavodoxin	-5.19	6.67	0.04	DVU0599	carbon starvation protein A, putative	2.11	2.77	0.82
DVU2652	hypothetical protein	3.07	4.67	2.12	DVU1592	methyl-accepting chemotaxis protein	-1.47	2.51	0.09
DVU2655	D-alanyl-D-alanine carboxypeptidase	0.75	1.17	0.95	DVU1593	chemotaxis protein ( <i>cheY-4</i> )	-1.61	2.62	0.12
					DVU1594	chemotaxis protein ( <i>cheA-3</i> )	-1.63	2.63	0.10
					DVU0806	chemotaxis protein CheY1, putative	-2.59	3.01	0.74


## PHENOTYPIC DIFFERENCES BETWEEN *D. VULGARIS* MUTANTS



Mutant	Description
<i>cooL</i>	transposon-interrupted mutant of [NiFe] hydrogenase small subunit contained within the CO-induced hydrogenase operon
<i>Hmc</i>	high-molecular weight cytochrome deletion mutant ( <i>hmcBCDE</i> )
<i>Hyd</i>	[Fe] hydrogenase deletion mutant ( <i>hydAB</i> )
<i>Hyn</i>	[NiFe] hydrogenase deletion mutant ( <i>hynAB</i> )

## ENERGETIC OVERVIEW

- Lactate → pyruvate + H<sub>2</sub>
- Energetically unfavorable unless pyruvate and/or H<sub>2</sub> kept at low concentrations.
- *Coo* acts to produce hydrogen (likely consumed by the methanogen) and translocate H<sup>+</sup> or Na<sup>+</sup>
- *Hmc* serves to keep pyruvate concentration low, allowing continued lactate oxidation.



- Up-regulated (log R = 2+)
- Up-regulated (log R = 1.5 - 2)
- Up-regulated (log R = 1.0 - 1.5)
- Up-regulated (log R = 0.75 - 1.0)
- Statistically unchanged

## FUTURE WORK

- Develop syntrophic growth pathway on other compounds, such as ethanol.
- Refine metabolite interaction model between *D. vulgaris* and *M. maripaludis* LL using transcriptional, proteome and phenotypic analyses.
- Explore salt, nitrate, pH, etc., stress response of coculture.
- Compare responses with other organisms, especially other *Desulfovibrio* strains.

## ACKNOWLEDGMENT

ESPP is part of the Virtual Institute for Microbial Stress and Survival supported by the U.S. Department of Energy, Office of Biological and Environmental Research, Genomics:GTL Program through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U.S. Department of Energy. The *hmc*, *hyd* and *hyn* mutants were kindly donated by Dr. G. Voordouw, U. of Calgary. *M. maripaludis* LL kindly donated by Dr. J. Leigh, U. of Washington.