

UC Davis

UC Davis Previously Published Works

Title

Enhancing knowledge of chemical exposures and fate in honey bee hives: Insights from colony structure and interactions

Permalink

<https://escholarship.org/uc/item/3x852036>

Journal

The Science of The Total Environment, 916

ISSN

0048-9697

Authors

Encerrado-Manriquez, Angela M

Pouv, Amara K

Fine, Julia D

et al.

Publication Date

2024-03-01

DOI

10.1016/j.scitotenv.2024.170193

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Review

Enhancing knowledge of chemical exposures and fate in honey bee hives: Insights from colony structure and interactions



Angela M. Encerrado-Manriquez^a, Amara K. Pouv^{a,b}, Julia D. Fine^c, Sascha C.T. Nicklisch^{a,*}

^a Department of Environmental Toxicology, University of California-Davis, Davis, CA 95616, USA


^b Department of Fisheries, Animal, and Veterinary Science, University of Rhode Island, Kingston, RI 02881, USA

^c Invasive Species and Pollinator Health Research Unit, USDA-ARS, 3026 Bee Biology Rd., Davis, CA 95616, USA

HIGHLIGHTS

- Honey bees are unintentionally exposed to various chemicals in their environment.
- Social structure and food sharing facilitates chemical transfer between members.
- Limited sensitive analytical techniques restrict studying cumulative impacts.
- New tracing and AI can track chemical transfers in hives to identify exposures.

GRAPHICAL ABSTRACT



ARTICLE INFO

Editor: Jay Gan

Keywords:

Honey bees
Sublethal exposures
Synergistic toxicity
Trophallaxis
Maternal offloading
Chemical diffusion
Chemical mixtures
Artificial intelligence

ABSTRACT

Honey bees are unintentionally exposed to a wide range of chemicals through various routes in their natural environment, yet research on the cumulative effects of multi-chemical and sublethal exposures on important caste members, including the queen bee and brood, is still in its infancy. The hive's social structure and food-sharing (trophallaxis) practices are important aspects to consider when identifying primary and secondary exposure pathways for residential hive members and possible chemical reservoirs within the colony. Secondary exposures may also occur through chemical transfer (maternal offloading) to the brood and by contact through possible chemical diffusion from wax cells to all hive members. The lack of research on peer-to-peer exposures to contaminants and their metabolites may be in part due to the limitations in sensitive analytical techniques for monitoring chemical fate and dispersion. Combined application of automated honey bee monitoring and modern chemical trace analysis techniques could offer rapid progress in quantifying chemical transfer and accumulation within the hive environment and developing effective mitigation strategies for toxic chemical co-exposures. To enhance the understanding of chemical fate and toxicity within the entire colony, it is crucial to consider both the intricate interactions among hive members and the potential synergistic effects arising from combinations of chemical and their metabolites.

* Corresponding author at: University of California, Davis College of Agricultural and Environmental Sciences, Department of Environmental Toxicology, 4117 Meyer Hall, USA.

E-mail address: nicklisch@ucdavis.edu (S.C.T. Nicklisch).

<https://doi.org/10.1016/j.scitotenv.2024.170193>

Received 19 October 2023; Received in revised form 13 January 2024; Accepted 13 January 2024

Available online 24 January 2024

0048-9697/© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

1.1. Economic importance of managed bees

Honey bees (*Apis mellifera*) are one of the best monitored and most widely used commercial crop pollinators worldwide. The practice of managing honey bees has been developing for millennia to obtain commercial products such as honey, wax, and propolis (Kritsky, 2017). It was not until 1851 that the invention of the movable frame hive gave rise to the existence of modern beekeeping practices and the extension of the use of honey bee colonies for commercial agriculture (Oxley and Oldroyd, 2010). Pollination services increase essential crop production by up to 90 % while also providing an annual income source for beekeepers (Porto et al., 2020). According to the 2022 USDA Annual Strategic Pollinator Priorities Report, honey bees pollinate over 100 commercially grown crops in the U.S. and increase their total value by tens of billions of dollars annually (USDA, 2022a). While there are increasing efforts to identify additional wild insect species with the potential to be managed for crop production (Klein et al., 2018; Osterman et al., 2021), honey bees still remain the most economically valuable pollinators for global crop monocultures (Khalifa et al., 2021; Klein et al., 2018). However, the continuous expansion and geographic concentration of pollinator-dependent crop cultivation, such as the almond industry in California, has demanded an increase in the honey bee colony movements across states, ultimately exposing bees to multiple agrochemicals within a short period of time (Bond et al., 2021; USDA, 2022b).

1.2. Multi-chemical and chemical mixture exposure as emerging threats to colony health

Multiple categories of stressors, including biological (parasites, pathogens), dietary (poor nutrition), climate (warmer, dryer), and man-made factors (pesticides, miticides, biocides) (Daisley et al., 2020), can have negative effects on the populations of both commercial and wild bee species, compromising their productivity and survival. The issue of pesticide exposure is particularly complex in commercial beekeeping operations, as they are often migratory (Alger et al., 2018; Simone-Finstrom et al., 2016; Traynor et al., 2016). Colonies are moved regionally and even nationally during massive monoculture pollination events in the U.S. (Bond et al., 2021). This constant annual migration exposes bee colonies to different types of crop pesticides, hive medications, and environmental chemicals within the span of only a few months (Sanchez-Bayo et al., 2016; Traynor et al., 2016). Managed honey bees live in a well-coordinated colony environment with their hives acting as superorganisms with intricate communication and food-sharing mechanisms. This makes chemical exposures a community problem, rather than an individual one, with hive-wide health implication. To begin to understand the dynamic nature and effects of multi-chemical co-exposures on colony health, it is important to first develop sensitive chemical analysis techniques to detect and monitor chemical residue levels and their possible metabolites within the hive environment.

1.3. Technical challenges to study chemical pathways and fate within the colony

Most studies on detection of trace chemicals in the hive environment have focused on quantifying pesticides in products of human use and consumption, such as wax, honey, and pollen (Calatayud-Vernich et al., 2018; Murcia-Morales et al., 2022). However, only a few studies have focused on identifying levels of pesticide residues as they are processed, transferred, and stored within different organismal and physical checkpoints of the honey bee colony (Murcia Morales et al., 2020). A major limitation when analyzing chemical residues in hive members and matrices is the fact that standard analytical methodologies, such as

liquid chromatography (LC-) or gas chromatography (GC-) coupled with mass spectrometry (MS), are often not sensitive enough to detect trace levels of chemicals and their metabolites as they are moved and transformed within the hive environment (Charlton and Jones, 2007; Wu et al., 2021). Furthermore, chemical compounds need to be isolated through optimized extraction techniques, such as Quick Easy Cheap Effective Rugged Safe (QuEChERS) and solid-liquid extraction protocols (Barganska et al., 2018; Barganska et al., 2014; Calatayud-Vernich et al., 2016), which are in turn limited to sample size or amount. When evaluating small samples such as eggs (~0.0001 g) or larvae (~0.002–0.2 g) (Taber and Roberts, 1963; Zoltowska et al., 2011), sample sizes can be prohibitively small. Yet bees are highly sensitive to pesticides even at sublethal concentrations and exposure to low-dose mixtures of these chemicals could promote synergistic effects on honey bee behavior and toxicity when acting the same molecular targets (Nicklisch et al., 2016; Nicklisch and Hamdoun, 2020; Taenzler et al., 2023; Wade et al., 2019). To develop a more holistic understanding of multi-chemical stressors in honey bees, it is important to apply modern tracking and quantification techniques to better elucidate the single and multi-chemical fate pathways, to identify chemical sinks within the honey bee hive environment, and to monitor and quantify how social interactions and caste structure can influence chemical exposures.

In this review, we synthesize current knowledge on sources and types of multi-chemical exposure experienced by commercial honey bees, the observed toxic effects on individual members and the impacts on colony health. We provide updates on the emerging information on possible chemical contaminant sinks and transfer from within-hive structures and food. We further shed light on how caste and developmental stage of the hive members can influence both chemical exposure and elimination pathways. We additionally highlight the importance of two unique but understudied features of the honey bee superorganism that can determine chemical fate within the colony, trophallaxis and maternal off-loading. The overall goal of this review is to emphasize the need to consider social structure and colony interactions to advance our understanding of chemical fate in bee hives and for mitigating the impacts of multi-chemical stressor to honey bees.

2. Discussion

2.1. Routes of honey bee exposures to chemical mixtures

2.1.1. Primary routes of exposure

Direct exposure of honey bees to pesticides and environmental chemicals can occur outside or within the hive (Fig. 1). Forager bees, that collect food and material for hive maintenance and colony food supply, are regularly exposed to multiple chemicals in their environment (Johnson, 2015; Vannette et al., 2015). Thereby, exposure can typically occur via three different routes: i) orally through consumption of contaminated food (pollen, nectar) or water; ii) via direct contact (e.g., cuticle, antenna) with pesticides applied to plants or other surfaces or sequestered in wax comb; and iii) via air exposure when pesticides are applied as aerial sprays and drift onto bees or through active and passive air influx into colonies from nearby treated agricultural areas (Benuszak et al., 2017; Calatayud-Vernich et al., 2018; Johnson, 2015; Krupke et al., 2012; Murcia Morales et al., 2020; Peters et al., 2019; Southwick and Moritz, 1987; Zhu et al., 2015).

Within the hive, physical hotspots and reservoirs of chemical contaminants primarily include stored contaminated food and various hive matrices, such as propolis or wax. The systemic transfer of pesticides to pollen, nectar, beebread, and wax has been widely reported in the literature and poses a significant threat to honey bees (Alder et al., 2006; Benuszak et al., 2017; Calatayud-Vernich et al., 2018; Chauzat and Faucon, 2007; Dively and Kamel, 2012; Gunes et al., 2021; Mullin et al., 2010; Ostiguy et al., 2019; Sanchez-Bayo and Goka, 2014; Stoner and Eitzer, 2012; Stoner and Eitzer, 2013; Traynor et al., 2016; Traynor et al., 2021). This is particularly important for developing larvae, which,

as one of the most vulnerable members in the hive, cannot relocate from contaminated wax cells and rely on the food produced by the nurse bees (Murcia Morales et al., 2020). The levels of pesticides detected in royal jelly are often relatively low, and there appears to be minimal transfer from worker bee diet to the glandular secretions they produce to feed to larvae. For instance, worker bees exposed in the field to a pollen-honey diet fortified with a cocktail of 13 commonly used pesticides produced royal jelly with the majority of the detected concentrations of each active ingredient below 1 µg/kg (Böhme et al., 2017). However, in recent field studies, it was observed that pesticide residues are higher in royal jelly collected from natural comb compared to artificial queen cups, suggesting that there is some diffusion of pesticide residues from wax into the larval diet (Alkassab et al., 2022). Wu and coworkers showed that worker bees reared in pesticide-contaminated wax have shorter lifespans, which may be partly due to this diffusion and direct contact with the pesticide-laden wax (Wu et al., 2011). This highlights the importance of rotating comb to reduce pesticide contamination in wax and the need to assess long-term exposure and sublethal effects of pesticides on caste differentiation and queen bee development, which are still not fully understood (Böhme et al., 2017; Calatayud-Vernich et al., 2018; Payne et al., 2019; Schmuck et al., 2001).

2.1.2. Secondary and tertiary routes of exposure

The complex dynamics within honey bee castes foster numerous biological intersections, creating various points where hive inhabitants engage in the processing and exchange of food. As such, food contaminant transfer and accumulation should be studied in the context of a colony “superorganism” that collectively absorbs, distributes, metabolizes, and eliminates (ADME) these chemicals. Indeed, pesticide residues have been detected in several hive members, such as foragers and larvae (Calatayud-Vernich et al., 2018; Kasiotis et al., 2014; Murcia Morales et al., 2020). These studies revealed that food sources consumed by all hive members (such as bee bread and nectar) were contaminated with a cocktail of pesticides (Table S1).

2.1.2.1. Trophallaxis. The unique process of food exchange through mouth-to-mouth regurgitation is a significant route of secondary exposure to pesticides for in-hive bees, including foragers, nurses, MABs, and drones. The queen bee and brood are likely to experience tertiary exposure since their food is first processed by nurse bees. As primary

consumers of contaminated food, worker bees may dilute ingested pesticides across hive members involved in food handling (depending on the number of organisms in the food-sharing chain) (Böhme et al., 2017; Sponsler and Johnson, 2017). Moreover, the vectorial transport of contaminated food toward the queen bee creates a potential avenue for pesticide bioaccumulation in this vital individual.

While the impacts of peer-to-peer trophallaxis on viral transmission and spread have been well-documented (Chen et al., 2006; de Miranda et al., 2015; Dolezal et al., 2016; Posada-Florez et al., 2021; Ullah et al., 2021), studies on chemical residue levels across all hive members and on possible chemical transfer between bees are patchy and often do not consider continuous biomonitoring of chemical and/or metabolite pathways from the flower (via worker bees) to the reproductive hive members (i.e., queen, drone) and bee products for human consumption (i.e., honey). For instance, to our knowledge, almost no studies have been conducted to determine pesticide, hive medicine, and other chemical residues in queen or drone bees (nor the manner of their transfer). Furthermore, the levels of ubiquitous environmental chemicals, such as the ICES-7 PCB compounds and other persistent, bioaccumulative, and toxic (PBT) chemicals, have so far only been studied in bee products of human consumption, including pollen, propolis and honey (Table S1) (Tang, 2013; Webster et al., 2013; Weisbrod et al., 2007).

Such a holistic approach toward sensitive multi-chemical tracing within a colony would allow for discriminating the extent to which interindividual trophallaxes of contaminated food and contact exposures could provide a biological pathway to mix and distribute toxic chemicals and their metabolites throughout the colony. Likewise, queen bee retinue behavior in worker bees, including grooming and trophallaxis, could facilitate toxic chemical and metabolite bioaccumulation in this longest-living hive member and have detrimental impacts on brood development and survival.

2.1.2.2. Maternal offloading. Another aspect of superorganism ADME is the possibility of maternal offloading via the rapid elimination of accumulated contaminants from the queen bee. This process has been extensively characterized in marine fish and birds and involves the passive transfer of xenobiotic chemicals (including pesticides) through lipids from the fat storages of the reproductive female to nourish and support offspring development (Chynel et al., 2021; Jouanneau et al.,




Fig. 1. Routes of honey bee exposure to mixtures of pesticides and environmental chemicals. VOCs = Volatile organic compounds; PM = particulate matter.

2022; Lyons and Adams, 2015; Lyons et al., 2013a; Lyons et al., 2013b; Mull et al., 2013). Honey bee larvae, consisting mostly of fat tissue, could serve as an ideal deposition site for hydrophobic chemicals originating from the queen bee's body (Bishop, 1961). Maternal environment influences traits in insects, including honey bees, such as the onset of diapause, egg production and size, and pupal and adult weight (Amiri et al., 2020; Fox, 1993; Mousseau and Dingle, 1991; Rossiter, 1991). However, the effects of maternal pesticide exposure on metrics such as disease susceptibility, behavior, and longevity of adult worker offspring are not known. Most evidence of maternal offloading in insects exists in the form of egg and immature insect toxicity data generated by exposing

reproductive adults (Fine, 2020; Fine et al., 2017b; Hodgson et al., 2011; Joseph, 2019; Medina et al., 2010; Trostanetsky and Kostyukovsky, 2008). Information pertaining directly to the amount of a pesticide that deposits in embryos is scarce, likely owing to the difficulty of performing quantitative assessments on small samples such as insect eggs. However, one study, using isotopically labeled pesticides, demonstrated that the insect growth disruptors pyriproxyfen and diflubenzuron are deposited into the eggs of the predatory lacewing bug (*Chrysoperla carnea*) following maternal exposure (Medina et al., 2002). This work generated precise quantitative data relating exposure scenarios to tissue-specific residue concentrations and physiological outcomes. Similar




Fig. 2. Hive member interactions determine chemical movement and fate within the honey bee superorganism. (A) Honey bee hive members and possible pathways of chemical contaminant deposition: Individual bee food deposition or consumption (white arrow), interindividual transfer across castes (purple arrows), passive absorption by the brood (green arrow), and active elimination by the queen (pink arrow). (B) List of possible first-hand oral (O), contact (C), and unknown (U) exposure routes for each hive member and bee product or food source. MAB = middle-aged bees.

information about chemical residue levels in honey bees could be valuable, where the outcome and fate of an entire colony is dependent on a single reproductive queen.

2.2. Hive structure and interactions determine chemical exposure risk

The colony structure of honey bees is characterized by a close-knit eusocial community where individuals share food, labor, and information (Fig. 2 and Table S2). This complex and coordinated behavior evolved through a caste system, typically formed by a single queen bee, multiple worker bees (i.e., nurse, forager, and middle-aged bees (MAB)), drones, and brood (including egg, larvae, and pupa) (Johnson, 2010). Acknowledging the unique social structures and food-sharing practices within the bee hive is important to comprehensively track chemical exposure pathways and identify biological and physical reservoirs of chemical contaminants. This requires considering factors such as caste-specific routes of exposure, differences in metabolic enzymes among bees, physical contacts and feeding behavior within the hive, and the storage sites of contaminated food and bee products (Barascou et al., 2022; Medrzycki et al., 2015). These factors can significantly influence the movement, degradation, and potential accumulation of pesticides and other chemicals within the hive, ultimately impacting the overall health and well-being of the colony.

2.2.1. Honey bee development and hive member interactions

Forager honey bees are the oldest of all worker bees (Fig. 2 and Table S2). Depending on the time of the year, foragers can live from 20 days (summer) to up to three months (winter) (Johnson, 2010). Forager honey bees venture outside of the hive and collect goods such as nectar, pollen, propolis, and water, which will serve the hive mainly as food sources. When forager bees return to the hive, they pass the collected goods to middle-aged bees (MABs), another type of worker bee, whose age ranges from 12 to 21 days old. At the entrance of the hive, MABs take nectar and pollen for storage, water for hive temperature control, and propolis for fixing gaps within the hive combs. MABs can also be in charge of handling comb building and hive defense. Deeper inside the hive at the brood comb region, nurse bees perform queen retinue behaviors and brood care. The life span of nurse bees ranges between 4 and 12 days, during which time they utilize gland secretions and hive-stored goods such as pollen, nectar, honey, and water to produce brood and queen bee food (Berenbaum and Liao, 2019) and share processed food via trophallaxis, the hive's main food-sharing mechanism. It is thought that queen bees are fed with royal jelly (though the full composition and extent of the queen bee's diet has not been well studied), while the worker brood are fed a mixture of primarily gland secretions, bee bread, and honey (though again, the specifics of these mixtures have not been established) (Berenbaum and Liao, 2019; Free, 1959; Haydak, 1943; von Planta, 1888; Wueppenhorst et al., 2022). The brood consist of eggs laid by the queen (~1500 per day), larvae (at different stages), and pupae (Remolina and Hughes, 2008). Brood are the youngest hive members and are taken care of by nurse bees until they become adult bees and replace them in their role. Finally, drones, or male bees are produced only to mate with virgin queens from other colonies to spread the colony's genes. During the summer, drones feed on stored goods and attempt to mate until they die. By winter any remaining drone will be removed from the colony to reduce the burden to the hive resources. Depending on the developmental stage and caste differentiation, each hive member can be exposed to multiple natural and synthetic chemicals in their foraging environment or within the hive.

2.3. Effects of multi-chemical exposure on colony health and survival

2.3.1. Synergistic toxicity of chemical mixtures

Managed and wild bees are continuously exposed to a wide-ranging mixture of natural compounds as well as man-made in-hive medicine, pesticides, and environmental chemicals. Once entering the organism,

these complex mixtures of chemicals can interact with cells and biomolecules to directly exert harmful effects. These chemicals further interact with one another, creating a complex web of interactions that indirectly contribute to their overall impact on bee health. If the chemicals act on the same molecular target, such as receptors, transporters or enzymes, the interactions can be antagonistic (combined effect is less than the sum of its individual effects), additive (combined effect is the sum of its individual effects), or synergistic (combined effect is greater than the sum of its individual effect) (Nicklisch and Hamdoun, 2020). In a recent literature review on the effects of binary mixtures of pesticides on honey bee, wild bee, and solitary bee toxicity, it was noted that additive and synergistic toxicity were the most common interactions, representing 17 % and 72 % of reported data, respectively (Carnesecchi et al., 2019). One historical example of the synergistic toxicity observed in laboratory and field studies is the interaction between ergosterol biosynthesis inhibiting (EBI) fungicides and neonicotinoids or pyrethroids caused by the fungicide's inhibition of the cytochrome P450 monooxygenase system (Meled et al., 1998; Pilling et al., 1995; Pilling and Jepson, 1993; Tosi and Nieh, 2019). While enhanced toxicity of multi-pesticide combinations toward bees has been thoroughly described and tested in field and laboratory assays (Berenbaum and Johnson, 2015; Carnesecchi et al., 2019; Pal et al., 2022; Siviter et al., 2021; Wade et al., 2019; Wang et al., 2021b; Zhu et al., 2017a; Zhu et al., 2017b), intended and unintended co-exposures to mixtures of pesticide-adjuvant, pesticide-in-hive medicine, and pesticide-environmental chemicals have been less appreciated.

2.3.1.1. Intended chemical mixtures

2.3.1.1.1. *Tank mixing of pesticides and adjuvants.* Pesticides are often applied in mixtures (known as tank mixing) to control multiple pests and diseases (e.g., an aphid infestation alongside a fungal infection) while reducing the number of trips over a field (Holloway and Western, 2003). Although intentional tank mixing of agrochemicals can be controlled and monitored by applicators, the possible synergistic toxicities of these chemical mixtures on non-target pollinators are not well understood. Toxicity to bees can also increase when pesticides are combined with adjuvants, a common agricultural practice of adding additional compounds like surfactants and oils to boost pesticide absorption and distribution with an eye toward improving their efficacy (Green, 2000; Hazen, 2000; Mullin et al., 2016). In a comparative study of over 1300 wax, pollen, and bee samples from hives across the US and Canada, Mullin and coworkers revealed a notable finding: only 70 % of the targeted pesticide active ingredients were detected in the beehive samples, whereas 100 % of the other sought-after formulation ingredients were present (Mullin et al., 2015). The ubiquitous and concurrent detection of pesticide contaminants and formulation ingredients in the hive environment could facilitate toxic pesticide bioaccumulation and toxicity in bees. Indeed, a 2015 field and laboratory study showed that the broad-spectrum insecticide Fipronil with the added spray surfactant and adjuvant, Sylgard, increased honey bee mortality up to 100 % within 24 h of application (Mayer and Lunden, 2015). Similar adjuvants typically considered biologically inert have also been shown in laboratory studies to promote viral pathogenicity in chronically exposed honey bee larvae (Fine et al., 2017a), enhance viral replication in adults, and impair queen oviposition rates (Fine et al., 2023). More recently, laboratory tests on 30 tank mixtures of different pesticide-adjuvant combinations showed that 50 % of the combinations increased honey bee mortality (Wernecke et al., 2021). In another study, Walker et al. demonstrated that the adjuvant *Dyne-Acmin*, when combined with the fungicide tilt and the insecticide altacor, can result in elevated toxicity levels, attributed to increases in overall toxicity of the pesticide mixture (Walker et al., 2022).

Adjuvants can be added to the pesticide formulation as "inert" ingredients, in which case they are classified as formulation adjuvants, or they can be added separately on top of other formulated products in a

tank mix, in which case they are classified as spray adjuvants. Each adjuvant can add a specific characteristic to the formulation or tank mix, therefore multiple adjuvants can be used in the same application to achieve several different functions (reduce drift, increase penetration, increase solubility, increase attachment, increase stability) (Hock, 2022; Wu et al., 2023). Adjuvants are thought to lack inherent pesticidal properties, and federal law exempts spray adjuvants from EPA pesticide registration, leaving them largely unregulated. However, this rule varies by state, and in California the CDPR does require spray adjuvants to be registered as pesticides prior to sale and use (Cox and Zeiss, 2022). Unlike in the EU, where toxicity testing of active ingredients and adjuvants is required to have a ‘representative formulation’, in the US toxicity testing of pesticide products does not involve inert ingredients such as adjuvants and mainly focuses on active ingredients (Straw et al., 2022). In addition, given the lack of scrutiny and toxicological testing of these “inert” formulation ingredients for lethal and sublethal effects on the reproductive Queen and vulnerable hive members, such as the bee brood, more research is needed to better predict the effects of various pesticide-adjuvant interactions on colony health and survival.

2.3.1.1.2. In-hive medicine. There is a growing body of evidence that in-hive medications applied to treat bee infections and hive infestations can be detrimental to bee health when bees are pre-exposed to other agrochemicals (Mullin et al., 2010; Rangel and Tarry, 2016; Traver et al., 2018; Traynor et al., 2016). It has been suggested that such unintended co-exposures of agricultural pesticides affect bee colony health directly (by causing acute additive or synergistic toxicity in bees) or indirectly (by sensitizing bees to other xenobiotics). For instance, in a laboratory assay, bees pre-treated with oxytetracycline (OTC), an antibiotic treatment for foulbrood diseases, and sequentially exposed to the acaricide coumaphos, a therapy for mite infestation, experienced a 44 % increase in mean mortality (Hawthorne and Dively, 2011). Likewise, bees fed with OTC and the acaricide tau-fluvalinate exhibited a 33 % increase in mortality. In another laboratory study, researchers found that orally administering fumagillin (an antibiotic used to treat *Nosema* within hives) together with tau-fluvalinate (an acaricide used to treat varroa mite infestations) can decrease the median LD₅₀ by 2-fold compared to sucrose syrup control treatment in exposed honey bees (Johnson et al., 2013).

As such, a precautionary approach accounting for potential synergies between in-hive medications and agrochemicals is prudent to ensure thriving honey bee colonies. Ongoing monitoring of pesticides exposures and a carefully timed application of in-hive medications to avoid simultaneous pesticide exposure may help mitigate interactive toxicity.

2.3.1.2. Unintended chemicals mixtures

2.3.1.2.1. Foodborne toxicities. Arguably the primary route of honey bee exposure to toxic xenobiotics is through ingestion of contaminated pollen and nectar, typically a result of the widespread application, systemic distribution, and environmental persistence of modern insecticides such as neonicotinoids applied to growing crops (Bonmatin et al., 2015; Stoner and Eitzer, 2012; Tosi et al., 2018). Exposures to multiple pesticides at lethal and sublethal concentrations may cause synergistic toxicity across hive members and contribute to colony loss (Biddinger et al., 2013; Farooqui, 2013; Iwasa et al., 2004; Johnson et al., 2010; Levine and Borgert, 2018; Lu et al., 2014; Magal et al., 2019; Magal et al., 2020; Pilling et al., 1995; Pilling and Jepson, 1993; Spurgeon et al., 2016; Thompson et al., 2014; Williamson et al., 2013; Williamson and Wright, 2013). For instance, Zhu and coworkers showed in two separate laboratory studies that when honey bees were exposed to eight different insecticides (imidacloprid, acephate, λ -cyhalothrin, oxamyl, tetraconazole, glyphosate, sulfoxaflor, and clothianidin) in binary combinations, the mixtures showed synergistic toxicity that increased mortality between 15 and 26 % (Zhu et al., 2017a; Zhu et al., 2017b). Notably, when bees were exposed to a mixture of all eight pesticides, the mortality increased to 100 % (Zhu et al., 2017b). Johnson

and coworkers performed a similar laboratory study using pairwise combinations of six different acaricides, eight fungicides, three antimicrobials, and three detoxification enzyme inhibitors (Johnson et al., 2013). The group reported synergistic toxicity as median lethal dose (LD₅₀) and observed the highest toxicities for the acaricide tau-fluvalinate in combination with the fungicide prochloraz (LD₅₀ = 0.01 mg/bee), the model CYP450 inhibitor piperonyl butoxide or PBO (LD₅₀ = 0.01 mg/bee), and the acaricide coumaphos (LD₅₀ = 0.78 mg/bee). Tau-fluvalinate also increased toxicity in honey bees pre-treated with different EBI fungicides, including myclobutanil (LD₅₀ = 0.121 mg/bee) and metconazole (LD₅₀ = 0.452 mg/bee).

2.3.1.2.2. Waterborne contaminants. Unintended co-exposures to water-soluble environmental chemicals, such as *Per*- and polyfluoroalkyl substances (PFAS), is another understudied research area and possible threat to honey bee health. PFAS are a group of persistent organic pollutants (POP) that are ubiquitously found in the environment and are recognized by the EPA as contaminant of emerging concern (CEC) in surface waters (Gluge et al., 2020; Miner et al., 2021; Podder et al., 2021). PFAS are regularly detected in aquatic environments, including wastewater, agricultural runoff waters, rainwater, groundwater, and receiving waters such as rivers, ponds, and lakes (Dauchy et al., 2019; Lin et al., 2014). These forever chemicals have also been shown to move from irrigated soil through roots into plants, including flowering crop plants known to be pollinated by commercial and solitary bees (Navarro et al., 2017; Xu et al., 2022). Honey bees need water as a dietary salt source, as a thinner to prevent honey crystallization, and for thermal homeostasis of the hive during the warm summer months (Butler, 1940; Lau and Nieh, 2016; Nicolson, 2009; Stabentheiner et al., 2021). As such, water consumption could be a significant route of PFAS exposure to honey bees. Indeed, PFAS compounds have been shown to bioaccumulate in different aquatic and terrestrial insects and can be transferred to their predators (Koch et al., 2021; Lan et al., 2020). Recently, Sontter and coworkers showed in a laboratory assay that the major water contaminant perfluorooctanesulfonic acid (PFOS) can accumulate in honey bees and increase mortality, reduce colony defense and maintenance behaviors, and even halt brood development (Sontter et al., 2021). Furthermore, Surma and coworkers showed the presence of perfluoroalkyl substance in honey from various European countries, indicating that bees who undergo the intricate process of producing and storing this natural product within the hive, come into contact with PFAS (da Silva et al., 2016; Surma et al., 2016).

2.3.1.2.3. Airborne toxicities. Bees can also be unintentionally exposed to airborne pesticides, heavy metals, micro- and nano-plastics (also present in water) (Alma et al., 2023; Buteler et al., 2022; Edo et al., 2021), and air pollutant mixtures due to wind drift or when bees are foraging on neighboring crop fields with varying foliar spray applications (Botias et al., 2015; Davis and Williams, 1990; Krupke et al., 2012; Macri et al., 2021; Marzaro et al., 2011; Moffett et al., 1972; Thimmegowda et al., 2020; Ucar and Hall, 2001; Zaric et al., 2018; Zhu et al., 2015). It has been suggested that both airborne particles such as respirable suspended particulate matter (RSPM) and volatile agrochemicals interfere with pest insect and pollinator sensory physiology, chemical and pheromone signaling, and behavior (Gate et al., 1995; Jacobson, 1966; McFrederick et al., 2009). Smoke has been traditionally used by beekeepers to reversibly interfere with the bee's sense of smell and subdue the alarm response (Gage et al., 2018; Visscher et al., 1995). But chemical air pollutant mixtures, such as agriculture and horticulture pesticides, diesel exhaust, and other volatile organic compounds (VOCs) have been shown in field studies to sensitize pollinators to additional stressors and affect crucial pollinator foraging behavior and reproduction (Reitmayer et al., 2019; Samuelson et al., 2018).

2.3.1.2.4. Micro- and nanoplastics. Bees can also incorporate ubiquitous micro- and nanoplastics (MNPs) from a variety of media in the environment and deliver it to multiple hive matrices and members (Alma et al., 2023; Edo et al., 2021). MNP toxicity depends not only the particle size, but also on the type of polymer, and the presence of other

chemicals (Al Naggar et al., 2021). For instance, in recent laboratory assays, micro particles of polystyrene have been shown to accumulate in honey bee tissues, and to promote adverse health effects (i.e., decrease gut microbiome diversity, changes in oxidative damage, impaired detoxification and immunity, and reduced lipid metabolism gene expression) when in combination with other pollutants or existing diseases (Alma et al., 2023; Buteler et al., 2022; Deng et al., 2021; Wang et al., 2021a). Further research is needed to better characterize the effects of MNP exposure on a colony level and to elucidate possible synergistic effects with MNP-adsorbed or co-exposed chemicals in the hive environment.

2.4. Sublethal pesticide exposures

The effects of repeated exposures of commercial and wild bee species to low levels of pesticides have traditionally been underestimated. Sublethal exposure has been shown to interfere with the colony's survival, performance, winterization, behavior, development, and physiology (Barron, 2015; Bryden et al., 2013; Dively et al., 2015; El Hassani et al., 2008; Fine, 2020; Hatjina et al., 2013; Ingram et al., 2015; Lu et al., 2014; Negi et al., 2022; Williamson et al., 2014; Wu et al., 2011; Wu et al., 2022; Xavier et al., 2015). Specifically, delayed larval development and adult emergence, downregulation of detoxification enzymes, and shortened worker bee longevity have been identified as added stressors that impact a colony's fitness and survival (Fisher et al., 2021; Glass et al., 2021; Smodis Skerl et al., 2009; Tome et al., 2020; Wu et al., 2011).

2.4.1. Learning and memory impairment

Low-dose exposures have also been shown to cause direct neurological effects in bees, including decreased memory and learning performance. For instance, honey bee exposure to sublethal concentrations of the neonicotinoids thiamethoxam and thiacloprid in field studies caused an increase in mortality and affected the ability of forager bees to navigate home (Henry et al., 2012; Tison et al., 2016). Thiamethoxam was also shown in two other field studies to alter honey bee and bumble bee feeding rates, motor skills, and visual learning (Laycock et al., 2014; Ludicke and Nieh, 2020). As a result, sublethal studies triggered a re-evaluation of thiamethoxam which led to its eventual banning in Europe for application to flowering plants (Stokstad, 2018). Acute laboratory exposure of bees to sub-lethal doses of the organophosphate in-hive acaricide coumaphos impaired short-term (10 min) and long-term (24 h) memory in response to a conditioned olfactory stimulus (Williamson et al., 2013), while environmentally-relevant subchronic treatment of bees over four days with either coumaphos or the neonicotinoid imidacloprid resulted in degraded olfactory learning and memory (Williamson et al., 2014). Using the proboscis extension response (PER) assay, Decourtye and coworkers found that worker bees exposed in the laboratory to sublethal concentrations of fipronil, deltamethrin, endosulfan, and prochloraz showed reduced learning performance (Decourtye et al., 2005). Since associative learning of odor, color, and shape of flowers with essential food sources such as pollen and nectar is an important behavior of foraging bees (Hammer and Menzel, 1995), the disruption of this vital sense through chronic exposure to sublethal concentrations of pesticides could both threaten colony survival and disturb the pollination services in the ecosystem.

2.4.2. Disruption of chemical communication

Laboratory assays examining low-level pesticide exposure have also demonstrated that chronic exposure can interfere with the production and quality of queen mandibular pheromone (QMP), thereby impacting communication between the hive and queen bee, affecting labor, queen-rearing suppression, and colony cohesion (Bortolotti and Costa, 2014; Milone and Tarpy, 2021; Rangel and Tarpy, 2016; Walsh et al., 2020). Laboratory pesticide exposures also affect chemically guided worker bee retinue behaviors toward the queen, which usually promotes

attenuating, grooming, and trophallaxis (Litsey et al., 2021). Indirect effects of low-level pesticide exposures in the laboratory have also been shown to cause glandular degeneration and impact the nutritional quality and quantity of royal jelly produced by nurse bees (Milone et al., 2021), which can have dire consequences for adequate provisioning of queen bees and healthy colony maintenance.

2.4.3. Disturbance of gut microbiome

Exposure to low levels and mixtures of agricultural pesticides have also been shown to affect gut bacterial colonization of larval, newly emerged, and adult honey bees (Almasri et al., 2022; Motta and Moran, 2020; Motta et al., 2018; Vazquez et al., 2023). Furthermore, exposure to commonly used in-hive medicine has been shown to affect honey bee microbiome structure and function (Kakumanu et al., 2016). Altering the bee gut microbiota, including core species like *Snodgrassella alvi*, *Gilliamella apicola*, *Lactobacillus Firm-4*, *Lactobacillus Firm-5*, and *Bifidobacterium asteroides*, can have detrimental effects on bee development, metabolism, immunity, and behavior (Daisley et al., 2020; Kwong and Moran, 2016; Motta and Moran, 2023). However, to date only a few pesticides have been tested for their effects on honey bee and wild bee microbiomes at either sub-lethal or environmentally relevant concentrations, and more research is needed to better understand the impacts on the host's immune system and overall health (Hotchkiss et al., 2022).

2.4.4. Cumulative multi-stressor effects

Finally, sublethal pesticide exposure to larval and adult bees already battling pathogens may also have synergistic detrimental effects. For instance, Doublet and coworkers found in laboratory assays that honey bee larvae exposed to sublethal doses of the neonicotinoid thiacloprid in conjunction with Black Queen Cell Virus (BQCV) have a lower survival rate (65 %) than the control (90 %) or exposure to just thiacloprid (80 %) or the virus (85 %) (Doublet et al., 2015). In a similar way, Pettis and coworkers showed in field studies with three brood generations that sublethal exposure to the pesticide imidacloprid sensitized honey bee colonies toward increased infections with the *Nosema* gut pathogen (Pettis et al., 2012).

Chronic exposure to multiple agricultural chemicals, even at low sublethal doses, can negatively impact honey bee health. At the colony level, these sublethal effects can reduce productivity and increase susceptibility to disease and parasites, ultimately leading to higher mortality. While testing myriad potential mixtures of real-world residues in a laboratory setting poses challenges related to scope and analytical detection limits, the integration of innovative computer-aided approaches for individual bee monitoring, coupled with sensitive trace chemical analysis techniques will be essential for developing a holistic understanding of sublethal and toxic effects resulting from chemical mixtures within bee hives.

2.5. Toward improved understanding of chemical fate and toxicity in bee colonies

Pesticides in the U.S. are regulated by the Environmental Protection Agency (EPA) to ensure the safety of humans, pollinators, and other non-target organisms (EPA, 2014; EPA, 2016a, 2016b). A plethora of pesticides have been approved for U.S. agricultural crops, often applied in mixtures to combat several pests at once. Globally, according to the FAOSTAT statistics (<https://www.fao.org/faostat>) of the Food and Agriculture Organization (FAO) of the United Nations, between 1990 and 2021 herbicides were the largest applied pesticide class (by tons), followed by fungicides in second place, and insecticides in third. In contrast to Europe, where in 2021 fungicides were the most applied pesticide class, in the US herbicides continue to dominate the market for agricultural use. The pesticide classes most associated with honey bee and other pollinator toxicity are neonicotinoids, pyrethroids, carbamates, organophosphates, and the phenylpyrazole Fipronil (Cressey, 2017; Johnson et al., 2010; Pisa et al., 2015; Sanchez-Bayo, 2014;

Sanchez-Bayo and Goka, 2014).

Historically, most evaluations of pesticide toxicity toward bees are based only on tests evaluating the effects of direct contact and oral exposures of adult bees or larvae to a single compound (Maus et al., 2022). Furthermore, assessments of pesticide hazards toward honey bee colonies by calculating a Hazard Quotient (HQ) or Risk Quotient (RQ) from pesticide application rate divided by the acute adult bee LD₅₀ values often produce poor correlation results between field application rates and actual pesticide residues reaching the individual colony members (Carlson et al., 2022; Drummond et al., 2018; Mu et al., 2022; Thompson, 2021). These methods were initially developed for foliar spray applications and assume that the combined total bee exposure from all routes (oral, contact, air) is directly related to the pesticide application rate. Due to a lack of residue-level validation studies, predicting effects of agrochemical exposure using these methods is hampered by the fact that dietary and contact exposure pathways during bee developmental stages are different (i.e., capped brood versus roaming adult bees).

The eusocial system within a honey bee hive forms another under-explored dimension of developmental and caste-specific exposure risks and interindividual chemical transfer and mixing that needs to be considered when evaluating effects of agrochemicals. Furthermore, within the hive, wax functions as the structural foundation for both present and future hive occupants, serving not only as their habitat but also as a repository for stored food. Its hydrophobic nature makes it an ideal sink for lipophilic environmental pollutants. However, there is a limited body of research exploring wax as a possible route of developmental exposure and the toxicity of contaminated wax across the diverse members of the hive. A major limitation to performing these studies is the fact that chemical concentrations in small samples such as eggs and larvae are typically below the detection limit of standard analytical methods. In addition, it is technically still challenging to determine the chemical-specific diffusion or transfer rates from the wax into larvae and other hive members to develop estimates of exposure (Alkassab et al., 2022; Kast and Kilchenmann, 2022; Murcia Morales et al., 2020; Wilmart et al., 2021).

Modern isotope tracing analysis techniques, such as accelerated mass spectrometry (AMS), that use stable radiolabeled chemicals to detect trace levels of multiple chemical compounds and their metabolites, can lower the detection capabilities to quintillion levels (10⁻¹⁸ mol) and sample size to milligrams (Enright et al., 2016; Kutschera, 2016; Nnane and Tao, 2005). This technique has shown its capability to do accurate quantification of parent pesticide and metabolite distribution in Argentinian ant colonies (Hooper-Bui et al., 2015). However, to our knowledge, to date no AMS studies have been performed on honey bees. An additional challenge to track chemical contaminant flow between hive members is the need to clearly identify and monitor individual bees and their interactions.

Artificial intelligence (AI)-based automated monitoring of insects using computer vision systems (Bjerger et al., 2021; Gernat et al., 2023; van Klink et al., 2022) is a promising, emerging technology in the field of animal behavior studies and could provide a new avenue to study parent compound and metabolite transfers within and across honey bee hive members. Such AI-guided systems could facilitate monitoring the interactivity of individual bees, including trophallaxis and behavior, and – in combination with the sensitive AMS technology – could aid in the prediction of contaminated food flow, deposition, and the effects on colony survival.

3. Conclusions

Honey bees are exposed to complex chemical mixtures in their natural habitat that can significantly threaten colony health and survival. Although this article focuses on the honey bee hive superorganism and its interactions with chemicals, it is worth remembering that chemical exposure is not exclusive of commercial pollinators and more research needs to be done in unmanaged pollinator species, such as wild bumble

bees and solitary bees, where the species survival can be entirely dependent on one individual. Levels and composition of these chemical mixtures may vary during the year and over time due to crop-specific pesticide application schedules and chemical degradation rates. Identifying levels and the metabolic fate of trace amounts of chemicals as they are processed, transferred, and stored within different organismal and physical checkpoints of the hive will be necessary to identify and apply appropriate mitigation strategies such as comb replacement (for physical sinks) or requeening (for biological sinks).

Based on the available literature, sublethal impacts on factors such as learning, immunity, and colony productivity can have substantial consequences for bee health at the colony level. Additionally, synergistic interactions between the multiple active and inert ingredients in commercial pesticide formulations and tank mixes may alter toxicity but are not fully captured in single active ingredient assays. To enhance protection of bees from pesticide-related harm, more research investigating sublethal endpoints across exposure scenarios at the individual and colony levels is needed. Continuing development of field-realistic chronic toxicity protocols, evaluation of pesticide mixtures, and increased use of semi-field and field studies can help ensure the intricate effects of toxicants are fully considered.

A fundamental understanding of the combined effects of the chemical mixtures on bees is crucial for addressing the complex and evolving challenges posed by multiple chemical exposures in their natural environment. This will require adopting modern tracing analysis techniques and artificial intelligence-based monitoring to better understand possible cumulative and synergistic effects of chemical mixtures, especially within the hive's social structure at field realistic concentrations. This comprehensive approach aims to achieve a fundamental understanding of chemical fate in hives for safeguarding honey bee health and colony survival in the face of diverse and changing chemical threats.

CRedit authorship contribution statement

Angela M. Encerrado-Manriquez: Writing – review & editing, Writing – original draft, Visualization, Methodology, Data curation, Conceptualization. **Amara K. Pouv:** Writing – review & editing, Writing – original draft, Conceptualization. **Julia D. Fine:** Writing – review & editing, Writing – original draft, Validation, Resources, Investigation, Funding acquisition, Conceptualization. **Sascha C.T. Nicklisch:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Resources, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization.

Declaration of competing interest

The authors declared no competing interests in this work.

Data availability

Data will be made available on request.

Acknowledgements

We thank Charles Mitter for carefully reading the manuscript and providing insightful discussions. The research reported in this publication was supported by a grant from the 2022-2025 PAM-Costco USA Scholarship. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.

Funding

This work was supported by the NIFA-USDA (CA-D-ETX-2526-H) to S.C.T.N and by USDA NACA (58-2030-3-034) to J.D.F. and S.C.T.N.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2024.170193>.

References

- Al Naggar, Y., Brinkmann, M., Sayes, C.M., Al-Kahtani, S.N., Dar, S.A., El-Seedi, H.R., et al., 2021. Are honey bees at risk from microplastics? *Toxics* 9.
- Albero, B., Miguel, E., García-Valcárcel, A.I., 2022. Acaricide residues in beeswax. In: *Implications in Honey, Brood and Honeybee and Influence of Landscape Around Colonies*.
- Alder, L., Greulich, K., Kempe, G., Vieth, B., 2006. Residue analysis of 500 high priority pesticides: better by GC-MS or LC-MS/MS? *Mass Spectrom. Rev.* 25, 838–865.
- Alger, S.A., Burnham, P.A., Lamas, Z.S., Brody, A.K., Richardson, L.L., 2018. Home sick: impacts of migratory beekeeping on honey bee (*Apis mellifera*) pests, pathogens, and colony size. *PeerJ* 6, e5812.
- Alkassab, A.T., Bischoff, G., Thorbahn, D., Frommberger, M., Pistorius, J., 2022. Transfer of xenobiotics from contaminated beeswax into different bee matrices under field conditions and the related exposure probability. *Chemosphere* 307, 135615.
- Alma, A.M., de Groot, G.S., Buteler, M., 2023. Microplastics incorporated by honeybees from food are transferred to honey, wax and larvae. *Environ. Pollut.* 320, 121078.
- Almasri, H., Liberti, J., Brunet, J.L., Engel, P., Belzunces, L.P., 2022. Mild chronic exposure to pesticides alters physio-logical markers of honey bee health without perturbing the core gut microbiota. *Sci. Rep.* 12, 4281.
- Amiri, E., Le, K., Melendez, C.V., Strand, M.K., Tarpay, D.R., Rueppell, O., 2020. Egg-size plasticity in *Apis mellifera*: honey bee queens alter egg size in response to both genetic and environmental factors. *J. Evol. Biol.* 33, 534–543.
- Arceo-Gómez, G., Jameel, M.I., Ashman, T.L., 2018. Effects of heterospecific pollen from a wind-pollinated and pesticide-treated plant on reproductive success of an insect-pollinated species. *Am. J. Bot.* 105, 836–841.
- Barascou, L., Sene, D., Le Conte, Y., Alaux, C., 2022. Pesticide risk assessment: honeybee workers are not all equal regarding the risk posed by exposure to pesticides. *Environ. Sci. Pollut. Res. Int.* 29, 90328–90337.
- Barganska, Z., Slebioda, M., Namiesnik, J., 2014. Determination of pesticide residues in honeybees using modified QUCHEERS sample work-up and liquid chromatography-tandem mass spectrometry. *Molecules* 19, 2911–2924.
- Barganska, Z., Konieczka, P., Namiesnik, J., 2018. Comparison of two methods for the determination of selected pesticides in honey and honeybee samples. *Molecules* 23.
- Barker, R.J., Lehner, Y., Kunzman, M.R., 1980. Pesticides and honey bees: nectar and pollen contamination in alfalfa treated with dimethoate. *Arch. Environ. Contam. Toxicol.* 9, 125–133.
- Barron, A.B., 2015. Death of the bee hive: understanding the failure of an insect society. *Curr Opin Insect Sci* 10, 45–50.
- Benuszak, J., Laurent, M., Chauzat, M.P., 2017. The exposure of honey bees (*Apis mellifera*; Hymenoptera: Apidae) to pesticides: room for improvement in research. *Sci. Total Environ.* 587–588, 423–438.
- Berenbaum, M.R., Johnson, R.M., 2015. Xenobiotic detoxification pathways in honey bees. *Curr Opin Insect Sci* 10, 51–58.
- Berenbaum, M.R., Liao, L.H., 2019. Honey bees and environmental stress: toxicologic pathology of a super-organism. *Toxicol. Pathol.* 47, 1076–1081.
- Bergero, M., Bosco, L., Giacomelli, A., Angelozzi, G., Perugini, M., Merola, C., 2021. Agrochemical contamination of honey and bee bread collected in the Piedmont Region, Italy. *Environments* 8.
- Beyer, M., Lenouvel, A., Guignard, C., Eickermann, M., Clermont, A., Kraus, F., Hoffmann, L., 2018. Pesticide residue profiles in bee bread and pollen samples and the survival of honeybee colonies—a case study from Luxembourg. *Environ. Sci. Pollut. Res. Int.* 25, 32163–32177.
- Biddinger, D.J., Robertson, J.L., Mullin, C., Frazier, J., Ashcraft, S.A., Rajotte, E.G., et al., 2013. Comparative toxicities and synergism of apple orchard pesticides to *Apis mellifera* (L.) and *Osmia cornifrons* (Radoszkow-ski). *PLoS One* 8, e72587.
- Bishop, G.H., 1961. Growth rates of honey bee larva. *J. Exp. Zool.* 146, 11–20.
- Bjerge, K., Mann, H.M.R., Høye, T.T., Sankey, T., Ahumada, J., 2021. Real-time insect tracking and monitoring with computer vision and deep learning. *Remote Sensing in Ecology and Conservation* 8, 315–327.
- Böhme, F., Bischoff, G., Zebitz, C.P.W., Rosenkranz, P., Wallner, K., 2017. From field to food—will pesticide-contaminated pollen diet lead to a contamination of royal jelly? *Apidologie* 49, 112–119.
- Bond, J.K., Hitaj, C., Smith, D., Hunt, K., Perez, A., Ferreira, G., 2021. Honey Bees on the Move: From Pollination to Honey Production and Back.
- Bonmatin, J.M., Moineau, I., Charvet, R., Fleche, C., Colin, M.E., Bengsch, E.R., 2003. A LC/APCI-MS/MS method for analysis of imidacloprid in soils, in plants, and in pollens. *Anal. Chem.* 75, 2027–2033.
- Bonmatin, J.M., Marchand, P.A., Charvet, R., Moineau, I., Bengsch, E.R., Colin, M.E., 2005. Quantification of imidacloprid uptake in maize crops. *J. Agric. Food Chem.* 53, 5336–5341.
- Bonmatin, J.M., Giorio, C., Girolami, V., Goulson, D., Kreutzweiser, D.P., Krupke, C., et al., 2015. Environmental fate and exposure; neonicotinoids and fipronil. *Environ. Sci. Pollut. Res. Int.* 22, 35–67.
- Bortolotti, L., Costa, C., 2014. Chemical communication in the honey bee society. In: *Mucignat-Caretta, C. (Ed.), Neurobiology of Chemical Communication*. CRC Press/Taylor & Francis © 2014 by Taylor & Francis Group, LLC, Boca Raton (FL).
- Botias, C., David, A., Horwood, J., Abdul-Sada, A., Nicholls, E., Hill, E., Goulson, D., 2015. Neonicotinoid residues in wildflowers, a potential route of chronic exposure for bees. *Environ. Sci. Technol.* 49, 12731–12740.
- Bryden, J., Gill, R.J., Mitton, R.A., Raine, N.E., Jansen, V.A., 2013. Chronic sublethal stress causes bee colony failure. *Ecol. Lett.* 16, 1463–1469.
- Buteler, M., Alma, A.M., Stadler, T., Gingold, A.C., Manattini, M.C., Lozada, M., 2022. Acute toxicity of microplastic fibers to honeybees and effects on foraging behavior. *Sci. Total Environ.* 822, 153320.
- Butler, C.G., 1940. The choice of drinking water by the honeybee. *J. Exp. Biol.* 17, 253–261.
- Byrne, F.J., Visscher, P.K., Leimkuehler, B., Fischer, D., Grafton-Cardwell, E.E., Morse, J. G., 2014. Determination of exposure levels of honey bees foraging on flowers of mature citrus trees previously treated with imidacloprid. *Pest Manag. Sci.* 70, 470–482.
- Calatayud-Vernich, P., Calatayud, F., Simo, E., Pico, Y., 2016. Efficiency of QUCHEERS approach for determining 52 pesticide residues in honey and honey bees. *MethodsX* 3, 452–458.
- Calatayud-Vernich, P., Calatayud, F., Simo, E., Pico, Y., 2018. Pesticide residues in honey bees, pollen and beeswax: assessing beehive exposure. *Environ. Pollut.* 241, 106–114.
- Calatayud-Vernich, P., Calatayud, F., Simo, E., Pascual Aguilar, J.A., Pico, Y., 2019. A two-year monitoring of pesticide hazard in-hive: high honey bee mortality rates during insecticide poisoning episodes in apiaries located near agricultural settings. *Chemosphere* 232, 471–480.
- Carlson, E.A., Melathopoulos, A., Sagili, R., 2022. The value of hazard quotients in honey bee (*Apis mellifera*) ecotoxicology: a review. *Front. Ecol. Evol.* 10.
- Carnesecchi, E., Svendsen, C., Lasagni, S., Grech, A., Quignot, N., Amzal, B., et al., 2019. Investigating combined toxicity of binary mixtures in bees: Meta-analysis of laboratory tests, modelling, mechanistic basis and implications for risk assessment. *Environ. Int.* 133, 105256.
- Charlton, A.J., Jones, A., 2007. Determination of imidazole and triazole fungicide residues in honeybees using gas chromatography-mass spectrometry. *J. Chromatogr. A* 1141, 117–122.
- Chauzat, M.P., Faucon, J.P., 2007. Pesticide residues in beeswax samples collected from honey bee colonies (*Apis mellifera* L.) in France. *Pest Manag. Sci.* 63, 1100–1106.
- Chauzat, M.P., Martel, A.C., Cougoule, N., Porta, P., Lachaize, J., Zeggane, S., et al., 2011. An assessment of honeybee colony matrices, *Apis mellifera* (Hymenoptera: Apidae) to monitor pesticide presence in continental France. *Environ. Toxicol. Chem.* 30, 103–111.
- Chen, Y.P., Pettis, J.S., Collins, A., Feldlaufer, M.F., 2006. Prevalence and transmission of honeybee viruses. *Appl. Environ. Microbiol.* 72, 606–611.
- Chynel, M., Munsch, C., Bely, N., Heas-Moisan, K., Pollono, C., Jaquemet, S., 2021. Legacy and emerging organic contaminants in two sympatric shark species from Reunion Island (Southwest Indian Ocean): levels, profiles and maternal transfer. *Sci. Total Environ.* 751, 141807.
- Cox, C., Zeiss, M., 2022. Health, pesticide adjuvants, and inert ingredients: California case study illustrates need for data access. *Environ. Health Perspect.* 130, 85001.
- Cressey, D., 2017. The bitter battle over the world's most popular insecticides. *Nature* 551, 156–158.
- da Silva, P.M., Gauche, C., Gonzaga, L.V., Costa, A.C., Fett, R., 2016. Honey: chemical composition, stability and authenticity. *Food Chem.* 196, 309–323.
- Daisley, B.A., Chmiel, J.A., Pitek, A.P., Thompson, G.J., Reid, G., 2020. Missing microbes in bees: how systematic depletion of key symbionts erodes immunity. *Trends Microbiol.* 28, 1010–1021.
- Dauchy, X., Boiteux, V., Colin, A., Bach, C., Rosin, C., Munoz, J.F., 2019. Poly- and perfluoroalkyl substances in runoff water and wastewater sampled at a firefighter training area. *Arch. Environ. Contam. Toxicol.* 76, 206–215.
- David, A., Botias, C., Abdul-Sada, A., Goulson, D., Hill, E.M., 2015. Sensitive determination of mixtures of neonicotinoid and fungicide residues in pollen and single bumblebees using a scaled down QUCHEERS method for exposure assessment. *Anal. Bioanal. Chem.* 407, 8151–8162.
- David, A., Botias, C., Abdul-Sada, A., Nicholls, E., Rotheray, E.L., Hill, E.M., Goulson, D., 2016. Widespread contamination of wildflower and bee-collected pollen with complex mixtures of neonicotinoids and fungicides commonly applied to crops. *Environ. Int.* 88, 169–178.
- Davis, B.N., Williams, C.T., 1990. Buffer zone widths for honeybees from ground and aerial spraying of insecticides. *Environ. Pollut.* 63, 247–259.
- de Miranda, J.R., Bailey, L., Ball, B.V., Blanchard, P., Budge, G.E., Chejanovsky, N., et al., 2015. Standard methods for virus research in *Apis mellifera*. *J. Apic. Res.* 52, 1–56.
- de Souza, A.P.F., Rodrigues, N.R., Fernández-Alba, A.R., Reyes Reyes, F.G., 2022. Occurrence of pesticide residues in Brazilian *Apis mellifera* beeswax by gas chromatography-tandem mass spectrometry and pesticide hazard evaluation. *J. Apic. Res.* 1–7.
- Decourtye, A., Devillers, J., Genecque, E., Le Menach, K., Budzinski, H., Cluzeau, S., Pham-Delegue, M.H., 2005. Comparative sublethal toxicity of nine pesticides on olfactory learning performances of the honeybee *Apis mellifera*. *Arch. Environ. Contam. Toxicol.* 48, 242–250.
- Deng, Y., Jiang, X., Zhao, H., Yang, S., Gao, J., Wu, Y., et al., 2021. Microplastic polystyrene ingestion promotes the susceptibility of honeybee to viral infection. *Environ. Sci. Technol.* 55, 11680–11692.

- Dively, G.P., Kamel, A., 2012. Insecticide residues in pollen and nectar of a cucurbit crop and their potential exposure to pollinators. *J. Agric. Food Chem.* 60, 4449–4456.
- Dively, G.P., Embrey, M.S., Kamel, A., Hawthorne, D.J., Pettis, J.S., 2015. Assessment of chronic sublethal effects of imidacloprid on honey bee colony health. *PLoS One* 10, e0118748.
- Dolezal, A.G., Hendrix, S.D., Scavo, N.A., Carrillo-Tripp, J., Harris, M.A., Wheelock, M.J., et al., 2016. Honey bee viruses in wild bees: viral prevalence, loads, and experimental inoculation. *PLoS One* 11, e0166190.
- Doublet, V., Labarussias, M., de Miranda, J.R., Moritz, R.F., Paxton, R.J., 2015. Bees under stress: sublethal doses of a neonicotinoid pesticide and pathogens interact to elevate honey bee mortality across the life cycle. *Environ. Microbiol.* 17, 969–983.
- Drummond, F.A., Ballman, E.S., Eitzer, B.D., Du Clos, B., Dill, J., 2018. Exposure of honey bee (*Apis mellifera* L.) colonies to pesticides in pollen, a statewide assessment in Maine. *Environ. Entomol.* 47, 378–387.
- Edo, C., Fernandez-Alba, A.R., Vejsnaes, F., van der Steen, J.J.M., Fernandez-Pinas, F., Rosal, R., 2021. Honeybees as active samplers for microplastics. *Sci. Total Environ.* 767, 144481.
- El Agrebi, N., Traynor, K., Wilmart, O., Tosi, S., Leinartz, L., Danneels, E., et al., 2020. Pesticide and veterinary drug residues in Belgian beeswax: occurrence, toxicity, and risk to honey bees. *Sci. Total Environ.* 745, 141036.
- El Hassani, A.K., Dacher, M., Gary, V., Lambin, M., Gauthier, M., Armengaud, C., 2008. Effects of sublethal doses of acetamiprid and thiamethoxam on the behavior of the honeybee (*Apis mellifera*). *Arch. Environ. Contam. Toxicol.* 54, 653–661.
- Enright, H.A., Malfatti, M.A., Zimmermann, M., Ognibene, T., Henderson, P., Turteltaub, K.W., 2016. Use of accelerator mass spectrometry in human health and molecular toxicology. *Chem. Res. Toxicol.* 29, 1976–1986.
- EPA, 2014. Guidance for Assessing Pesticide Risks to Bees.
- EPA, 2016a. Guidance on Exposure and Effects Testing for Assessing Risks to Bees.
- EPA, 2016b. Process for Requiring Exposure and Effects Testing for Assessing Risks to Bees During Registration and Registration Review.
- Erdogru, Ö., 2007. Levels of selected pesticides in honey samples from Kahramanmaraş, Turkey. *Food Control* 18, 866–871.
- Farooqui, T., 2013. A potential link among biogenic amines-based pesticides, learning and memory, and colony collapse disorder: a unique hypothesis. *Neurochem. Int.* 62, 122–136.
- Fine, J.D., 2020. Evaluation and comparison of the effects of three insect growth regulators on honey bee queen oviposition and egg eclosion. *Ecotoxicol. Environ. Saf.* 205, 111142.
- Fine, J.D., Cox-Foster, D.L., Mullin, C.A., 2017a. An inert pesticide adjuvant synergizes viral pathogenicity and mortality in honey bee larvae. *Sci. Rep.* 7, 40499.
- Fine, J.D., Mullin, C.A., Frazier, M.T., Reynolds, R.D., 2017b. Field residues and effects of the insect growth regulator novaluron and its major co-formulant N-methyl-2-pyrrolidone on honey bee reproduction and development. *J. Econ. Entomol.* 110, 1993–2001.
- Fine, J.D., Cox-Foster, D.L., Moor, K.J., Chen, R., Avalos, A., 2023. Trisiloxane surfactants negatively affect reproductive behaviors and enhance viral replication in honey bees. *Environ. Toxicol. Chem.* 43, 222–233.
- Fisher 2nd, A., DeGrandi-Hoffman, G., Smith, B.H., Johnson, M., Kaftanoglu, O., Cogley, T., et al., 2021. Colony field test reveals dramatically higher toxicity of a widely-used mitotic fungicide on honey bees (*Apis mellifera*). *Environ. Pollut.* 269, 115964.
- Fox, C.W., 1993. The influence of maternal age and mating frequency on egg size and offspring performance in *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Oecologia* 96, 139–146.
- Free, J.B., 1959. The transfer of food between the adult members of a honeybee community. *Bee World* 40, 193–201.
- Gage, S.L., Ahumada, F., Rivera, A., Graham, H., DeGrandi-Hoffman, G., 2018. Smoke conditions affect the release of the venom droplet accompanying sting extension in honey bees (Hymenoptera: Apidae). *J. Insect Sci.* 18.
- Gate, I.M., McNeill, S., Ashmore, M.R., 1995. Effects of air pollution on the searching behaviour of an insect parasitoid. *Water Air Soil Pollut.* 85, 1425–1430.
- Gérez, N., Pérez-Parada, A., Cesio, M.V., Heinzen, H., 2017. Occurrence of pesticide residues in candies containing bee products. *Food Control* 72, 293–299.
- Gernat, T., Jagla, T., Jones, B.M., Middendorf, M., Robinson, G.E., 2023. Automated monitoring of honey bees with barcodes and artificial intelligence reveals two distinct social networks from a single affiliative behavior. *Sci. Rep.* 13, 1541.
- Giroud, B., Vaucher, A., Vulliet, E., Wiest, L., Bulete, A., 2013. Trace level determination of pyrethroid and neonicotinoid insecticides in bee bread using acetonitrile-based extraction followed by analysis with ultra-high-performance liquid chromatography-tandem mass spectrometry. *J. Chromatogr. A* 1316, 53–61.
- Glass, J.R., Fisher 2nd, A., Fewell, J.H., DeGrandi-Hoffman, G., Ozturk, C., Harrison, J.F., 2021. Consumption of field-realistic doses of a widely used mitotic fungicide reduces thorax mass but does not negatively impact flight capacities of the honey bee (*Apis mellifera*). *Environ. Pollut.* 274, 116533.
- Gluge, J., Scheringer, M., Cousins, I.T., DeWitt, J.C., Goldenman, G., Herzke, D., et al., 2020. An overview of the uses of per- and polyfluoroalkyl substances (PFAS). *Environ. Sci. Process Impacts* 22, 2345–2373.
- Green, J., 2000. Adjuvant outlook for pesticides. *Pestic. Outlook* 11, 196–199.
- Gunes, M.E., Sari, M.F., Esen, F., 2021. Organochlorine pesticides in honeybee, pollen and honey in Bursa, Turkey. *Food Addit. Contam. Part B Surveill.* 14, 126–132.
- Hammer, M., Menzel, R., 1995. Learning and memory in the honeybee. *J. Neurosci.* 15, 1617–1630.
- Hatjina, F., Papaefthimiou, C., Charistos, L., Dogaroglu, T., Bouga, M., Emmanouil, C., Arnold, G., 2013. Sublethal doses of imidacloprid decreased size of hypopharyngeal glands and respiratory rhythm of honeybees in vivo. *Apidologie* 44, 467–480.
- Hawthorne, D.J., Dively, G.P., 2011. Killing them with kindness? In-hive medications may inhibit xenobiotic efflux transporters and endanger honey bees. *PLoS One* 6, e26796.
- Haydak, M.H., 1943. Larval food and development of castes in the honeybee. *J. Econ. Entomol.* 36, 778–792.
- Hazen, J.L., 2000. Adjuvants—terminology, classification, and chemistry. *Weed Technol.* 14, 773–784.
- Hb Kanga, L., 2019. Pesticide residues in conventionally and organically managed apiaries in south and North Florida. *Current Investigations in Agriculture and Current Research* 7.
- Henry, M., Beguin, M., Requier, F., Rollin, O., Odoux, J.F., Aupinel, P., et al., 2012. A common pesticide decreases foraging success and survival in honey bees. *Science* 336, 348–350.
- Henry, M., Cerrutti, N., Aupinel, P., Decourtye, A., Gayrard, M., Odoux, J.F., et al., 2015. Reconciling laboratory and field assessments of neonicotinoid toxicity to honeybees. *Proc. Biol. Sci.* 282.
- Hock, W., 2022. Spray Adjuvants.
- Hodgson, E.W., Pitts-Singer, T.L., Barbour, J.D., 2011. Effects of the insect growth regulator, novaluron on immature alfalfa leafcutting bees, *Megachile rotundata*. *J. Insect Sci.* 11, 43.
- Holloway, P.J., Western, N.M., 2003. Tank-mix adjuvants and pesticide residues: some regulatory and quantitative aspects. *Pest Manag. Sci.* 59, 1237–1244.
- Hooper-Bui, L.M., Kwok, E.S., Buchholz, B.A., Rust, M.K., Eastmond, D.A., Vogel, J.S., 2015. Insecticide transfer efficiency and lethal load in Argentine ants. *Nucl. Inst. Methods Phys. Res. B* 361, 665–669.
- Hotchkiss, M.Z., Poulain, A.J., Forrest, J.R.K., 2022. Pesticide-induced disturbances of bee gut microbiotas. *FEMS Microbiol. Rev.* 46.
- Ingram, E.M., Augustin, J., Ellis, M.D., Siegfried, B.D., 2015. Evaluating sub-lethal effects of orchard-applied pyrethroids using video-tracking software to quantify honey bee behaviors. *Chemosphere* 135, 272–277.
- Issa, M.M., S, M.T., El-Marsafy, A.M., MMH, Khalil, Ismail, E.H., 2020. Acetonitrile-ethyl acetate based method for the residue analysis of 373 pesticides in beeswax using LC-MS/MS and GC-MS/MS. *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* 1145, 122106.
- Iwasa, T., Motoyama, N., Ambrose, J.T., Roe, R.M., 2004. Mechanism for the differential toxicity of neonicotinoid insecticides in the honey bee, *Apis mellifera*. *Crop Prot.* 23, 371–378.
- Jacobson, M., 1966. Chemical insect attractants and repellents. *Annu. Rev. Entomol.* 11, 403–422.
- Jiang, J., Ma, D., Zou, N., Yu, X., Zhang, Z., Liu, F., Mu, W., 2018. Concentrations of imidacloprid and thiamethoxam in pollen, nectar and leaves from seed-dressed cotton crops and their potential risk to honeybees (*Apis mellifera* L.). *Chemosphere* 201, 159–167.
- Johnson, B.R., 2010. Division of labor in honeybees: form, function, and proximate mechanisms. *Behav. Ecol. Sociobiol.* 64, 305–316.
- Johnson, R.M., 2015. Honey bee toxicology. *Annu. Rev. Entomol.* 60, 415–434.
- Johnson, R.M., Ellis, M.D., Mullin, C.A., Frazier, M., 2010. Pesticides and honey bee toxicity - USA. *Apidologie* 41, 312–331.
- Johnson, R.M., Dahlgren, L., Siegfried, B.D., Ellis, M.D., 2013. Acaricide, fungicide and drug interactions in honey bees (*Apis mellifera*). *PLoS One* 8, e54092.
- Joseph, S.V., 2019. Transovarial effects of insect growth regulators on *Stephanitis pyrioides* (Hemiptera: Tingidae). *Pest Manag. Sci.* 75, 2182–2187.
- Jouanneau, W., Leandri-Breton, D.J., Corbeau, A., Herzke, D., Moe, B., Nikiforov, V.A., et al., 2022. A bad start in life? Maternal transfer of legacy and emerging poly- and perfluoroalkyl substances to eggs in an Arctic seabird. *Environ. Sci. Technol.* 56, 6091–6102.
- Kakumanu, M.L., Reeves, A.M., Anderson, T.D., Rodrigues, R.R., Williams, M.A., 2016. Honey bee gut microbiome is altered by in-hive pesticide exposures. *Front. Microbiol.* 7, 1255.
- Kasiotis, K.M., Anagnostopoulos, C., Anastasiadou, P., Machera, K., 2014. Pesticide residues in honeybees, honey and bee pollen by LC-MS/MS screening: reported death incidents in honeybees. *Sci. Total Environ.* 485–486, 633–642.
- Kast, C., Kilchenmann, V., 2022. An in vitro model for assessing the toxicity of pesticides in beeswax on honey bee larvae. *Chemosphere* 287, 132214.
- Khalifa, S.A.M., Elshafiey, E.H., Shetaia, A.A., El-Wahed, A.A.A., Algethami, A.F., Musharraf, S.G., et al., 2021. Overview of bee pollination and its economic value for crop production. *Insects* 12.
- Klein, A.M., Boreux, V., Fornoff, F., Mupepele, A.C., Pufal, G., 2018. Relevance of wild and managed bees for human well-being. *Curr Opin Insect Sci* 26, 82–88.
- Koch, A., Jonsson, M., Yeung, L.W.Y., Karrman, A., Ahrens, L., Ekblad, A., Wang, T., 2021. Quantification of biotransfer of per- and polyfluoroalkyl substances from the aquatic to the terrestrial environment via emergent insects. *Environ. Sci. Technol.* 55, 7900–7909.
- Kritsky, G., 2017. Beekeeping from antiquity through the middle ages. *Annu. Rev. Entomol.* 62, 249–264.
- Krupke, C.H., Hunt, G.J., Eitzer, B.D., Andino, G., Given, K., 2012. Multiple routes of pesticide exposure for honey bees living near agricultural fields. *PLoS One* 7, e29268.
- Kutschera, W., 2016. Accelerator mass spectrometry: state of the art and perspectives. *Advances in Physics: X* 1, 570–595.
- Kwong, W.K., Moran, N.A., 2016. Gut microbial communities of social bees. *Nat. Rev. Microbiol.* 14, 374–384.
- Lan, Z., Yao, Y., Xu, J., Chen, H., Ren, C., Fang, X., et al., 2020. Novel and legacy per- and polyfluoroalkyl substances (PFASs) in a farmland environment: soil distribution and biomonitoring with plant leaves and locusts. *Environ. Pollut.* 263, 114487.

- Larson, J.L., Redmond, C.T., Potter, D.A., 2015. Mowing mitigates bioactivity of neonicotinoid insecticides in nectar of flowering lawn weeds and turfgrass guttation. *Environ. Toxicol. Chem.* 34, 127–132.
- Lau, P.W., Nieh, J.C., 2016. Salt preferences of honey bee water foragers. *J. Exp. Biol.* 219, 790–796.
- Laurent, F.M., Rathahao, E., 2003. Distribution of [(14)C]imidacloprid in sunflowers (*Helianthus annuus* L.) following seed treatment. *J. Agric. Food Chem.* 51, 8005–8010.
- Laycock, I., Cotterell, K.C., O'Shea-Wheller, T.A., Cresswell, J.E., 2014. Effects of the neonicotinoid pesticide thiamethoxam at field-realistic levels on microcolonies of *Bombus terrestris* worker bumble bees. *Ecotoxicol. Environ. Saf.* 100, 153–158.
- Lentola, A., David, A., Abdul-Sada, A., Tapparo, A., Goulson, D., Hill, E.M., 2017. Ornamental plants on sale to the public are a significant source of pesticide residues with implications for the health of pollinating insects. *Environ. Pollut.* 228, 297–304.
- Levine, S.L., Borgert, C.J., 2018. Review and recommendations on criteria to evaluate the relevance of pesticide interaction data for ecological risk assessments. *Chemosphere* 209, 124–136.
- Lin, A.Y., Panchangam, S.C., Tsai, Y.T., Yu, T.H., 2014. Occurrence of perfluorinated compounds in the aquatic environment as found in science park effluent, river water, rainwater, sediments, and biotissues. *Environ. Monit. Assess.* 186, 3265–3275.
- Litsey, E.M., Chung, S., Fine, J.D., 2021. The behavioral toxicity of insect growth disruptors on *Apis mellifera* Queen care. *Front. Ecol. Evol.* 9.
- Lu, C.S., Warchol, K.M., Callahan, R.A., 2014. Sub-lethal exposure to neonicotinoids impaired honey bees winterization before proceeding to colony collapse disorder. *Bulletin of Insectology* 67, 125–130.
- Ludicke, J.C., Nieh, J.C., 2020. Thiamethoxam impairs honey bee visual learning, alters decision times, and increases abnormal behaviors. *Ecotoxicol. Environ. Saf.* 193, 110367.
- Lyons, K., Adams, D.H., 2015. Maternal offloading of organochlorine contaminants in the yolk-sac placental scalloped hammerhead shark (*Sphyrna lewini*). *Ecotoxicology* 24, 553–562.
- Lyons, K., Carlisle, A., Preti, A., Mull, C., Blasius, M., O'Sullivan, J., et al., 2013a. Effects of trophic ecology and habitat use on maternal transfer of contaminants in four species of young of the year lamniform sharks. *Mar. Environ. Res.* 90, 27–38.
- Lyons, K., Lowe, C.G., Gillanders, B.M., 2013b. Mechanisms of maternal transfer of organochlorine contaminants and mercury in the common thresher shark (*Alopias vulpinus*). *Can. J. Fish. Aquat. Sci.* 70, 1667–1672.
- Macri, I.N., Vazquez, D.E., Pagano, E.A., Zavala, J.A., Farina, W.M., 2021. Evaluating the impact of post-emergence weed control in honeybee colonies located in different agricultural surroundings. *Insects* 12.
- Magal, P., Webb, G.F., Wu, Y., 2019. An environmental model of honey bee colony collapse due to pesticide contamination. *Bull. Math. Biol.* 81, 4908–4931.
- Magal, P., Webb, G.F., Wu, Y., 2020. A spatial model of honey bee colony collapse due to pesticide contamination of foraging bees. *J. Math. Biol.* 80, 2363–2393.
- Manning, R., 2018. Chemical residues in beebread, honey, pollen and wax samples collected from bee hives placed on canola crops in Western Australia. *J. Apic. Res.* 57, 696–708.
- Marti, J.N.G., Kilchenmann, V., Kast, C., 2022. Evaluation of pesticide residues in commercial Swiss beeswax collected in 2019 using ultra-high performance liquid chromatographic analysis. *Environ. Sci. Pollut. Res. Int.* 29, 32054–32064.
- Marzaro, M., Vivan, L., Targa, A., Mazzon, L., Mori, N., Greatti, M., et al., 2011. Lethal aerial powdering of honey bees with neonicotinoids from fragments of maize seed coat. *Bulletin of Insectology* 64, 119–126.
- Maus, C., Alix, A., Schmehl, D.R., 2022. Assessing the impact of pesticides on pollinators. Promoting Pollination and Pollinators in Farming 183–224.
- Mayer, D.F., Lunden, J.D., 2015. Field and laboratory tests of the effects of fipronil on adult female bees of *Apis mellifera*, *Megachile rotundata* and *Nomia melanderi*. *J. Apic. Res.* 38, 191–197.
- McFrederick, Q.S., Fuentes, J.D., Roulston, T., Kathilankal, J.C., Lerdau, M., 2009. Effects of air pollution on biogenic volatiles and ecological interactions. *Oecologia* 160, 411–420.
- Medina, P., Smaghe, G., Budia, F., Del Estal, P., Tirry, L., Viñuela, E., 2002. Significance of penetration, excretion, and transovarial uptake to toxicity of three insect growth regulators in predatory lacewing adults. *Arch. Insect Biochem. Physiol.* 51, 91–101.
- Medina, P., Budia, F., Tirry, L., Smaghe, G., Viñuela, E., 2010. Compatibility of Spinosad, Tebufenozide and Azadirachtin with eggs and pupae of the predator *Chrysoperla carnea* (Stephens) under laboratory conditions. *Biocontrol Sci. Tech.* 11, 597–610.
- Medrzycki, P., Giffard, H., Aupinel, P., Belzunces, L.P., Chauzat, M.-P., Clafén, C., et al., 2015. Standard methods for toxicology research in *Apis mellifera*. *J. Apic. Res.* 52, 1–60.
- Meled, M., Thrasylvoulou, A., Belzunces, L.P., 1998. Seasonal variations in susceptibility of *Apis mellifera* to the synergistic action of prochloraz and deltamethrin. *Environ. Toxicol. Chem.* 17, 2517–2520.
- Milone, J.P., Tarpy, D.R., 2021. Effects of developmental exposure to pesticides in wax and pollen on honey bee (*Apis mellifera*) queen reproductive phenotypes. *Sci. Rep.* 11, 1020.
- Milone, J.P., Chakrabarti, P., Sagili, R.R., Tarpy, D.R., 2021. Colony-level pesticide exposure affects honey bee (*Apis mellifera* L.) royal jelly production and nutritional composition. *Chemosphere* 263, 128183.
- Miner, K.R., Clifford, H., Taruscio, T., Potocki, M., Solomon, G., Ritari, M., et al., 2021. Deposition of PFAS 'forever chemicals' on Mt. Everest. *Sci. Total Environ.* 759, 144421.
- Mititelu, M., Udeanu, D., Nedeleescu, M., Neacsu, S., Nicoara, A., Oprea, E., Ghica, M., 2022. Quality control of different types of honey and propolis collected from Romanian accredited beekeepers and consumer's risk assessment. *Crystals* 12.
- Moffett, J.O., Morton, H.L., Macdonald, R.H., 1972. Toxicity of some herbicidal sprays to honey bees. *J. Econ. Entomol.* 65, 32–36.
- Motta, E.V.S., Moran, N.A., 2020. Impact of glyphosate on the honey bee gut microbiota: effects of intensity, duration, and timing of exposure. *mSystems* 5.
- Motta, E.V.S., Moran, N.A., 2023. The honeybee microbiota and its impact on health and disease. *Nat. Rev. Microbiol.* <https://doi.org/10.1038/s41579-023-00990-3>.
- Motta, E.V.S., Raymann, K., Moran, N.A., 2018. Glyphosate perturbs the gut microbiota of honey bees. *Proc. Natl. Acad. Sci. U. S. A.* 115, 10305–10310.
- Mousseau, T.A., Dingle, H., 1991. Maternal effects in insect life histories. *Annu. Rev. Entomol.* 36, 511–534.
- Mu, H., Wang, K., Yang, X., Xu, W., Liu, X., Ritsema, C.J., Geissen, V., 2022. Pesticide usage practices and the exposure risk to pollinators: a case study in the North China Plain. *Ecotoxicol. Environ. Saf.* 241, 113713.
- Mull, C.G., Lyons, K., Blasius, M.E., Winkler, C., O'Sullivan, J.B., Lowe, C.G., 2013. Evidence of maternal offloading of organic contaminant residues in white sharks (*Carcharodon carcharias*). *PLoS One* 8, e62886.
- Mullin, C.A., Frazier, M., Frazier, J.L., Ashcraft, S., Simonds, R., Vanengelsdorp, D., Pettis, J.S., 2010. High levels of miticides and agrochemicals in North American apiaries: implications for honey bee health. *PLoS One* 5, e9754.
- Mullin, C.A., Chen, J., Fine, J.D., Frazier, M.T., Frazier, J.L., 2015. The formulation makes the honey bee poison. *Pestic. Biochem. Physiol.* 120, 27–35.
- Mullin, C.A., Fine, J.D., Reynolds, R.D., Frazier, M.T., 2016. Toxicological risks of agrochemical spray adjuvants: organosilicone surfactants may not be safe. *Front. Public Health* 4, 92.
- Murcia Morales, M., Gomez Ramos, M.J., Parrilla Vazquez, P., Diaz Galiano, F.J., Garcia Valverde, M., Gamiz Lopez, V., et al., 2020. Distribution of chemical residues in the beehive compartments and their transfer to the honeybee brood. *Sci. Total Environ.* 710, 136288.
- Murcia-Morales, M., Heinzen, H., Parrilla-Vázquez, P., Gómez-Ramos, M.D., Fernández-Alba, A.R., 2022. Presence and distribution of pesticides in apicultural products: a critical appraisal. *TrAC Trends Anal. Chem.* 146.
- Navarro, I., de la Torre, A., Sanz, P., Porcel, M.A., Pro, J., Carbonell, G., Martínez, M.L., 2017. Uptake of perfluoroalkyl substances and halogenated flame retardants by crop plants grown in biosolids-amended soils. *Environ. Res.* 152, 199–206.
- Negi, B.R., Sharma, H.K., Thakur, M., Prasad, H., 2022. Lethal and sublethal effects of thiamethoxam, a neonicotinoid molecule, on colony performance of *A. mellifera*. *Environ. Sci. Pollut. Res. Int.* 29, 10826–10837.
- Nicklisch, S.C.T., Hamdoun, A., 2020. Disruption of small molecule transporter systems by Transporter-Interfering Chemicals (TICs). *FEBS Lett.* 594, 4158–4185.
- Nicklisch, S.C., Rees, S.D., McGrath, A.P., Gokirmak, T., Bonito, L.T., Vermeer, L.M., et al., 2016. Global marine pollutants inhibit P-glycoprotein: environmental levels, inhibitory effects, and cocrystal structure. *Sci. Adv.* 2, e1600001.
- Nicolson, S.W., 2009. Water homeostasis in bees, with the emphasis on sociality. *J. Exp. Biol.* 212, 429–434.
- Nnane, I.P., Tao, X., 2005. Drug metabolism | isotope studies. *Encyclopedia of Analytical Science* 311–316.
- Nozal, M.J., Imaz, E., Bernal, J.L., Nieto, J.L., Higes, M., Bernal, J., 2021. An optimized extraction procedure for determining acaricide residues in foundation sheets of beeswax by using gas chromatography-mass spectrometry. *Agronomy* 11.
- Orantes-Bermejo, F.J., Pajuelo, A.G., Megías, M.M., Fernández-Piñar, C.T., 2015. Pesticide residues in beeswax and beebread samples collected from honey bee colonies (*Apis mellifera* L.) in Spain. Possible implications for bee losses. *J. Apic. Res.* 49, 243–250.
- Osterman, J., Aizen, M.A., Biesmeijer, J.C., Bosch, J., Howlett, B.G., Inouye, D.W., et al., 2021. Global trends in the number and diversity of managed pollinator species. *Agric. Ecosyst. Environ.* 322.
- Ostiguy, N., Drummond, F.A., Aronstein, K., Eitzer, B., Ellis, J.D., Spivak, M., Sheppard, W.S., 2019. Honey bee exposure to pesticides: a four-year nationwide study. *Insects* 10.
- Oxley, P.R., Oldroyd, B.P., 2010. The Genetic Architecture of Honeybee Breeding 83–118.
- Page Jr., R.E., 2013. *The Spirit of the Hive*. Harvard University Press, Cambridge, MA and London, England.
- Page Jr., R.E., Peng, C.Y., 2001. Aging and development in social insects with emphasis on the honey bee, *Apis mellifera* L. *Exp. Gerontol.* 36, 695–711.
- Pal, E., Almasri, H., Paris, L., Diogon, M., Pioz, M., Cousin, M., et al., 2022. Toxicity of the pesticides imidacloprid, difenoconazole and glyphosate alone and in binary and ternary mixtures to winter honey bees: effects on survival and antioxidative defenses. *Toxics* 10.
- Pareja, L., Colazzo, M., Perez-Parada, A., Niell, S., Carrasco-Letelier, L., Besil, N., et al., 2011. Detection of pesticides in active and depopulated beehives in Uruguay. *Int. J. Environ. Res. Public Health* 8, 3844–3858.
- Payne, A.N., Walsh, E.M., Rangel, J., 2019. Initial exposure of wax foundation to agrochemicals causes negligible effects on the growth and winter survival of incipient honey bee (*Apis mellifera*) colonies. *Insects* 10.
- Peters, J.M., Peleg, O., Mahadevan, L., 2019. Collective ventilation in honeybee nests. *J. R. Soc. Interface* 16, 20180561.
- Pettis, J.S., vanEngelsdorp, D., Johnson, J., Dively, G., 2012. Pesticide exposure in honey bees results in increased levels of the gut pathogen *Nosema*. *Naturwissenschaften* 99, 153–158.
- Pilling, E.D., Jepson, P.C., 1993. Synergism between EBI fungicides and a pyrethroid insecticide in the honeybee (*Apis mellifera*). *Pestic. Sci.* 39, 293–297.
- Pilling, E.D., Bromleychallenor, K.A.C., Walker, C.H., Jepson, P.C., 1995. Mechanism of synergism between the pyrethroid insecticide λ -cyhalothrin and the imidazole fungicide prochloraz, in the honeybee (*Apis mellifera* L.). *Pestic. Biochem. Physiol.* 51, 1–11.

- Pisa, L.W., Amaral-Rogers, V., Belzunces, L.P., Bonmatin, J.M., Downs, C.A., Goulson, D., et al., 2015. Effects of neonicotinoids and fipronil on non-target invertebrates. *Environ. Sci. Pollut. Res. Int.* 22, 68–102.
- Podder, A., Sadmani, A., Reinhart, D., Chang, N.B., Goel, R., 2021. Per and poly-fluoroalkyl substances (PFAS) as a contaminant of emerging concern in surface water: a transboundary review of their occurrences and toxicity effects. *J. Hazard. Mater.* 419, 126361.
- Pohorecka, K., Skubida, P., Miszczyk, A., Semkiw, P., Sikorski, P., Zagibajto, K., et al., 2012. Residues of neonicotinoid insecticides in bee collected plant materials from oilseed rape crops and their effect on bee colonies. *Journal of Apicultural Science* 56, 115–134.
- Pohorecka, K., Szczesna, T., Witek, M., Miszczyk, A., Sikorski, P., 2017. The exposure of honey bees to pesticide residues in the hive environment with regard to winter colony losses. *Journal of Apicultural Science* 61, 105–125.
- Porto, R.G., de Almeida, R.F., Cruz-Neto, O., Tabarelli, M., Viana, B.F., Peres, C.A., Lopes, A.V., 2020. Pollination ecosystem services: a comprehensive review of economic values, research funding and policy actions. *Food Security* 12, 1425–1442.
- Posada-Florez, F., Lamas, Z.S., Hawthorne, D.J., Chen, Y., Evans, J.D., Ryabov, E.V., 2021. Pupal cannibalism by worker honey bees contributes to the spread of deformed wing virus. *Sci. Rep.* 11, 8989.
- Raimets, R., Bontsutsnaja, A., Bartkevics, V., Pugajeva, I., Kaart, T., Puusepp, L., et al., 2020. Pesticide residues in beehive matrices are dependent on collection time and matrix type but independent of proportion of foraged oilseed rape and agricultural land in foraging territory. *Chemosphere* 238, 124555.
- Rangel, J., Tarpy, D.R., 2016. The combined effects of miticides on the mating health of honey bee (*Apis mellifera*) queens. *J. Apic. Res.* 54, 275–283.
- Reitmayer, C.M., Ryalls, J.M.W., Farthing, E., Jackson, C.W., Girling, R.D., Newman, T.A., 2019. Acute exposure to diesel exhaust induces central nervous system stress and altered learning and memory in honey bees. *Sci. Rep.* 9, 5793.
- Remolina, S.C., Hughes, K.A., 2008. Evolution and mechanisms of long life and high fertility in queen honey bees. *Age (Dordr.)* 30, 177–185.
- Rossiter, M.C., 1991. Environmentally-based maternal effects: a hidden force in insect population dynamics? *Oecologia* 87, 288–294.
- Rozko, M.L., Kaminska, M., Szymczyk, K., Jedrzejczak, R., 2016. Levels of selected persistent organic pollutants (PCB, PBDE) and pesticides in honey bee pollen sampled in Poland. *PLoS One* 11, e0167487.
- Samuelson, A.E., Gill, R.J., Brown, M.J.F., Leadbeater, E., 2018. Lower bumblebee colony reproductive success in agricultural compared with urban environments. *Proc. Biol. Sci.* 285.
- Sanchez-Bayo, F., 2014. Environmental science. The trouble with neonicotinoids. *Science* 346, 806–807.
- Sanchez-Bayo, F., Goka, K., 2014. Pesticide residues and bees—a risk assessment. *PLoS One* 9, e94482.
- Sanchez-Bayo, F., Goulson, D., Pennacchio, F., Nazzi, F., Goka, K., Desneux, N., 2016. Are bee diseases linked to pesticides? - a brief review. *Environ. Int.* 89-90, 7–11.
- Schaad, E., Fracheboud, M., Droz, B., Kast, C., 2023. Quantification of pesticides in bee bread collected from honey bee colonies in an agricultural environment in Switzerland. *Environ. Sci. Pollut. Res. Int.* 30, 56353–56367.
- Schmuck, R., Schoning, R., Stork, A., Schramel, O., 2001. Risk posed to honeybees (*Apis mellifera* L., Hymenoptera) by an imidacloprid seed dressing of sunflowers. *Pest Manag. Sci.* 57, 225–238.
- Seeley, T.D., 2009. *The Wisdom of the Hive*. Harvard University Press.
- Simone-Finstrom, M., Li-Byarlay, H., Huang, M.H., Strand, M.K., Rueppell, O., Tarpy, D.R., 2016. Migratory management and environmental conditions affect lifespan and oxidative stress in honey bees. *Sci. Rep.* 6, 32023.
- Simek, I., Kuzukiran, O., Yurdakok-Dikmen, B., Sireli, U.T., Beykaya, M., Filazi, A., 2021. Comparison of selected lipophilic compound residues in honey and propolis. *J. Food Compos. Anal.* 102.
- Siviter, H., Bailes, E.J., Martin, C.D., Oliver, T.R., Koricheva, J., Leadbeater, E., Brown, M.J.F., 2021. Agrochemicals interact synergistically to increase bee mortality. *Nature* 596, 389–392.
- Smodis Skerl, M.L., Velikonja Bolta, S., Basa Cesnik, H., Gregorc, A., 2009. Residues of pesticides in honeybee (*Apis mellifera carnica*) bee bread and in pollen loads from treated apple orchards. *Bull. Environ. Contam. Toxicol.* 83, 374–377.
- Sontner, C.A., Rader, R., Stevenson, G., Stavert, J.R., Wilson, S.C., 2021. Biological and behavioral responses of European honey bee (*Apis mellifera*) colonies to perfluorooctane sulfonate exposure. *Integr. Environ. Assess. Manag.* 17, 673–683.
- Southwick, E.E., Moritz, R.F.A., 1987. Social control of air ventilation in colonies of honey bees, *Apis mellifera*. *J. Insect Physiol.* 33, 623–626.
- Sponsler, D.B., Johnson, R.M., 2017. Mechanistic modeling of pesticide exposure: the missing keystone of honey bee toxicology. *Environ. Toxicol. Chem.* 36, 871–881.
- Spurgeon, D., Hesketh, H., Lahive, E., Svendsen, C., Baas, J., Robinson, A., et al., 2016. Chronic oral lethal and sub-lethal toxicities of different binary mixtures of pesticides and contaminants in bees (*Apis mellifera*, *Osmia bicornis* and *Bombus terrestris*). EFSA Supporting Publications 13.
- Stabentheiner, A., Kovac, H., Mandl, M., Kafer, H., 2021. Coping with the cold and fighting the heat: thermal homeostasis of a superorganism, the honeybee colony. *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* 207, 337–351.
- Stokstad, E., 2018. European Union Expands Ban of Three Neonicotinoid Pesticides, 2023.
- Stoner, K.A., Eitzer, B.D., 2012. Movement of soil-applied imidacloprid and thiamethoxam into nectar and pollen of squash (*Cucurbita pepo*). *PLoS One* 7, e39114.
- Stoner, K.A., Eitzer, B.D., 2013. Using a hazard quotient to evaluate pesticide residues detected in pollen trapped from honey bees (*Apis mellifera*) in Connecticut. *PLoS One* 8, e77550.
- Straw, E.A., Thompson, L.J., Leadbeater, E., Brown, M.J.F., 2022. 'Inert' ingredients are understudied, potentially dangerous to bees and deserve more research attention. *Proc. Biol. Sci.* 289, 20212353.
- Surma, M., Zielinski, H., Piskula, M., 2016. Levels of contamination by perfluoroalkyl substances in honey from selected European countries. *Bull. Environ. Contam. Toxicol.* 97, 112–118.
- Taber, S., Roberts, W.C., 1963. Egg weight variability and its inheritance in the honey bee. *Ann. Entomol. Soc. Am.* 56, 473–476.
- Taenzler, V., Weyers, A., Maus, C., Ebeling, M., Levine, S., Cabrera, A., et al., 2023. Acute toxicity of pesticide mixtures to honey bees is generally additive, and well predicted by concentration addition. *Sci. Total Environ.* 857, 159518.
- Tang, H.P.O., 2013. Recent development in analysis of persistent organic pollutants under the Stockholm convention. *TrAC Trends Anal. Chem.* 45, 48–66.
- Thimmegowda, G.G., Mullen, S., Sottolare, K., Sharma, A., Mohanta, S.S., Brockmann, A., et al., 2020. A field-based quantitative analysis of sublethal effects of air pollution on pollinators. *Proc. Natl. Acad. Sci. U. S. A.* 117, 20653–20661.
- Thompson, H.M., 2021. The use of the Hazard Quotient approach to assess the potential risk to honeybees (*Apis mellifera*) posed by pesticide residues detected in bee-relevant matrices is not appropriate. *Pest Manag. Sci.* 77, 3934–3941.
- Thompson, H.M., Fryday, S.L., Harkin, S., Milner, S., 2014. Potential impacts of synergism in honeybees (*Apis mellifera*) of exposure to neonicotinoids and sprayed fungicides in crops. *Apidologie* 45, 545–553.
- Tison, L., Hahn, M.L., Holtz, S., Rossner, A., Greggers, U., Bischoff, G., Menzel, R., 2016. Honey bees' behavior is impaired by chronic exposure to the neonicotinoid thiacloprid in the field. *Environ. Sci. Technol.* 50, 7218–7227.
- Tome, H.V.V., Schmehl, D.R., Wedde, A.E., Godoy, R.S.M., Ravaiano, S.V., Guedes, R.N.C., et al., 2020. Frequently encountered pesticides can cause multiple disorders in developing worker honey bees. *Environ. Pollut.* 256, 113420.
- Tong, Z., Duan, J., Wu, Y., Liu, Q., He, Q., Shi, Y., et al., 2018. A survey of multiple pesticide residues in pollen and bee bread collected in China. *Sci. Total Environ.* 640-641, 1578–1586.
- Tosi, S., Nieh, J.C., 2019. Lethal and sublethal synergistic effects of a new systemic pesticide, flupyradifurone (Sivanto(R)), on honeybees. *Proc. Biol. Sci.* 286, 20190433.
- Tosi, S., Costa, C., Vesco, U., Quaglia, G., Guido, G., 2018. A 3-year survey of Italian honey bee-collected pollen reveals widespread contamination by agricultural pesticides. *Sci. Total Environ.* 615, 208–218.
- Traver, B.E., Feazel-Orr, H.K., Catalfamo, K.M., Brewster, C.C., Fell, R.D., 2018. Seasonal effects and the impact of in-hive pesticide treatments on parasite, pathogens, and health of honey bees. *J. Econ. Entomol.* 111, 517–527.
- Traynor, K.S., Pettis, J.S., Tarpy, D.R., Mullin, C.A., Frazier, J.L., Frazier, M., vanEngelsdorp, D., 2016. In-hive pesticide exposure: assessing risks to migratory honey bees from in-hive pesticide contamination in the eastern United States. *Sci. Rep.* 6, 33207.
- Traynor, K.S., Tosi, S., Rennie, K., Steinhauer, N., Forsgren, E., Rose, R., et al., 2021. Pesticides in honey bee colonies: establishing a baseline for real world exposure over seven years in the USA. *Environ. Pollut.* 279, 116566.
- Trostanetsky, A., Kostyukovsky, M., 2008. Note: Transovarial activity of the chitin synthesis inhibitor novaluron on egg hatch and subsequent development of larvae of *Tribolium castaneum*. *Phytoparasitica* 36, 38–41.
- Ucar, T., Hall, F.R., 2001. Windbreaks as a pesticide drift mitigation strategy: a review. *Pest Manag. Sci.* 57, 663–675.
- Ullah, A., Tlak Gajger, I., Majoros, A., Dar, S.A., Khan, S., Kalimullah, et al., 2021. Viral impacts on honey bee populations: a review. *Saudi. Aust. J. Biol. Sci.* 28, 523–530.
- Umsza-Guez, M.A., Silva-Beltran, N.P., Machado, B.A.S., Balderrama-Carmona, A.P., 2021. Herbicide determination in Brazilian propolis using high pressure liquid chromatography. *Int. J. Environ. Health Res.* 31, 507–517.
- USDA, 2022a. Annual Strategic Pollinator Priorities Report.
- USDA, 2022b. California Almond Acreage Report.
- van Klink, R., August, T., Bas, Y., Bodesheim, P., Bonn, A., Fossoy, F., et al., 2022. Emerging technologies revolutionise insect ecology and monitoring. *Trends Ecol. Evol.* 37, 872–885.
- Vannette, R.L., Mohamed, A., Johnson, B.R., 2015. Forager bees (*Apis mellifera*) highly express immune and detoxification genes in tissues associated with nectar processing. *Sci. Rep.* 5, 16224.
- Vazquez, D.E., Villegas Martinez, L.E., Medina, V., Latorre-Estivalis, J.M., Zavala, J.A., Farina, W.M., 2023. Glyphosate affects larval gut microbiota and metamorphosis of honey bees with differences between rearing procedures. *Environ. Pollut.* 334, 122200.
- Visscher, P.K., Vetter, R.S., Robinson, G.E., 1995. Alarm pheromone perception in honeybees is decreased by smoke (Hymenoptera, Apidae). *J. Insect Behav.* 8, 11–18.
- von Planta, A., 1888. Ueber den Futtersaft der Biene. *bchm* 12, 327–354.
- Wade, A., Lin, C.H., Kurkul, C., Regan, E.R., Johnson, R.M., 2019. Combined toxicity of insecticides and fungicides applied to California almond orchards to honey bee larvae and adults. *Insects* 10.
- Walker, E.K., Brock, G.N., Arvidson, R.S., Johnson, R.M., 2022. Acute toxicity of fungicide-insecticide-adjunct combinations applied to almonds during bloom on adult honey bees. *Environ. Toxicol. Chem.* 41, 1042–1053.
- Waller, G.D., Barker, R.J., 1979. Effects of dimethoate on honey bee colonies. *J. Econ. Entomol.* 72, 549–551.
- Walsh, E.M., Sweet, S., Knap, A., Ing, N., Rangel, J., 2020. Queen honey bee (*Apis mellifera*) pheromone and reproductive behavior are affected by pesticide exposure during development. *Behav. Ecol. Sociobiol.* 74.
- Wang, J., Kliks, M.M., Jun, S., Li, Q.X., 2010. Residues of polybrominated diphenyl ethers in honeys from different geographic regions. *J. Agric. Food Chem.* 58, 3495–3501.

- Wang, X., Wang, Z., Di, S., Xue, X., Jin, Y., Qi, P., et al., 2020. Determination of 14 lipophilic pesticide residues in raw propolis by selective sample preparation and gas chromatography–tandem mass spectrometry. *Food Anal. Methods* 13, 1726–1735.
- Wang, K., Li, J., Zhao, L., Mu, X., Wang, C., Wang, M., et al., 2021a. Gut microbiota protects honey bees (*Apis mellifera* L.) against polystyrene microplastics exposure risks. *J. Hazard. Mater.* 402, 123828.
- Wang, Y., Zhu, Y.C., Li, W., Yao, J., Reddy, G.V.P., Lv, L., 2021b. Binary and ternary toxicological interactions of clothianidin and eight commonly used pesticides on honey bees (*Apis mellifera*). *Ecotoxicol. Environ. Saf.* 223, 112563.
- Wang, F., Wang, Y., Li, Y., Zhang, S., Shi, P., Li-Byarlay, H., Luo, S., 2022. Pesticide residues in beebread and honey in *Apis cerana cerana* and their hazards to honey bees and human. *Ecotoxicol. Environ. Saf.* 238, 113574.
- Webster, L., Roose, L., Bersuder, P., Kotterman, M., Haarich, M., Vorkamp, K., 2013. Determination of polychlorinated biphenyls (PCBs) in sediment and biota. In: *ICES Techniques in Marine Environmental Science (TIMES)*.
- Weisbrod, A.V., Burkhard, L.P., Arnot, J., Mekenyan, O., Howard, P.H., Russom, C., et al., 2007. Workgroup report: review of fish bioaccumulation databases used to identify persistent, bioaccumulative, toxic substances. *Environ. Health Perspect.* 115, 255–261.
- Wernecke, A., Eckert, J.H., Forster, R., Kurlemann, N., Odemer, R., 2021. Inert agricultural spray adjuvants may increase the adverse effects of selected insecticides on honey bees (*Apis mellifera* L.) under laboratory conditions. *Journal of Plant Diseases and Protection* 129, 93–105.
- Williamson, S.M., Wright, G.A., 2013. Exposure to multiple cholinergic pesticides impairs olfactory learning and memory in honeybees. *J. Exp. Biol.* 216, 1799–1807.
- Williamson, S.M., Baker, D.D., Wright, G.A., 2013. Acute exposure to a sublethal dose of imidacloprid and coumaphos enhances olfactory learning and memory in the honeybee *Apis mellifera*. *Invertebr. Neurosci.* 13, 63–70.
- Williamson, S.M., Willis, S.J., Wright, G.A., 2014. Exposure to neonicotinoids influences the motor function of adult worker honeybees. *Ecotoxicology* 23, 1409–1418.
- Wilmart, O., Legreve, A., Scippo, M.L., Reybroeck, W., Urbain, B., de Graaf, D.C., et al., 2021. Honey bee exposure scenarios to selected residues through contaminated beeswax. *Sci. Total Environ.* 772, 145533.
- Wu, J.Y., Anelli, C.M., Sheppard, W.S., 2011. Sub-lethal effects of pesticide residues in brood comb on worker honey bee (*Apis mellifera*) development and longevity. *PLoS One* 6, e14720.
- Wu, C., Liu, X., He, M., Dong, F., Wu, X., Xu, J., Zheng, Y., 2021. Quantitative determination of pyriproxyfen and its metabolite residues in bee products of China using a modified QuEChERS approach with UPLC-MS/MS. *Ecotoxicol. Environ. Saf.* 220, 112388.
- Wu, X., Liao, C., He, X., Zhang, L., Yan, W., Zeng, Z., 2022. Sublethal fluvalinate negatively affect the development and flight capacity of honeybee (*Apis mellifera* L.) workers. *Environ. Res.* 203, 111836.
- Wu, W.Y., Liao, L.H., Lin, C.H., Johnson, R.M., Berenbaum, M.R., 2023. Effects of pesticide-adjuvant combinations used in almond orchards on olfactory responses to social signals in honey bees (*Apis mellifera*). *Sci. Rep.* 13, 15577.
- Wueppenhorst, K., Eckert, J.H., Steinert, M., Erler, S., 2022. What about honey bee jelly? Pesticide residues in larval food jelly of the Western honey bee *Apis mellifera*. *Sci. Total Environ.* 850, 158095.
- Xavier, V.M., Message, D., Picanco, M.C., Chediak, M., Santana, P.A., Ramos, R.S., Martins, J.C., 2015. Acute toxicity and sublethal effects of botanical insecticides to honey bees. *J. Insect Sci.* 15.
- Xu, B., Qiu, W., Du, J., Wan, Z., Zhou, J.L., Chen, H., et al., 2022. Translocation, bioaccumulation, and distribution of perfluoroalkyl and polyfluoroalkyl substances (PFASs) in plants. *iScience* 25, 104061.
- Zaric, N.M., Deljanin, I., Iljjevic, K., Stanisavljevic, L., Ristic, M., Grzetic, I., 2018. Assessment of spatial and temporal variations in trace element concentrations using honeybees (*Apis mellifera*) as bioindicators. *PeerJ* 6, e5197.
- Zhu, Y.C., Adamczyk, J., Rinderer, T., Yao, J., Danka, R., Luttrell, R., Gore, J., 2015. Spray toxicity and risk potential of 42 commonly used formulations of row crop pesticides to adult honey bees (Hymenoptera: Apidae). *J. Econ. Entomol.* 108, 2640–2647.
- Zhu, Y.C., Yao, J., Adamczyk, J., Luttrell, R., 2017a. Feeding toxicity and impact of imidacloprid formulation and mixtures with six representative pesticides at residue concentrations on honey bee physiology (*Apis mellifera*). *PLoS One* 12, e0178421.
- Zhu, Y.C., Yao, J., Adamczyk, J., Luttrell, R., 2017b. Synergistic toxicity and physiological impact of imidacloprid alone and binary mixtures with seven representative pesticides on honey bee (*Apis mellifera*). *PLoS One* 12, e0176837.
- Zoltowska, K., Frączek, R., Lipiński, Z., 2011. Hydrolases of developing worker brood and newly emerged worker of *Apis mellifera carnica*. *Journal of Apicultural Science* 55, 27–36.