

UCLA

UCLA Previously Published Works

Title

The Effects of Collaborative Care Training on Case Managers' Perceived Depression-Related Services Delivery

Permalink

<https://escholarship.org/uc/item/3xp74139>

Journal

Psychiatric Services, 68(2)

ISSN

1075-2730

Authors

Landry, Craig M
Jackson, Aurora P
Tang, Lingqi
et al.

Publication Date

2017-02-01

DOI

10.1176/appi.ps.201500550

Peer reviewed



HHS Public Access

Author manuscript

Psychiatr Serv. Author manuscript; available in PMC 2019 January 06.

Published in final edited form as:

Psychiatr Serv. 2017 February 01; 68(2): 123–130. doi:10.1176/appi.ps.201500550.

The Effects of Collaborative Care Training on Case Managers' Perceived Depression Related Services Delivery

Craig Martin Landry,

UCLA Semel Institute - Department of Psychiatry, 10920 Wilshire Blvd., Suite 300 Los Angeles, CA 90024-6505, Los Angeles, California 90024-6505

Aurora P. Jackson,

UCLA Luskin School of Public Affairs - Social Welfare, Los Angeles, California

Lingqi Tang,

UCLA – Psychiaatry, Los Angeles, California

Jeanne Miranda,

University of California, Los Angeles - Psychiatry, Center for Health Services and Society, Los Angeles, California

Bowen Chung,

UCLA Robert Wood Johnson Clinical Scholars Program, David Geffen School of Medicine - Department of Psychiatry, 911 Broxton Avenue 3rd Floor, Los Angeles, California 90024

Felica Jones,

Health African American Families II (HAAF), Los Angeles, California

Michael K. Ong, and

Shields for Families, Los Angeles, California

Kenneth Wells

UCLA - Psychiatry and Biobehavioral Sciences, 10920 Wilshire Blvd. Suite # 300, Los Angeles, California 90024

Abstract

Objective—This study examined the effects of a Community Engagement and Planning (CEP) depression quality improvement intervention across social service agencies relative to Resources for Services (RS) technical assistance, on training participation and service delivery by primarily non-licensed, ethnically diverse case managers in two low-income communities in Los Angeles. Whether CEP is more effective than (RS) at engaging mostly non-licensed case managers in depression quality improvement training and improving depression competencies or time spent providing community services is unknown.

Methods—Cluster-randomized trial with program-level assignment to CEP or RS for training providers in quality improvement training for depression. Within 84 health and community-based

Disclosures: None of the authors of this paper have any conflicts of interest

Previous presentations: There have been no previous presentations of this material.

programs eligible for the provider outcomes study, staff with patient contact were invited to participate in trainings and baseline and one-year follow-up surveys; 117 case managers (59 RS, 58 CEP) completed follow-up from 52 programs. Outcomes are time spent in community settings, use of depression case management and problem solving practices (primary) and depression knowledge and attitudes and work environment support (secondary).

Results—CEP relative to RS increased case managers' participation in depression trainings, time spent in community settings and use of problem solving therapeutic approaches (each $p < .05$).

Conclusions—CEP relative to RS for depression quality improvement increased primarily non-licensed, ethnically diverse case managers' training participation, time in community settings and use of problem solving skills, offering a model for including case managers in community-wide depression improvement efforts.

Introduction

Depression is common across race/ethnic and socioeconomic groups (1), affects health, cost and productivity outcomes (2, 3) and is a leading cause of disability (4-6). African Americans relative to whites, report lower lifetime prevalence of major depression but have more severe symptoms (7, 8), use fewer services and terminate treatment early (7, 9, 10). African Americans and Latinos are less likely than whites to receive evidence-based depression care and have worse outcomes (8,9,12-14). While evidenced-based practices for depression exist, their implementation in community settings has been limited (11, 12). Quality improvement (QI) programs for depression based on the collaborative care model improve depressive symptoms, quality of life, and social outcomes and reduce ethnic/racial outcome disparities (13). However, these programs are often not implemented in under-resourced communities, where there is low availability of specialty care and historical distrust of services (14, 15). Depressed individuals in such settings may seek support from alternative sectors not included in healthcare QI efforts. In these settings, non-licensed providers may be primary supports for depressed individuals. Although the Affordable Care Act (ACA) provides incentives for health homes to utilize case managers (16, 17), such roles in collaborative care are typically filled by licensed providers (18, 19). Little is known about the effects of implementing depression QI on implementation outcomes for non-licensed providers, such as the use of therapeutic practices (20).

This study addresses this gap by analyzing data from Community Partners in Care (CPIC), a cluster-randomized trial of two depression QI implementation conditions, community engagement and planning (CEP) to support networks of health and community-based agencies versus technical assistance to individual agencies (RS). Following community feedback, health and community-based programs were included as sites for depression QI programs (9), with non-licensed and licensed case managers, given community awareness of mental health professional shortages in under-resourced communities (21, 22) and to increase provider diversity and community trust. Chung et al., reported that CEP relative to RS increased staff participation in depression trainings among estimated eligible staff (23). However, change in practice outcomes for predominantly non-licensed case managers was not reported. We sought to replicate findings of increased training participation under CEP over RS among case-managers participating in a provider outcomes sub-study and to

examine intervention effects on case-manager use of depression case management, therapeutic problem solving and on time spent delivering community services as primary outcomes; and depression knowledge, attitudes and perceived work-environment support, as secondary outcomes. We hypothesized that case-managers in programs assigned to CEP relative to RS would participate more in trainings, spend more time providing community services, and report higher use of depression case management and therapeutic services. We thought that there might be greater effects of CEP on improving depression knowledge and attitudes and work environment support, owing to a greater focus in CEP on network development.

Methods

Design

Data are from the provider sub-study of Community Partners in Care, a group-level randomized trial implemented using community-partnered participatory research (CPPR) (24, 25), that supports academic and community partners as equal decision makers in research design and implementation (26). As described elsewhere (23, 27, 28), the study was conducted in South Los Angeles of roughly 1.5 million population and Hollywood-Metro Los Angeles of roughly 500K population, having high rates of unemployment, homelessness and lack of insurance (29).

The interventions represent two ways of implementing a depression QI training based on the collaborative care model (30-33), adapted for licensed and non-licensed providers (34) from health and community services programs. Toolkits for case managers included guidelines for case management, depression screening, care coordination, outreach strategies, problem-solving therapy, behavioral management, and activation skills. Materials were introduced in kick-off conferences prior to enrollment and randomization and available to both conditions in hardcopy, flash drives and a website (<http://www.communitypartnersincare.org/community-engagement/cep/>). Interventions encouraged but did not require use of these resources by eligible providers.

Community Engagement and Planning (CEP)

CEP invited participating program administrators to attend 2-hour, bi-monthly Council meetings for 4 months, to adapt QI toolkits to their community and collaborate as a network, following a workbook based on principles of CPPR. Councils were asked to develop and implement a written plan for toolkit adaptation, training and monitoring, supported by \$15,000 from the study. Final plans featured conferences, follow-up with programs, telephone and webinar supervision for CBT and case-management and innovations such as provider self-care and depression book clubs.

CEP case-manager trainings were co-led by academic and community leaders. Eligible providers were invited to half or all-day conferences with follow-up and make-up sessions at individual programs. The 6-hour case-manager trainings reviewed the study's purpose, defined terms and presented resources. Participants were taught client engagement and outreach, depression screening with the PHQ-9, behavioral activation, making referrals and

problem solving, reinforced by role-play. Participation in other components (e.g., medication management) was encouraged. Case manager telephone supervision (2-3 sessions per community) was offered.

Resources for Services (RS)

The RS intervention offered outreach and technical assistance to individual programs using a “train-the-trainer” approach. A team of psychiatrists, nurse care manager, CBT trainer, QI expert, support staff and community-engagement specialist offered twelve 90-120 minute webinars on team management, CBT, care management and patient education plus visits to primary care sites on medication management. Case-managers received four 1-2 hour webinars per community,. Topics included client engagement, depressive symptom recognition, screenings using the PHQ-9, making referrals and use of problem-solving strategies.

Participants

Using County lists and community nominations, 97 of 149 agencies were confirmed as potentially eligible based on financial stability, serving adults or parents of children and permitting oversampling of community-prioritized subgroups (i.e., homeless, seniors, African Americans, and substance abuse programs). We recruited 50 agencies (52%) and enumerated 177 programs offering mental health; primary-care/public health; substance abuse; social services (homeless-serving, prisoner re-entry, family preservation); and “community-trusted” programs, including faith-based, park community centers, hair salons, and exercise clubs. Of these, 11 refused evaluation and of the remainder, 122 were confirmed as potentially eligible (serving ≥ 15 clients per week, having ≥ 1 staff, not focused primarily on psychotic disorders or home services) and randomized to CEP or RS, stratified by community and sector (35). At site visits to confirm participation, 8 refused and 114 were confirmed as eligible of which 95 (83%) enrolled (Appendix 1),. Program administrators were informed of intervention status by letter prior to screening. Participating and nonparticipating programs were from comparable neighborhoods by zip-code-level data (36).

Provider outcomes sub-study

In eligible programs having more than one staff (N=84), providers with direct patient contact including volunteers were invited to participate in the provider sub-study through agency presentations and recruitment packets distributed by administrators with telephone follow-up and study site visits. The goal was to achieve a baseline sample of 300 providers and a one-year follow-up sample of 200. From a pool of 370 consented providers at baseline, 326 (88%) from 80 of 84 eligible programs completed baseline surveys. New providers were permitted to enter after baseline. At follow-up, 297 were eligible (at the agency and alive) and 237 (80%) participated from 82 of 84 eligible programs. After verbal and written consent, the sample for this sub-study included 117 primarily non-licensed case-managers (58 CEP, 59 RS) from 52 programs. Internal Review Boards of RAND, UCLA, and participating agencies approved this study, registered as a clinical trial post baseline enrollment (clinical trials.gov NCT01699789).

Measures

The main independent variable is program intervention status (CEP or RS). Service sector (health or community) was assigned for each program. Other measures are from provider baseline and follow-up surveys, and for training participation, sign-in logs. Standardized alpha coefficients were calculated for scales with ≥ 2 items.

Sample Characteristics

Case-manager characteristics include age in years, sex, education, race/ethnicity and license status.

Primary Outcomes

Training participation: Attendance at trainings was assessed by sign-in logs or self-reported at follow-up, including study-provided and in-house trainings.

Depression Care Techniques

We report the mean of 9 items (response, 5 point scale from 1=never to 5=always $\alpha = .929$) for how often respondents performed the following tasks for people with symptoms of depression in the past 6 months: Encourage positive thinking, discuss costs of alternative mental health treatments, encourage pleasurable activities, discuss ways to improve social skills, determine depression treatment preferences, Recommend ways to take care of one's self, reframe or clarify the individual's problems, discuss benefits of treatments, help the individual feel better about his/her life. Case-managers not providing services for depressed clients were set to a response of "never".

Depression Case Management

We report the mean of 5 items (5-point response scale from 1=never to 5=always $\alpha = .917$) of how often respondents performed the following tasks if providing services for depressed clients in the past 6 months: Explain what depression is, ask the individual what he or she thinks depression is, ask about prior treatment, make a referral ask about barriers to depression care. Case-managers not providing services for depressed clients were set to a response of "never".

Community Services Provision

Respondents were asked to indicate on a six-point ranked scale how many hours in a typical week they spent providing services to individuals in community settings (0=0 hours, 1=1-10, 2=11-20, 3=21-30, 4=31-40, 5=more than 40 hours),

Secondary outcomes

Perceived depression knowledge was assessed by 3 items adapted from Partners in Care (37), (5-point scale from 1=strongly agree to 5=strongly disagree $\alpha = .626$) with the following statements: Depression is a medical condition, depression runs in families, depression can cause physical changes like aches and pains.

Perceived depression skill was assessed by the mean of 7 items on a 4-point scale from 1=not at all skilled to 4=very skilled $\alpha = .890$) for: Case finding, depression screening with a standardized instrument, educating individuals or families about depression, depression counseling, referring to mental health specialty care; providing social support for depression (support groups); Engaging in community outreach for depression.

Personal depression stigma was assessed by the mean of 3 items adapted from Link's Devaluation and Discrimination scale (38, 39): I have no patience with a person who is always feeling 'blue' or depressed; I would be embarrassed if people thought I was depressed; and most people think less of a person who has been depressed. Responses were on a 5-point scale from 1=strongly agree to 5=strongly disagree $\alpha = .579$.

Perceived system barriers were assessed by a count of 4 dichotomized items, with response of limited a great deal versus limited somewhat or not limited, on the extent to which optimal depression care services were limited in the past 6 months by: treatment difficult to obtain, MH professionals not available, poor reimbursement, limited insurance or other benefits, other barriers.

Analysis Plan

We conducted intent-to-treat comparative effectiveness analyses with the provider sub-study case-manager sample having one-year follow-up data. We used logistic regression models for dichotomous and multiple linear regression models for continuously scaled variables, adjusted for baseline status of the dependent variable, sector (formal vs. informal) and provider type (licensed vs. non-licensed). There was no baseline status for training participation. We compared case manager baseline characteristics by intervention status.

We used an extended hot-deck technique to impute missing values for nonresponse, using five imputed datasets for baseline and follow-ups, and multiple imputation inference for all analyses (37, 38). To control for potential nonresponse bias, we created nonresponse weights on one-year follow-up respondents for attrition to present eligible sample. The final weights included predictors that were significant ($p < 0.10$) for either CEP or RS groups: service sector (formal vs. informal), education, and baseline perception of depression attitude and skill. The reciprocal of the predicted response probability was used as the attrition weight for each participant. Five versions of the weight were created corresponding to imputed datasets (39, 40). Significance of comparisons by intervention status is based on regression coefficients. Results are presented for linear regression as between-group differences and for logistic regression as odds ratios (OR) with 95% confidence intervals (CI). We illustrate average results for an intervention group adjusted for all covariates using standardized predictions generated from the fitted regression model (40). To account for client clustering within programs, the variance estimation was based on the Taylor Series Linearization Method (41) All analyses were conducted using SUDAAN Version 11.0.0 (Software for the Statistical Analysis of Correlated Data at <http://www.rti.org/sudaan/>) with the design specification of sampling with-replacement in the first stage of sample selection (programs), accounting for attrition weights.

For primary outcomes, to account for multiple comparisons, we calculated the False Discovery Rate (FDR)(42) comparing observed significance findings with expected order statistics from a uniform distribution (42).

Results

Demographic and Descriptive Characteristics

As shown in Table 1, 55 (47%) of case managers self-identified as Hispanic; 52 (45%) African American; 7 (6%) non-Hispanic white; and 3 (2%) Asian Pacific Islander. The mean age was 43.6 years and 71% were female. Over 70% were non-licensed, with over half (62%) working in health sectors. Except for personal stigma of depression (CEP greater stigma than RS, $p < .05$), there were no significant baseline differences by intervention status.

Outcomes

As shown in Table 2, 27% of case managers in RS and 74% in CEP participated in CPIC-sponsored trainings (OR=7.78, 95% CI=2.90–20.89 $p < .001$). CEP case-managers reported more of depression-care techniques relative to RS (mean score of 3.1 versus 2.8; difference=.32, 95% CI=.03–.61, $p < .05$). CEP case-managers relative to RS reported greater community services provision (mean score of 1.2. versus .7; difference .51, 95% CI =.13–.89 $p < .05$. Findings remained significant applying the FDR for multiple comparisons. Intervention differences in depression case management were not statistically significant.

As shown in Table 3, there were no significant intervention differences at the .05 in perceived depression knowledge, skill, depression stigma, or system barriers to services, as secondary outcomes.

Discussion

We examined effects of two interventions to implement depression QI across under-resourced communities, CEP for supporting networks and RS for technical assistance to individual agencies, on case-manager outcomes. Case managers were predominantly non-licensed and ethnic minority, a resource for expanding workforce diversity. We found that case-managers in CEP programs were nearly 3 times more likely than those in RS to participate in trainings, confirming findings for eligible providers (23). Compared to RS, CEP case managers reported increased hours delivering community services and greater use of therapeutic problem-solving skills for depression (43). However, we found no intervention differences in case-management tasks, which may be standard competencies of case managers or easier to influence with technical assistance. The stronger CEP effect on increasing use of therapeutic strategies may be due to the novel nature of those skills for case managers, plus role modeling and supervision in CEP. Contrary to expectations, we did not find significant intervention effects on attitudes or perceived skill. In CEP relative to RS greater knowledge and system support were of borderline significance, suggesting areas for future research.

Others have noted that information given prior to training can improve self-efficacy and motivation (44), which can increase participation (45). In CPIC, providers in both

interventions could participate in conferences prior to randomization, for the same priming. From a social-cognitive theory (SCT) perspective, RS supported knowledge exchange while CEP supported in-person guidance, skills-building (self-efficacy), modeling (observational learning), peer support, networking and collaboration. Under self-efficacy theory (46) individuals believing in their ability to perform successfully are more likely to initiate and maintain behavior change; while those with less self-efficacy may avoid the task (47). Torrey et al (48) found that clinicians are motivated to change practice if the change is perceived as clinically helpful and reinforced through observation, supervision and feedback—features of CEP. The CEP focus on administrator involvement and community feedback may build collective efficacy (49, 50) to motivate an expansion of case manager roles.

Future research should focus on ways to further build and sustain capacity for depression services, as required for medical homes (16, 17), given that CEP supported a predominantly ethnic minority case-manager workforce in under-resourced communities. While we do not comment on the link of provider change to client outcomes, we previously reported that CEP relative to RS improved client mental health-related quality of life and reduced behavioral health hospitalizations over 6-12 months, with significance sensitive to methods (23, 51). In addition, it will be important to determine if intervention differences in provider outcomes result from differences in training participation.

Several limitations should be acknowledged. Data are from provider self-report in two communities, meriting replication with service records. Many of the survey questions were worded in one direction. The higher the score the more appropriate the answer. The case-manager sample was modest for observing small effects that are typical for public health implementation strategies. There was attrition at follow-up primarily due to turnover, which may be expected in under-resourced communities, accounted for with weights. Knowledge and attitude measures may require further development to capture intervention framework (52, 53) such as the Quality Enhancement Research initiative (Queri) framework (54) to assess intervention adoption and sustainability.

This study suggests that depression QI programs can feasibly include predominantly non-licensed, ethnically-diverse case managers; and that CEP relative to RS increases participation in trainings, time spent by case managers in delivering community services and use of therapeutically-oriented problem solving skills. As one partner noted: “this offers hope for under-resourced communities.” It may be important to identify policy mechanisms to sustain such efforts through certification as health workers, formal partnerships, and funding by HRSA, SAMSHA, CMS, health plans or foundations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the RAND Corporation, UCLA Semel Institute and the Los Angeles County Departments of Mental Health, Public Health and Health Services for institutional support. We thank the 95 participating healthcare and community-based agencies, the CPIC Council and members of the Association of Clinical and Translational

Science Team Science Award for CPIC (2014) and Campus-Community Partnerships for Health 2015 Annual Award (<https://ccph.memberclicks.net/annual-award-winner-2015>).

Grant Support: This study was funded by Grants 64244 from the Robert Wood Johnson Foundation, R01MD007721 from the National Institute on Minority Health and Health Disparities and the parent CPIC study by R01MH078853, P30MH082760, and P30MH068639 from the National Institute of Mental Health; CMCH-12-97088 from the California Community Foundation, G08LM011058 from the National Library of Medicine, and UL1TR000124 from the NIH/National Center For Advancing Translational Science UCLA CTSI.

References

1. Bruce ML, Smith W, Miranda J, et al. Community-Based Interventions. *Mental Health Services Research*. 4:205–13.2002; [PubMed: 12558005]
2. Lubner P, Hollenberg J, Williams-Russo P, et al. Diagnosis, Treatment, Comorbidity and Resource Utilization of Depressed Patients in a General Medical Practice. *International Journal Psychiatry in Medicine*. 30:1–13.2000;
3. Greenberg PE. The Economic Burden of Depression in the United States: How Did It Change Between 1990 and 2000? *Journal of Clinical Psychiatry*. 64:1465–75.2003; [PubMed: 14728109]
4. Organization, WH. Depression in Fact Sheet Number 369. WH, O, editor World Health Organization; 2009.
5. Mark, TL, Shern, DL, Bagalman, JE. , et al. Ranking America's Mental Health: An Analysis of Depression Across the States. Alexandria, VA: Mental Health America; 2007.
6. Kessler CR, Chiu WTC, Demler O, et al. Prevalence, severity, and comorbidity of Twelve-month DSM-IV disorders in the National Comorbidity Survey Replication (NCS-R). *Archives of General Psychiatry*. 6:617–27.2005;
7. Williams DR, Neighbors H, Nesse R, et al. Prevalence and distribution of major depressive disorder in African Americans, Caribbean Blacks, and Non-Hispanic Whites. *Archives of General Psychiatry*. 2007
8. Riolo SA, Nguyen TANH, Greden JF, et al. Prevalence of Depression by Race/Ethnicity: Findings From the National Health and Nutrition Examination Survey III. *American Journal of Public Health*. 95:998–1000.2005; [PubMed: 15914823]
9. Wells K. Building Wellness Pilot Application. 2006
10. Wells K. The design of Partners in Care: evaluating the cost-effectiveness of improving care for depression in primary care. *Social Psychiatry and Psychiatric Epidemiology*. 34:20–9.1999; [PubMed: 10073117]
11. Proctor EK, Landsverk J, Aarons G, et al. Implementation Research in Mental Health Services: an Emerging Science with Conceptual, Methodological, and Training Challenges. *Admin Policy Mental Health*. 36:2009;
12. Glasgow RE, Lichtenstein E, Marcus AC. Why Don't We See More Translation of Health Promotion Research to Practice? Rethinking the Efficacy to Effectiveness. *Transition American Journal of Public Health*. 93:1261–7.2003; [PubMed: 12893608]
13. Wells K, Sherbourne C, Miranda J, et al. The cumulative effects of quality improvement for depression on outcome disparities over 9 years: results from a randomized, controlled group-level trial. *Med Care*. 45:1052–9.2007; [PubMed: 18049345]
14. Suite D, Labril R, Primm A, et al. Beyond Misdiagnosis, Misunderstanding and Mistrust: Relevance of the Historical Perspective in the Medical and Mental Health Treatment of People of Color. *Journal of the National Medical Association*. 99:2007;
15. Miranda J, McGuire T, Williams D, et al. Mental Health in the Context of Health Disparities. *American Journal of Psychiatry*. 165:2008;
16. DeSilva M, Samele C, Saxena S, et al. Policy Actions To Achieve Integrated Community-Based Mental Health Services. *Health Affairs*. 33:1595–602.2014; [PubMed: 25201664]
17. Katzen, A, Morgan, M. Affordable Care Act Opportunities for Community Health Workers. Boston: Center for Health Law & Policy Innovation Harvard Law School; 2014.
18. Gilbody S, Bower P, Fletcher J, et al. Collaborative Care for Depression. *Arch Internal Medicine*. 166:2314–21.2006;

19. Taylor EF, Machta RM, Meyers DS, et al. Enhancing the Primary Care Team to Provide Redesigned Care: The Roles of Practice Facilitators and Care Managers. *Annals of Fam Med.* 11:80–3.2013;
20. Fixsen, DL, Naoom, SF, Blase, KA. , et al. *Implementation Research: A synthesis of the literature.* Tampa Florida: University of South Florida, Florida Mental Health Institute; 2005.
21. Rivelli, SK, Shirley, KG. *Prevalence of Psychiatric Symptom/Syndromes in Medical Settings in Integrated Care in Psychiatry, 2014, Redefining the Role of Mental Health Professional in the Medical Setting.* New York: Springer Science + Business Media; 2014.
22. Boer, P, Wiersma, D, Russo, S. , et al. *Cochrane Database of Systematic Reviews 2005.* John Wiley & Sons Ltd; 2005. Paraprofessionals for anxiety and depressive disorders.
23. Chung B, Ngo VK, Ong MK, et al. Participations in Training for Depression Care Quality Improvement: A Randomized Trial of Community Engagement or Technical Support. *Psychiatric Services.* 66:831–9.2015; [PubMed: 25930037]
24. Jones L, Wells K. Strategies for Academic and Clinician Engagement in Community-Participatory Partnered Research. *JAMA.* 297:24–31.2007;
25. Wells K, Jones L. Commentary: “Research” in Community-Partnered, Participatory Research. *JAMA.* 302:320–1.2009; [PubMed: 19602693]
26. Wallerstein N, Duran B. Using Community- Based Participatory Research to Address Health Disparities. *Health Promotion Practice.* 7:312.2006; [PubMed: 16760238]
27. Chung B, Ong M, Ettner SL, et al. 12-Month Outcomes of Community Engagement Versus Technical Assistance to Implement Depression Collaborative Care: A Partnered, Cluster, Randomized, Comparative-Effectiveness Trial. *Annals of Internal Medicine.* 1612014;
28. Wells KB, Jones L, Chung B, et al. Community-Partnered Cluster-Randomized Comparative Effectiveness Trial of Community Engagement and Planning or Resources for Services to Address Depression Disparities. *Journal of General Internal Medicine.* 28:1268–78.2013; [PubMed: 23649787]
29. US Census Bureau. Los Angeles County DoHS. LA County Department of Health Services Key Health Indicators. Los Angeles County C; 2009.
30. Wells K, Sherbourne C, Schoenbaum M, et al. Impact of Disseminating Quality Improvement Programs for Depression in Managed Primary Care: A Randomized Controlled Trial. *Journal of American Medical Association.* 283:212–20.2000;
31. Miranda J, Naihua D, Sherbourne C, et al. Improving Care for Minorities: Can Quality Improvement Interventions Improve Care and Outcomes For Depressed Minorities? Results of a Randomized, Controlled Trial. *Health Services Research.* 38:613–30.2003; [PubMed: 12785564]
32. Unutzer J, Katon W, Callahan CM, et al. Collaborative Care Management of Late-Life Depression in the Primary Care Setting A Randomized Controlled Trial. *Journal of American Medical Association.* 2892002;
33. Springgate B, Allen C, Jones C, et al. Rapid Community Participatory Assessment of Health Care in Post Storm New Orleans. *American Journal of Preventative Medicine.* 372009;
34. Wennerstrom A, Vannoy S, Allen C, et al. Community Based Participatory Development of a Community Health Worker Mental Health Outreach Role to Extend Collaborative Care in Post Katrina New Orleans. *Ethnicity and Disease.* 212011;
35. Murray, DM. *Design and Analysis of Group-Randomized Trials.* New York: Oxford University Press; 1998.
36. Wells K, Jones L, Chung B, et al. Community-Partnered Cluster-Randomized Comparative Effectiveness Trial of Community Engagement and Planning or Resources for Services to Address Depression Disparities. *Journal of General Internal Medicine.* 28:1268–78.2013; [PubMed: 23649787]
37. Wells KB. The design of Partners in Care: evaluating the cost-effectiveness of improving care for depression in primary care. *Soc Psychiatry Psychiatry Epidemiol.* 34:20–9.1999;
38. Link BG, Struening ELRM, Phelan PJC, et al. On Stigma and Its Consequences: Evidence from a Longitudinal Study of Men with Dual Diagnoses of Mental Illness and Substance Abuse. *Journal of Health and Social Behavior.* 38:177–90.1997; [PubMed: 9212538]

39. Link BG, Struening EL, Rahav M, et al. On Stigma and Its Consequences: Evidence from a Longitudinal Study of Men with Dual Diagnoses of Mental Illness and Substance Abuse. *Journal of Health and Social Behavior*. 38:177–90.1997; [PubMed: 9212538]
40. Korn, E, Graubard, B. *Analysis of Health Surveys*. Hoboken New Jersey: John Wiley & sons-Intersciences; 1999.
41. Binder D. On the Variances of Asymptotically Normal Estimators from Complex Surveys. *International Statistical Review / Revue Internationale de Statistique*. 51:279–92.1983;
42. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society*. 57:289–300.1995;
43. Nezu AM. Problem Solving and Behavior Therapy Revisited. *Behavior Therapy*. 35:1–33.2004;
44. Wei-Tao T. Effects of Training Framing, General Self-efficacy and Training Motivation on Trainees' Training Effectiveness. *Personnel Review*. 35:51–65.2004;
45. Carlson DS, Bozeman DP, Kacmar K, Michele, et al. Training Motivation In Organizations: An Analysis of Individual-Level Antecedents. *Journal of Managerial Issues*. 12:271–87.2000;
46. Bandura, A. *Self Efficacy the exercise of Control*. NY,NY: W.H. Freeman; 1997.
47. Parle M, Maguire P, Heaven C. The Development of A Training Model to Improve Health Professionals Skills, Self-Efficacy and Outcome Expectancies When Communicating with Cancer Patients. *Social Science Medicine*. 44:231–40.1997; [PubMed: 9015875]
48. Torrey WC, Drake RE, Dixson L, et al. Implementing Evidenced-Based Practices for Persons With Severe Mental Illnesses. *Psychiatric Services*. 52:45–50.2000; 1.
49. Ballesteros-Fernandez R, Nicolas-Diaz J, Bandura A. Determinants and Structural Relation of Personal Efficacy to Collective Efficacy. *Applied Psychology: An International Review* 51. :107–25.2002
50. Bandura A. Exercise of Human Agency through Collective Efficacy. *Current Directions in Psychological Sciences*. 9:75–8.2000;
51. Wells, K, Tang, L. Report on the survey method for the CPIC Provider on-line Survey. Landry, C, editor2013.
52. Ajzen I. The Theory of Planned Behavior. *Organization Behavior and Human Decision Processes* 50. :179–211.1991
53. Montano, DE, Kasprzyk, D. Theory of Reasoned Action, Theory of Planned Behavior, and the Intergrated Behavioral Model. In: Glanz, K, Rimer, BK, Viswanath, K, editors *Health Behavior and Health Education: Theory, Research, and Practice*. San Francisco: Josey Bass; 2008.
54. Rubenstein LV, Mittman BS, Yano EM, et al. From Understanding Health Care Provider Behavior to Improving Health Care: The QUERI Framework for Quality Improvement. *Medical Care*. 38:129–141.2000;

Table 1
Baseline characteristics of 117 case managers in outcomes analysis, by intervention group

Characteristic	Overall (N=117)		RS (N=59)		CEP (N=58)		Test	
	No.	%	No.	%	No.	%	χ^2	df p
Healthcare screening sector (vs. community sector)	69	62	40	70	29	54	.9	1 .343
Non-licensed	87	73	45	76	42	70	.17	1 .676
Age (M \pm SD)	43.6 \pm 1.6		43.4 \pm 2.5		43.7 \pm 2.1		.01	1 .922
Female sex	82	71	38	65	44	76	.97	1 .325
Race							.36	3 .948
Hispanic	55	47	26	45	29	49		
Black or African American	52	45	27	46	25	45		
Non-Hispanic white	7	6	4	7	3	5		
Other (API/Native/other)	3	2	2	3	1	2		
Some college or above	105	90	53	90	52	90	0	1 .985
Depression care techniques ^b (M \pm SD)	2.6 \pm .1		2.5 \pm .2		2.7 \pm .2		.38	1 .537
Depression case management ^b (M \pm SD)	2.5 \pm .1		2.6 \pm .2		2.5 \pm .2		.46	1 .499
Community services provision ^c (M \pm SD)	1.4 \pm .2		1.1 \pm .2		1.6 \pm .3		3.07	1 .08
Perception of depression knowledge ^d (M \pm SD)	2.1 \pm .1		2 \pm .1		2.1 \pm .1		.45	1 .502
Perception of depression skill ^e (M \pm SD)	2.2 \pm .1		2.3 \pm .1		2.2 \pm .1		.39	1 .533
Personal depression stigma ^f (M \pm SD)	3.8 \pm .1		3.7 \pm .1		4.0 \pm .1		5.17	1 .023
Number of system barriers ^g (M \pm SD)	.8 \pm .1		.9 \pm .2		.7 \pm .2		.3	1 .582

^aData were multiply imputed and weighted for attrition to present eligible sample; Chi-square test was used for a comparison between the two groups accounting for the design effect of the cluster randomization. Percentages may not add to 100% because of rounding.

^bPossible scores range from 1 to 5, with higher scores indicating greater use depression care techniques.

^cPossible scores range from 0 to 5, with higher scores indicating greater hours providing services.

^dPossible scores range from 1 to 5, with lower scores indicating greater depression knowledge.

^ePossible scores range from 1 to 4, with higher scores indicating greater perception of skills.

^fPossible scores range from 1 to 5, with higher scores indicating less depression stigma.

^gPossible scores range from 0 to 4, with lower scores indicating fewer number of barriers from a list of 4.

Table 2
Case manager training participation and care practices at follow-up by intervention group status

Variable	Unadjusted Estimates [†]				Adjusted Analysis [‡]											
	RS	CEP	RS	CEP	RS	CEP	OR	95% CI	t	df	p	P-FDR				
Analytic N	No.	%	No.	%	Estimate	Estimate	Estimate	95% CI	Estimate	95% CI	OR	95% CI	Estimate	95% CI	OR	95% CI
Training participation %	112	15	27	42	74	<.001	27	15.9 – 41.8	73.9	57.9 – 85.4	7.78	2.9 – 20.89	4.2	50	<.001	<.001
		Mean		Mean							Difference					
Depression care techniques ^a (mean)	85	2.5 ± 1.1	3 ± 1.1	.04	2.8	.04	2.8	2.5 – 3	3.1	2.9 – 3.3	.32	.03 – .61	2.2	57	.031	.042
Depression case management ^a (mean)	97	2.8 ± 1.3	2.8 ± 1.2	.893	2.8	.893	2.8	2.5 – 3.1	2.9	2.6 – 3.1	.05	-.30 – .41	.3	43	.767	.767
Community services provision ^b (mean)	116	.6 ± .9	1.3 ± 1.4	.003	.7	.003	.7	.4 – .9	1.2	0.9 – 1.5	.51	.13 – .89	2.7	62	.009	.018

[†]Raw data without weighting or imputation; plus-minus values are mean ± SD.

[‡]Adjusted analyses used multiply imputed data (N=117); data were weighted to account for attrition to present eligible sample; a logistic regression model for the binary variable (training participation) and linear regression models for continuous variables adjusted for baseline status of the dependent variable, sector (formal vs. informal), and provider type (licensed vs. non-licensed) and accounted for the design effect of the cluster randomization. p-FDR=adjusted p value from the False Discovery Rate procedure.

^aPossible scores range from 1 to 5, with higher scores indicating greater use depression care techniques.

^bPossible scores range from 0 to 5, with higher scores indicating greater hours providing services.

Table 3
Case manager knowledge and attitudes at follow-up by intervention group status

	Unadjusted Estimates [†]			Adjusted Analysis [‡]									
	RS	CEP	p	RS	Estimate	95% CI	CEP	Estimate	95% C	CEP vs RS	t	df	p
Perceived depression knowledge ^a	104	2 ± .7	2.1 ± .9	.366	2	1.8 – 2.2	2.1	1.8 – 2.4	.09	-.25 – .42	.52	65	.605
Perception of depression skill ^b	97	2.2 ± .7	2.3 ± .7	.709	2.3	2.2 – 2.5	2.4	2.2 – 2.6	.07	-.15 – .3	.67	29	.505
Personal depression Stigma ^c	112	3.8 ± .7	4 ± .6	.061	3.8	3.7 – 3.9	3.9	3.8 – 4	.08	-.1 – .25	.92	35	.366
Perceived system barriers ^d	57	.7 ± 1.2	.1 ± .3	.008	.7	.4 – .9	.3	.1 – .6	-.33	-.7 – .05	-1.76	41	.085

[†]Raw data without weighting or imputation; plus-minus values are mean ± SD.

[‡]Adjusted analyses used multiply imputed data (N=117); data were weighted to account for attrition to present eligible sample; linear regression models for continuous variables adjusted for baseline status of the dependent variable, sector (formal vs. informal), and provider type (licensed vs. non-licensed) and accounted for the design effect of the cluster randomization. PFDR=adjusted p value from the False Discovery Rate procedure.

^aPossible scores range from 1 to 4, with higher scores indicating greater perception of skills.

^bPossible scores range from 1 to 5, with higher scores indicating less depression stigma.

^cPossible scores range from 0 to 4, with lower scores indicating fewer number of barriers from a list of 4.