UC Irvine

UC Irvine Previously Published Works

Title

Identity of ergosterol '5 β ,8 β -peroxide'

Permalink

<https://escholarship.org/uc/item/3zz7p346>

Journal

Journal of the Chemical Society, Chemical Communications, 0(15)

ISSN

0022-4936

Authors

Fisch, MH
Ernst, R
Flick, BH
et al.

Publication Date

1973

DOI

10.1039/C39730000530

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Identity of Ergosterol '5 β ,8 β -Peroxide'

By MICHAEL H. FISCH,*† ROBERT ERNST, BRIGITTA H. FLICK, and JOSEPH ARDITTI

(Department of Developmental and Cell Biology, University of California, Irvine, California 92664)

and DEREK H. R. BARTON, PHILIP D. MAGNUS,* and IAN D. MENZIES

(Chemistry Department, Imperial College, London SW7)

Summary Ergosterol '5 β ,8 β -peroxide' has been identified as 9,11-dehydroergosterol peroxide.

THE recent report¹ that ergosterol peroxide (I), long accepted as being a single, pure compound, is in fact a mixture of the 5 α ,8 α - and 5 β ,8 β -endo-peroxides (I) and (II) in the ratio 84:16 requires amendment.

Samples of ergosterol peroxide (I) prepared at Imperial College from recrystallised ergosterol by photo-oxidation (singlet oxygen) and aminium radical-cation catalysed oxygenation² (triplet oxygen) contained only ergosterol peroxide (I). No trace of a second compound could be found by 220 MHz n.m.r. spectroscopy.

A comparison of pure ergosterol peroxide (I), the claimed mixture [(I) + (II)] and 9,11-dehydroergosterol peroxide (III) is instructive.

	Ergosterol peroxide (I)	[(I) + (II)] (84:16)	(III)
M.p. ..	176–178°	167–171°	161–165°
[α] _D ^{21.5} (CHCl ₃)	–32.9°	–13.5°	+80°

A mixture of 16% (III) and 84% (I), prepared from the pure compounds, had [α]_D –14.2°. Similarly the n.m.r. spectrum of 16% (III) and 84% (I) is identical with that reported for [(I) + (II)]. Acetylation of a specimen of the ergosterol peroxide [(I) + (II)] prepared earlier¹ at Irvine by oxidation of ergosterol (singlet oxygen) and p.l.c. of the

product against ergosterol peroxide acetate and 9,11-dehydroergosterol peroxide acetate (AgNO₃-SiO₂ plates) demonstrated the presence of 9,11-dehydroergosterol peroxide acetate. The mass spectrum of [(I) + (II)] shows a strong $M-2$ 428 m/e corresponding in relative intensity to approximately 16% of (III).

These data conclusively show that the supposed ergosterol 5 β ,8 β -peroxide (II) is in reality the long known 9,11-dehydroergosterol peroxide.

(Received, 1st June 1973; Com. 784.)

† Present address: Witco Chemical Corporation, Technical Center, P.O. Box 110, Oakland, New Jersey 07436.

¹ J. Arditti, R. Ernst, M. H. Fisch, and B. H. Flick, *J.C.S. Chem. Comm.*, 1972, 1217.

² D. H. R. Barton, G. Leclercq, P. D. Magnus, and I. D. Menzies, *J.C.S. Chem. Comm.*, 1972, 446.