

Lawrence Berkeley National Laboratory

Recent Work

Title

COULOMB EXCITATION INTO THE BACKBEND REGION OF IDTr

Permalink

<https://escholarship.org/uc/item/42k9m0g1>

Author

Lee, I.Y.

Publication Date

1976-05-01

0 0 0 0 4 5 0 4 4 3 4

Submitted to Physical Review Letters

LBL-5031
Preprint c.1

RECEIVED
LIBRARY
DOCS


COULOMB EXCITATION INTO THE BACKBEND
REGION OF ^{164}Er

I. Y. Lee, D. Cline, R. S. Simon, P. A. Butler,
P. Colombani, M. W. Guidry, F. S. Stephens,
R. M. Diamond, N. R. Johnson, and E. Eichler

May 1976

Prepared for the U. S. Energy Research and
Development Administration under Contract W-7405-ENG-48

For Reference
Not to be taken from this room



LBL-5031
c.1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

COULOMB EXCITATION INTO THE BACKBEND REGION OF $^{164}\text{Er}^*$

I. Y. Lee, D. Cline,[†] R. S. Simon,[‡] P. A. Butler,[§] P. Colombani,^{||}
M. W. Guidry, F. S. Stephens and R. M. Diamond

Nuclear Science Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

and

N. R. Johnson and E. Eichler
Oak Ridge National Laboratory**
Oak Ridge, Tennessee 37830

ABSTRACT

Multiple Coulomb excitation has been used to excite previously unknown members of the ground state band above the known backbend in ^{164}Er . The ground-band $B(E2)$ values obey the rigid rotor relation to within $\pm 25\%$. A two band mixing analysis shows that the two intersecting bands have remarkably small interaction matrix elements at the backbend i.e., < 40 keV. This weak band interaction is predicted by the rotation-alignment model to be a general feature.

The discovery¹ of backbending (an anomalous behavior of the moment of inertia at high spin in nuclear rotational bands) has stimulated an intensive theoretical investigation of this phenomenon.²⁻⁶ Present experimental evidence^{3,4} suggests that backbending is caused by the intersection of the ground-state rotational band with a second rotational band possessing an appreciably larger moment of inertia. Two possibilities have emerged for the most likely nature of this second band. The Coriolis antipairing⁵ model considers it to be a band for which the pairing has collapsed while the rotation-alignment⁶ model attributes the band to two quasiparticles which are aligned with the rotating core by the Coriolis force. Observation of additional levels and a determination of the interaction matrix elements between the intersecting bands can shed considerable light on the structure of the bands.

Previously, backbending has been studied exclusively using (HI,xn) reactions to populate highly excited high-spin states which subsequently deexcite by γ -ray cascades into the yrast sequence of states. In contrast, multiple Coulomb excitation specifically excites those collective bands which are strongly coupled to the ground state and thus is a complementary probe of the backbending phenomenon. In addition, Coulomb excitation can be used to study neutron-rich nuclei which cannot be reached by (HI,xn) reactions. The present paper describes the first case where states through a reasonably sharp backbend region have been Coulomb excited. The nucleus ^{164}Er has been studied because the high spin yrast states up to spin 18^+ have been seen previously^{7,8} via the $^{164}\text{Dy}(\alpha,4n)$ reaction and because ^{164}Er is one of the few stable isotopes known to backbend sharply.

Beams of 612 MeV and 547 MeV ^{136}Xe ions from the LBL SuperHILAC were used to bombard a 1.34 mg/cm^2 self-supporting metallic foil of ^{164}Er . The isotopic enrichment was 73.6%. Three silicon detectors were used to detect scattered Xe ions at angles of 65° , 77° and 90° in coincidence with deexcitation γ -rays observed in a Ge(Li) detector located at -30° to the incident beam. The Ge(Li) detector was placed in the average recoil direction where the Doppler shift is a maximum, 8%, and the Doppler broadening is a minimum. A γ -ray energy resolution of $\leq 1\%$ (FWHM) was achieved. Four 7.6 cm by 7.6 cm NaI detectors, serving as a multiplicity filter, were placed around the target. The number of NaI detectors in coincidence was used to determine the multiplicity of each γ -ray transition observed in the Ge(Li) spectrum in coincidence with the scattered ions. The dependence of the γ -ray yields on the multiplicity distribution, on the bombarding energy and on the projectile scattering angle provided three independent measures of the location of each deexcitation gamma transition in the nuclear decay scheme. A γ -ray spectrum is shown in the upper section of Fig. 1. The unmarked γ -ray lines are due to Coulomb excitation of the $^{166,168}\text{Er}$ contaminants and also to excited target nuclei which recoil into the silicon detectors and exhibit a small Doppler shift.

The $^{164}\text{Dy}(\alpha,4n)^{164}\text{Er}$ reaction was studied, in addition to the Coulomb excitation, to search for weak branching at the backbend. A 10 mg/cm^2 self-supporting metallic foil, enriched to 93% in ^{164}Dy , was bombarded with a 51 MeV α -particle beam from the LBL 88" Cyclotron. Two 50 cm^3 coaxial Ge(Li) detectors, with energy resolution of 2.3 keV FWHM at 1.1 MeV, were used and both

singles and coincident γ -ray spectra were accumulated. The lower part of Fig. 1 shows the coincidence spectrum gated by transitions originating from states with $I \geq 12$.

The decay scheme derived from the present work is shown in Fig. 2. The yrast sequence up to spin 18^+ has been seen in earlier work where spin assignments were made on the basis of γ -ray angular distribution data.^{7,8} The present work supports these previous results. In addition, the $^{164}\text{Dy}(\alpha,4n)$ reaction clearly shows that an incompletely resolved 707 keV self-coincident doublet feeds into the yrast 14^+ state. This unresolved doublet, which has not been seen previously, was strongly excited by Coulomb excitation suggesting E2 character. The observed yield of this doublet is 1.5 times the calculated yield for Coulomb excitation of the ground band 16^+ state but is in agreement with the predicted sum of the yields of the $18^+ \rightarrow 16^+$ and $16^+ \rightarrow 14^+$ transitions if rigid rotor $B(E2)$ values are assumed. Thus this doublet is presumed to deexcite the 18^+ and 16^+ members of the ground band. Neither the 14^+ nor the 20^+ members of the second band was located in the present work. However, the transitions involving these states could have been masked by transitions in ^{162}Er excited by $^{162}\text{Dy}(\alpha,4n)$ since the 506 keV ($10^+ \rightarrow 8^+$) transition in ^{162}Er and the ($18^+ \rightarrow 16^+$) in ^{164}Er coincide.

Above the $14^+ \rightarrow 12^+$ transition, the discontinuity in the spacing between the ground-band transition energies is a striking feature of the Coulomb excitation spectrum shown in Fig. 1. The measured yields of these ground-band transitions were compared with calculations using the Winther-deBoer⁹ semiclassical Coulomb excitation code. An axially symmetric rigid rotor was assumed with $\langle 0||M(E2)||2 \rangle = 2.315$ e.b. taken

from an α -particle Coulomb excitation measurement,¹⁰ and with $\langle 0 || M(E4) || 4 \rangle = 0.2 \text{ e.b.}^2$ taken from systematics.¹¹ The ratio of experimental yields for adjacent ground band transitions agreed with the calculated ratio to better than $\pm 15\%$. The systematic uncertainties involved in using this code are expected to be less than $\pm 20\%$ from comparison with experimental yields for high-spin ground band states in other strongly deformed nuclei.^{12,13} Thus the ground-band $B(E2)$ values obey the rigid rotor relation to within $\leq 25\%$. Unfortunately the Coulomb excitation of the second band was difficult to observe because the $16^{+'} \rightarrow 14^{+}$ transition was unresolved from the strong $12^{+} \rightarrow 10^{+}$ transition, and the yrast $18^{+'} \rightarrow 16^{+}$ transition is predicted to be weak. The Coulomb excitation data places an upper limit on the ratio $B(E2; 14 \rightarrow 16') / B(E2; 14 \rightarrow 16)$ of ≤ 0.4 . On a two band mixing picture this ratio should be the same as the ratio $B(E2; 16' \rightarrow 14) / B(E2; 16' \rightarrow 14')$ if both bands have the same intrinsic quadrupole moment. This second ratio is given by the branching ratio for deexcitation of the $16'$ state. Systematics would suggest that the $16' \rightarrow 14'$ transition energy falls between 380 keV and 480 keV. No such transition was observed and the upper limit for branching to a 14^{+} state is ≤ 0.25 from the $^{164}\text{Dy}(\alpha, 4n)$ reaction data. This sets a lower limit of $B(E2; 16' \rightarrow 14) / B(E2; 16' \rightarrow 14') \geq 0.5$.

A conventional backbending plot of these results is shown in Fig. 3. The $N = 96$ isotones ^{166}Yb , ^{168}Hf and ^{170}W also exhibit very similar backbending and the upper band has about the same moment of inertia and excitation energy in all these nuclei.¹⁴ Below the backbend the moment of inertia in the ground band increases slightly with increasing spin,

presumably due to the influence of Coriolis antipairing.

The ratio of the intra- to interband $B(E2)$ values at the band intersection directly determines the interaction strength when only two bands are interacting, provided that the level energies are known and the bands have the same intrinsic quadrupole moments. The Coulomb excitation and branching ratio data suggest that the ratio $B(E2;16' \rightarrow 14)/B(E2;16' \rightarrow 14') \approx 0.45$ which leads to an average interaction matrix element of 38 keV for these states if the 14^+ states splitting is 130 keV. In addition, the unperturbed ground band 14^+ and 16^+ states fall on an extension of the line through the lower spin states on a backbending plot, as indicated by the dashed line in Fig. 3, provided that the interaction matrix elements are taken to be 38 keV. This interaction predicts a 24% reduction in the ratio $B(E2;16 \rightarrow 14)/B(E2;14 \rightarrow 12)$ for the ground band which is within the experimental limit given by the Coulomb excitation yields. The γ -ray branching ratio at the backbend has been measured^{15,16} in two other nuclei, the $N = 90$ isotones ^{154}Gd and ^{156}Dy . A similar analysis gives an average interaction matrix element for the 16^+ and 18^+ states of 23.5 ± 1.5 keV in ^{154}Gd and 8.5 ± 1.5 keV for the 16^+ state in ^{156}Dy which is consistent with the values previously reported.^{15,16}

The energy for the 18^+ state given by the smooth extrapolation in Fig. 3 lies 27 keV above the experimental energy. The two quasiparticle-plus-rotor model suggests additional bands occur in this energy region and the above shift could be due to the intersection of the ground band with one of these additional bands. Such behavior would result in a rapid loss of identity of the ground band at higher spin values.

Band interaction matrix elements of less than 40 keV at the backband are remarkably small, i.e., they are nearly two orders of magnitude smaller than might be expected for Coriolis matrix elements at these spins. However, this behavior can be understood in the rotation-alignment model. Calculations with the two-quasiparticle plus rotor model^{6,17} show that the aligned two $i_{13/2}$ quasineutron eigenfunctions for the yrast states become localized around $J = 12$ and $R = I - 12$ with increasing spin I . On the other hand the zero-quasiparticle ground band has $I = R$ for a fully paired state. The Coriolis force does not couple states with differing core rotation R and thus the two bands interact only via the overlap of weak components in the wavefunctions. This overlap becomes progressively smaller with increasing spin due to the increased localization in R space of the aligned states. Two calculations within this model^{6,17,18} suggest that the interaction is ≤ 140 keV and is nearly constant for $10 < I < 22$. However, the assumptions made in these calculations may not be adequate for accurately reproducing the interband interaction strength. A more complete Hartree-Fock-Bogoliubov calculation by Mang¹⁹ also predicts a small interaction strength.

This first example of Coulomb excitation through a known backband illustrates the power of this technique which can be used on many nuclei that cannot be excited by (HI, xn) reactions. The ground band $B(E2)$ values have been measured in ^{164}Er and follow the rigid-rotor relation to within $\pm 25\%$ throughout the backband. The band intersecting the ground band in ^{164}Er is closely similar to the bands seen in ^{154}Gd and ^{156}Dy which shows that this type of behavior is not peculiar to the

90-neutron region. The $B(E2)$ data and the level energies in all three nuclei are consistent with a two band mixing model having a remarkably weak interaction strength at the backbend, i.e., <40 keV. This behavior is reasonably well described by the rotation-alignment model.

FOOTNOTES AND REFERENCES

* This work was done with support from the U. S. Energy Research and Development Administration. Any conclusions or opinions expressed in this report represent solely those of the authors and not necessarily those of the Lawrence Berkeley Laboratory nor of the U. S. Energy Research and Development Administration.

[†] Partially supported by the N.S.F. Permanent address: Nuclear Structure Research Laboratory, University of Rochester, Rochester, NY 14627.

[‡] On leave from Sektion Physik der Ludwig-Maximilians - Universität München, 8046 Garching, Germany; sponsored by the Bundesministerium für Forschung und Technologie.

[§] U.K.S.R.C./N.A.T.O. fellow.

^{||} Permanent address: Institut de Physique Nucléaire, 91406 Orsay, France.


** Operated by Union Carbide Co. for the U. S. Energy Research and Development Administration.

1. A. Johnson, H. Ryde and J. Sztarkier, Phys. Lett. 34B, 605 (1971).
2. A. Johnson and Z. Szymański, Phys. Rep. 7C, 182 (1973).
3. R. A. Sorensen, Rev. Mod. Phys. 45, 353 (1973).
4. F. S. Stephens, Rev. Mod. Phys. 47, 43 (1975).
5. B. R. Mottelson and J. G. Valatin, Phys. Rev. Lett. 5, 511 (1960).
6. F. S. Stephens and R. S. Simon, Nucl. Phys. A183, 257 (1972).
7. M. V. Banaschik, C. Günther, H. Hübel, A. C. Rester, G. Nowicki and J. J. Pinajian, Nucl. Phys. A222, 459 (1974).
8. W. F. Davidson, R. M. Lieder, H. Beuscher, A. Neskakis, G. A. Varley, J. C. Willmott, F. Kearns and J. C. Lisle, J. Phys. G 3, 199 (1976).
9. A. Winther and J. deBoer, in Coulomb Excitation (Academic Press, NY, 1966), p. 303.
10. R. Ronnigen, private communication (1975).
11. K. A. Erb, J. E. Holden, I. Y. Lee, J. X. Saladin and T. K. Saylor, Phys. Rev. Lett. 29, 1010 (1972).

12. D. Ward, P. Colombani, I. Y. Lee, P. A. Butler, R. S. Simon, R. M. Diamond and F. S. Stephens, to be published in Nucl. Phys.
13. M. W. Guidry, P. A. Butler, P. Colombani, I. Y. Lee, D. Ward, R. M. Diamond, F. S. Stephens, E. Eichler, N. R. Johnson and R. Sturm, to be published in Nucl. Phys.
14. R. O. Sayer, J. S. Smith, III and W. T. Milner, Atomic and Nuclear Data Tables 15, 85 (1975).
15. T. L. Khoo, F. M. Bernthal, J. S. Boyno and R. A. Warner, Phys. Rev. Lett. 31, 1146 (1973).
16. H. R. Andrews, D. Ward, R. L. Graham and J. S. Geiger, Nucl. Phys. A217, 141 (1974).
17. C. Flaum and D. Cline, Proc. of the Symp. on High Spin States, Jülich 1975, Phys. Rev. (1976), in press.
18. The interaction between the zero and two quasiparticle bands given in Ref. 6 is too large due to an incorrect phase convention used for the zero-quasiparticle ground band. This error has little effect on the other calculated properties and was corrected in the present calculations.
19. H. J. Mang, private communication (1976).


FIGURE CAPTIONS

- Fig. 1. Coincidence γ -ray spectra for ^{164}Er . The upper spectrum is for the excitation of ^{164}Er by ^{136}Xe . The lower spectrum is for the sum of the coincidence spectra gated by the transitions from states with spin ≥ 12 fed by the $^{164}\text{Dy}(\alpha,4n)$ reaction.
- Fig. 2. Level scheme of ^{164}Er .
- Fig. 3. Plot of the moment of inertia vs the square of the angular velocity for ^{164}Er . The dashed line calculates a smooth extrapolation of the line through the lower spin states.



XBL 763-2513A


Fig. 1



^{164}Er

XBL 763-2514 A

Fig. 2



XBL 763-2515A

Fig. 3

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720