Lawrence Berkeley National Laboratory

Lawrence Berkeley National Laboratory

Title

IMPLICATIONS OF THE VIRIAL THEOREM ON THE DESCRIPTION OF WEAK BONDS

Permalink

<https://escholarship.org/uc/item/43n3x38f>

Author

Winn, John S.

Publication Date

1981

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Molecular Research Division

RECEIVED
LAWRENCE
BERKELEY LABORATORY

MAR 5 1981

Submitted to the Journal of Chemical Physics LIBRARY AND
DOCUMENTS SECTION

IMPLICATIONS OF THE VIRIAL THEOREM ON THE
DESCRIPTION OF WEAK BONDS

John S. Winn

January 1981

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782.*

LBL-12241
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

IMPLICATIONS OF THE VIRIAL THEOREM
ON THE DESCRIPTION OF WEAK BONDS

By

John S. Winn*
*Department of Chemistry and
Materials and Molecular Research Division
Lawrence Berkeley Laboratories
University of California
Berkeley, CA 94720*

This research was supported by the National Science Foundation and by the Division of Chemical Sciences, Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract W-7405-ENG-48.

---

*Alfred P. Sloan Research Fellow

Abstract

The origins of and variations in bonding among weakly bound diatomics are discussed in terms of the virial theorem as applied to Born-Oppenheimer total energy curves. These curves are represented by accurate and flexible analytic functions. Expressions are given for the relations among the parameters of these functions and the electron average kinetic energy, $\langle T(R) \rangle$, and the total molecular potential energy, $\langle V(R) \rangle$. The relatively brief distance through which weak bonding occurs (as opposed to the relative distances for chemical bonding) is discussed, as is the origin of bond saturation through a homologous series of weakly bound diatomics.

Application of the virial theorem to the total energy of a diatomic molecule within the Born-Oppenheimer approximation yields an instructive and familiar¹ description of the physical origin of bonding between two atoms. If $W(R)$ is the Born-Oppenheimer total energy of the atoms as a function of separation, R , then the virial theorem relates the average kinetic energy of the electrons, $\langle T(R) \rangle$, and the average Coulombic potential energy of the diatomic, $\langle V(R) \rangle$, to $W(R)$ via the relations

$$\langle T(R) \rangle = -W(R) - R \frac{dW(R)}{dR} \quad (1)$$

and

$$\langle V(R) \rangle = 2W(R) + R \frac{dW(R)}{dR} \quad (2)$$

These three functions are shown in Fig. 1 for the ground state of H_2 , using the ab initio data of Kołos and Wolniewicz².

The distance scale is measured in multiples of the H_2 equilibrium bond length (i.e., the distance, R_e , at which $W(R)$ is a minimum), and the energy scale is measured in multiples of the H_2 bond energy (i.e., $W(R=\infty) - W(R_e)$).

Constants have been added to each curve so that each goes to

zero as $R \rightarrow \infty$

As the atoms approach, the potential energy rises (electrons are moving away from nuclei) and the kinetic energy falls (as delocalization begins). In the vicinity of $R/R_e=2$, this trend reverses. The kinetic energy increases as the electronic wavefunction is localized further, raising the momentum, but the potential energy falls, as charge is now brought nearer both nuclei. Only at $R/R_e < 0.5$ does nuclear repulsion cause the potential energy to increase and contribute to the total repulsion energy.

These general features apply to any diatomic in a bound state (with the exception¹ of purely ionic attractions). To construct and interpret $\langle T(R) \rangle$ and $\langle V(R) \rangle$, one needs only $W(R)$. It has been shown³ that $W(R)$ for many weakly bound diatomics can be given accurately by the Thakkar analytic function

$$W(R) = e_o \lambda^2 \left[1 + \sum_{n=2}^{\infty} e_n \lambda^n \right] + W(R_e) \quad (3)$$

where $\lambda = 1 - (R_e/R)^p$ and p is a positive constant. This function has been shown to represent the true $W(R)$ throughout the bound region of the molecule, including the long-range attractive tail, in many cases.

In Fig. 2, we show $W(R)$ for Ar_2 , using Eq. (3), as well as the $\langle T(R) \rangle$ and $\langle V(R) \rangle$ functions derived from it.⁴ These functions are plotted in a reduced coordinate system analogous to that used in Fig. 1. All three curves are dramatically different in qualitative appearance from those for H_2 .

Note that chemical bonding (i.e. H_2) can be attributed to a gradual decrease in $\langle V \rangle$ throughout a wide portion of the total energy well. Weak bonding can be attributed to the decrease in electron kinetic energy over the attractive tail, but, in the vicinity of R_e , the roles of $\langle T \rangle$ and $\langle V \rangle$ in promoting binding are rapidly reversed.

The behavior of $\langle T \rangle$ and $\langle V \rangle$ at long range is, for Ar_2 , a manifestation of the dispersion interaction. Admixing excited state character to form the wavefunction descriptive of the induced dipole-induced dipole attraction necessarily increases $\langle V \rangle$ and decreases $\langle T \rangle$. (Recall that, for an atom, $W = -\langle T \rangle < 0$.) What this analysis shows rather clearly is the persistence of this behavior as R decreases towards R_e .

One may take the position of the maximum in $\langle V \rangle$ or of the minimum in $\langle T \rangle$ as a lower bound to the inner limit of classical long range attraction. Expressions for these distances (which are slightly different) can be found from

Eq. (1) only by numerical methods, except for the special case $W(R) = e_o \lambda^2 + W(R_e)$. This is a simple Lennard-Jones (p,2p) function. For this case, one finds

$$d\langle T \rangle / dR = 0 \quad \text{when } R/R_e = \left(\frac{2p-1}{p-1} \right)^{1/p} \quad (4)$$

and

$$d\langle V \rangle / dR = 0 \quad \text{when } R/R_e = \left(\frac{2p-2}{p-2} \right)^{1/p} \quad (5)$$

Both of these distances are larger than the location of the inflection point in $W(R)$, which occurs at $R/R_e = ((2p+1)/(p+1))^{1/p}$. Since p is a property of $W(R_e)$ and can be calculated³ from derivatives of $W(R)$ at R_e , from observed equilibrium spectroscopic constants, or, as shown below, from first principles, Eq's (4) and (5) provide unambiguous criteria by which one may locate with certainty the "end" of long range attraction as R is decreased towards R_e . These relations should be of use in checking the somewhat arbitrary decisions made to locate this point in schemes using piecewise analytic representations⁵ of $W(R)$.

Several simple and informative relations pertaining to $\langle V(R_e) \rangle$ and $\langle T(R_e) \rangle$ can be derived from Eq's (1)-(3). The well-known fact that the decrease in $\langle V \rangle$ at R_e is twice the increase in $\langle T \rangle$ at R_e is, of course, general, and does not depend on the functional form of $W(R)$. Using the following

definitions for derivations of W and for p:

$$k_e \equiv \left(\frac{d^2 W(R)}{dR^2} \right)_{R_e},$$

$$l_e \equiv \left(\frac{d^3 W(R)}{dR^3} \right)_{R_e},$$

and

$$p = -l_e R_e / 3k_e - 1,$$

one finds

$$\left(\frac{d\langle V \rangle}{dR} \right)_{R_e} = - \left(\frac{d\langle T \rangle}{dR} \right)_{R_e} = k_e R_e,$$

$$\left(\frac{d^2 \langle T \rangle}{dR^2} \right)_{R_e} = 3pk_e,$$

and

$$\left(\frac{d^2 \langle V \rangle}{dR^2} \right)_{R_e} = (1-3p)k_e = k_e - \left(\frac{d^2 \langle T \rangle}{dR^2} \right)_{R_e}.$$

Also, for Eq. (3),

$$k_e = 2p^2 e_o (1 + \sum_n e_n) / R_e^2.$$

It is helpful to consider these expressions in reduced coordinates so that the relative variations in $\langle T \rangle$ and $\langle V \rangle$

across the bound portion of $W(R)$ can be compared. Thus, when distances are scaled by R_e and energies by D_e , as in the figures, the behavior of $\langle T \rangle$ and $\langle V \rangle$ at $R/R_e=1$ can be related directly to p , viz., $-d\langle T \rangle/dR \sim p^2$, $d^2\langle T \rangle/dR^2 \sim p^3$, etc.

For chemically bound species, p 's are found, from spectroscopic data,⁶ to lie in the range 0.6 to 3.0. In contrast, p 's for weakly bound species³ are found throughout the range 3.1 to >6 . Thus, weak bonds may be characterized in the vicinity of $R/R_e=1$ as having $\langle T \rangle$ ($\langle V \rangle$) increase (decrease) at relatively much greater rates than do chemically bound species in general. It is a somewhat surprising outcome of this analysis that weak bonding is a relatively more localized interaction than chemical bonding.⁷

The reason Eq. (3), the Thakkar function, works as well as it does (or, equivalently, fails rather miserably for strongly bound diatomics) appears to involve two factors. The first is the persistence of long range behavior to R values close to R_e (see Fig. 2) in many weakly bound molecules⁸, allowing Eq. (3) to reproduce the long range tail with a power series lead by a term in R^{-p} with p close to the anticipated integral value.

The second is the physical similarity of repulsive interactions for weakly bound molecules: the Pauli repulsion of two (usually) spherical atoms. Binding, as noted above, is relatively less localized in covalent, ionic, or other strong interactions, and the electron distribution changes less rapidly with R in the vicinity of R_e . Eq. (3) accomodates the relatively abrupt repulsion of weakly bound species through the interrelation of p to k_e and the derivatives of $\langle T \rangle$ and $\langle V \rangle$ noted above.

It is also possible, given the relationship of p to derivatives of $W(R)$, to express p from first principles. Let $\psi(r, R)$ be the exact Born-Oppenheimer wavefunction for the diatomic, given as a function of all electron coordinates, r , and, parametrically, as a function of internuclear distance R . First, from the Hellmann-Feynman theorem⁹, the internuclear force is given by

$$-\frac{dW(R)}{dR} = -\int \psi^* \frac{\partial V}{\partial R} \psi d\tau = -\langle \partial V / \partial R \rangle. \quad (6)$$

Clinton¹⁰ has derived an expression for k_e from the virial theorem definition of $\langle V \rangle$;

$$k_{eR_e} = \left(\int \frac{\partial \psi^*}{\partial R} \psi V d\tau \right)_{R_e}.$$

From the differentiation of Eq.(6), one has

$$k_e = \left(\int \frac{\partial \psi^* \psi}{\partial R} \frac{\partial V}{\partial R} d\tau \right)_{R_e} + \left(\int \psi^* \psi \frac{\partial^2 V}{\partial R^2} d\tau \right)_{R_e}$$

From further differentiation of $\langle V \rangle$, one has

$$\frac{d^2 \langle V \rangle}{dR^2} = \int \frac{\partial^2 \psi^* \psi}{\partial R^2} V d\tau + 2 \int \frac{\partial \psi^* \psi}{\partial R} \frac{\partial V}{\partial R} d\tau + \int \psi^* \frac{\partial^2 V}{\partial R^2} \psi d\tau$$

When this expression is evaluated at R_e and combined with the expression for $(d^2 \langle V \rangle / dR^2)_{R_e}$ given above, one finds

$$\left[\int \frac{\partial^2 \psi^* \psi}{\partial R^2} V d\tau \right]_{R_e} + \left[\int \frac{\partial \psi^* \psi}{\partial R} \frac{\partial V}{\partial R} d\tau \right]_{R_e} = -3pk_e = - \left(\frac{d^2 \langle T \rangle}{dR^2} \right)_{R_e}$$

Thus p can be evaluated from first principles. This expression is also more transparent than the perturbation expression for p given by Thakkar.¹¹

Finally, we note that this expression can be used to find the spectroscopic vibration-rotation interaction constant, α_e , (or more precisely, the Dunham energy term,³ Y_{11}). This constant leads the expansion of the rotational constant as a function of vibrational quantum number, v , via

$$B_v = B_e - \alpha_e (v + \frac{1}{2}).$$

From the spectroscopic definition³ of p ,

$$p = \frac{\alpha_e \omega_e}{6B_e^2},$$

and the familiar expressions for ω_e and B_e in terms of k_e , R_e , ψ , and molecular masses, one can evaluate α_e from first principles.

One is used to expressing the anharmonic shape of a potential function through the first vibrational correction $\omega_e x_e$ (or Y_{20}). It is important to remember that α_e not only measures the same effect, but it does so in a more direct fashion. The constant α_e describes the lengthening (if $\alpha_e > 0$) of the average internuclear separation as the vibrational quantum number is increased.¹²

In conclusion, we apply the results of this analysis to a qualitative view of an aspect of weak bonding which is, at first sight, somewhat unexpected and puzzling. Given the periodic variation of polarizabilities throughout the Periodic Table, one might expect the binding energies of the homologous series Ne_2 , NeAr , NeKr , and NeXe to increase uniformly throughout the series. Instead, the binding energy increases¹³ from Ne_2 to NeAr , but then saturates at a virtually constant value from NeAr to NeXe . Similarly, the greater polarizability of Na when compared to Ar might lead one to predict NaAr to be more strongly bound than Ar_2 .

In fact, NaAr exhibits 40% of the binding energy of Ar_2 .

Since any of these binding energies is a trivial fraction of the total molecular energy, only very slight and perhaps subtle changes to the separated atom wavefunctions occur near R_e . Consider first the Ne containing rare gas diatomics. From the measured¹³ $W(R)$'s, one deduces p values ranging from 5.3 for Ne_2 to 5.6, 6.0, and 6.9 for NaAr, NeKr, and NeXe, respectively. This uniform increase in p implies a uniform (and more pronounced) increase in the rates of change of $\langle T \rangle$ and $\langle V \rangle$ near R_e as Ne is paired with successively larger atoms. The smaller (less polarizable) Ne finds its electrons confined to an even smaller region of space at the onset of repulsive overlap, and the degree of confinement will be greater the larger the other atom happens to be. This will raise the total electron kinetic energy at a rate governed by the identity of the less polarizable atom, in analogy with shortening the length of the box containing a particle in an already short box.

Similar arguments would seem to apply to the alkali metal-rare gas diatomics. The large bond lengths found for these species (i.e., $R_e = 5 \text{ \AA}$ for NaAr) imply a repulsive interaction at a distance governed by the spatially diffuse outer s electron of the alkali. However, p for NaAr is³

~ 4.3 , indicating that the single s electron is, as expected, less effective in raising the kinetic energy of the outer Ar electrons. The near equality of the NeAr-NeXe binding energies and the "anomalously small" binding of NaAr should be regarded as somewhat fortuitous outcomes. But the variations in the detailed shapes of $W(R)$ near R_e can be understood in terms of the effects of electron density variations in controlling the repulsive part of $\langle T(R) \rangle$.

To summarize, the availability of accurate energy functions for many weakly bound diatomics permits the analysis of weak bonding in terms of the physical consequences of the virial theorem and the detailed shape of $W(R)$ functions near R_e . The relatively more abrupt onset of repulsion in weak bonds (compared to true chemical bonds) is traced to the abrupt end of long range attractions and consequent rapid increase in the total kinetic energy of the diatomic's electrons. Periodic variations in non-bonding electron distributions alter the rate of rise of this repulsion in a way which can be measured by the parameters describing the shape of the total energy function near R_e . These parameters can be directly related to expectation values of the molecular Hamiltonian or to various derivatives of them.

ACKNOWLEDGMENT

This research was supported by the National Science Foundation and by the Division of Chemical Sciences, Office of Basic Energy Sciences, U. S. Department of Energy, under contract W-7405-Eng-48.

REFERENCES

1. J. C. Slater, *J. Chem. Phys.* 1, 687 (1933).
2. W. Kołos and L. Wolniewicz, *J. Chem. Phys.*, 43, 2429 (1965).
3. J. H. Goble, D. C. Hartman, and J. S. Winn, *J. Chem. Phys.* 67, 4206 (1977); J. H. Goble and J. S. Winn, *J. Chem. Phys.* 70, 2051 (1979); J. H. Goble and J. S. Winn, *J. Chem. Phys.* 70, 2058 (1979).
4. J. H. Goble, S. M. Welsh, and J. S. Winn (to be published).
5. See, for example, R. W. Bickes, Jr., and R. B. Bernstein, *J. Chem. Phys.* 66, 2408 (1977), or J. M. Parson, P. E. Siska, and Y. T. Lee, *J. Chem. Phys.* 56, 1511 (1972).
6. K. P. Huber and G. Herzberg, *Constants of Diatomic Molecules*, (van Nostrand Reinhold, New York, 1979).
7. Of course, R_e is generally larger for weakly bound diatomics than for chemically bound diatomics, and the real distances over which bonding occurs is correspondingly larger.
8. Important exceptions to this behavior include the alkaline earth diatomics.
9. H. Hellmann, *Einführung in die Quantenchemie* (Lepizig, 1937), p. 285; R. P. Feynman, *Phys. Rev.* 56, 340 (1939).
10. W. L. Clinton, *J. Chem. Phys.* 33, 1603 (1960). See also P. Phillipson, *J. Chem. Phys.* 39, 3010 (1963).
11. A. J. Thakkar, *J. Chem. Phys.* 62, 1693 (1975). See also R. G. Parr and R. J. White, *J. Chem. Phys.* 49, 1059 (1968).
12. Strictly speaking, the expansion of B_v about B_e should be given a classical or semiclassical^v interpretation here. Thus, B_e is B_v for the fictitious level $v = -1/2$, and an expansion in powers of $(v + 1/2)$ should be either mass scaled or reduced to a classical limit of infinite reduced mass.
13. C. Y. Ng, Y. T. Lee, and J. A. Barker, *J. Chem. Phys.* 61, 1996 (1974).

Figure Captions

Figure 1. The total energy, W , the average electron kinetic energy, T , and the average diatomic potential energy, V , of H_2 , measured in units of the H_2 bond energy, plotted as a function of internuclear distance, measured in units of the equilibrium H_2 bond length. Data from Ref. 2.

Figure 2. As Fig. 1, but for Ar_2 . Data from Ref. 4.

XBL 811-7881

Figure 1

XBL 811-7882

Figure 2