

UCSF

UC San Francisco Previously Published Works

Title

Thalamocortical circuits in generalized epilepsy: Pathophysiologic mechanisms and therapeutic targets.

Permalink

<https://escholarship.org/uc/item/43n7j8nh>

Authors

Lindquist, Britta E
Timbie, Clare
Voskobiynyk, Yuliya
[et al.](#)

Publication Date

2023-06-01

DOI

10.1016/j.nbd.2023.106094

Peer reviewed



Published in final edited form as:

Neurobiol Dis. 2023 June 01; 181: 106094. doi:10.1016/j.nbd.2023.106094.

Thalamocortical circuits in generalized epilepsy: Pathophysiologic mechanisms and therapeutic targets

Britta E. Lindquist^{a,b,d}, Clare Timbie^{a,c,d}, Yuliya Voskobiynyk^{a,d}, Jeanne T. Paz^{a,d,e,*}

^aGladstone Institute of Neurological Disease, United States of America

^bUCSF Department of Neurology, Division of Neurocritical Care, United States of America

^cUCSF Department of Neurology, Division of Pediatric Epilepsy, United States of America

^dUCSF Department of Neurology, United States of America

^eKavli Institute for Fundamental Neuroscience, UCSF, United States of America

Abstract

Generalized epilepsy affects 24 million people globally; at least 25% of cases remain medically refractory. The thalamus, with widespread connections throughout the brain, plays a critical role in generalized epilepsy. The intrinsic properties of thalamic neurons and the synaptic connections between populations of neurons in the nucleus reticularis thalami and thalamocortical relay nuclei help generate different firing patterns that influence brain states. In particular, transitions from tonic firing to highly synchronized burst firing mode in thalamic neurons can cause seizures that rapidly generalize and cause altered awareness and unconsciousness. Here, we review the most recent advances in our understanding of how thalamic activity is regulated and discuss the gaps in our understanding of the mechanisms of generalized epilepsy syndromes. Elucidating the role of the thalamus in generalized epilepsy syndromes may lead to new opportunities to better treat pharmaco-resistant generalized epilepsy by thalamic modulation and dietary therapy.

Keywords

Absence epilepsy; Burst firing; Generalized spike-and-wave discharge; Oscillation; Thalamus; Genetic generalized epilepsy; Idiopathic generalized epilepsy

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

*Corresponding author at: Gladstone Institutes, 1650 Owens St, San Francisco, CA 94158, United States of America.

jeanne.paz@gladstone.ucsf.edu (J.T. Paz).

Author contributions

Britta E Lindquist: conceptualization, original draft

Clare Timbie: original draft

Yuliya Voskobiynyk: original draft

Jeanne T Paz: conceptualization; supervision; review and editing of draft manuscript

Declaration of Competing Interest

None. The authors have no competing interests to declare.

1. Introduction

Epilepsy comprises a broad category of diseases occurring across the lifespan, united by a common tendency toward unprovoked seizures (Fisher et al., 2017). Generalized epilepsy disorders affect 24 million individuals worldwide (GBDN Collaborators, 2019). Unlike focal epilepsy, which remains localized to specific parts of the brain, generalized epilepsy involves the whole brain. The thalamus is thought to be a key contributor to generalized epilepsy because it has widespread connections to the cortex and other brain regions.

Most, if not all, generalized epilepsy syndromes have a genetic or presumed genetic etiology. Thus, the latest ILAE Task Force classified these disorders as genetic generalized epilepsies (GGEs). GGE is characterized by generalized seizure semiologies (myoclonic, absence, myoclonic-tonic-clonic, and/or generalized tonic-clonic) and generalized discharges at 2.5–5.5 Hz on electrographic recordings outside of seizures (Hirsch et al., 2022). GGE includes the defined clinical syndromes known as idiopathic generalized epilepsies (IGEs); some developmental epileptic encephalopathies (DEEs) also meet the criteria of GGE (Hirsch et al., 2022; Wirrell et al., 2022). IGEs and DEEs differ in etiology—largely unknown genetic factors in IGE vs. known causal mutations for many DEEs—and in clinical presentation—age of onset, comorbidities, and response to treatment (Fig. 1). In addition, a number of IGEs have been modeled in animals, whereas many DEEs lack validated models. Therefore, much of our review summarizes work on IGEs, as animal models of IGEs have established the foundation for our understanding of intrathalamic circuits and long-range projections between the thalamus and other structures in generalized epilepsy. Nevertheless, we also draw parallels with generalized DEEs, as the presence of generalized seizure semiologies in DEEs suggests there may be overlap in pathogenic mechanisms with IGEs.

The aim of this review is to summarize our current knowledge about mechanisms of GGEs and to identify remaining gaps in our understanding of pathogenesis and potential treatment targets. First, we provide an overview of the epidemiology and clinical features of GGEs, contrasting IGEs and DEEs (Fig. 1). Then, we summarize the functional neuroanatomy of intrathalamic and thalamo-cortico-thalamic circuits and their modulation by local metabolic conditions, inputs from other brain regions, and systemic factors, highlighting evidence implicating these pathways in GGE. We also report recent preclinical work on thalamocortical mechanisms in DEEs, which will be important in identifying treatment targets for syndromes that currently lack highly effective medical therapies. Finally, we discuss unresolved questions in the field relevant to the pathogenesis of GGEs and emerging treatment approaches targeting the thalamus.

2. Key features of idiopathic generalized epilepsies (IGEs)

There are four key IGE syndromes: childhood absence epilepsy, juvenile absence epilepsy, juvenile myoclonic epilepsy, and generalized tonic-clonic seizures alone. These syndromes differ in their age of onset, time course, prognosis, and major seizure types. Despite this heterogeneity, pharmacologic treatments largely overlap, including ethosuximide for absence seizures and lamotrigine, valproic acid, or levetiracetam for convulsive (i.e.,

generalized tonic-clonic) seizures (Beydoun and D'Souza, 2012; Mastroianni et al., 2021). However, a large portion of IGEs are refractory to these approaches and require better treatments or new drug targets. Below we review the common clinical features and pathophysiologic mechanisms of these four IGE syndromes, current drug targets, and the unresolved gaps that need to be addressed to generate better treatments.

2.1. Childhood absence epilepsy

2.1.1. Epidemiology—Childhood absence epilepsy (CAE) occurs in 6.3–8.0 children per 100,000 and is more common in girls (60–75% of cases) than in boys (ILAE, 1989; Wirrell et al., 1996). CAE accounts for approximately 18% of epilepsy in school-aged children, with onset typically at 4–10 years of age (Blom et al., 1978; Cavazzuti, 1980; Olsson, 1988; ILAE, 1989; Loiseau et al., 1990; Wirrell et al., 1996; Grosso et al., 2005; Valentin et al., 2007; Ma et al., 2011; Hirsch et al., 2022).

2.1.2. Features and natural history—CAE clinical features include daily typical absence seizures associated with 2.5–4-Hz generalized spike-and-wave discharges (SWDs), which can be reliably provoked by hyperventilation (Salvati and Beenhakker, 2019), and an interictal (between seizures) generalized spike-and-wave pattern, often occurring during drowsiness and sleep. Typical absence seizures are characterized by abrupt onset and loss of awareness lasting 3–20 seconds, followed by an instant return to normal activity with possible confusion (Panayiotopoulos et al., 1989b; Wirrell et al., 1996; Sadleir et al., 2006; Sadleir et al., 2009; Kessler et al., 2017; Seneviratne et al., 2017; Elmali et al., 2020). The seizures are accompanied by inert facial expression, staring, behavioral arrest (interruption of activity), and sometimes manual automatisms and eye involvement. Incontinence and loss of postural control are rare (Kessler et al., 2017; Elmali et al., 2020). Antecedent febrile seizures occur in 10–15% of children with CAE (Livingston et al., 1965; Dieterich et al., 1985; Marini et al., 2003). Generalized tonic-clonic seizures may occur later during adolescence, and myoclonic seizures, by definition, do not occur in CAE patients (Wirrell et al., 1996; Ma et al., 2011; Elmali et al., 2020).

Children with CAE show typical development; however, subtle learning difficulties and attention deficit hyperactivity disorder (ADHD) might be present, with an increased risk for depression and anxiety (Wirrell et al., 1997; Austin et al., 2001; Hermann et al., 2007; Caplan et al., 2008; Shinnar et al., 2017; Abarategui et al., 2018). CAE is usually drug-responsive (Table 1); first-line agents for treatment are ethosuximide, valproic acid, and lamotrigine. Sixty percent of patients achieve spontaneous remission in early adolescence. The remainder continue to require treatment or develop other generalized epilepsy syndromes; in one longitudinal study, 7% required medication after 12–17 years of follow-up (Wirrell et al., 1996; Trinka et al., 2004; Grosso et al., 2005; Valentin et al., 2007; Callenbach et al., 2009; Morse et al., 2019). Why remission occurs in some patients but not others is not well understood.

2.2. Juvenile absence epilepsy

2.2.1. Epidemiology—Juvenile absence epilepsy (JAE) is less common than CAE, accounting for 2.4–3.1% of child and adolescent epilepsy, with onset typically at 9–13 years of age (Berg et al., 1999; Jallon and Latour, 2005; Wirrell et al., 2011).

2.2.2. Features and natural history—JAE clinical features include both absence seizures, with 3–Hz generalized SWDs, and generalized tonic-clonic seizures. A history of febrile seizures is noted in 6–33% of patients (Marini et al., 2004; Asadi-Pooya et al., 2013; Asadi-Pooya and Homayoun, 2020). JAE has fewer (less than daily) absence seizures, and seizures are associated with less loss of awareness than in CAE; seizures are also provoked by drowsiness. Seizure duration in JAE is 5–30 s, during which patients can usually respond to commands but cannot perform complex tasks (Trinka et al., 2004; Beghi et al., 2006). Similar to CAE, absence seizures in JAE are accompanied by an unreactive facial expression and motor automatisms. With seizure offset, EEG activity immediately returns to normal (Beghi et al., 2006; Elmali et al., 2020). Absence status epilepticus, generalized tonic-clonic seizures, and myoclonic seizures occur in 20%, >90%, and 0% of patients, respectively (Agathonikou et al., 1998; Trinka et al., 2004).

Children with JAE develop without major cognitive impairment; however, learning problems, ADHD, depression, and anxiety are prominent comorbidities (Marini et al., 2004; Henkin et al., 2005; Prassouli et al., 2007; Asadi-Pooya et al., 2013; Abarategui et al., 2018; Asadi-Pooya and Homayoun, 2020). JAE is usually drug-responsive, though some patients require lifelong treatment (Trinka et al., 2004; Table 1). Ethosuximide as a monotherapy is not recommended due to high likelihood of generalized tonic-clonic seizures for which ethosuximide is ineffective; rather, broad-spectrum antiseizure medications for generalized epilepsies should be used (Trinka et al., 2004; Vorderwülbecke et al., 2017; Healy et al., 2018; Hirsch et al., 2022).

2.3. Juvenile myoclonic epilepsy

2.3.1. Epidemiology—Juvenile myoclonic epilepsy (JME) is common, with a prevalence of 1–3 per 10,000 people, accounting for 9.3% of all epilepsies (Juul-Jensen and Foldspang, 1983; Syvertsen et al., 2015; Syvertsen et al., 2017). The typical age at onset is 10–24 years, females are represented slightly more than males, and 5–15% of cases evolve from CAE (Juul-Jensen and Foldspang, 1983; Wirrell et al., 1996; Syvertsen et al., 2015; Syvertsen et al., 2017).

2.3.2. Features and natural history—JME clinical features include myoclonic seizures, usually occurring within the first hour of awakening. Seizures can be triggered by sleep deprivation (Hirsch et al., 2022). Electroencephalography features 3–3.5-Hz interictal spike-and-wave and polyspike-and-wave signatures (Elmali et al., 2020). Generalized tonic-clonic seizures (>90% of patients) are common, and febrile seizures occur in 4–5% of patients (Jain et al., 1998). In one third of patients, absence seizures occur, less than daily, with subtle impairment to awareness (Panayiotopoulos et al., 1989a; Yacubian, 2017).

Patients with JME develop typically, and intellectual disability is rare (Wandschneider et al., 2012; Iqbal et al., 2015; Abarrategui et al., 2018; Sezikli et al., 2018; Almane et al., 2019; Chawla et al., 2021). Some cognitive domains might be altered (e.g., executive function), and depression, anxiety, impulsivity, and social or psychiatric problems are common (Wandschneider et al., 2012; de Araujo Filho and Yacubian, 2013; Iqbal et al., 2015; Abarrategui et al., 2018; Sezikli et al., 2018; Almane et al., 2019; Syvertsen et al., 2019; Gama et al., 2020; Taura et al., 2020; Chawla et al., 2021). JME is responsive to anti-seizure medications in 65–92% of patients, though it often requires lifelong treatment (Yacubian, 2017; see Table 1). Myoclonic seizures may be more difficult to control than generalized tonic-clonic seizures: sodium channel blockers often aggravate both myoclonic and absence seizures in JME, and lamotrigine (which has multiple mechanisms of action, including sodium channel blockade) may aggravate myoclonic seizures in some patients (Geithner et al., 2012; Senf et al., 2013; Höfler et al., 2014; Yacubian, 2017; Stevelink et al., 2019; Zhang et al., 2019; Pietrafusa et al., 2021; Hirsch et al., 2022). The mechanisms by which sodium channel blockers exacerbate seizures are not well understood but may relate to inhibition of Na_v1.1 channels on GABAergic neurons implicated in generalized epilepsies (Catterall, 2014).

2.4. Generalized tonic-clonic seizures alone

2.4.1. Epidemiology—Generalized tonic-clonic seizures alone (GTCA) accounts for one-third of all adolescent-onset IGEs (Vorderwülbecke et al., 2017). The typical age at onset is 10–25 years, later than in JAE or JME (Beghi et al., 2006; Vorderwülbecke et al., 2017).

2.4.2. Features and natural history—GTCA clinical features include generalized tonic-clonic seizures and interictal 3–5.5-Hz generalized spike-and-wave or polyspike-and-wave signatures. Seizures usually occur within 2 hours of awakening but can happen at other times. Seizures are infrequent, occurring yearly or less, and are sometimes triggered by sleep deprivation, fatigue, or alcohol (Holtkamp et al., 2014).

Patients' cognition is grossly normal interictally; however, specific cognitive domains might be altered (e.g., attention or decision-making) (Abarrategui et al., 2018). Antiseizure medications, including valproic acid, lamotrigine, levetiracetam, topiramate, perampanel, and zonisamide, are used to treat seizures in GTCA (Beydoun and D'Souza, 2012; Rohrer et al., 2016). Many patients with GTCA require medication throughout their adult lives.

In summary, IGEs share key clinical features, including seizures with impaired awareness and increased susceptibility surrounding sleep-wake transitions. While most IGEs are managed with medications, current therapies have off-target effects (for instance, sleep perturbations), and a substantial fraction of cases are refractory to treatment, motivating the pursuit of mechanistic investigations in animal models that will aid the discovery of more effective drug targets.

The factors that provoke seizures provide valuable clues to underlying mechanisms. For example, absence seizures provoked by hyperventilation suggest pH-dependent processes,

and the association between drowsiness and seizure susceptibility may be linked to high adenosine tone and low acetylcholine and norepinephrine levels during these periods. In addition, to understand why some drugs are ineffective or exacerbate a particular seizure type, it is important to investigate their effects on specific cells and brain circuits. In the sections that follow, we highlight recent developments linking clinical observations with mechanistic studies.

3. Contrasts and parallels between IGEs and DEEs

DEEs are a heterogeneous group of disorders characterized by early-onset, often severe seizures and developmental impairment, and encephalopathy that tends to worsen with increasing seizure burden (reviewed in Guerrini et al., 2023). While some DEEs are caused by focal structural changes and have a largely focal EEG pattern, others include generalized seizure types and display generalized interictal patterns on EEG, consistent with GGEs (Wirrell et al., 2022; Guerrini et al., 2023).

The latest ILAE Task Force on Nosology and Definitions for epilepsy syndromes defined 18 syndromes as DEEs, subcategorized by age of onset (Guerrini et al., 2023; for detailed review, see Guerrini et al., 2023). Here we highlight those DEEs that meet criteria for GGE (such as epilepsy with myoclonic-atonic seizures), and also those that prominently feature generalized seizure types (such as Dravet syndrome), because they are the most likely to have any mechanistic overlap with IGEs. There is also genetic overlap between DEEs with generalized features and IGEs, including mutations in *GABRA1* and *GABRG2* associated with both familial IGE syndromes and Dravet syndrome (Helbig, 2015).

The neonatal- and infantile-onset DEEs with generalized features include clinically defined syndromes with multiple etiologies (such as early infantile developmental and epileptic encephalopathy (EIDEE) or Dravet syndrome), and a growing list of etiology-specific syndromes such as those with an identified genetic cause (i.e., *KCNQ2* and glucose transporter-1 deficiency (Glu1DS)). Childhood-onset DEEs with generalized features include clinically defined syndromes with multiple etiologies (Lennox-Gastaut Syndrome (LGS), epilepsy with myoclonic-atonic seizures, spike-and-wave activation in sleep). DEEs with generalized features that present at variable ages include progressive myoclonic epilepsy, which has multiple genetic and metabolic causes such as mitochondrial disease (myoclonus epilepsy and ragged red fibers), and lysosome- and glycogen-storage disorders (including Lafora disease and Unverricht-Lundborg disease). This is not an exhaustive list of the DEEs, and we direct the reader to a recent review for a more detailed enumeration of the genetic associations identified so far (Guerrini et al., 2023).(Guerrini et al., 2023).

3.1. Epidemiology

The true prevalence of DEEs is not well established, although it has been estimated that the overall annual incidence of monogenic epilepsies is around 1 per 2100 live births (Symonds et al., 2019; Aledo-Serrano et al., 2021).

3.2. Features and natural history

DEE clinical features encompass a broad spectrum of severity, seizure types, and developmental comorbidities. Currently, most therapies for DEEs target individual symptoms such as seizures and not the underlying disease mechanisms (with a few notable exceptions, such as the ketogenic diet for Glu1DS). By contrast with IGEs, which are largely drug-responsive, seizure control is elusive for many individuals with DEEs, even when anti-seizure medications are optimized for the underlying etiology. First-line treatments for IGEs such as ethosuximide, valproate, and lamotrigine are often ineffective in DEEs and can cause neuropsychiatric side effects, adding to comorbid developmental impairment (Strzelczyk and Schubert-Bast, 2022). By contrast, there are medications with specific indications for DEEs that are not typically used in IGEs, including cannabidiol and fenfluramine for Dravet syndrome, and ganaxolone for CDKL5 (Elkommos and Mula, 2022). Despite combination therapy (i.e., >50% of patients with LGS take three or more anti-seizure medications concurrently (Spoor et al., 2021), many patients with DEEs still experience untreated symptoms and turn to nonspecific treatments such as the ketogenic diet (Sondhi et al., 2020). In DEE patients in whom seizure control is achieved, developmental impairments and other comorbidities often remain severe (McTague et al., 2016; Stamberger et al., 2016).

4. Animal models of generalized epilepsy

Foundational work on mechanisms of GGEs was made possible by studying animal models with generalized SWDs and behavioral arrest that recapitulate absence epilepsy. SWDs are an electrographic pattern composed of a bilaterally synchronous spike and after-going slow wave; in humans, absence seizures are associated with 2.5–5 Hz SWDs while in animal models of IGE, SWDs tend to occur faster, at 5–11 Hz (Jafarian et al., 2020; Hirsch et al., 2021; Hirsch et al., 2022).

Animal models of IGEs were initially obtained by inbreeding rat and mouse strains and screening them for features of IGEs, including absence-type seizures and SWDs. The resulting models include Genetic Absence Epilepsy Rat from Strasbourg (GAERS, Marescaux et al., 1992), Wistar Albino Glaxo Rijswijk rats (WAG/Rij), *tottering*, *lethargic*, *coloboma*, and *stargazer* (reviewed in Crunelli and Leresche, 2002b, and in Maheshwari and Noebels, 2014). Eventually, the disease-conferring loci were identified in these strains by forward genetics. As molecular biology technologies advanced and reverse genetics became possible, epileptic mice (and, to a lesser degree, rats) were generated with targeted mutations in pathways implicated in IGEs (for example, *Gabrg2* missense (R43Q) mutation, encoding the GABA-A γ_2 subunit (Petrou and Reid, 2012); *Gabra1* knock-out, encoding the GABA-A α_{1A} subunit (Arain et al., 2012); *Cacna1g* knock-out, encoding the T-type Ca²⁺ channel (Kim et al., 2001); *Hcn2* missense (R591E, T592A) mutation, ablating cAMP sensitivity (Hammelmann et al., 2019). These models reproduce the seizure semiology (behavioral arrest) and electroencephalographic pattern (generalized SWDs) common to CAE and JAE, although the SWD frequency is faster in rodents than in humans, and persists beyond the development window seen in humans.

Nevertheless, important gaps remain in animal models of GGEs. The rodent models of JME (Ding and Gallagher, 2016; McCarthy et al., 2020) and GTCA (Harada et al., 2013) have not been as exhaustively characterized as the absence epilepsy models, and could yield additional important insights into how pathogenic mechanisms differ between absence and convulsive seizures. Moreover, developing animal models of DEEs is challenging due to the sheer number and variety of insults that are known to cause clinical disease. For example, Dravet syndrome can be caused by thousands of mutations in numerous genes (He et al., 2022). A Dravet syndrome model induced by a mutation in *Scn1a* may not translate to the same clinical syndrome as induced with a *Gabra1* mutation and vice versa (He et al., 2022). Many etiology-specific DEEs now have animal models in which patient-derived mutations have been introduced, but the resulting animal phenotype and seizure repertoire do not always phenocopy the human disease (for review of genetic models see Wang and Frankel, 2021).

5. Neuroanatomical basis for the role of the thalamus in generalized epilepsy

Here we summarize the functional neuroanatomy of the thalamus as it relates to normal physiology and its disruption in GGEs. We describe the intrinsic properties of thalamocortical (TC) and nucleus reticularis thalami (nRT) neurons and their modulation by larger regulatory loops connecting thalamus and cortex.

5.1. Neuronal components of intrathalamic circuits underlying adaptive and maladaptive rhythmic oscillations

There are two functionally distinct populations of neurons in the thalamus implicated in generalized epilepsy: TC neurons and nRT neurons.

TC neurons are intrinsically oscillatory, due in part to the alternating activation and deactivation of hyperpolarization-activated cyclic nucleotide-gated cation channels (HCN) and TWIK-related acid-sensitive K^+ (TASK) channels (Meuth et al., 2006), interacting with T-type Ca^{2+} channels (Coulter et al., 1989; Tisone et al., 2021). TC neurons in the ventrobasal (VB) thalamus are glutamatergic projection neurons that innervate neocortical neurons in layer 4 and receive glutamatergic input from corticothalamic (CT) neurons in neocortical layers 5 and 6 (Sherman and Guillery, 1996; Fig. 2A and discussed below). The glutamate signal from CT neurons acts on two types of receptors expressed on the TC neurons' surface: ionotropic glutamate receptors are essential to the flow of information through the thalamocortical circuit (Koerner et al., 1996; Paz et al., 2011; Barad et al., 2012), and metabotropic glutamate receptors contribute to the regulation of the firing mode of TC neurons (McCormick and von Krosigk, 1992; Bertaso et al., 2008; Cheong et al., 2009).

The activity of TC neurons is influenced by GABAergic inputs from nRT neurons, and $GABA_A$ and $GABA_B$ receptor activation in the thalamocortical circuit has long been implicated in the generation of SWDs (Coulter et al., 1990b; Hosford et al., 1997; Kim et al., 1997).

The nRT, composed of GABAergic neurons spatially arranged as a capsule around and between projection nuclei, forms an important input to TC neurons (Fig. 2A). Like TC neurons, nRT neurons are also intrinsically oscillatory (Steriade et al., 1987; Fuentealba et al., 2005) due to the presence of low-voltage-activated T-type Ca^{2+} channels, Ca^{2+} -activated small conductance potassium (SK) channels (Cueni et al., 2008), and Ca^{2+} -activated cation TRPM4 channels (O'Malley et al., 2020).

TC and CT neurons send glutamatergic collaterals to the nRT (Fig. 2A), which also expresses both ionotropic and metabotropic glutamate receptors. The inhibition caused by the CT->nRT->TC pathway is referred to as feedforward inhibition, whereas inhibition from the TC->nRT->TC pathway is referred to as feedback inhibition (for review see Paz and Huguenard, 2015a). Fast AMPA-receptor (AMPA)-mediated currents are critical for CT-nRT-TC feedforward inhibition. Evidence for the importance of this pathway in epilepsy comes from the *stargazer* mouse, which lacks an auxiliary AMPAR subunit and displays absence epilepsy (Bats et al., 2007; Lacey et al., 2012). AMPARs that include the GluR4 subunit are particularly enriched on nRT neurons, opposite CT presynaptic elements, and loss of GluR4 is sufficient to cause loss of CT-nRT-TC feedforward inhibition and absence epilepsy (Paz et al., 2011). Both feedforward and feedback inhibition can result in a powerful GABAergic input to TC neurons, promoting a recurrent cycle of hyperpolarization followed by a burst of Na^+ action potentials; the mechanisms underlying burst firing and their disruption in SWDs are discussed below.

5.2. Tonic and burst firing in the intrathalamic circuit

TC and nRT neurons operate in two modes: burst firing (predominant during sleep, seizures, and anesthesia) and tonic firing (during wakefulness).

The tonic firing mode allows TC neurons to integrate and relay activity between cortical regions and the subcortical regions to which they are connected (Whitmire et al., 2021). Tonic firing occurs when the TC membrane is at resting potential (~60 mV). With T-type Ca^{2+} channels inactivated, the TC neurons generate Na^+ -channel-dependent action potentials in response to excitatory stimuli.

The burst firing mode allows the thalamus to entrain wide regions of the brain into hypersynchronous activity, as occurs normally during sleep. Burst firing occurs when the TC membrane is hyperpolarized (Sorokin et al., 2017; Whitmire et al., 2021). Hyperpolarization of TC neurons, which can result from GABAergic inputs from the nRT, relieves the inactivation of T-type Ca^{2+} channels and activates HCN channels. These conditions permit TC cells to fire a rebound Ca^{2+} spike with an overriding burst of high-frequency Na^+ -mediated action potentials (Fig. 2B, C). The volley of high-frequency action potentials from TC neurons excites distant targets throughout the brain. CT projection neurons then send glutamatergic inputs back to the thalamus (Fig. 2A).

Key to the burst firing mode in both TC and nRT neurons, T-type Ca^{2+} channels voltage-activate in response to low voltage depolarization (-50 mV), inactivate with further depolarization (-30 mV), and require hyperpolarization (-90 mV and below) to relieve voltage inactivation, as was demonstrated in acutely dissociated rat TC neurons (Coulter

et al., 1989). Enhanced T-type Ca^{2+} currents are implicated in many rodent models of absence epilepsy. For example, T-type Ca^{2+} current is augmented in nRT in GAERS rats, resulting in more powerful hyperpolarizing output onto TC neurons (Tsakiridou et al., 1995). T-type Ca^{2+} current is also augmented in the *tottering*, *stargazer*, *lethargic*, *coloboma*, and *Cacna1a*^{-/-} models (Ernst et al., 2009). Thus, monogenic mouse models of epilepsy, such as those with deficient or dysfunctional P/Q Ca^{2+} channels ($\text{Ca}_v2.1 \alpha_{1A}$), may cause epilepsy via a compensatory increase in T-type Ca^{2+} channels (Zhang et al., 2002; Song et al., 2004; Zhang et al., 2004). Deletion of functional α_{1G} -containing T-type Ca^{2+} channels on TC neurons is protective against SWDs (Kim et al., 2001; Miao et al., 2020), and overexpression of this subunit is sufficient to cause an absence epilepsy phenotype in mice (Ernst et al., 2009). T-type Ca^{2+} channels are one of the targets of ethosuximide, which is effective in treating absence seizures in IGE (Huguenard, 1999; Table 1).

In spite of these observations, other lines of evidence call into question the primacy of TC T-type Ca^{2+} channels: 1) ethosuximide is a relatively weak T-type Ca^{2+} channel blocker (Coulter et al., 1990a), and it has been suggested that its anti-absence effects could be mediated by modifying other channels (Crunelli and Leresche, 2002a); 2) a recent absence epilepsy model generated by deletion of P/Q channels in adult mice failed to show the expected increases in T-type Ca^{2+} current or burst-firing (Miao et al., 2020); 3) It has been observed that some TC neurons can remain in tonic firing mode during SWDs (Pinault et al., 1998; McCafferty et al., 2018).

In the GAERS rat model of absence epilepsy, nRT neurons fire in burst mode during SWDs, as measured using either intracellular (Slaght et al., 2002) or extracellular recordings (McCafferty et al., 2018). The burst firing in nRT neurons is critical for SWDs *in vivo* (McCafferty et al., 2018) and in SWD-like activity in brain slices from GAERS rats (Cain et al., 2018). Burst firing mode is also observed in TC neurons in the motor thalamus during SWDs in GAERS rats (Paz et al., 2007). However, burst firing in single TC neurons is not commonly observed during SWDs in the VB thalamus (McCafferty et al., 2018). Although the results between brain regions seem variable as to the proportion of burst firing in TC cells, there is consensus that universal burst firing in every single TC neuron is not necessary for the expression of SWDs, but that switching the firing of a population of TC neurons from synchronized to non-synchronized is sufficient to interrupt SWDs in both the rat and mouse models of absence epilepsy (Sorokin et al., 2017).

Activation of GABA receptors in TC neurons can be pro-epileptic, because GABA_A and GABA_B receptors hyperpolarize TC neurons with differing time courses and promote post-hyperpolarization rebound firing (Kim et al., 1997). Consistent with this idea, enhancing the bioavailability of endogenous GABA with reuptake inhibitors (D'Amore et al., 2015), loss-of-function mutations in *Slc6a1* (a mutation encoding GABA transporter GAT-1 and implicated in a generalized DEE, epilepsy with myoclonic-atonic seizures) (Pirttimaki et al., 2013), or administering GABA pro-drug GHB (Godschalk et al., 1977; Liu et al., 1991), can all provoke SWDs. Burst firing in TC neurons is key to understanding the mechanism by which clinical GABA reuptake inhibitors, such as tiagabine and vigabatrin, which provoke absence status in patients with IGE (Knake et al., 1999).

In contrast to the seizure-promoting effects of GABA receptor activation on TC neurons, GABA_A receptor activation in the nRT is antiepileptic. This effect probably reflects lateral inhibition between nRT neurons, which desynchronizes GABAergic output to TC neurons and decreases the duration of bursts in thalamic slices (Sohal et al., 2000; Beenhakker and Huguenard, 2009); loss of lateral inhibition is a putative pathogenic mechanism of GGE in *Scn8a*^{+/-} mice (Makinson et al., 2017). Indeed, overactivity of nRT cells, which results in increased GABAergic output to TC neurons, is implicated in several animal models of IGE and DEE (GAERS Tsakiridou et al., 1995, and Bessaih et al., 2006, *Scn1a*^{+/-} Ritter-Makinson et al., 2019). Thus, the therapeutic effect of benzodiazepines such as clonazepam on absence seizures is thought to be due to enhancement of α 3-containing GABA_A receptor current within the nRT (Sohal et al., 2003; Schofield et al., 2009).

The difficulty in selectively promoting anti-epileptic vs. pro-epileptic actions of GABA within the thalamus underlines the need for cell-type specific targeting of therapeutics in GGEs.

5.3. The role of thalamo-cortico-thalamic loops in spike-and-wave discharges

SWDs result from synchronized oscillations in thalamo-cortico-thalamic loops. Bursting TC neurons cause excitatory postsynaptic potentials in neocortical neurons, which generates a time-locked extracellular negative potential seen as a “spike” on a surface electroencephalogram (Staak and Pape, 2001; Terlau et al., 2020). The spike is followed by a slow wave representing complex intracortical polysynaptic excitatory and inhibitory postsynaptic potentials (Terlau et al., 2020).

Controversy has persisted for decades regarding the brain region—cortex or thalamus—that instigates absence seizures in susceptible animals (Meeren et al., 2002; Manning et al., 2004; Polack et al., 2009). Multi-electrode arrays in GAERS rats demonstrated that precursor activity could be recorded in the somatosensory cortex, before the appearance of SWDs (Seidenbecher et al., 1998). Similar studies in WAG/Rij rats showed that SWDs appear earliest in the peri-oral region of the somatosensory cortex, then rapidly generalize to other cortical regions (Meeren et al., 2002; Polack et al., 2009). These observations gave rise to the “cortical focus theory,” which has accumulated additional evidence from network approaches such as cross-correlation, phase synchronization analysis and Granger causality (Luttjohann and van Luijtelaar, 2012; Luttjohann et al., 2013; Sysoeva et al., 2016; Luttjohann and Pape, 2019). The early involvement of the somatosensory cortex was demonstrated for both absence seizures and myoclonic seizures, in a mouse model of JME (Ding and Gallagher, 2016).

One caveat to the interpretation of cortically recorded SWDs is that standard electrode arrays lack the spatial resolution to resolve which neuronal population(s) generate the signals. Thus, local field recordings could reflect the postsynaptic responses of layer IV neurons (the target of thalamocortical inputs), layer II/III (thalamocortical and corticocortical inputs), or layer VI neurons (which receive a mix of both thalamocortical and intracolumnar cortical input), while unit activity recordings could indicate intracortical (layer II/III) or corticofugal (layer V & VI) neuronal spikes. Thus, single-electrode techniques cannot distinguish a cortical focus firing autonomously vs a cortical focus driven

by thalamocortical stimuli. Efforts to disentangle these possibilities might be enhanced by current source density analysis. Terlau et al. (2020) have recently described depth profiles in the somatosensory cortex and mapped current sources and sinks during SWDs in GAERS rats using high-density multi-contact electrodes. The same approach could be used to characterize the precursor activity and initiation of SWD in the somatosensory cortical focus (Luttjohann and van Luijtelaar, 2012; Luttjohann and Pape, 2019). Furthermore, the cortical focus theory should be evaluated in animal models of generalized DEEs, as the existence of a cortical focus, if confirmed, could inform clinical approaches to treatment such as placement of electrodes for responsive neurostimulation or other neuromodulatory interventions.

The current consensus is that thalamus and cortex are both needed for the expression of generalized seizures. Intervening at any point in the thalamo-cortico-thalamic loop is likely to interrupt SWDs as long as a critical mass of neurons is affected. Thus, absence seizures in animal models can be attenuated by ablative lesions of the nRT (Banerjee and Snead 3rd, 1994) or an infusion of glutamate receptor antagonists into thalamic relay nuclei (Seidenbecher and Pape, 2001; Terlau et al., 2020), but also by lesioning the ipsilateral cortex (Scicchitano et al., 2015) or silencing it with lidocaine, tetrodotoxin or cortical spreading depression (Vergnes and Marescaux, 1992; Sitnikova and van Luijtelaar, 2004; Polack et al., 2009). Wherever the seizures start, they almost instantaneously generalize throughout the bilateral thalamo-cortico-thalamic loops and cause unconsciousness by disrupting the tonic firing mode required for awake awareness (Paz and Huguenard, 2015a; Paz and Huguenard, 2015b; Sorokin et al., 2017).

Thus, the thalamus remains a potent therapeutic target because it is a key gatekeeper to seizure maintenance, whether the SWDs started there or not. Indeed, the first real-time optogenetic disruption of thalamic output during SWDs showed that switching the output of the thalamus from bursting to tonic firing mode is sufficient to instantaneously interrupt SWDs and behavioral absences in multiple rodent models of absence epilepsy (Sorokin et al., 2017). This on-demand switch in the firing mode of TC neurons can be achieved simply by manipulating the membrane voltage of TC neurons: by expressing the inhibitory halorhodopsin in TC neurons of various rodent models of absence epilepsy – WAG/Rij rats and stargazer mice – were able to cause a short hyperpolarization of TC neurons with a brief yellow light pulse delivered in the thalamus with an optical fiber, which caused rebound burst firing in TC neurons and an SWD accompanied by a behavioral arrest. In contrast, by expressing the excitatory stabilized step function opsin in TC neurons in these rodent models, Sorokin et al. (2017) were able to depolarize TC neurons with a brief blue light pulse delivered to the thalamus, which caused a switch from bursting to tonic firing mode, instantaneously ending the SWD and the associated behavioral absence. Thus, depolarizing TC neurons is a powerful way to interrupt the high-frequency burst firing in TC neurons, which is due to the fact that depolarization inactivates T-type Ca^{2+} channels in these neurons. These first real-time optogenetic perturbations of thalamic firing mode revealed that synchronized output from the thalamus is *necessary* to maintain an SWD. Whether single TC neurons fire in bursts or not, their collective output from the thalamus is “bursty” during SWDs because the output of a mass of TC neurons is highly synchronized, which is sufficient to cause an SWD in mice prone to absence epilepsy.

Emerging literature is beginning to clarify how mode switching is accomplished endogenously and where those processes can go wrong and contribute to GGEs.

6. Intrinsic properties and synaptic inputs that influence firing states of intrathalamic circuits in healthy states and epilepsy

Several factors regulate thalamic neuron membrane potential and, thus, their firing mode.

6.1. Metabotropic glutamate receptors in TC and nRT neurons

Metabotropic glutamate receptors on TC neurons can switch these cells' firing mode from tonic to burst firing. For example, group I metabotropic glutamate receptors (mGluR1 and mGluR5) activate phospholipase C (PLC), which activates protein kinase C (PKC) and modifies the function of T- and L-type Ca^{2+} channels in TC neurons (Cheong et al., 2009). In the absence of this downstream signaling, PLC $\beta 4$ knockout animals demonstrate increased voltage sensitivity of T-type Ca^{2+} channels, more easily generate bursts of spikes, and develop spontaneous absence seizures, which can be rescued by PKC activation (Cheong et al., 2009). Disrupting the subcellular targeting of mGluR7, a group III metabotropic glutamate receptor coupled to the $\text{G}_{i/o}$ second messenger system, and expressed presynaptically in both nRT and TC neurons, decreases neurotransmitter release at multiple synapses in the intrathalamic circuit, and also leads to absence epilepsy (Okamoto et al., 1994; Bertaso et al., 2008; Tassin et al., 2016).

6.2. Hyperpolarization-activated cyclic nucleotide-gated nonselective (HCN) in TC neurons

HCN channels carry a nonselective (depolarizing) current that activates at hyperpolarized membrane potentials and inactivates at depolarized membrane potentials (He et al., 2014). This property makes HCN channels well-positioned to stabilize the resting membrane potential of TC neurons. In the presence of cAMP, the voltage sensitivity of the channel shifts to more depolarized potentials, such that the TC neuron resting membrane potential is maintained above the threshold for T-type Ca^{2+} channel deinactivation, favoring tonic firing over burst firing (Budde et al., 2005; Zobeiri et al., 2019).

Hyperpolarization-activated cAMP nucleotide-gated (HCN) channels have been implicated in generalized epilepsy by genetic and pharmacologic studies (Ludwig et al., 2003; Budde et al., 2005; David et al., 2018; Hammelmann et al., 2019; Nishitani et al., 2019; Iacone et al., 2021). HCN channels' cAMP sensitivity provides a candidate mechanism for state-dependent susceptibility to SWDs during drowsiness and sleep, as neuromodulators linked to wakefulness (norepinephrine $\alpha 2$ and β receptors, acetylcholine mAChRs 2/4) and sleep pressure (adenosine receptors A1, 2A/B, 3) are coupled to G_i and G_s proteins that control adenylyl cyclase activity. HCN channels are sensitive to intracellular pH, and their voltage sensitivity is shifted by lamotrigine (Table 1). HCN channels make an attractive target for rational drug therapy if their modulation can reduce burst-firing in drowsy states without disrupting neuronal function during alert wakefulness.

6.3. TWIK-related potassium (TREK) and TWIK-related acid-sensitive potassium (TASK) channels in TC neurons

Together with HCN channels, TREK and TASK channels on TC neurons contribute to resting membrane potential and facilitate switching between tonic and burst firing modes (Meuth et al., 2006; Bista et al., 2012; Bista et al., 2015). TASK and TREK channels contribute to the background potassium leak current (Enyedi and Czirjak, 2010). When this leak current is inhibited, as by Gq-coupled receptors (Bista et al., 2012, Bista et al., 2015), resting membrane potential shifts upward, and T-type Ca^{2+} channels are maintained in an inactivated state, thus favoring tonic firing. Tonic firing is thereby favored over burst firing (Meuth et al., 2006; Bista et al., 2012). Conversely, under conditions of extracellular or intracellular alkalinization, TASK channels open (Niemeyer et al., 2010), hyperpolarizing the resting membrane potential and favoring burst firing. TASK channel mutations have not, to our knowledge, been reported in generalized epilepsy cohorts (Kananura et al., 2002), but TASK and TREK channels should be explored as potential therapeutic targets for much-needed treatments.

6.4. Ca^{2+} -activated potassium channels in nRT neurons

Ca^{2+} activated small conductance K^{+} currents (SK) in nRT reticular thalamic neurons regulate GABAergic input to TC neurons (Ying and Goldstein, 2005; Kleiman-Weiner et al., 2009), thus influencing firing mode in the thalamocortical circuit (Wong et al., 2013). SK channels are critical for the afterhyperpolarization (Debarbieux et al., 1998; Zaman et al., 2011) in nRT neurons following each low threshold Ca^{2+} spike. This afterhyperpolarization relieves voltage inactivation of T-type Ca^{2+} channels, restoring conditions required to initiate the next burst (Debarbieux et al., 1998, Zaman et al., 2011). The role of these channels in epilepsy pathogenesis is less well understood, but diminished SK current and hyperactive nRT output are seen in brain slices from mice with a model of Dravet syndrome induced by *Scn1a* deficiency (Ritter-Makinson et al., 2019). Further study is needed on the role of SK currents in epilepsy to inform anti-seizure medication development.

6.5. Non-neuronal modulators of intrathalamic circuits: thalamic glia

Astrocytes are an important non-neuronal cell type that regulates ion concentrations and neurotransmitter levels, stores glycogen, and provides metabolic support for neurons during high energy demands of seizures (Walz, 2000; Pirttimaki et al., 2013; Crunelli et al., 2015; Philippot et al., 2021; Dienel et al., 2023). More than just “support cells,” astrocytes demonstrate intrinsic oscillations, carry Ca^{2+} waves, release signals of their own, and have been implicated in the pathogenesis of some types of epilepsy (Parri and Crunelli, 2001; Cho et al., 2022).

Of particular relevance to the pathophysiology of GGEs, thalamic astrocytes are key modifiers of thalamo-cortico-thalamic oscillations; they regulate glutamate and GABA synthesis (Bryant et al., 2009) and clearance (Beenhakker and Huguenard, 2010; Pirttimaki et al., 2013), thus shaping the kinetics of synaptic neurotransmission. GABA reuptake, in particular, has been implicated in absence epilepsy: for instance, reduced GAT-1 function in thalamic astrocytes prolongs neuronal GABA_A and GABA_B receptor activation and promotes SWDs (Pirttimaki et al., 2013).

Interactions between astrocytes and neurons in the development of generalized seizures could be bidirectional. For instance, in various IGE models (Akbar et al., 1998; Dutuit et al., 2000; Dutuit et al., 2002; Cavdar et al., 2019; Kose et al., 2022; Thompson et al., 2022), as well as in some models of DEEs (Scn8a epileptic encephalopathy Thompson et al., 2022, Dravet syndrome Uchino et al., 2023), thalamic astrocytes proliferate and take on a reactive phenotype – downregulating glutamate and GABA transporters, and decreasing potassium uptake – while spontaneous seizures develop. Experimental work has now extended these observations and established a role for thalamic astrocytes in thalamocortical circuit physiology and generalized seizures. Thus, optogenetic depolarization of astrocytes in the VB relay thalamus prolongs SWDs in GAERS rats (Ozgur et al., 2022), whereas ablating glial function (with fluorocitrate or a glutamine shuttle toxin) disrupts intrathalamic oscillations elicited in brain slices from normal rats (Yang and Cox, 2011).

Electrolyte homeostasis by astrocytes may contribute to seizures in the Noda epileptic rat, a model of GTCA that shows decreased astrocyte expression of *Kir4.1* in thalamus and amygdala (Harada et al., 2013). In the special case of Lafora disease, a rare glycogen storage disorder causing progressive myoclonus epilepsy with neurodevelopmental impairment, toxic accumulation of glycogen in astrocytes results in impaired neurotransmitter clearance (Munoz-Ballester et al., 2016; Munoz-Ballester et al., 2019), and the disease can be rescued by astrocyte-specific deletion of glycogen synthase (Duran et al., 2021); whether manipulations of thalamic astrocytes are sufficient to rescue the epileptic phenotype is not known and should be investigated.

In conclusion, thalamic astrocytes represent an important therapeutic target for GGEs that is not currently targeted by existing anti-seizure medications, and should be researched further.

7. Neuromodulation in the thalamus

Thalamic modulation occurs through neuromodulators released by a heterogeneous group of afferents, and through systemic drivers (Sherman and Guillery, 1998; Varela, 2014). Examples of thalamic modulators include local purinergic, afferent noradrenergic, cholinergic, serotonergic, dopaminergic, and histaminergic projections, and systemic carbon dioxide and blood pH (CO₂/pH), among others. Here, we focus on purinergic, noradrenergic, cholinergic, and CO₂/pH modulation (Fig. 3) due to their prominent role in vigilance state changes controlled by the thalamus.

7.1. Norepinephrine

The adrenergic neuromodulator norepinephrine (NE) has potent and long-lasting effects on thalamocortical circuits and is released from NE axons to all thalamic nuclei (McCormick et al., 1991; Pérez-Santos et al., 2021) under conditions of alert wakefulness (Berridge et al., 2012). NE axons project from the locus coeruleus (LC) in the brainstem reticular formation (Simpson et al., 1997) and the A5 noradrenergic region in the brainstem (Swanson and Hartman, 1975; Morrison and Foote, 1986; Byrum and Guyenet, 1987; De Lima and Singer, 1987; Simpson et al., 1997; Vogt et al., 2008; Varela, 2014).

In acute thalamic slices from guinea pig and cat brains, norepinephrine application evokes a slow depolarization, reduced burst firing, and increased tonic activity by decreasing K^+ leak current (likely via TASK and TREK channels) and changing voltage sensitivity of I_h (likely reflecting HCN channels) which remains active at resting membrane potential and makes it more difficult to switch to burst mode (McCormick and Prince, 1988; Pape and McCormick, 1989; McCormick and Pape, 1990; Varela, 2014). In addition, norepinephrine increases synaptic strength in thalamocortical synapses (Varela, 2013).

In vivo, LC stimulation increases thalamocortical response to whisker stimulation (Moxon et al., 2007; Devilbiss and Waterhouse, 2011), and phasic LC stimulation sensitizes TC neurons to low-level stimuli (Devilbiss and Waterhouse, 2011). However, 20–40% of TC neurons show suppression of their response to whisker stimulation when LC is stimulated (Moxon et al., 2007; Devilbiss and Waterhouse, 2011), congruent with *in vivo* iontophoretic delivery of noradrenergic agonists inhibiting TC neurons (Grasso et al., 2006).

Taken together, these data support an antiepileptic role of endogenous NE, though this has not yet been exploited therapeutically (reviewed in Giorgi et al., 2004).

7.2. Acetylcholine

Acetylcholine, released by widespread nerve terminals arising from cell bodies in brainstem nuclei (pedunculo pontine nucleus (PPT), laterodorsal tegmentum (LDT), parabigeminal nucleus (PBG)), and the basal forebrain (BF), is high during alert wakefulness when absence seizures and SWDs are least likely to occur. Acetylcholine powerfully switches the firing patterns of thalamic neurons, inhibiting nRT neurons and activating tonic firing in TC projection neurons (Ben-Ari et al., 1976). Acetylcholine effects are mediated through Gq-coupled muscarinic M1 and M3 receptors, which close TASK and TREK channels (Bista et al., 2012), thus decreasing membrane conductance and depolarizing TC cells above the threshold of T-type channel deinactivation (McCormick and von Krosigk, 1992; Broicher et al., 2008). Importantly, acetylcholine mimics like pilocarpine (an M3 receptor agonist) provoke convulsive seizures (Levesque et al., 2021), but may exert contrasting effects on absence seizures due to intrathalamic signaling. Similar to norepinephrine, potential cholinergic antiepileptic mechanisms have not yet been explored therapeutically, but will have to be targeted to the appropriate patient population, and to the correct brain structure and cell type.

7.3. Adenosine

The purinergic neuromodulator adenosine regulates neural activity via four subtypes of G-protein coupled receptors. Adenosine is a low-energy metabolite released into the extracellular milieu by equilibrative and concentrative transporters (Lovatt et al., 2012; Wall and Dale, 2013) and under some conditions in the form of ATP by neurons (Pankratov et al., 2007) and glia (Newman, 2004; Pascual et al., 2005; Wall and Dale, 2013). A1 receptors are widely expressed in the brain, and A1R activation inhibits synaptic transmission and hyperpolarizes the membrane potential via the activation of K^+ channels (Dunwiddie and Masino, 2001; Borea et al., 2018; Wall et al., 2022).

Adenosine has a well-established role as an endogenous anti-epileptic neuromodulator in focal and convulsive seizures (Weltha et al., 2019). Adenosine accumulates during seizures and provides negative feedback to terminate bursts of neuronal activity and delay the next event (During and Spencer, 1992; Boison, 2008). The role of adenosine in the thalamocortical circuit and generalized seizures is less clear.

In the thalamus, A1Rs are highly expressed, particularly in the VB thalamus and the nRT, supporting a role for adenosine signaling in modulating oscillations in thalamic microcircuit and TC neurons. A recent study showed that endogenous adenosine is released during thalamic oscillations and acts via A1 receptors to feedback and reduce thalamic oscillatory activity in acute thalamic slices from mice (Wall et al., 2022).

However, adenosine also accumulates with sleep pressure and drowsiness, which may contribute to its observed pro-epileptic role in absence seizures. Exogenous agonists of adenosine receptors increase SWDs, while antagonists (i.e., caffeine) decrease SWDs in rodent models of absence epilepsy (Ilbay et al., 2001; Ates et al., 2004; Germé et al., 2015; Dede et al., 2019).

7.4. CO₂/pH

Voluntary hyperventilation provokes SWDs in 90% of patients with absence epilepsy and can be useful as a bedside diagnostic maneuver (Holowach et al., 1962; Panayiotopoulos, 1999; Caplan et al., 2008; Glauser et al., 2013; Caplan, 2015; Salvati and Beenhakker, 2019). This physiological response to hyperventilation is caused by the intimate relationship between carbon dioxide (CO₂) and blood pH. After CO₂ is formed as a waste product of aerobic respiration, CO₂ diffuses out of tissues into blood plasma and then red blood cells. Plasma CO₂ is in equilibrium with carbonic acid (H₂CO₃), bicarbonate ions (HCO₃⁻), and hydrogen ions (H⁺) (Svichar and Chesler, 2003). These free H⁺ ions determine blood plasma pH. Hyperventilation causes CO₂ plasma to drop, promoting the reverse formation of H₂CO₃ from HCO₃⁻ and H⁺ ions. This decrease of H⁺ ions in the blood is called respiratory alkalosis, which was shown to directly provoke SWDs in seizure-prone rats (Salvati et al., 2022). Importantly, absence seizures are induced by hyperventilation at a low metabolic state and not during exercise, during which increased breathing does not induce respiratory alkalosis (Esquivel et al., 1991; Salvati and Beenhakker, 2019).

There is well-established evidence that the thalamus is responsible for hyperventilation-induced absence seizures. In 1965, hyperventilation-induced SWDs were shown to be abolished by severing connections between the cortex and subcortical structures, and in 1967, the thalamus was shown to be the responsible pH sensor (Sherwin, 1965; Sherwin, 1967). TC neurons express HCN and TASK channels which are pH-sensitive (Soltesz et al., 1991; Meuth et al., 2006). Their respective currents, I_h and I_{TASK} , are blocked by extracellular acidification (Meuth et al., 2006; Biel et al., 2009). However, since the I_h is depolarizing (reversal potential is around -20 mV) and I_{TASK} is hyperpolarizing (reversal potential is around -70 mV), the net result of extracellular acidification is minimal, and neither the resting membrane potential nor firing properties of TC neurons change much during extracellular acidification (Meuth et al., 2006). Thus, it remains unclear how TC neurons are responsible for hyperventilation-induced absence seizures (Salvati and

Beenhakker, 2019). The possible proposed mechanisms include, but are not limited to, 1) unbalanced HCN and TASK1/3 channel expression in TC sub-populations (Salvati and Beenhakker, 2019); 2) differences in HCN and TASK 1/3 channel responses to extracellular alkalization (Salvati and Beenhakker, 2019); or 3) TC neurons' modulation by nitric oxide (NO) produced by vasculature in response to blood pH changes (Najarian et al., 2000; Yang and Cox, 2007; Yang and Cox, 2008), 4) cell-specific response of thalamic nuclei to hyperventilation (Salvati et al., 2022). Interestingly, the most recent study to date on thalamic pH-sensitivity determined that hypoxia-induced hyperventilation consistently activates neurons within the intralaminar nuclei (Salvati et al., 2022). pH effects may be particularly potent because the intralaminar nuclei – along with other nonspecific thalamic nuclei – have such broad connections throughout the cortex.

8. Inputs to the thalamus from the basal ganglia, cerebellum, and limbic system are disrupted in generalized epilepsy

The thalamus is extensively connected to other brain regions. Here we focus on the afferent connections that modulate the activity of the thalamus and are implicated in the pathogenesis of seizures (Fig. 3), as well as the outputs from the thalamus to cortex that form thalamo-cortico-thalamic loops which are activated in SWDs. Thalamic projections play an important role in numerous other cortical and subcortical circuits, which are beyond the scope of this review (reviewed in Castro-Alamancos and Connors, 1997, and Haber and McFarland, 2001).

8.1. The role of the basal ganglia in regulating thalamic circuits in generalized epilepsy

Parallel direct and indirect pathways through the basal ganglia form inhibitory inputs to TC neurons in circuits critical for the control of behavior and implicated in the pathogenesis of neurodegenerative, movement, and neuropsychiatric disorders (Graybiel, 2000). Within these circuits, inhibitory projections from the globus pallidus interna (GPi) and substantia nigra pars reticulata (SNr) are organized topographically in the thalamic mediodorsal (MD), ventral anterior (VA) and ventral lateral (VL) nuclei (Haber and Calzavara, 2009). In turn, TC neurons in MD innervate the prefrontal cortex, while TC neurons in VA and VL largely target motor and premotor cortices (Haber and Calzavara, 2009).

The role of increased inhibitory output from SNr to the thalamus in the pathogenesis of generalized seizures has been demonstrated in the GAERS rat model of absence epilepsy. In freely behaving GAERS rats, multi-unit recordings of SNr neurons revealed a change in firing properties time-locked with spontaneous SWDs, with an increase in firing rates and a switch to burst firing (Deransart et al., 2003). The impact of this change in the firing pattern of nigral neurons was further elucidated through recordings of ventral medial thalamic neurons in GAERS rats, which also show an increase in burst firing during SWDs, and demonstrate tonic firing with disruption of SWDs when glutamatergic signaling in SNr is blocked (Paz et al., 2007).

Alterations in basal ganglia-thalamic pathways have also been implicated in secondary seizure generalization in focal epilepsy (reviewed in Gale, 1992). Patients with recent

focal to bilateral tonic-clonic seizures have shown decreased pallidal–thalamic interaction through the indirect pathway in measures of resting state functional connectivity, compared with patients with exclusively focal seizures or only remote history of focal to bilateral tonic-clonic seizures (He et al., 2020).

8.2. Altered cerebellothalamic pathways in generalized epilepsy

The efferent pathways from the cerebellum form a major excitatory input to thalamic neurons, originating in the projection neurons of the deep cerebellar nuclei, and altered intrinsic cerebellar firing has been implicated in generalized epilepsy (reviewed in Streng and Krook-Magnuson, 2021). Cerebellar projection neuron activity is largely modulated by inhibitory input from Purkinje neurons. In rodent models of absence epilepsy, subsets of Purkinje neurons and cerebellar nuclei projection neurons show firing patterns phase-locked to SWDs (Kandel and Buzsáki, 1993; Kros et al., 2015), and genetic manipulation of cerebellar neurons can cause SWDs (Schwitalla et al., 2022). Pharmacologic or optogenetic modulation of cerebellar projection neurons has shown that decreasing cerebellar projection neuron output leads to an increase in SWDs, while conversely, increasing firing reduces SWDs (Kros et al., 2015; Eelkman Rooda et al., 2021).

8.3. Limbic-thalamic pathways implicated in generalized epilepsy

The epileptogenic role of limbic pathways involving the hippocampus and amygdala have been well-defined through clinical data and animal models of focal temporal lobe epilepsy but may also be disrupted in generalized epilepsy (reviewed in Nayel et al., 1991, Jefferys et al., 2012). Specifically, the connections between limbic cortical and subcortical regions with the nRT and limbic thalamic nuclei (including anterior and midline nuclei) are likely to be altered in some GGEs (reviewed in Onat et al., 2013).

The strongest evidence for the role of limbic pathways in generalized epilepsy comes from models of atypical absence epilepsy. This has been studied through a mouse model generated through a mutation in the R1 subunit of the metabotropic GABA_B receptor (Wang et al., 2009). SWDs were recorded in vivo from the midline thalamus, hippocampal CA1 and nRT; the nRT and midline thalamus were necessary for the generation of SWDs (Wang et al., 2009). Though not directly implicated in epileptogenesis, there is also evidence for disrupted limbic pathways in typical absence epilepsy models: data from WAG/Rij rats suggests a lower threshold to trigger the spread of SWDs to limbic pathways, and GAERS rats are resistant to secondary generalized seizures after amygdala kindling (E kazan et al., 2002; Tolmacheva et al., 2004). Further strengthening the link between alterations in the limbic system and generalized epilepsy is the presence of comorbid mood disorders. Neuropsychiatric comorbidities, including anxiety and depression, are common in patients with IGE, and anxious and depressive behaviors have also been described in corresponding animal models (Gruenbaum et al., 2021).

9. Mechanistic overlap between seizures and sleep

Additional clues to the pathogenesis of IGEs have come from studying the mechanisms of sleep. The thalamocortical circuits underlying abnormal rhythmic activity in generalized

seizures are also thought to generate the physiologic rhythms of sleep (reviewed in Beenhakker and Huguenard, 2009). Specifically, sleep spindles, brief bursts of rhythmic waveforms seen on scalp EEG during non-REM sleep, are generated through recurrent loops between the nRT and TC neurons (Steriade et al., 1985; Contreras et al., 1997). Disconnection experiments have demonstrated that similar to the pathways generating SWDs, the circuit connecting cortex, nRT, and thalamocortical neurons is necessary to generate synchronized spindles within the thalamus and cortex (Steriade et al., 1985, Contreras et al., 1997). Specifically, excitatory corticothalamic projections targeting the inhibitory nRT drive burst firing in thalamocortical neurons, and ablating the cortical input to the thalamus or the inhibitory nRT input to thalamocortical neurons disrupts spindle synchronization in healthy cats (Contreras et al., 1997).

In addition to sharing underlying pathways, sleep spindles and SWDs can arise in similar brain states. SWDs are often observed at sleep-wake transitions and can be seen in non-REM sleep, similar to the temporal pattern of sleep spindles (Kostopoulos, 2000). Further supporting the mechanistic links between sleep spindles and generalized seizures, there is evidence that spindle oscillations directly transform into SWDs in feline and rodent models of epilepsy (reviewed in Kostopoulos, 2000, Halasz et al., 2002). Lesional studies targeting these pathways have demonstrated that ablating the nRT and thalamocortical relay nuclei suppresses both sleep spindles and SWDs ipsilateral to the lesion (Meeren et al., 2009). Further work is needed to determine the relationship between sleep spindles and SWDs in humans, but there is evidence that sleep spindles are disrupted in patients with focal and generalized epilepsy (Drake et al., 1991; Tezer et al., 2014; Kramer et al., 2021).

Further supporting the link between sleep and epilepsy, sleep is often disrupted in patients with epilepsy, and anti-seizure medications commonly cause disordered sleep, including insomnia or daytime sleepiness (reviewed in Liguori et al., 2021). These differential effects on sleep do not clearly cluster by drug mechanism, and additional research is needed to identify potential mechanisms underlying these side effects and reduce or eliminate them in newer anti-seizure treatments. Data from functional imaging implicates alterations in thalamic metabolism in sleep disorders, including hypersomnia and insomnia (Nofzinger, 2005). Given the key role of the thalamus in both sleep and epilepsy, alterations in thalamocortical circuits likely play a role in these symptoms in patients with epilepsy.

10. Discussion: unresolved questions and opportunities

Our understanding of TC circuits in GGEs has been advanced by genetic, histologic, anatomic, electrophysiological, pharmacological, and molecular approaches, as summarized above. However, important questions for the field remain unanswered.

10.1. Thalamic circuits in developmental and epileptic encephalopathies

There is limited research on the role of the TC circuit in developmental and epileptic encephalopathies (DEEs). Lennox-Gastaut Syndrome (LGS), characterized by slow spike-and-wave and commonly including generalized seizure types, is an area of research using surgical ablation and functional modulation (Velasco et al., 2006). Intracranial recordings from patients with LGS show ictal discharges in the centromedian thalamic nucleus (CM)

during generalized seizures, including tonic, tonic-clonic, and myoclonic seizures (Velasco et al., 1991). In Dravet syndrome resulting from *SCN1A* mutations (predominantly affecting $Na_v1.1$ on inhibitory neurons), an association has been demonstrated between loss of SK current, hyperactivity in nRT neurons, and downstream enhancement of TC neuron bursting (Ritter-Makinson et al., 2019; Studtmann et al., 2022). $Na_v1.2$, by contrast, is predominantly expressed on excitatory cells, and mutations in the *SCN2A* gene encoding $Na_v1.2$ are also associated with DEEs with predominantly focal seizures (Wolff et al., 2017), though animals carrying these mutations have absence-type seizures (Ogiwara et al., 2018). The mechanisms remain unknown. Further studies are needed to investigate the role of TC neuron disruption in the pathogenesis of these syndromes and the many other syndromes caused by other channelopathies (*SCN8A*, *SCN1B*, etc.) and to discover new therapeutic options for these conditions.

10.2. Dietary approaches may reveal untapped mechanisms

The ketogenic diet is effective for a wide range of epilepsies and represents an important tool in the clinical management of pharmaco-resistant seizures (Kinsman et al., 1992; Martin-McGill et al., 2020). A few randomized clinical trials (Kim et al., 2016; Sondhi et al., 2020) and innumerable clinical case reports suggest that ketogenic diets improve GGEs, including typical (Clemens et al., 2013) and atypical absence epilepsy (Ross et al., 1985), juvenile myoclonic epilepsy (Kverneland et al., 2015), and LGS (Caraballo et al., 2011), Dravet syndrome (Caraballo et al., 2005), and epilepsy with myoclonic-atonic seizures (Wiemer-Kruel et al., 2017). The ketogenic diet is pleiotropic, and the mechanisms of action so far identified include the provision of ketone bodies as an alternative metabolic substrate (Klepper, 2008), shifting the glutamate-glutamine equilibrium (Melo et al., 2006; Yudkoff et al., 2007), decreasing quantal content of glutamatergic vesicles (Juge et al., 2010), acidification of the brain parenchyma (Choudhary et al., 2021) and activation of ATP-sensitive potassium channels (Fogle et al., 2016). The extent to which these mechanisms act on TC circuits is not well understood, but the pH sensitivity of TASK and HCN channels provides one possible link.

10.3. Therapeutic thalamic stimulation in refractory epilepsy

Thalamic stimulation with deep brain stimulation (DBS) and responsive neurostimulation (RNS) are emerging treatments for refractory generalized epilepsy syndromes, and the efficacy of these treatments further supports a central role for the thalamus in generalized epilepsies. Chronic stimulation with DBS in the centromedian (CM) thalamic nucleus has been shown to reduce the frequency of generalized seizures, including absence and generalized tonic-clonic seizures in patients with GGE, multifocal epilepsy, or LGS (Velasco et al., 1987; Velasco et al., 2006; Dalic et al., 2022). There is also evidence for improvement of cognitive symptoms in patients with LGS after CM neurostimulation (Velasco et al., 2006). This effect on generalized epilepsy through stimulating CM is thought to be mediated by the widespread cortical connections of CM, including prefrontal, orbitofrontal, insular, motor, and sensory cortices (reviewed in Velasco et al., 2021). In contrast, targeting the anterior thalamic nucleus, with its largely temporal and frontal lobe connections, reduces seizure frequency in refractory temporal lobe epilepsy (Fisher et al., 2010; Salanova et al., 2021).

Closed-loop neurostimulation with RNS has also been used to target thalamic nuclei in patients with refractory epilepsy, recording thalamic ictal discharges and delivering stimulation in response to seizure detection. In case reports of patients with LGS, RNS targeting bilateral CM reduced the frequency of generalized seizures, including tonic, atonic, myoclonic, and generalized tonic-clonic seizures (Kwon et al., 2020; Welch et al., 2021).

In addition to modifying thalamic network activity with RNS or DBS, thalamic ablation has also been studied as a treatment for refractory epilepsy (Aguado-Carrillo et al., 2021). In a case series of patients with refractory generalized or multifocal epilepsy, bilateral radiofrequency ablation of CM significantly reduced the frequency of generalized seizures and was well tolerated, with only transient feeding difficulties after surgery (Aguado-Carrillo et al., 2021). Together, these results support the key role of the thalamus, and specifically the broad reciprocal cortical connections of CM, in generalized seizures and open new avenues for treating generalized epilepsy through thalamic modulation.

Further refinement of thalamic targets for neurostimulation is needed to improve treatment options in refractory epilepsy. Hypothesis-driven intracranial recordings from thalamic nuclei may help guide patient selection and improve outcomes as this technique becomes more widespread (Chaitanya et al., 2020). Furthermore, additional work in preclinical models is needed to study the therapeutic impact of modulating other thalamic nuclei. Potential candidates include the mediodorsal nucleus, with its widespread frontal connections, or nRT, with its key role in modulating thalamocortical loops.

10.4. Charting a path to future therapeutic breakthroughs

A number of technical and conceptual barriers should be addressed to improve treatments for GGE. One conceptual challenge is the remarkable clinical heterogeneity in the causes, expression, and treatments of the GGE syndromes. On the technical side, our understanding of the pathophysiology of generalized seizures would benefit from more widespread implementation of optogenetics and miniaturized probes and amplifiers that can record electrophysiologic signals in awake, behaving animals with high spatial and temporal resolution.

The rodent strains with susceptibility to absence epilepsy, which have served as powerful prototype disease models for the IGEs, have limited utility for the study of DEEs. Monogenic knock-in and knock-out genetic mouse models are helpful for identifying mechanisms and screening treatments for DEEs, but these approaches remain hampered by their low throughput. DEEs tend to be rare diseases, with only a handful of patients sharing any given mutation – it would be a staggering task to develop a unique mouse strain for every pathogenic human mutation. High-throughput assays and personalized medicine approaches should be leveraged where possible.

Current pharmacologic approaches to epilepsy treatment do not take full advantage of our knowledge about pathophysiological mechanisms. More selective T-type Ca^{2+} channel antagonists are needed. Notable gaps exist in medication classes that target reactive astrocytes, shift membrane potential, influence second messenger systems downstream of metabotropic glutamate receptors, adenosine receptors, adrenergic receptors, and muscarinic

acetylcholine receptors, and modulate emerging targets such as HCN and TASK channels. As highlighted above, we suggest that broader use of direct thalamic stimulation may expand therapeutic options for GGEs.

11. Conclusion

The unique physiologic properties of thalamic neurons and the widespread cortical and subcortical connections of the thalamus are dysregulated in generalized epilepsy syndromes and animal models of epilepsy. However, a more nuanced understanding of the relationship between TC circuit dysfunction and epilepsy is needed to identify new pharmacologic, dietary, gene therapy, surgical, and neuromodulation therapeutic targets to treat refractory generalized epilepsy. In particular, more studies are required to dissect thalamocortical cellular heterogeneity in health and epilepsy through single-cell and spatial transcriptomics. The rapid development of research tools to address these questions, including molecular and genetic models combined with optogenetics and advancements in recording techniques, makes this a pivotal time for the field.

Acknowledgements

We thank Dr. Rose Creed, PhD, and the members of Dr. Alexandra Nelson's lab for expert discussions. We thank Francoise Chanut and Kathryn Claiborn for editorial guidance.

Funding

JTP is supported by the National Institutes of Health grants R01NS121287; R01MH121446, Department of Defense grant EP190020, SLC6A1 Connect; Gladstone Institutes; and American Epilepsy Society.

BEL and CT are supported by the National Institute of Neurological Disorders and Stroke, National Institutes of Health, through UCSF grant number 5R25NS070680-13.

YV is supported by the FamilieSCN2A Foundation Action Potential grant and the National Institute of Neurological Disorders and Stroke, National Institutes of Health grant number 1F32NS127998-01.

The funding agencies had no role in the preparation of this review. The contents of this publication are solely the responsibility of the authors and do not necessarily represent the views of the funding agencies.

Data availability

No data were used for the research described in the article.

Abbreviations:

| | |
|-------------|--|
| AMPA | α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid |
| CAE | childhood absence epilepsy |
| cAMP | cyclic adenosine monophosphate |
| CT | corticothalamic |
| DBS | deep brain stimulation |
| DEE | developmental and epileptic encephalopathy |

| | |
|----------------|---|
| EEG | electroencephalogram |
| GABA | γ -aminobutyric acid |
| GAERS | genetic absence epilepsy rat from Strasbourg |
| GGE | genetic generalized epilepsy |
| GPe | globus pallidus pars externa |
| GTCA | generalized tonic clonic seizures alone |
| HCN | hyperpolarization-activated cyclic nucleotide-gated channel |
| IGE | idiopathic generalized epilepsy |
| JAE | juvenile absence epilepsy |
| JME | juvenile myoclonic epilepsy |
| LGS | Lennox-Gastaut syndrome |
| NE | norepinephrine |
| NMDA | N-methyl-D-aspartic acid |
| nRT | nucleus reticularis thalami |
| RNS | responsive neurostimulation |
| SWD | spike-and-wave discharge |
| TASK | TWIK-related acid-sensitive potassium channel |
| TC | thalamocortical |
| TREK | TWIK-related potassium channel |
| WAG/Rij | Wistar Albino Glaxo rats from Rijswijk. |

References

- Abarrategui B, Parejo-Carbonell B, García García ME, Di Capua D, García-Morales I, 2018. The cognitive phenotype of idiopathic generalized epilepsy. *Epilepsy Behav* 89, 99–104. [PubMed: 30408705]
- Agathonikou A, Panayiotopoulos CP, Giannakodimos S, Koutroumanidis M, 1998. Typical absence status in adults: diagnostic and syndromic considerations. *Epilepsia* 39 (12), 1265–1276. [PubMed: 9860061]
- Aguado-Carrillo G, Velasco AL, Saucedo-Alvarado PE, Cuellar-Herrera M, Trejo-Martínez D, Navarro-Olvera JL, Heres-Becerril SD, Barrera-Domínguez ED, Velasco-Campos F, 2021. Radiofrequency ablation of the centromedian thalamic nucleus in the treatment of drug-resistant epilepsy. *Epilepsy Behav. E&B* 114 (Pt A), 107560.
- Akbar MT, Rattray M, Williams RJ, Chong NW, Meldrum BS, 1998. Reduction of GABA and glutamate transporter messenger RNAs in the severe-seizure genetically epilepsy-prone rat. *Neuroscience* 85 (4), 1235–1251. [PubMed: 9681960]

- Aledo-Serrano A, Sánchez-Alcudia R, Toledano R, García-Morales I, Beltrán-Corbellini Á, del Pino I, Gil-Nagel A, 2021. Developmental and epileptic encephalopathies after negative or inconclusive genetic testing: What is next? *J. Transl. Genet. Genom* 5 (4), 443–455.
- Almane DN, Jones JE, McMillan T, Stafstrom CE, Hsu DA, Seidenberg M, Hermann BP, Oyegbile TO, 2019. The timing, nature, and range of neurobehavioral comorbidities in Juvenile Myoclonic epilepsy. *Pediatr. Neurol* 101, 47–52. [PubMed: 31122836]
- Arain FM, Boyd KL, Gallagher MJ, 2012. Decreased viability and absence-like epilepsy in mice lacking or deficient in the GABAA receptor alpha1 subunit. *Epilepsia* 53 (8), e161–e165. [PubMed: 22812724]
- Asadi-Pooya AA, Homayoun M, 2020. Idiopathic (genetic) generalized epilepsies with absences: clinical and electrographic characteristics and seizure outcome. *Neurol. Sci* 41 (12), 3677–3682. [PubMed: 32504278]
- Asadi-Pooya AA, Emami M, Sperling MR, 2013. A clinical study of syndromes of idiopathic (genetic) generalized epilepsy. *J. Neurol. Sci* 324 (1–2), 113–117. [PubMed: 23149265]
- Ates N, Sahin D, Ilbay G, 2004. Theophylline, a methylxanthine derivative, suppresses absence epileptic seizures in WAG/Rij rats. *Epilepsy Behav* 5 (5), 645–648. [PubMed: 15380114]
- Austin JK, Harezlak J, Dunn DW, Huster GA, Rose DF, Ambrosius WT, 2001. Behavior problems in children before first recognized seizures. *Pediatrics* 107 (1), 115–122. [PubMed: 11134444]
- Banerjee PK, Snead OC 3rd, 1994. Thalamic mediodorsal and intralaminar nuclear lesions disrupt the generation of experimentally induced generalized absence-like seizures in rats. *Epilepsy Res* 17 (3), 193–205. [PubMed: 8013443]
- Barad Z, Shevtsova O, Arbutnott GW, Leitch B, 2012. Selective loss of AMPA receptors at corticothalamic synapses in the epileptic stargazer mouse. *Neuroscience* 217, 19–31. [PubMed: 22609941]
- Bats C, Groc L, Choquet D, 2007. The interaction between Stargazin and PSD-95 regulates AMPA receptor surface trafficking. *Neuron* 53 (5), 719–734. [PubMed: 17329211]
- Beenhakker MP, Huguenard JR, 2009. Neurons that fire together also conspire together: Is normal sleep circuitry hijacked to generate epilepsy? *Neuron* 62 (5), 612–632. [PubMed: 19524522]
- Beenhakker MP, Huguenard JR, 2010. Astrocytes as gatekeepers of GABAB receptor function. *J. Neurosci* 30 (45), 15262–15276. [PubMed: 21068331]
- Beghi M, Beghi E, Cornaggia CM, Gobbi G, 2006. Idiopathic generalized epilepsies of adolescence. *Epilepsia* 47 (Suppl. 2), 107–110. [PubMed: 17105478]
- Ben-Ari Y, Dingledine R, Kanazawa I, Kelly JS, 1976. Inhibitory effects of acetylcholine on neurones in the feline nucleus reticularis thalami. *J. Physiol* 261 (3), 647–671. [PubMed: 978591]
- Berg AT, Levy SR, Testa FM, Shinnar S, 1999. Classification of childhood epilepsy syndromes in newly diagnosed epilepsy: interrater agreement and reasons for disagreement. *Epilepsia* 40 (4), 439–444. [PubMed: 10219269]
- Berridge CW, Schmeichel BE, España RA, 2012. Noradrenergic modulation of wakefulness/arousal. *Sleep Med. Rev* 16 (2), 187–197. [PubMed: 22296742]
- Bertaso F, Zhang C, Scheschonka A, de Bock F, Fontanaud P, Marin P, Haganir RL, Betz H, Bockaert J, Fagni L, Lerner-Natoli M, 2008. PICK1 uncoupling from mGluR7a causes absence-like seizures. *Nat. Neurosci* 11 (8), 940–948. [PubMed: 18641645]
- Bessaih T, Bourgeois L, Badiu CI, Carter DA, Toth TI, Ruano D, Lambolez B, Crunelli V, Leresche N, 2006. Nucleus-specific abnormalities of GABAergic synaptic transmission in a genetic model of absence seizures. *J. Neurophysiol* 96 (6), 3074–3081. [PubMed: 16971676]
- Beydoun A, D'Souza J, 2012. Treatment of idiopathic generalized epilepsy - a review of the evidence. *Expert. Opin. Pharmacother* 13 (9), 1283–1298. [PubMed: 22568658]
- Biel M, Wahl-Schott C, Michalakis S, Zong X, 2009. Hyperpolarization-activated cation channels: from genes to function. *Physiol. Rev* 89 (3), 847–885. [PubMed: 19584315]
- Bista P, Meuth SG, Kanyshkova T, Cerina M, Pawlowski M, Ehling P, Landgraf P, Borsotto M, Heurteaux C, Pape HC, Baukrowitz T, Budde T, 2012. Identification of the muscarinic pathway underlying cessation of sleep-related burst activity in rat thalamocortical relay neurons. *Pflugers Arch* 463 (1), 89–102. [PubMed: 22083644]

- Bista P, Cerina M, Ehling P, Leist M, Pape HC, Meuth SG, Budde T, 2015. The role of two-pore-domain background K(+) (K(2)p) channels in the thalamus. *Pflugers Arch* 467 (5), 895–905. [PubMed: 25346156]
- Blom S, Heijbel J, Bergfors PG, 1978. Incidence of epilepsy in children: a follow-up study three years after the first seizure. *Epilepsia* 19 (4), 343–350. [PubMed: 100316]
- Boison D, 2008. The adenosine kinase hypothesis of epileptogenesis. *Prog. Neurobiol* 84 (3), 249–262. [PubMed: 18249058]
- Borea PA, Gessi S, Merighi S, Vincenzi F, Varani K, 2018. Pharmacology of adenosine receptors: the state of the art. *Physiol. Rev* 98 (3), 1591–1625. [PubMed: 29848236]
- Broicher T, Wetschurck N, Munsch T, Coulon P, Meuth SG, Kanyshkova T, Seidenbecher T, Offermanns S, Pape HC, Budde T, 2008. Muscarinic ACh receptor-mediated control of thalamic activity via G(q)/G (11)-family G-proteins. *Pflugers Arch* 456 (6), 1049–1060. [PubMed: 18350314]
- Bryant AS, Li B, Beenhakker MP, Huguenard JR, 2009. Maintenance of thalamic epileptiform activity depends on the astrocytic glutamate-glutamine cycle. *J. Neurophysiol* 102 (5), 2880–2888. [PubMed: 19741104]
- Budde T, Caputi L, Kanyshkova T, Staak R, Abrahamczik C, Munsch T, Pape HC, 2005. Impaired regulation of thalamic pacemaker channels through an imbalance of subunit expression in absence epilepsy. *J. Neurosci* 25 (43), 9871–9882. [PubMed: 16251434]
- Byrum CE, Guyenet PG, 1987. Afferent and efferent connections of the A5 noradrenergic cell group in the rat. *J. Comp. Neurol* 261 (4), 529–542. [PubMed: 2440916]
- Cain SM, Tyson JR, Choi HB, Ko R, Lin PJC, LeDue JM, Powell KL, Bernier LP, Rungta RL, Yang Y, Cullis PR, O'Brien TJ, MacVicar BA, Snutch TP, 2018. Cav 3.2 drives sustained burst-firing, which is critical for absence seizure propagation in reticular thalamic neurons. *Epilepsia* 59 (4), 778–791. 10.1111/epi.14018. [PubMed: 29468672]
- Callenbach PM, Bouma PA, Geerts AT, Arts WF, Stroink H, Peeters EA, van Donselaar CA, Peters AC, Brouwer OF, 2009. Long-term outcome of childhood absence epilepsy: Dutch study of epilepsy in childhood. *Epilepsy Res* 83 (2–3), 249–256. [PubMed: 19124226]
- Caplan R, 2015. A call to arms against “The phantom of Epilepsy”. *Epilepsia* 56 (11), 1710–1711. [PubMed: 26415884]
- Caplan R, Siddarth P, Stahl L, Lanphier E, Vona P, Gurbani S, Koh S, Sankar R, Shields WD, 2008. Childhood absence epilepsy: behavioral, cognitive, and linguistic comorbidities. *Epilepsia* 49 (11), 1838–1846. [PubMed: 18557780]
- Caraballo RH, Cersosimo RO, Sakr D, Cresta A, Escobal N, Fejerman N, 2005. Ketogenic diet in patients with Dravet syndrome. *Epilepsia* 46 (9), 1539–1544. [PubMed: 16146451]
- Caraballo R, Vaccarezza M, Cersosimo R, Rios V, Soraru A, Arroyo H, Agosta G, Escobal N, Demartini M, Maxit C, Cresta A, Marchione D, Carniello M, Panico L, 2011. Long-term follow-up of the ketogenic diet for refractory epilepsy: multicenter Argentinean experience in 216 pediatric patients. *Seizure* 20 (8), 640–645. [PubMed: 21763159]
- Castro-Alamancos MA, Connors BW, 1997. Thalamocortical synapses. *Prog. Neurobiol* 51 (6), 581–606. [PubMed: 9175158]
- Catterall WA, 2014. Sodium channels, inherited epilepsy, and antiepileptic drugs. *Annu. Rev. Pharmacol. Toxicol* 54, 317–338. [PubMed: 24392695]
- Cavazzuti GB, 1980. Epidemiology of different types of epilepsy in school age children of Modena, Italy. *Epilepsia* 21 (1), 57–62. [PubMed: 6766394]
- Cavdar S, Kuvvet Y, Sur-Erdem I, Ozgur M, Onat F, 2019. Relationships between astrocytes and absence epilepsy in rat: an experimental study. *Neurosci. Lett* 712, 134518. [PubMed: 31560994]
- Chaitanya G, Romeo AK, Ilyas A, Irannejad A, Toth E, Elsayed G, Bentley JN, Riley KO, Pati S, 2020. Robot-assisted stereoelectroencephalography exploration of the limbic thalamus in human focal epilepsy: implantation technique and complications in the first 24 patients. *Neurosurg. Focus* 48 (4), E2.
- Chawla T, Chaudhry N, Puri V, 2021. Cognitive dysfunction in Juvenile Myoclonic Epilepsy (JME) - A tertiary care center study. *Ann. Indian Acad. Neurol* 24 (1), 40–50. [PubMed: 33911378]

- Cheong E, Zheng Y, Lee K, Lee J, Kim S, Sanati M, Lee S, Kim YS, Shin HS, 2009. Deletion of phospholipase C beta4 in thalamocortical relay nucleus leads to absence seizures. *Proc. Natl. Acad. Sci. U. S. A* 106 (51), 21912–21917. [PubMed: 19955421]
- Cho FS, Vainchtein ID, Voskobiynyk Y, Morningstar AR, Aparicio F, Higashikubo B, Ciesielska A, Broekaart DWM, Anink JJ, van Vliet EA, Yu X, Khakh BS, Aronica E, Molofsky AV, Paz JT, 2022. Enhancing GAT-3 in thalamic astrocytes promotes resilience to brain injury in rodents. *Sci. Transl. Med* 14 (652), eabj4310.
- Choudhary A, Mu C, Barrett KT, Charkhand B, Williams-Dyjur C, Marks WN, Shearer J, Rho JM, Scantlebury MH, 2021. The link between brain acidosis, breathing and seizures: a novel mechanism of action for the ketogenic diet in a model of infantile spasms. *Brain Commun* 3 (4), fcab189.
- Clemens Z, Kelemen A, Fogarasi A, Toth C, 2013. Childhood absence epilepsy successfully treated with the paleolithic ketogenic diet. *Neurol. Ther* 2 (1–2), 71–76. [PubMed: 26000218]
- Collaborators GBDN, 2019. Global, regional, and national burden of neurological disorders, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 18 (5), 459–480. [PubMed: 30879893]
- Contreras D, Destexhe A, Sejnowski TJ, Steriade M, 1997. Spatiotemporal patterns of spindle oscillations in cortex and thalamus. *J. Neurosci* 17 (3), 1179–1196. [PubMed: 8994070]
- Coulter DA, Huguenard JR, Prince DA, 1989. Calcium currents in rat thalamocortical relay neurones: kinetic properties of the transient, low-threshold current. *J. Physiol* 414, 587–604. [PubMed: 2607443]
- Coulter DA, Huguenard JR, Prince DA, 1990a. Differential effects of petit mal anticonvulsants and convulsants on thalamic neurones: calcium current reduction. *Br. J. Pharmacol* 100 (4), 800–806. [PubMed: 2169941]
- Coulter DA, Huguenard JR, Prince DA, 1990b. Differential effects of petit mal anticonvulsants and convulsants on thalamic neurones: GABA current blockade. *Br. J. Pharmacol* 100 (4), 807–813. [PubMed: 2119843]
- Crunelli V, Leresche N, 2002a. Block of thalamic T-Type Ca(2+) channels by ethosuximide is not the whole story. *Epilepsy Curr* 2 (2), 53–56. [PubMed: 15309166]
- Crunelli V, Leresche N, 2002b. Childhood absence epilepsy: genes, channels, neurons and networks. *Nat. Rev. Neurosci* 3 (5), 371–382. [PubMed: 11988776]
- Crunelli V, Carmignoto G, Steinhauser C, 2015. Novel astrocyte targets: new avenues for the therapeutic treatment of epilepsy. *Neuroscientist* 21 (1), 62–83. [PubMed: 24609207]
- Cueni L, Canepari M, Lujan R, Emmenegger Y, Watanabe M, Bond CT, Franken P, Adelman JP, Luthi A, 2008. T-type Ca²⁺ channels, SK2 channels and SERCAs gate sleep-related oscillations in thalamic dendrites. *Nat. Neurosci* 11 (6), 683–692. [PubMed: 18488023]
- Dalic LJ, Warren AEL, Bulluss KJ, Thevathasan W, Roten A, Churilov L, Archer JS, 2022. DBS of thalamic centromedian nucleus for Lennox-Gastaut Syndrome (ESTEL Trial). *Ann. Neurol* 91 (2), 253–267. [PubMed: 34877694]
- D'Amore V, von Randow C, Nicoletti F, Ngomba RT, van Luijtelaar G, 2015. Anti-absence activity of mGlu1 and mGlu5 receptor enhancers and their interaction with a GABA reuptake inhibitor: effect of local infusions in the somatosensory cortex and thalamus. *Epilepsia* 56 (7), 1141–1151. [PubMed: 26040777]
- David F, Carcak N, Furdan S, Onat F, Gould T, Meszaros A, Di Giovanni G, Hernandez VM, Chan CS, Lorincz ML, Crunelli V, 2018. Suppression of hyperpolarization-activated cyclic nucleotide-gated channel function in thalamocortical neurons prevents genetically determined and pharmacologically induced absence seizures. *J. Neurosci* 38 (30), 6615–6627. [PubMed: 29925625]
- de Araujo Filho GM, Yacubian EM, 2013. Juvenile myoclonic epilepsy: psychiatric comorbidity and impact on outcome. *Epilepsy Behav* 28 (Suppl. 1), S74–S80. [PubMed: 23756487]
- De Lima AD, Singer W, 1987. The brainstem projection to the lateral geniculate nucleus in the cat: identification of cholinergic and monoaminergic elements. *J. Comp. Neurol* 259 (1), 92–121. [PubMed: 2884241]

- Debarbieux F, Brunton J, Charpak S, 1998. Effect of bicuculline on thalamic activity: a direct blockade of IAHP in reticularis neurons. *J. Neurophysiol* 79 (6), 2911–2918. [PubMed: 9636097]
- Dede F, Karadenizli S, Ozsoy OD, Eraldemir FC, Sahin D, Ates N, 2019. Antagonism of adenosinergic system decrease SWD occurrence via an increment in thalamic NFkB and IL-6 in absence epilepsy. *J. Neuroimmunol* 326, 1–8. [PubMed: 30423516]
- Deransart C, Hellwig B, Heupel-Reuter M, eger J-F, Heck D, Lüicking CH, 2003. Single-unit analysis of substantia nigra pars reticulata neurons in freely behaving rats with genetic absence epilepsy. *Epilepsia* 44 (12), 1513–1520. [PubMed: 14636321]
- Devilbiss DM, Waterhouse BD, 2011. Phasic and tonic patterns of locus coeruleus output differentially modulate sensory network function in the awake rat. *J. Neurophysiol* 105 (1), 69–87. [PubMed: 20980542]
- Dienel GA, Gillinder L, McGonigal A, Borges K, 2023. Potential new roles for glycogen in epilepsy. *Epilepsia* 64 (1), 29–53. 10.1111/epi.17412. [PubMed: 36117414]
- Dieterich E, Doose H, Baier WK, Fichsel H, 1985. Longterm follow-up of childhood epilepsy with absences. II. Absence-epilepsy with initial grand mal. *Neuropediatrics* 16 (3), 155–158. [PubMed: 3930989]
- Ding L, Gallagher MJ, 2016. Dynamics of sensorimotor cortex activation during absence and myoclonic seizures in a mouse model of juvenile myoclonic epilepsy. *Epilepsia* 57 (10), 1568–1580. [PubMed: 27573707]
- Drake ME, Pakalnis A, Padamadan H, Weate SM, Cannon PA, 1991. Sleep spindles in Epilepsy. *Clin. Electroencephalogr* 22 (3), 144–149. [PubMed: 1879053]
- Dunwiddie TV, Masino SA, 2001. The role and regulation of adenosine in the central nervous system. *Annu. Rev. Neurosci* 24, 31–55. [PubMed: 11283304]
- Duran J, Hervera A, Markussen KH, Varea O, Lopez-Soldado I, Sun RC, Del Rio JA, Gentry MS, Guinovart JJ, 2021. Astrocytic glycogen accumulation drives the pathophysiology of neurodegeneration in Lafora disease. *Brain* 144 (8), 2349–2360. [PubMed: 33822008]
- During MJ, Spencer DD, 1992. Adenosine: a potential mediator of seizure arrest and postictal refractoriness. *Ann. Neurol* 32 (5), 618–624. [PubMed: 1449242]
- Dutuit M, Didier-Bazes M, Vergnes M, Mutin M, Conjard A, Akaoka H, Belin MF, Touret M, 2000. Specific alteration in the expression of glial fibrillary acidic protein, glutamate dehydrogenase, and glutamine synthetase in rats with genetic absence epilepsy. *Glia* 32 (1), 15–24. [PubMed: 10975907]
- Dutuit M, Touret M, Szymocha R, Nehlig A, Belin MF, Didier-Bazes M, 2002. Decreased expression of glutamate transporters in genetic absence epilepsy rats before seizure occurrence. *J. Neurochem* 80 (6), 1029–1038. [PubMed: 11953453]
- Eelkman Rooda OHJ, Kros L, Faneyte SJ, Holland PJ, Gornati SV, Poelman HJ, Jansen NA, Tolner EA, van den Maagdenberg A, De Zeeuw CI, Hoebeek FE, 2021. Single-pulse stimulation of cerebellar nuclei stops epileptic thalamic activity. *Brain Stimul* 14 (4), 861–872. [PubMed: 34022430]
- Elkommos S, Mula M, 2022. Current and future pharmacotherapy options for drug-resistant epilepsy. *Expert. Opin. Pharmacother* 23 (18), 2023–2034. [PubMed: 36154780]
- Elmali AD, Auvin S, Bast T, Rubboli G, Koutroumanidis M, 2020. How to diagnose and classify idiopathic (genetic) generalized epilepsies. *Epileptic Disord* 22 (4), 399–420. [PubMed: 32782228]
- Enyedi P, Czirjak G, 2010. Molecular background of leak K⁺ currents: two-pore domain potassium channels. *Physiol. Rev* 90 (2), 559–605. [PubMed: 20393194]
- Ernst WL, Zhang Y, Yoo JW, Ernst SJ, Noebels JL, 2009. Genetic enhancement of thalamocortical network activity by elevating alpha 1g-mediated low-voltage-activated calcium current induces pure absence epilepsy. *J. Neurosci* 29 (6), 1615–1625. [PubMed: 19211869]
- E kaza E, Onat FY, Aker R, Oner G, 2002. Resistance to propagation of amygdaloid kindling seizures in rats with genetic absence epilepsy. *Epilepsia* 43 (10), 1115–1119. [PubMed: 12366723]
- Esquivel EE, Pitt MC, Boyd SG, 1991. EEG findings in hypomelanosis of Ito. *Neuropediatrics* 22 (4), 216–219. [PubMed: 1775218]

- Fisher R, Salanova V, Witt T, Worth R, Henry T, Gross R, Oommen K, Osorio I, Nazzaro J, Labar D, Kaplitt M, Sperling M, Sandok E, Neal J, Handforth A, Stern J, DeSalles A, Chung S, Shetter A, Bergen D, Bakay R, Henderson J, French J, Baltuch G, Rosenfeld W, Youkilis A, Marks W, Garcia P, Barbaro N, Fountain N, Bazil C, Goodman R, McKhann G, Krishnamurthy K Babu, Papavassiliou S, Epstein C, Pollard J, Tonder L, Grebin J, Coffey R, Graves N, S. S. Group, 2010. Electrical stimulation of the anterior nucleus of thalamus for treatment of refractory epilepsy. *Epilepsia* 51 (5), 899–908. [PubMed: 20331461]
- Fisher RS, Cross JH, French JA, Higurashi N, Hirsch E, Jansen FE, Lagae L, Moshe SL, Peltola J, Roulet Perez E, Scheffer IE, Zuberi SM, 2017. Operational classification of seizure types by the International League Against Epilepsy: position paper of the ILAE commission for classification and terminology. *Epilepsia* 58 (4), 522–530. [PubMed: 28276060]
- Fogle KJ, Hertzler JI, Shon JH, Palladino MJ, 2016. The ATP-sensitive K channel is seizure protective and required for effective dietary therapy in a model of mitochondrial encephalomyopathy. *J. Neurogenet* 30 (3–4), 247–258. [PubMed: 27868454]
- Fuentealba P, Timofeev I, Bazhenov M, Sejnowski TJ, Steriade M, 2005. Membrane bistability in thalamic reticular neurons during spindle oscillations. *J. Neurophysiol* 93 (1), 294–304. [PubMed: 15331618]
- Gale K, 1992. Subcortical structures and pathways involved in convulsive seizure generation. *J. Clin. Neurophysiol* 9 (2), 264–277. [PubMed: 1350593]
- Gama AP, Taura M, Alonso NB, Sousa AM, Noffs M, Yacubian EM, Guilhoto LM, 2020. Impulsiveness, personality traits and executive functioning in patients with juvenile myoclonic epilepsy. *Seizure* 82, 125–132. [PubMed: 33068959]
- Geithner J, Schneider F, Wang Z, Berneiser J, Herzer R, Kessler C, Runge U, 2012. Predictors for long-term seizure outcome in juvenile myoclonic epilepsy: 25–63 years of follow-up. *Epilepsia* 53 (8), 1379–1386. [PubMed: 22686598]
- Germé K, Faure JB, Koning E, Nehlig A, 2015. Effect of caffeine and adenosine receptor ligands on the expression of spike-and-wave discharges in Genetic Absence Epilepsy Rats from Strasbourg (GAERS). *Epilepsy Res* 110, 105–114. [PubMed: 25616462]
- Giorgi FS, Pizzanelli C, Biagioni F, Murri L, Fornai F, 2004. The role of norepinephrine in epilepsy: from the bench to the bedside. *Neurosci. Biobehav. Rev* 28 (5), 507–524. [PubMed: 15465138]
- Glauser TA, Cnaan A, Shinnar S, Hirtz DG, Dlugos D, Masur D, Clark PO, Adamson PC, 2013. Ethosuximide, valproic acid, and lamotrigine in childhood absence epilepsy: initial monotherapy outcomes at 12 months. *Epilepsia* 54 (1), 141–155. [PubMed: 23167925]
- Godschalk M, Dzoljic MR, Bonta IL, 1977. Slow wave sleep and a state resembling absence epilepsy induced in the rat by gamma-hydroxybutyrate. *Eur. J. Pharmacol* 44 (2), 105–111. [PubMed: 407087]
- Grasso C, Li Volsi G, Licata F, Ciranna L, Santangelo F, 2006. Aminergic control of neuronal firing rate in thalamic motor nuclei of the rat. *Arch. Ital. Biol* 144 (3–4), 173–196. [PubMed: 16977832]
- Graybiel AM, 2000. The basal ganglia. *Curr. Biol. CB* 10 (14), R509–R511. [PubMed: 10899013]
- Grosso S, Galimberti D, Vezzosi P, Farnetani M, Di Bartolo RM, Bazzotti S, Morgese G, Balestri P, 2005. Childhood absence epilepsy: evolution and prognostic factors. *Epilepsia* 46 (11), 1796–1801. [PubMed: 16302860]
- Gruenbaum BF, Sandhu MRS, Bertasi RAO, Bertasi TGO, Schonwald A, Kurup A, Gruenbaum SE, Freedman IG, Funaro MC, Blumenfeld H, Sanacora G, 2021. Absence seizures and their relationship to depression and anxiety: evidence for bidirectionality. *Epilepsia* 62 (5), 1041–1056. [PubMed: 33751566]
- Guerrini R, Conti V, Mantegazza M, Balestrini S, Galanopoulou AS, Benfenati F, 2023. Developmental and epileptic encephalopathies: from genetic heterogeneity to phenotypic continuum. *Physiol. Rev* 103 (1), 433–513. [PubMed: 35951482]
- Haber SN, Calzavara R, 2009. The cortico-basal ganglia integrative network: the role of the thalamus. *Brain Res. Bull* 78 (2–3), 69–74. [PubMed: 18950692]
- Haber S, McFarland NR, 2001. The place of the thalamus in frontal cortical-basal ganglia circuits. *Neurosci. Rev. J. Bring. Neurobiol. Neurol. Psychiatry* 7 (4), 315–324.

- Halasz P, Terzano MG, Parrino L, 2002. Spike-wave discharge and the microstructure of sleep-wake continuum in idiopathic generalised epilepsy. *Neurophysiol. Clin* 32 (1), 38–53. [PubMed: 11915485]
- Hammelmann V, Stieglitz MS, Hulle H, Le Meur K, Kass J, Brummer M, Gruner C, Rotzer RD, Fenske S, Hartmann J, Zott B, Luthi A, Spahn S, Moser M, Isbrandt D, Ludwig A, Konnerth A, Wahl-Schott C, Biel M, 2019. Abolishing cAMP sensitivity in HCN2 pacemaker channels induces generalized seizures. *JCI Insight* 4 (9).
- Harada Y, Nagao Y, Shimizu S, Serikawa T, Terada R, Fujimoto M, Okuda A, Mukai T, Sasa M, Kurachi Y, Ohno Y, 2013. Expressional analysis of inwardly rectifying Kir4.1 channels in Noda epileptic rat (NER). *Brain Res* 1517, 141–149. [PubMed: 23603404]
- He C, Chen F, Li B, Hu Z, 2014. Neurophysiology of HCN channels: from cellular functions to multiple regulations. *Prog. Neurobiol* 112, 1–23. [PubMed: 24184323]
- He X, Chaitanya G, Asma B, Caciagli L, Bassett DS, Tracy JJ, Sperling MR, 2020. Disrupted basal ganglia-thalamocortical loops in focal to bilateral tonic-clonic seizures. *England* 143.
- He Z, Li Y, Zhao X, Li B, 2022. Dravet syndrome: Advances in etiology, clinical presentation, and treatment. *Epilepsy Res* 188, 107041. [PubMed: 36368227]
- Healy L, Moran M, Singhal S, O'Donoghue MF, Alzoubidi R, Whitehouse WP, 2018. Relapse after treatment withdrawal of antiepileptic drugs for Juvenile Absence Epilepsy and Juvenile Myoclonic Epilepsy. *Seizure* 59, 116–122. [PubMed: 29807291]
- Helbig I, 2015. Genetic causes of generalized epilepsies. *Semin. Neurol* 35 (3), 288–292. [PubMed: 26060908]
- Henkin Y, Sadeh M, Kivity S, Shabtai E, Kishon-Rabin L, Gadoth N, 2005. Cognitive function in idiopathic generalized epilepsy of childhood. *Dev. Med. Child Neurol* 47 (2), 126–132. [PubMed: 15707236]
- Hermann B, Jones J, Dabbs K, Allen CA, Sheth R, Fine J, McMillan A, Seidenberg M, 2007. The frequency, complications and aetiology of ADHD in new onset paediatric epilepsy. *Brain* 130 (Pt 12), 3135–3148. [PubMed: 17947336]
- Hirsch LJ, Fong MWK, Leitinger M, LaRoche SM, Beniczky S, Abend NS, Lee JW, Wusthoff CJ, Hahn CD, Westover MB, Gerard EE, Herman ST, Haider HA, Osman G, Rodriguez-Ruiz A, Maciel CB, Gilmore EJ, Fernandez A, Rosenthal ES, Claassen J, Husain AM, Yoo JY, So EL, Kaplan PW, Nuwer MR, van Putten M, Sutter R, Drislane FW, Trinka E, Gaspard N, 2021. American clinical neurophysiology society's standardized critical care EEG terminology: 2021 version. *J. Clin. Neurophysiol* 38 (1).
- Hirsch E, French J, Scheffer IE, Bogacz A, Alsaadi T, Sperling MR, Abdulla F, Zuberi SM, Trinka E, Specchio N, Somerville E, Samia P, Riney K, Nabbout R, Jain S, Wilmshurst JM, Auvin S, Wiebe S, Perucca E, Moshe SL, Tinuper P, Wirrell EC, 2022. ILAE definition of the Idiopathic Generalized Epilepsy Syndromes: position statement by the ILAE task force on nosology and definitions. *Epilepsia* 63 (6), 1475–1499. [PubMed: 35503716]
- Höfler J, Unterberger I, Döbesberger J, Kuchukhidze G, Walser G, Trinka E, 2014. Seizure outcome in 175 patients with juvenile myoclonic epilepsy—a long-term observational study. *Epilepsy Res* 108 (10), 1817–1824. [PubMed: 25443450]
- Holowach J, Thurston DL, O'Leary JL, 1962. Petit mal epilepsy. *Pediatrics* 30, 893–901. [PubMed: 13961432]
- Holtkamp M, Kowski AB, Merkle H, Janz D, 2014. Long-term outcome in epilepsy with grand mal on awakening: forty years of follow-up. *Ann. Neurol* 75 (2), 298–302. [PubMed: 24395517]
- Hosford DA, Wang Y, Cao Z, 1997. Differential effects mediated by GABAA receptors in thalamic nuclei in lh/lh model of absence seizures. *Epilepsy Res* 27 (1), 55–65. [PubMed: 9169291]
- Huguenard JR, 1999. Neuronal circuitry of thalamocortical epilepsy and mechanisms of antiabsence drug action. *Adv. Neurol* 79, 991–999. [PubMed: 10514881]
- Iacone Y, Morais TP, David F, Delicata F, Sandle J, Raffai T, Parri HR, Weisser JJ, Bundgaard C, Klewe IV, Tamas G, Thomsen MS, Crunelli V, Lorincz ML, 2021. Systemic administration of ivabradine, a hyperpolarization-activated cyclic nucleotide-gated channel inhibitor, blocks spontaneous absence seizures. *Epilepsia* 62 (7), 1729–1743. [PubMed: 34018186]

- ILAE, 1989. Proposal for revised classification of epilepsies and epileptic syndromes. Commission on Classification and Terminology of the International League Against Epilepsy. *Epilepsia* 30 (4), 389–399. [PubMed: 2502382]
- Ilbay G, Sahin D, Karson A, Ates N, 2001. Effects of adenosine administration on spike-wave discharge frequency in genetically epileptic rats. *Clin. Exp. Pharmacol. Physiol* 28 (8), 643–646. [PubMed: 11473530]
- Iqbal N, Caswell H, Muir R, Cadden A, Ferguson S, Mackenzie H, Watson P, Duncan S, 2015. Neuropsychological profiles of patients with juvenile myoclonic epilepsy and their siblings: an extended study. *Epilepsia* 56 (8), 1301–1308. [PubMed: 26075864]
- Jafarian M, Esmail Alipour M, Karimzadeh F, 2020. Experimental models of absence epilepsy. *Basic Clin. Neurosci* 11 (6), 715–726. [PubMed: 33850609]
- Jain S, Padma MV, Puri A, Maheshwari MC, 1998. Juvenile myoclonic epilepsy: disease expression among Indian families. *Acta Neurol. Scand* 97 (1), 1–7. [PubMed: 9482671]
- Jallon P, Latour P, 2005. Epidemiology of idiopathic generalized epilepsies. *Epilepsia* 46 (Suppl. 9), 10–14.
- Jefferys JGR, Jiruska P, de Curtis M, Avoli M, 2012. Limbic network synchronization and temporal lobe Epilepsy. In: Noebels JL, Avoli M, Rogawski MA, Olsen RW, Delgado-Escueta AV (Eds.), *Jasper's Basic Mechanisms of the Epilepsies* National Center for Biotechnology Information (US), Bethesda (MD).
- Juge N, Gray JA, Omote H, Miyaji T, Inoue T, Hara C, Uneyama H, Edwards RH, Nicoll RA, Moriyama Y, 2010. Metabolic control of vesicular glutamate transport and release. *Neuron* 68 (1), 99–112. [PubMed: 20920794]
- Juul-Jensen P, Foldspang A, 1983. Natural history of epileptic seizures. *Epilepsia* 24 (3), 297–312. [PubMed: 6406213]
- Kananura C, Sander T, Rajan S, Preisig-Muller R, Grzeschik KH, Daut J, Derst C, Steinlein OK, 2002. Tandem pore domain K(+)-channel TASK-3 (KCNK9) and idiopathic absence epilepsies. *Am. J. Med. Genet* 114 (2), 227–229. [PubMed: 11857586]
- Kandel A, Buzsáki G, 1993. Cerebellar neuronal activity correlates with spike and wave EEG patterns in the rat. *Epilepsy Res* 16 (1), 1–9. [PubMed: 8243435]
- Kessler SK, Shinnar S, Cnaan A, Dlugos D, Conry J, Hirtz DG, Hu F, Liu C, Mizrahi EM, Moshe SL, Clark P, Glauser TA and G. Childhood Absence Epilepsy Study (2017). “Pretreatment seizure semiology in childhood absence epilepsy”. *Neurology* 89(7): 673–679. [PubMed: 28724582]
- Kim U, Sanchez-Vives MV, McCormick DA, 1997. Functional dynamics of GABAergic inhibition in the thalamus. *Science* 278 (5335), 130–134. [PubMed: 9311919]
- Kim D, Song I, Keum S, Lee T, Jeong MJ, Kim SS, McEnery MW, Shin HS, 2001. Lack of the burst firing of thalamocortical relay neurons and resistance to absence seizures in mice lacking alpha(1G) T-type Ca(2+) channels. *Neuron* 31 (1), 35–45. [PubMed: 11498049]
- Kim JA, Yoon JR, Lee EJ, Lee JS, Kim JT, Kim HD, Kang HC, 2016. Efficacy of the classic ketogenic and the modified Atkins diets in refractory childhood epilepsy. *Epilepsia* 57 (1), 51–58. [PubMed: 26662710]
- Kinsman SL, Vining EP, Quaskey SA, Mellits D, Freeman JM, 1992. Efficacy of the ketogenic diet for intractable seizure disorders: review of 58 cases. *Epilepsia* 33 (6), 1132–1136. [PubMed: 1464275]
- Kleiman-Weiner M, Beenhakker MP, Segal WA, Huguenard JR, 2009. Synergistic roles of GABAA receptors and SK channels in regulating thalamocortical oscillations. *J. Neurophysiol* 102 (1), 203–213. [PubMed: 19386752]
- Klepper J, 2008. Glucose transporter deficiency syndrome (GLUT1DS) and the ketogenic diet. *Epilepsia* 49 (Suppl. 8), 46–49. [PubMed: 19049586]
- Knake S, Hamer HM, Schomburg U, Oertel WH, Rosenow F, 1999. Tiagabine-induced absence status in idiopathic generalized epilepsy. *Seizure* 8 (5), 314–317. [PubMed: 10486298]
- Koerner C, Danober L, Boehrer A, Marescaux C, Vergnes M, 1996. Thalamic NMDA transmission in a genetic model of absence epilepsy in rats. *Epilepsy Res* 25 (1), 11–19. [PubMed: 8886657]
- Kose B, Ozkan M, Sur-Erdem I, Cavdar S, 2022. Does astrocyte gap junction protein expression differ during development in absence epileptic rats? *Synapse* 76 (3–4), e22225. [PubMed: 35137459]

- Kostopoulos GK, 2000. Spike-and-wave discharges of absence seizures as a transformation of sleep spindles: the continuing development of a hypothesis. *Clin. Neurophysiol* 111 (Suppl. 2), S27–S38. [PubMed: 10996552]
- Kramer MA, Stoyell SM, Chinappen D, Ostrowski LM, Spencer ER, Morgan AK, Emerton BC, Jing J, Westover MB, Eden UT, Stickgold R, Manoach DS, Chu CJ, 2021. Focal sleep spindle deficits reveal focal thalamocortical dysfunction and predict cognitive deficits in sleep activated developmental epilepsy. *J. Neurosci* 41 (8), 1816–1829. [PubMed: 33468567]
- Kros L, Eelkman Rooda OHJ, Spanke JK, Alva P, van Dongen MN, Karapatis A, Tolner EA, Strydis C, Davey N, Winkelman BHJ, Negrello M, Serdijn WA, Steuber V, van den Maagdenberg AMJM, De Zeeuw CI, Hoebeek FE, 2015. Cerebellar output controls generalized spike-and-wave discharge occurrence. *Ann. Neurol* 77 (6), 1027–1049. [PubMed: 25762286]
- Kverneland M, Selmer KK, Nakken KO, Iversen PO, Tauboll E, 2015. A prospective study of the modified Atkins diet for adults with idiopathic generalized epilepsy. *Epilepsy Behav* 53, 197–201. [PubMed: 26588588]
- Kwon C-S, Schupper AJ, Fields MC, Marcuse LV, La Vega-Talbott M, Panov F, Ghatan S, 2020. Centromedian thalamic responsive neurostimulation for Lennox-Gastaut epilepsy and autism. *Ann. Clin. Transl. Neurol* 7 (10), 2035–2040. [PubMed: 32860345]
- Lacey CJ, Bryant A, Brill J, Huguenard JR, 2012. Enhanced NMDA receptor-dependent thalamic excitation and network oscillations in stargazer mice. *J. Neurosci* 32 (32), 11067–11081. [PubMed: 22875939]
- Levesque M, Biagini G, de Curtis M, Gnatkovsky V, Pitsch J, Wang S, Avoli M, 2021. The pilocarpine model of mesial temporal lobe epilepsy: over one decade later, with more rodent species and new investigative approaches. *Neurosci. Biobehav. Rev* 130, 274–291. [PubMed: 34437936]
- Liguori C, Toledo M, Kothare S, 2021. Effects of anti-seizure medications on sleep architecture and daytime sleepiness in patients with epilepsy: a literature review. *Sleep Med. Rev* 60, 101559. [PubMed: 34710770]
- Liu Z, Snead OC 3rd, Vergnes M, Depaulis A, Marescaux C, 1991. Intrathalamic injections of gamma-hydroxybutyric acid increase genetic absence seizures in rats. *Neurosci. Lett* 125 (1), 19–21. [PubMed: 1857553]
- Livingston S, Torres I, Pauli LL, Rider RV, 1965. Petit mal epilepsy. Results of a prolonged follow-up study of 117 patients. *Jama* 194 (3), 227–232. [PubMed: 4953647]
- Loiseau J, Loiseau P, Guyot M, Duche B, Dartigues JF, Aublet B, 1990. Survey of seizure disorders in the French southwest. I. Incidence of epileptic syndromes. *Epilepsia* 31 (4), 391–396. [PubMed: 2369875]
- Lovatt D, Xu Q, Liu W, Takano T, Smith NA, Schnermann J, Tieu K, Nedergaard M, 2012. Neuronal adenosine release, and not astrocytic ATP release, mediates feedback inhibition of excitatory activity. *Proc. Natl. Acad. Sci. U. S. A* 109 (16), 6265–6270. [PubMed: 22421436]
- Ludwig A, Budde T, Stieber J, Moosmang S, Wahl C, Holthoff K, Langebartels A, Wotjak C, Munsch T, Zong X, Feil S, Feil R, Lancel M, Chien KR, Konnerth A, Pape HC, Biel M, Hofmann F, 2003. Absence epilepsy and sinus dysrhythmia in mice lacking the pacemaker channel HCN2. *EMBO J* 22 (2), 216–224. [PubMed: 12514127]
- Luttjohann A, Pape HC, 2019. Regional specificity of cortico-thalamic coupling strength and directionality during waxing and waning of spike and wave discharges. *Sci. Rep* 9 (1), 2100. [PubMed: 30765744]
- Luttjohann A, van Luijckelaar G, 2012. The dynamics of cortico-thalamo-cortical interactions at the transition from pre-ictal to ictal LFPs in absence epilepsy. *Neurobiol. Dis* 47 (1), 49–60. [PubMed: 22465080]
- Luttjohann A, Schoffelen JM, van Luijckelaar G, 2013. Peri-ictal network dynamics of spike-wave discharges: phase and spectral characteristics. *Exp. Neurol* 239, 235–247. [PubMed: 23124095]
- Ma X, Zhang Y, Yang Z, Liu X, Sun H, Qin J, Wu X, Liang J, 2011. Childhood absence epilepsy: electroclinical features and diagnostic criteria. *Brain and Development* 33 (2), 114–119. [PubMed: 20378290]

- Maheshwari A, Noebels JL, 2014. Monogenic models of absence epilepsy: windows into the complex balance between inhibition and excitation in thalamocortical microcircuits. *Prog. Brain Res* 213, 223–252. [PubMed: 25194492]
- Makinson CD, Tanaka BS, Sorokin JM, Wong JC, Christian CA, Goldin AL, Escayg A, Huguenard JR, 2017. Regulation of thalamic and cortical network synchrony by Scn8a. *Neuron* 93 (5), 1165–1179 e1166. [PubMed: 28238546]
- Manning JP, Richards DA, Leresche N, Crunelli V, Bowery NG, 2004. Cortical-area specific block of genetically determined absence seizures by ethosuximide. *Neuroscience* 123 (1), 5–9. [PubMed: 14667436]
- Marescaux C, Vergnes M, Depaulis A, 1992. Genetic absence epilepsy in rats from Strasbourg—a review. *J. Neural Transm. Suppl* 35, 37–69. [PubMed: 1512594]
- Marini C, Harkin LA, Wallace RH, Mulley JC, Scheffer IE, Berkovic SF, 2003. Childhood absence epilepsy and febrile seizures: a family with a GABA(A) receptor mutation. *Brain* 126 (Pt 1), 230–240. [PubMed: 12477709]
- Marini C, Scheffer IE, Crossland KM, Grinton BE, Phillips FL, McMahon JM, Turner SJ, Dean JT, Kivity S, Mazarib A, Neufeld MY, Korczyn AD, Harkin LA, Dibbens LM, Wallace RH, Mulley JC, Berkovic SF, 2004. Genetic architecture of idiopathic generalized epilepsy: clinical genetic analysis of 55 multiplex families. *Epilepsia* 45 (5), 467–478. [PubMed: 15101828]
- Martin-McGill KJ, Bresnahan R, Levy RG, Cooper PN, 2020. Ketogenic diets for drug-resistant epilepsy. *Cochrane Database Syst. Rev* 6 (6), CD001903. [PubMed: 32588435]
- Mastroianni G, Ascoli M, Gasparini S, Brigo F, Cianci V, Neri S, Russo E, Aguglia U, Ferlazzo E, 2021. Therapeutic approach to difficult-to-treat typical absences and related epilepsy syndromes. *Expert. Rev. Clin. Pharmacol* 14 (11), 1427–1433. [PubMed: 34289757]
- McCafferty C, David F, Venzi M, Lorincz ML, Delicata F, Atherton Z, Recchia G, Orban G, Lambert RC, Di Giovanni G, Leresche N, Crunelli V, 2018. Cortical drive and thalamic feed-forward inhibition control thalamic output synchrony during absence seizures. *Nat. Neurosci* 21 (5), 744–756. [PubMed: 29662216]
- McCarthy E, Shakil F, Saint Ange P, Morris Cameron E, Miller J, Pathak S, Greenberg DA, Veliskova J, Velisek L, 2020. Developmental decrease in parvalbumin-positive neurons precedes increase in flurothyl-induced seizure susceptibility in the Brd2(+/-) mouse model of juvenile myoclonic epilepsy. *Epilepsia* 61 (5), 892–902. [PubMed: 32301507]
- McCormick DA, Pape HC, 1990. Noradrenergic and serotonergic modulation of a hyperpolarization-activated cation current in thalamic relay neurones. *J. Physiol* 431, 319–342. [PubMed: 1712844]
- McCormick DA, Prince DA, 1988. Noradrenergic modulation of firing pattern in guinea pig and cat thalamic neurons, in vitro. *J. Neurophysiol* 59 (3), 978–996. [PubMed: 3367206]
- McCormick DA, von Krosigk M, 1992. Corticothalamic activation modulates thalamic firing through glutamate “metabotropic” receptors. *Proc. Natl. Acad. Sci. U. S. A* 89 (7), 2774–2778. [PubMed: 1313567]
- McCormick DA, Pape HC, Williamson A, 1991. Actions of norepinephrine in the cerebral cortex and thalamus: implications for function of the central noradrenergic system. *Prog. Brain Res* 88, 293–305. [PubMed: 1726028]
- McTague A, Howell KB, Cross JH, Kurian MA, Scheffer IE, 2016. The genetic landscape of the epileptic encephalopathies of infancy and childhood. *Lancet Neurol* 15 (3), 304–316. [PubMed: 26597089]
- Meeren HK, Pijn JP, Van Luijckelaar EL, Coenen AM, Lopes da Silva FH, 2002. Cortical focus drives widespread corticothalamic networks during spontaneous absence seizures in rats. *J. Neurosci* 22 (4), 1480–1495. [PubMed: 11850474]
- Meeren HK, Veening JG, Modersheim TA, Coenen AM, van Luijckelaar G, 2009. Thalamic lesions in a genetic rat model of absence epilepsy: dissociation between spike-wave discharges and sleep spindles. *Exp. Neurol* 217 (1), 25–37. [PubMed: 19416679]
- Melo TM, Nehlig A, Sonnewald U, 2006. Neuronal-glia interactions in rats fed a ketogenic diet. *Neurochem. Int* 48 (6–7), 498–507. [PubMed: 16542760]

- Meuth SG, Kanyshkova T, Meuth P, Landgraf P, Munsch T, Ludwig A, Hofmann F, Pape HC, Budde T, 2006. Membrane resting potential of thalamocortical relay neurons is shaped by the interaction among TASK3 and HCN2 channels. *J. Neurophysiol* 96 (3), 1517–1529. [PubMed: 16760342]
- Miao QL, Herlitz S, Mark MD, Noebels JL, 2020. Adult loss of *Cacna1a* in mice recapitulates childhood absence epilepsy by distinct thalamic bursting mechanisms. *Brain* 143 (1), 161–174. [PubMed: 31800012]
- Morrison JH, Foote SL, 1986. Noradrenergic and serotonergic innervation of cortical, thalamic, and tectal visual structures in Old and New World monkeys. *J. Comp. Neurol* 243 (1), 117–138. [PubMed: 3950077]
- Morse E, Giblin K, Chung MH, Dohle C, Berg AT, Blumenfeld H, 2019. Historical trend toward improved long-term outcome in childhood absence epilepsy. *Epilepsy Res* 152, 7–10. [PubMed: 30856420]
- Moxon KA, Devilbiss DM, Chapin JK, Waterhouse BD, 2007. Influence of norepinephrine on somatosensory neuronal responses in the rat thalamus: a combined modeling and in vivo multi-channel, multi-neuron recording study. *Brain Res* 1147, 105–123. [PubMed: 17368434]
- Munoz-Ballester C, Berthier A, Viana R, Sanz P, 2016. Homeostasis of the astrocytic glutamate transporter GLT-1 is altered in mouse models of Lafora disease. *Biochim. Biophys. Acta* 1862 (6), 1074–1083. [PubMed: 26976331]
- Munoz-Ballester C, Santana N, Perez-Jimenez E, Viana R, Artigas F, Sanz P, 2019. In vivo glutamate clearance defects in a mouse model of Lafora disease. *Exp. Neurol* 320, 112959. [PubMed: 31108086]
- Najarian T, Marrache AM, Dumont I, Hardy P, Beauchamp MH, Hou X, Peri K, Gobeil F Jr., Varma DR, Chemtob S, 2000. Prolonged hypercapnia-evoked cerebral hyperemia via K(+) channel- and prostaglandin E(2)-dependent endothelial nitric oxide synthase induction. *Circ. Res* 87 (12), 1149–1156. [PubMed: 11110772]
- Nayel M, Awad IA, Larkins M, Lüders H, 1991. Experimental limbic epilepsy: models, pathophysiologic concepts, and clinical relevance. *Cleve. Clin. J. Med* 58 (6), 521–530. [PubMed: 1752034]
- Newman EA, 2004. Glial modulation of synaptic transmission in the retina. *Glia* 47 (3), 268–274. [PubMed: 15252816]
- Niemeyer MI, Cid LP, Pena-Munzenmayer G, Sepulveda FV, 2010. Separate gating mechanisms mediate the regulation of K2P potassium channel TASK-2 by intra- and extracellular pH. *J. Biol. Chem* 285 (22), 16467–16475. [PubMed: 20351106]
- Nishitani A, Kunisawa N, Sugimura T, Sato K, Yoshida Y, Suzuki T, Sakuma T, Yamamoto T, Asano M, Saito Y, Ohno Y, Kuramoto T, 2019. Loss of HCN1 subunits causes absence epilepsy in rats. *Brain Res* 1706, 209–217. [PubMed: 30408474]
- Nofzinger EA, 2005. Functional neuroimaging of sleep. *Semin. Neurol* 25 (1), 9–18. [PubMed: 15798933]
- Ogiwara I, Miyamoto H, Tatsukawa T, Yamagata T, Nakayama T, Atapour N, Miura E, Mazaki E, Ernst SJ, Cao D, Ohtani H, Itohara S, Yanagawa Y, Montal M, Yuzaki M, Inoue Y, Hensch TK, Noebels JL, Yamakawa K, 2018. *Nav1.2* haplodeficiency in excitatory neurons causes absence-like seizures in mice. *Commun. Biol* 1, 96. [PubMed: 30175250]
- Okamoto N, Hori S, Akazawa C, Hayashi Y, Shigemoto R, Mizuno N, Nakanishi S, 1994. Molecular characterization of a new metabotropic glutamate receptor mGluR7 coupled to inhibitory cyclic AMP signal transduction. *J. Biol. Chem* 269 (2), 1231–1236. [PubMed: 8288585]
- Olsson I, 1988. Epidemiology of absence epilepsy. I. Concept and incidence. *Acta Paediatr. Scand* 77 (6), 860–866. [PubMed: 3144826]
- O'Malley JJ, Seibt F, Chin J, Beierlein M, 2020. TRPM4 conductances in thalamic reticular nucleus neurons generate persistent firing during slow oscillations. *J. Neurosci* 40 (25), 4813–4823. [PubMed: 32414784]
- Onat FY, van Luijtelaaar G, Nehlig A, Snead OC 3rd, 2013. The involvement of limbic structures in typical and atypical absence epilepsy. *Epilepsy Res* 103 (2–3), 111–123. [PubMed: 22989853]

- Ozgun M, Ozyurt MG, Arkan S, Cavdar S, 2022. The effects of optogenetic activation of astrocytes on spike-and-wave discharges in genetic absence epileptic rats. *Ann. Neurosci* 29 (1), 53–61. [PubMed: 35875425]
- Panayiotopoulos CP, 1999. Typical absence seizures and their treatment. *Arch. Dis. Child* 81 (4), 351–355. [PubMed: 10490445]
- Panayiotopoulos CP, Obeid T, Waheed G, 1989a. Absences in juvenile myoclonic epilepsy: a clinical and video-electroencephalographic study. *Ann. Neurol* 25 (4), 391–397. [PubMed: 2496640]
- Panayiotopoulos CP, Obeid T, Waheed G, 1989b. Differentiation of typical absence seizures in epileptic syndromes. A video EEG study of 224 seizures in 20 patients. *Brain* 112 (Pt 4), 1039–1056. [PubMed: 2505885]
- Pankratov Y, Lalo U, Verkhatsky A, North RA, 2007. Quantal release of ATP in mouse cortex. *J. Gen. Physiol* 129 (3), 257–265. [PubMed: 17325196]
- Pape HC, McCormick DA, 1989. Noradrenaline and serotonin selectively modulate thalamic burst firing by enhancing a hyperpolarization-activated cation current. *Nature* 340 (6236), 715–718. [PubMed: 2475782]
- Parri HR, Crunelli V, 2001. Pacemaker calcium oscillations in thalamic astrocytes in situ. *Neuroreport* 12 (18), 3897–3900. [PubMed: 11742206]
- Pascual O, Casper KB, Kubera C, Zhang J, Revilla-Sanchez R, Sul JY, Takano H, Moss SJ, McCarthy K, Haydon PG, 2005. Astrocytic purinergic signaling coordinates synaptic networks. *Science* 310 (5745), 113–116. [PubMed: 16210541]
- Paz JT, Huguenard JR, 2015a. Microcircuits and their interactions in epilepsy: is the focus out of focus? *Nat. Neurosci* 18 (3), 351–359. [PubMed: 25710837]
- Paz JT, Huguenard JR, 2015b. Optogenetics and epilepsy: past, present and future. *Epilepsy Curr* 15 (1), 34–38. [PubMed: 25678887]
- Paz JT, Chavez M, Saillet S, Deniau JM, Charpier S, 2007. Activity of ventral medial thalamic neurons during absence seizures and modulation of cortical paroxysms by the nigrothalamic pathway. *J. Neurosci* 27 (4), 929–941. [PubMed: 17251435]
- Paz JT, Bryant AS, Peng K, Fenno L, Yizhar O, Frankel WN, Deisseroth K, Huguenard JR, 2011. A new mode of corticothalamic transmission revealed in the *Gria4(-/-)* model of absence epilepsy. *Nat. Neurosci* 14 (9), 1167–1173. [PubMed: 21857658]
- Pérez-Santos I, Palomero-Gallagher N, Zilles K, Cavada C, 2021. Distribution of the noradrenaline innervation and adrenoceptors in the Macaque Monkey Thalamus. *Cereb. Cortex* 31 (9), 4115–4139. [PubMed: 34003210]
- Petrou S, Reid CA, 2012. The GABA(A)gamma2(R43Q) mouse model of human genetic epilepsy. In: Noebels JL, Avoli M, Rogawski MA, Olsen RW, Delgado-Escueta AV (Eds.), *Jasper's Basic Mechanisms of the Epilepsies* Bethesda (MD).
- Philippot C, Griemsmann S, Jabs R, Seifert G, Kettenmann H, Steinhauser C, 2021. Astrocytes and oligodendrocytes in the thalamus jointly maintain synaptic activity by supplying metabolites. *Cell Rep* 34 (3), 108642. [PubMed: 33472059]
- Pietrafusa N, La Neve A, de Palma L, Boero G, Luisi C, Vigeveno F, Specchio N, 2021. Juvenile myoclonic epilepsy: long-term prognosis and risk factors. *Brain and Development* 43 (6), 688–697. [PubMed: 33781581]
- Pinault D, Leresche N, Charpier S, Deniau JM, Marescaux C, Vergnes M, Crunelli V, 1998. Intracellular recordings in thalamic neurones during spontaneous spike and wave discharges in rats with absence epilepsy. *J. Physiol* 509 (Pt 2), 449–456. [PubMed: 9575294]
- Pirttimaki T, Parri HR, Crunelli V, 2013. Astrocytic GABA transporter GAT-1 dysfunction in experimental absence seizures. *J. Physiol* 591 (4), 823–833. [PubMed: 23090943]
- Polack PO, Mahon S, Chavez M, Charpier S, 2009. Inactivation of the somatosensory cortex prevents paroxysmal oscillations in cortical and related thalamic neurons in a genetic model of absence epilepsy. *Cereb. Cortex* 19 (9), 2078–2091. [PubMed: 19276326]
- Prassouli A, Katsarou E, Attilakos A, Antoniadou I, 2007. Learning difficulties in children with epilepsy with idiopathic generalized epilepsy and well-controlled seizures. *Dev. Med. Child Neurol* 49 (11), 874 author reply 874–875. [PubMed: 17979868]

- Ritter-Makinson S, Clemente-Perez A, Higashikubo B, Cho FS, Holden SS, Bennett E, Chkhaidze A, Eelkman Rooda OHJ, Cornet MC, Hoebeek FE, Yamakawa K, Cilio MR, Delord B, Paz JT, 2019. Augmented reticular thalamic bursting and seizures in *Scn1a*-Dravet syndrome. *Cell Rep* 26 (1), 54–64 e56. [PubMed: 30605686]
- Rohracher A, Brigo F, Höfler J, Kalss G, Neuray C, Dobesberger J, Kuchukhidze G, Leitinger M, Trinkka E, 2016. Perampanel for the treatment of primary generalized tonic-clonic seizures in idiopathic generalized epilepsy. *Expert. Opin. Pharmacother* 17 (10), 1403–1411. [PubMed: 27267634]
- Ross DL, Swaiman KF, Torres F, Hansen J, 1985. Early biochemical and EEG correlates of the ketogenic diet in children with atypical absence epilepsy. *Pediatr. Neurol* 1 (2), 104–108. [PubMed: 3939741]
- Sadleir LG, Farrell K, Smith S, Connolly MB, Scheffer IE, 2006. Electroclinical features of absence seizures in childhood absence epilepsy. *Neurology* 67 (3), 413–418. [PubMed: 16894100]
- Sadleir LG, Scheffer IE, Smith S, Carstensen B, Farrell K, Connolly MB, 2009. EEG features of absence seizures in idiopathic generalized epilepsy: impact of syndrome, age, and state. *Epilepsia* 50 (6), 1572–1578. [PubMed: 19243419]
- Salanova V, Sperling MR, Gross RE, Irwin CP, Vollhaber JA, Giftakis JE, Fisher RS, 2021. The SANTÉ study at 10 years of follow-up: effectiveness, safety, and sudden unexpected death in epilepsy. *Epilepsia* 62 (6), 1306–1317. [PubMed: 33830503]
- Salvati KA, Beenhakker MP, 2019. Out of thin air: hyperventilation-triggered seizures. *Brain Res* 1703, 41–52. [PubMed: 29288644]
- Salvati KA, Souza G, Lu AC, Ritger ML, Guyenet P, Abbott SB, Beenhakker MP, 2022. Respiratory alkalosis provokes spike-wave discharges in seizure-prone rats. *Elife* 11.
- Schofield CM, Kleiman-Weiner M, Rudolph U, Huguenard JR, 2009. A gain in GABAA receptor synaptic strength in thalamus reduces oscillatory activity and absence seizures. *Proc. Natl. Acad. Sci. U. S. A* 106 (18), 7630–7635. [PubMed: 19380748]
- Schwitalla JC, Pakusch J, Mucher B, Bruckner A, Depke DA, Fenzl T, De Zeeuw CI, Kros L, Hoebeek FE, Mark MD, 2022. Controlling absence seizures from the cerebellar nuclei via activation of the Gq signaling pathway. *Cell. Mol. Life Sci* 79 (4), 197. [PubMed: 35305155]
- Seicchitano F, van Rijn CM, van Luijteleaer G, 2015. Unilateral and bilateral cortical resection: effects on Spike-Wave discharges in a genetic absence epilepsy model. *PLoS One* 10 (8), e0133594. [PubMed: 26262879]
- Seidenbecher T, Pape HC, 2001. Contribution of intralaminar thalamic nuclei to spike-and-wave discharges during spontaneous seizures in a genetic rat model of absence epilepsy. *Eur. J. Neurosci* 13 (8), 1537–1546. [PubMed: 11328348]
- Seidenbecher T, Staak R, Pape HC, 1998. Relations between cortical and thalamic cellular activities during absence seizures in rats. *Eur. J. Neurosci* 10 (3), 1103–1112. [PubMed: 9753178]
- Seneviratne U, Hepworth G, Cook M, D'Souza W, 2017. Can EEG differentiate among syndromes in genetic generalized epilepsy? *J. Clin. Neurophysiol* 34 (3), 213–221. [PubMed: 27763964]
- Senf P, Schmitz B, Holtkamp M, Janz D, 2013. Prognosis of juvenile myoclonic epilepsy 45 years after onset: seizure outcome and predictors. *Neurology* 81 (24), 2128–2133. [PubMed: 24212391]
- Sezikli S, Pulat TA, Tekin B, Ak PD, Keskinkılıç C, Ataklı D, 2018. Frontal lobe cognitive functions and electroencephalographic features in juvenile myoclonic epilepsy. *Epilepsy Behav* 86, 102–107. [PubMed: 30017834]
- Sherman SM, Guillery RW, 1996. Functional organization of thalamocortical relays. *J. Neurophysiol* 76 (3), 1367–1395. [PubMed: 8890259]
- Sherman SM, Guillery RW, 1998. On the actions that one nerve cell can have on another: distinguishing “drivers” from “modulators”. *Proc. Natl. Acad. Sci. U. S. A* 95 (12), 7121–7126. [PubMed: 9618549]
- Sherwin I, 1965. Differential effects of hyperventilation on the excitability of intact and isolated cortex. *Electroencephalogr. Clin. Neurophysiol* 18, 599–607. [PubMed: 14296837]
- Sherwin I, 1967. Alterations in the non-specific cortical afference during hyperventilation. *Electroencephalogr. Clin. Neurophysiol* 23 (6), 532–538. [PubMed: 4169823]

- Shinnar RC, Shinnar S, Cnaan A, Clark P, Dlugos D, Hirtz DG, Hu F, Liu C, Masur D, Weiss EF, Glauser TA, 2017. Pretreatment behavior and subsequent medication effects in childhood absence epilepsy. *Neurology* 89 (16), 1698–1706. [PubMed: 28916534]
- Simpson KL, Altman DW, Wang L, Kirifides ML, Lin RC, Waterhouse BD, 1997. Lateralization and functional organization of the locus coeruleus projection to the trigeminal somatosensory pathway in rat. *J. Comp. Neurol* 385 (1), 135–147. [PubMed: 9268121]
- Sitnikova E, van Luijtelaar G, 2004. Cortical control of generalized absence seizures: effect of lidocaine applied to the somatosensory cortex in WAG/Rij rats. *Brain Res* 1012 (1–2), 127–137. [PubMed: 15158169]
- Slaght SJ, Leresche N, Deniau JM, Crunelli V, Charpier S, 2002. Activity of thalamic reticular neurons during spontaneous genetically determined spike and wave discharges. *J. Neurosci* 22 (6), 2323–2334. [PubMed: 11896171]
- Sohal VS, Huntsman MM, Huguenard JR, 2000. Reciprocal inhibitory connections regulate the spatiotemporal properties of intrathalamic oscillations. *J. Neurosci* 20 (5), 1735–1745. [PubMed: 10684875]
- Sohal VS, Keist R, Rudolph U, Huguenard JR, 2003. Dynamic GABA(A) receptor subtype-specific modulation of the synchrony and duration of thalamic oscillations. *J. Neurosci* 23 (9), 3649–3657. [PubMed: 12736336]
- Soltesz I, Lightowler S, Leresche N, Jassik-Gerschenfeld D, Pollard CE, Crunelli V, 1991. Two inward currents and the transformation of low-frequency oscillations of rat and cat thalamocortical cells. *J. Physiol* 441, 175–197. [PubMed: 1667794]
- Sondhi V, Agarwala A, Pandey RM, Chakrabarty B, Jauhari P, Lodha R, Toteja GS, Sharma S, Paul VK, Kossoff E, Gulati S, 2020. Efficacy of ketogenic diet, modified Atkins diet, and low glycemic index therapy diet among children with drug-resistant epilepsy: a randomized clinical trial. *JAMA Pediatr* 174 (10), 944–951. [PubMed: 32761191]
- Song I, Kim D, Choi S, Sun M, Kim Y, Shin HS, 2004. Role of the alpha1G T-type calcium channel in spontaneous absence seizures in mutant mice. *J. Neurosci* 24 (22), 5249–5257. [PubMed: 15175395]
- Sorokin JM, Davidson TJ, Frechette E, Abramian AM, Deisseroth K, Huguenard JR, Paz JT, 2017. Bidirectional control of generalized epilepsy networks via rapid real-time switching of firing mode. *Neuron* 93 (1), 194–210. [PubMed: 27989462]
- Spoor JKH, Greco T, Kamp MA, Faini S, Senft C, Dibue M, 2021. Quantifying the burden of disease in patients with Lennox Gastaut syndrome. *Epilepsy Behav. Rep* 16, 100508. [PubMed: 34917922]
- Staak R, Pape HC, 2001. Contribution of GABA(A) and GABA(B) receptors to thalamic neuronal activity during spontaneous absence seizures in rats. *J. Neurosci* 21 (4), 1378–1384. [PubMed: 11160409]
- Stamberger H, Nikanorova M, Willemsen MH, Accorsi P, Angriman M, Baier H, Benkel-Herrenbrueck I, Benoit V, Budetta M, Caliebe A, Cantalupo G, Capovilla G, Casara G, Courage C, Deprez M, Destree A, Dilella R, Erasmus CE, Fannemel M, Fjaer R, Giordano L, Helbig KL, Heyne HO, Klepper J, Kluger GJ, Lederer D, Lodi M, Maier O, Merckenschlager A, Michelberger N, Minetti C, Muhle H, Phalin J, Ramsey K, Romeo A, Schallner J, Schanze I, Shinawi M, Slegers K, Sterbova K, Syrbe S, Traverso M, Tzschach A, Uldall P, Van Coster R, Verhelst H, Viri M, Winter S, Wolff M, Zenker M, Zoccante L, De Jonghe P, Helbig I, Striano P, Lemke JR, Moller RS, Weckhuysen S, 2016. STXBP1 encephalopathy: a neurodevelopmental disorder including epilepsy. *Neurology* 86 (10), 954–962. [PubMed: 26865513]
- Steriade M, Deschênes M, Domich L, Mulle C, 1985. Abolition of spindle oscillations in thalamic neurons disconnected from nucleus reticularis thalami. *J. Neurophysiol* 54 (6), 1473–1497. [PubMed: 4087044]
- Steriade M, Domich L, Oakson G, Deschenes M, 1987. The deafferented reticular thalamic nucleus generates spindle rhythmicity. *J. Neurophysiol* 57 (1), 260–273. [PubMed: 3559675]
- Stevellink R, Koeleman BPC, Sander JW, Jansen FE, Braun KPJ, 2019. Refractory juvenile myoclonic epilepsy: a meta-analysis of prevalence and risk factors. *Eur. J. Neurol* 26 (6), 856–864. [PubMed: 30223294]

- Streng ML, Krook-Magnuson E, 2021. The cerebellum and epilepsy. *Epilepsy Behav. E&B* 121 (Pt B), 106909.
- Strzelczyk A, Schubert-Bast S, 2022. Psychobehavioural and cognitive adverse events of anti-seizure medications for the treatment of developmental and epileptic encephalopathies. *CNS Drugs* 36 (10), 1079–1111. [PubMed: 36194365]
- Studtmann C, Ladislav M, Topolski MA, Safari M, Swanger SA, 2022. Na(V)1.1 haploinsufficiency impairs glutamatergic and GABAergic neuron function in the thalamus. *Neurobiol. Dis* 167, 105672. [PubMed: 35219855]
- Svichar N, Chesler M, 2003. Surface carbonic anhydrase activity on astrocytes and neurons facilitates lactate transport. *Glia* 41 (4), 415–419. [PubMed: 12555208]
- Swanson LW, Hartman BK, 1975. The central adrenergic system. An immunofluorescence study of the location of cell bodies and their efferent connections in the rat utilizing dopamine-beta-hydroxylase as a marker. *J. Comp. Neurol* 163 (4), 467–505. [PubMed: 1100685]
- Symonds JD, Zuberi SM, Stewart K, McLellan A, O'Regan M, MacLeod S, Jollands A, Joss S, Kirkpatrick M, Brunklaus A, Pilz DT, Shetty J, Dorris L, Abu-Arafeh I, Andrew J, Brink P, Callaghan M, Cruden J, Diver LA, Findlay C, Gardiner S, Grattan R, Lang B, MacDonnell J, McKnight J, Morrison CA, Nairn L, Slean MM, Stephen E, Webb A, Vincent A, Wilson M, 2019. Incidence and phenotypes of childhood-onset genetic epilepsies: a prospective population-based national cohort. *Brain* 142 (8), 2303–2318. [PubMed: 31302675]
- Sysoeva MV, Luttjohann A, van Luijckelaar G, Sysoev IV, 2016. Dynamics of directional coupling underlying spike-wave discharges. *Neuroscience* 314, 75–89. [PubMed: 26633265]
- Syvertsen M, Nakken KO, Edland A, Hansen G, Hellum MK, Koht J, 2015. Prevalence and etiology of epilepsy in a Norwegian county-A population based study. *Epilepsia* 56 (5), 699–706. [PubMed: 25810072]
- Syvertsen M, Hellum MK, Hansen G, Edland A, Nakken KO, Selmer KK, Koht J, 2017. Prevalence of juvenile myoclonic epilepsy in people <30 years of age-A population-based study in Norway. *Epilepsia* 58 (1), 105–112. [PubMed: 27861775]
- Syvertsen M, Selmer K, Enger U, Nakken KO, Pal DK, Smith A, Koht J, 2019. Psychosocial complications in juvenile myoclonic epilepsy. *Epilepsy Behav* 90, 122–128. [PubMed: 30530133]
- Tassin V, Girard B, Chotte A, Fontanaud P, Rigault D, Kalinichev M, Perroy J, Acher F, Fagnia L, Bertaso F, 2016. Phasic and tonic mGlu7 receptor activity modulates the thalamocortical network. *Front. Neural Circuits* 10, 31. [PubMed: 27199672]
- Taura M, Gama AP, Sousa AVM, Noffs MHS, Alonso NB, Yacubian EM, Guilhoto LM, 2020. Dysfunctional personality beliefs and executive performance in patients with juvenile myoclonic epilepsy. *Epilepsy Behav* 105, 106958. [PubMed: 32097884]
- Terlau J, Yang JW, Khastkhodaei Z, Seidenbecher T, Luhmann HJ, Pape HC, Luttjohann A, 2020. Spike-wave discharges in absence epilepsy: segregation of electrographic components reveals distinct pathways of seizure activity. *J. Physiol* 598 (12), 2397–2414. [PubMed: 32144956]
- Tezer FI, Rémi J, Erbil N, Noachtar S, Saygi S, 2014. A reduction of sleep spindles heralds seizures in focal epilepsy. *Clin. Neurophysiol* 125 (11), 2207–2211. [PubMed: 24684944]
- Thompson JA, Miralles RM, Wengert ER, Wagley PK, Yu W, Wenker IC, Patel MK, 2022. Astrocyte reactivity in a mouse model of SCN8A epileptic encephalopathy. *Epilepsia Open* 7 (2), 280–292. [PubMed: 34826216]
- Tissone AI, Vidal VB, Nadal MS, Mato G, Amarillo Y, 2021. Differential contribution of the subthreshold operating currents I(T), I(h), and I(Kir) to the resonance of thalamocortical neurons. *J. Neurophysiol* 126 (2), 561–574. [PubMed: 34232785]
- Tolmacheva EA, van Luijckelaar G, Chepurinov SA, Kaminskij Y, Mares P, 2004. Cortical and limbic excitability in rats with absence epilepsy. *Epilepsy Res* 62 (2–3), 189–198. [PubMed: 15579307]
- Trinka E, Baumgartner S, Unterberger I, Unterrainer J, Luef G, Haberlandt E, Bauer G, 2004. Long-term prognosis for childhood and juvenile absence epilepsy. *J. Neurol* 251 (10), 1235–1241. [PubMed: 15503104]

- Tsakiridou E, Bertollini L, de Curtis M, Avanzini G, Pape HC, 1995. Selective increase in T-type calcium conductance of reticular thalamic neurons in a rat model of absence epilepsy. *J. Neurosci* 15 (4), 3110–3117. [PubMed: 7722649]
- Uchino K, Tanaka Y, Ikezawa W, Deshimaru M, Kubota K, Watanabe T, Katsurabayashi S, Iwasaki K, Hirose S, 2023. Astrocyte Ca(2+) signaling is facilitated in Scn1a(+/-) mouse model of Dravet syndrome. *Biochem. Biophys. Res. Commun* 643, 169–174. [PubMed: 36610382]
- Valentin A, Hindocha N, Osei-Lah A, Fisniku L, McCormick D, Asherson P, Moran N, Makoff A, Nashef L, 2007. Idiopathic generalized epilepsy with absences: syndrome classification. *Epilepsia* 48 (11), 2187–2190. [PubMed: 17666074]
- Varela C, 2013. The gating of neocortical information by modulators. *J. Neurophysiol* 109 (5), 1229–1232. [PubMed: 22956786]
- Varela C, 2014. Thalamic neuromodulation and its implications for executive networks. *Front. Neural Circuits* 8, 69. [PubMed: 25009467]
- Velasco F, Velasco M, Ogarrio C, Fanghanel G, 1987. Electrical stimulation of the centromedian thalamic nucleus in the treatment of convulsive seizures: a preliminary report. *Epilepsia* 28 (4), 421–430. [PubMed: 3497802]
- Velasco M, Velasco F, Alcalá H, Davila G, Diaz-de-Leon AE, 1991. Epileptiform EEG activity of the centromedian thalamic nuclei in children with intractable generalized seizures of the Lennox-Gastaut syndrome. *Epilepsia* 32 (3), 310–321. [PubMed: 1904342]
- Velasco AL, Velasco F, Jiménez F, Velasco M, Castro G, Carrillo-Ruiz JD, Fanghanel G, Boleaga B, 2006. Neuromodulation of the centromedian thalamic nuclei in the treatment of generalized seizures and the improvement of the quality of life in patients with Lennox-Gastaut syndrome. *Epilepsia* 47 (7), 1203–1212. [PubMed: 16886984]
- Velasco F, Saucedo-Alvarado PE, Reichrath A, Valdés-Quiroz H, Aguado-Carrillo G, Velasco AL, 2021. Centromedian nucleus and epilepsy. *J. Clin. Neurophysiol* 38 (6), 485–493. [PubMed: 34261113]
- Vergnes M, Marescaux C, 1992. Cortical and thalamic lesions in rats with genetic absence epilepsy. *J. Neural Transm. Suppl* 35, 71–83. [PubMed: 1512595]
- Vogt BA, Hof PR, Friedman DP, Sikes RW, Vogt LJ, 2008. Norepinephrinergic afferents and cytology of the macaque monkey midline, mediodorsal, and intralaminar thalamic nuclei. *Brain Struct. Funct* 212 (6), 465–479. [PubMed: 18317800]
- Vorderwülbecke BJ, Kowski AB, Kirschbaum A, Merkle H, Senf P, Janz D, Holtkamp M, 2017. Long-term outcome in adolescent-onset generalized genetic epilepsies. *Epilepsia* 58 (7), 1244–1250. [PubMed: 28464258]
- Wall MJ, Dale N, 2013. Neuronal transporter and astrocytic ATP exocytosis underlie activity-dependent adenosine release in the hippocampus. *J. Physiol* 591 (16), 3853–3871. [PubMed: 23713028]
- Wall MJ, Puddefoot K, Yin W, Bingham C, Seifi M, Swinny JD, Ngomba RT, 2022. Adenosine is released during thalamic oscillations to provide negative feedback control. *Neuropharmacology* 216, 109172. [PubMed: 35780977]
- Walz W, 2000. Role of astrocytes in the clearance of excess extracellular potassium. *Neurochem. Int* 36 (4–5), 291–300. [PubMed: 10732996]
- Wandschneider B, Thompson PJ, Vollmar C, Koepf MJ, 2012. Frontal lobe function and structure in juvenile myoclonic epilepsy: a comprehensive review of neuropsychological and imaging data. *Epilepsia* 53 (12), 2091–2098. [PubMed: 23106095]
- Wang W, Frankel WN, 2021. Overlaps, gaps, and complexities of mouse models of developmental and epileptic encephalopathy. *Neurobiol. Dis* 148, 105220. [PubMed: 33301879]
- Wang X, Stewart L, Cortez MA, Wu Y, Velazquez JL, Liu CC, Shen L, Snead OC 3rd, 2009. The circuitry of atypical absence seizures in GABA(B)R1a transgenic mice. *Pharmacol. Biochem. Behav* 94 (1), 124–130. [PubMed: 19666047]
- Welch WP, Hect JL, Abel TJ, 2021. Case report: responsive neurostimulation of the centromedian thalamic nucleus for the detection and treatment of seizures in pediatric primary generalized epilepsy. *Front. Neurol* 12, 656585. [PubMed: 33995254]

- Weltha L, Reemmer J, Boison D, 2019. The role of adenosine in epilepsy. *Brain Res. Bull* 151, 46–54. [PubMed: 30468847]
- Whitmire CJ, Liew YJ, Stanley GB, 2021. Thalamic state influences timing precision in the thalamocortical circuit. *J. Neurophysiol* 125 (5), 1833–1850. [PubMed: 33760642]
- Wiemer-Kruel A, Haberlandt E, Hartmann H, Wohlrab G, Bast T, 2017. Modified Atkins diet is an effective treatment for children with Doose syndrome. *Epilepsia* 58 (4), 657–662. [PubMed: 28229464]
- Wirrell EC, Camfield CS, Camfield PR, Gordon KE, Dooley JM, 1996. Long-term prognosis of typical childhood absence epilepsy: remission or progression to juvenile myoclonic epilepsy. *Neurology* 47 (4), 912–918. [PubMed: 8857718]
- Wirrell EC, Camfield CS, Camfield PR, Dooley JM, Gordon KE, Smith B, 1997. Long-term psychosocial outcome in typical absence epilepsy. Sometimes a wolf in sheep's clothing. *Arch. Pediatr. Adolesc. Med* 151 (2), 152–158. [PubMed: 9041870]
- Wirrell EC, Grossardt BR, Wong-Kissel LC, Nickels KC, 2011. Incidence and classification of new-onset epilepsy and epilepsy syndromes in children in Olmsted County, Minnesota from 1980 to 2004: a population-based study. *Epilepsy Res* 95 (1–2), 110–118. [PubMed: 21482075]
- Wirrell EC, Nabbout R, Scheffer IE, Alsaadi T, Bogacz A, French JA, Hirsch E, Jain S, Kaneko S, Riney K, Samia P, Snead OC, Somerville E, Specchio N, Trinka E, Zuberi SM, Balestrini S, Wiebe S, Cross JH, Perucca E, Moshe SL, Tinuper P, 2022. Methodology for classification and definition of epilepsy syndromes with list of syndromes: report of the ILAE Task Force on Nosology and Definitions. *Epilepsia* 63 (6), 1333–1348. [PubMed: 35503715]
- Wolff M, Johannesen KM, Hedrich UBS, Masnada S, Rubboli G, Gardella E, Lesca G, Ville D, Milh M, Villard L, Afenjar A, Chantot-Bastarud S, Mignot C, Lardennois C, Nava C, Schwarz N, Gerard M, Perrin L, Doummar D, Auvin S, Miranda MJ, Hempel M, Brilstra E, Knoers N, Verbeek N, van Kempen M, Braun KP, Mancini G, Biskup S, Hortnagel K, Docker M, Bast T, Loddenkemper T, Wong-Kissel L, Baumeister FM, Fazeli W, Striano P, Dilena R, Fontana E, Zara F, Kurlemann G, Klepper J, Thoene JG, Arndt DH, Deconinck N, Schmitt-Mechelke T, Maier O, Muhle H, Wical B, Finetti C, Bruckner R, Pietz J, Golla G, Jillella D, Linnet KM, Charles P, Moog U, Oiglane-Shlik E, Mantovani JF, Park K, Deprez M, Lederer D, Mary S, Scalais E, Selim L, Van Coster R, Lagae L, Nikanorova M, Hjalgrim H, Korenke GC, Trivisano M, Specchio N, Ceulemans B, Dorn T, Helbig KL, Hardies K, Stamberger H, de Jonghe P, Weckhuysen S, Lemke JR, Krageloh-Mann I, Helbig I, Kluger G, Lerche H, Moller RS, 2017. Genetic and phenotypic heterogeneity suggest therapeutic implications in SCN2A-related disorders. *Brain* 140 (5), 1316–1336. [PubMed: 28379373]
- Wong AY, Borduas JF, Clarke S, Lee KF, Beique JC, Bergeron R, 2013. Calcium influx through N-type channels and activation of SK and TRP-like channels regulates tonic firing of neurons in rat paraventricular thalamus. *J. Neurophysiol* 110 (10), 2450–2464. [PubMed: 24004531]
- Yacubian EM, 2017. Juvenile myoclonic epilepsy: challenges on its 60th anniversary. *Seizure* 44, 48–52. [PubMed: 27665373]
- Yang S, Cox CL, 2007. Modulation of inhibitory activity by nitric oxide in the thalamus. *J. Neurophysiol* 97 (5), 3386–3395. [PubMed: 17376843]
- Yang S, Cox CL, 2008. Excitatory and anti-oscillatory actions of nitric oxide in thalamus. *J. Physiol* 586 (15), 3617–3628. [PubMed: 18535092]
- Yang S, Cox CL, 2011. Attenuation of inhibitory synaptic transmission by glial dysfunction in rat thalamus. *Synapse* 65 (12), 1298–1308. [PubMed: 21656574]
- Ying SW, Goldstein PA, 2005. Propofol-block of SK channels in reticular thalamic neurons enhances GABAergic inhibition in relay neurons. *J. Neurophysiol* 93 (4), 1935–1948. [PubMed: 15563549]
- Yudkoff M, Daikhin Y, Melo TM, Nissim I, Sonnewald U, Nissim I, 2007. The ketogenic diet and brain metabolism of amino acids: relationship to the anticonvulsant effect. *Annu. Rev. Nutr* 27, 415–430. [PubMed: 17444813]
- Zaman T, Lee K, Park C, Paydar A, Choi JH, Cheong E, Lee CJ, Shin HS, 2011. Cav2.3 channels are critical for oscillatory burst discharges in the reticular thalamus and absence epilepsy. *Neuron* 70 (1), 95–108. [PubMed: 21482359]

- Zhang Y, Mori M, Burgess DL, Noebels JL, 2002. Mutations in high-voltage-activated calcium channel genes stimulate low-voltage-activated currents in mouse thalamic relay neurons. *J. Neurosci* 22 (15), 6362–6371. [PubMed: 12151514]
- Zhang Y, Vilaythong AP, Yoshor D, Noebels JL, 2004. Elevated thalamic low-voltage-activated currents precede the onset of absence epilepsy in the SNAP25-deficient mouse mutant coloboma. *J. Neurosci* 24 (22), 5239–5248. [PubMed: 15175394]
- Zhang Y, Chen J, Ren J, Liu W, Yang T, Zhou D, 2019. Clinical features and treatment outcomes of Juvenile myoclonic epilepsy patients. *Epilepsia Open* 4 (2), 302–308. [PubMed: 31168497]
- Zobeiri M, Chaudhary R, Blaich A, Rottmann M, Herrmann S, Meuth P, Bista P, Kanyshkova T, Luttjohann A, Narayanan V, Hundehege P, Meuth SG, Romanelli MN, Urbano FJ, Pape HC, Budde T, Ludwig A, 2019. The hyperpolarization-activated HCN4 channel is important for proper maintenance of oscillatory activity in the thalamocortical system. *Cereb. Cortex* 29 (5), 2291–2304. [PubMed: 30877792]




Fig. 1. Clinical differences and areas of overlap within GGE, between IGEs and generalized DEEs. CAE: childhood absence epilepsy, DEE: developmental and epileptic encephalopathy, GTCA: generalized tonic-clonic seizures alone, IGE: idiopathic generalized epilepsy, JAE: juvenile absence epilepsy, JME: juvenile myoclonic epilepsy. Blue: features associated with IGE, red: features associated with generalized DEEs.




Fig. 2. Simplified diagram of the major thalamocortical circuit and schematic representation of burst-firing and tonic-firing modes regulating thalamocortical output. **A.** Feedforward axons from TC neurons target middle cortical layers (e.g., a VB TC neuron targeting cortical layer IV), while corticothalamic (CT) pyramidal neurons in layers V and VI project to the thalamus. Both projection neurons also synapse on inhibitory neurons in the nucleus reticularis thalami (nRT). Intra-nRT lateral inhibitory projections are omitted for clarity. **B.** Schematic of tonic firing mode. During alert wakefulness, under conditions of high acetylcholine and high norepinephrine tone, TASK channels close (inhibited by Gq-coupled signaling cascades), and cAMP-bound HCN channels open at higher voltages (downstream of Gs-coupled signaling cascades), resulting in a relatively depolarized TC membrane potential. Consequently, T-type Ca^{2+} channels remain inactivated. Under these conditions, TC neurons fire Na^{+} -mediated action potentials in response to excitatory synaptic inputs. **C.** Schematic of burst firing observed when neurons are exposed to conditions of low acetylcholine and low norepinephrine tone: TASK channels are open, contributing to outward potassium “leak current,” and cAMP-unbound HCN channels require lower voltages to open; TC membrane potential is dynamically regulated around a relatively hyperpolarized set-point, and this relieves the inactivation of T-type Ca^{2+} channels. In these conditions, excitatory post-synaptic potentials (EPSPs) depolarize the membrane to the activation threshold of T-type Ca^{2+} channels, which elicits a low-threshold Ca^{2+} spike (LTS) and a burst of Na^{+} action potentials in nRT and variable responses in TC neurons, which can fire either bursts of action potentials or single action potentials during a SWD.

An intrathalamic network oscillation is generated by interactions between bursting nRT and TC neurons: nRT neuron firing leads to GABA-mediated inhibitory post-synaptic potentials (IPSPs) in TC neurons, leading to activation of HCN channels, resulting in membrane depolarization to the activation threshold of T-type Ca^{2+} channels. A subsequent LTS and train of Na^+ spikes lead to glutamate release from the TC neurons onto the nRT neurons and the next cycle of burst firing. In nRT, Ca^{2+} influx activates SK channels, resulting in afterhyperpolarization of nRT neurons which makes T-type Ca^{2+} available for another burst and perpetuates the cycle.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 3. Simplified diagram of the major glutamatergic, GABAergic, and neuromodulatory inputs to the thalamus. Projections from the cortex, basal ganglia, cerebellum, and brainstem converge on TC neurons. BF: basal forebrain, GPe: globus pallidus externa, GPI: globus pallidus interna, LC: locus coeruleus, LDT: laterodorsal tegmentum, nRT: nucleus reticularis thalami, PBG: parabigeminal nucleus, PPT: pedunculopontine tegmentum, STN: subthalamic nucleus, SNc: substantia nigra pars compacta, SNr: substantia nigra pars reticulata.

Table 1

Clinical and electrographic features differ between IGEs, though first-line medications and putative targets of current treatments overlap (Jallon and Latour, 2005; Mastroianni et al., 2021; Hirsch et al., 2022).

| IGE | Epidemiology | Seizure type and age of onset | Susceptible brain state(s) | Medications | Putative medication targets |
|------|---|--|---|---|---|
| CAE | Prevalence: 5–50 per 100,000 Accounts for 18% of children with epilepsy | Typical absence seizures, generalized tonic-clonic seizures may occur in adolescence | Hyperventilation | ETX | T-type Ca^{2+} channels |
| JAE | Prevalence: 10 per 100,000 Accounts for 2.4–3.1% of children and adolescents with epilepsy | Age of onset: 4–10 (range: 2–13) Typical absence seizures, generalized tonic-clonic seizures (>90%) | Drowsiness and sleep Upon waking | VPA LTG VPA | Na^+ , K^+ , and Ca^{2+} channels, GABA Na^+ channels, HCN channels Na^+ , K^+ , and Ca^{2+} channels, GABA |
| JME | Prevalence: 1–3 per 100,000 Accounts for 9.3% of all epilepsies | Age of onset: 9–13 (range: 8–20) Myoclonic seizures, generalized tonic-clonic seizures (>90%), absence seizures (33%) | Drowsiness and sleep Soon after waking | LTG ETX (adjunctive therapy for absence seizures) VPA | Na^+ channels, HCN channels T-type Ca^{2+} channels Na^+ , K^+ , and Ca^{2+} channels, GABA |
| GTCA | Prevalence: 1.8 per 100,000 Accounts for 33% of all adolescent epilepsies | Age of onset: 10–24 (range: 8–40) Generalized tonic-clonic seizures Age of onset: 10–25 (range: 5–40) | Increased risk with sleep deprivation and photic stimulation Sleep Soon after waking Increased risk with sleep deprivation, fatigue, and alcohol use, photic stimulation | LEV LTG CZP TPM, ZNS VPA LEV | Presynaptic glutamate release HCN channels, Na^+ channels GABA _A receptors Carbonic anhydrase Na^+ , K^+ , and Ca^{2+} channels, GABA Presynaptic glutamate release |
| | | | Sleep | LTG TPM, ZNS | HCN channels, Na^+ channels Carbonic anhydrase |

CZP: clonazepam, ETX: ethosuximide, LEV: levetiracetam, LTG: lamotrigine, TPM: topiramate, VPA: valproic acid, ZNS: zonisamide.