

Lawrence Berkeley National Laboratory

LBL Publications

Title

A Critical Look at Colloid Generation, Stability, and Transport in Redox-Dynamic Environments: Challenges and Perspectives.

Permalink

<https://escholarship.org/uc/item/4407m0g3>

Journal

ACS Earth and Space Chemistry, 8(4)

ISSN

2472-3452

Authors

Spielman-Sun, Eleanor
Boye, Kristin
Engel, Maya
et al.

Publication Date

2024-04-18

DOI

10.1021/acsearthspacechem.3c00255

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

A Critical Look at Colloid Generation, Stability, and Transport in Redox-Dynamic Environments: Challenges and Perspectives

Eleanor Spielman-Sun, Kristin Boye, Dipankar Dwivedi, Maya Engel, Aaron Thompson, Naresh Kumar,* and Vincent Noël



Cite This: *ACS Earth Space Chem.* 2024, 8, 630–653



Read Online


ACCESS |

Metrics & More

Article Recommendations

ABSTRACT: Colloid generation, stability, and transport are important processes that can significantly influence the fate and transport of nutrients and contaminants in environmental systems. Here, we critically review the existing literature on colloids in redox-dynamic environments and summarize the current state of knowledge regarding the mechanisms of colloid generation and the chemical controls over colloidal behavior in such environments. We also identify critical gaps, such as the lack of universally accepted cross-discipline definition and modeling infrastructure that hamper an in-depth understanding of colloid generation, behavior, and transport potential. We propose to go beyond a size-based operational definition of colloids and consider the functional differences between colloids and dissolved species. We argue that to predict colloidal transport in redox-dynamic environments, more empirical data are needed to parametrize and validate models. We propose that colloids are critical components of element budgets in redox-dynamic systems and must urgently be considered in field as well as lab experiments and reactive transport models. We intend to bring further clarity and openness in reporting colloidal measurements and fate to improve consistency. Additionally, we suggest a methodological toolbox for examining impacts of redox dynamics on colloids in field and lab experiments.

KEYWORDS: *redox, colloids, colloidal stability, experiment vs model, nutrient and contaminant mobility, colloid transport modeling*



1. INTRODUCTION

The existence of colloids and their role in nutrient and contaminant transport is well recognized in both oxic and anoxic environments. Despite the ubiquity of colloids in environmental systems, defining a “colloid” still remains a challenge. Colloids are broadly defined as particles that are mobile in water but not fully dissolved as solutes. Commonly reported colloids in the environment include clay minerals, metal (oxyhydr)oxides, bacteria, viruses, and organic macromolecules.^{1–3} Colloids exhibit unique properties and behaviors due to their immense intra- and interparticle chemical heterogeneity, making differentiating “colloids” from the conventional “dissolved” vs “solid” phases quite difficult.

Owing to their high specific surface area (SSA) and reactive surface functional group density, colloids are important carriers of critical substances, including organic compounds, macro- and micronutrients, heavy metals, and organic pollutants, and therefore, have influence over ground and surface water quality.^{4–10} The efficiency of colloidal transport relies on the stability of the suspension, with prolonged stability directly facilitating transport.⁵ Colloid-facilitated transport of nutrients and contaminants can sustain aqueous-phase concentrations

beyond thermodynamic solubility¹¹ and thus, in certain conditions, can enhance the transport of sparingly soluble contaminants.^{10,12–16} Despite these observations, the exact mechanisms of colloid generation, their biogeochemical behavior, and transport, particularly in redox-dynamic environments, remain elusive.

Recurring seasonal wetting and drying of soils and sediments drives redox processes at solid–water interfaces.^{17–22} Wetting (e.g., through rain/snowmelt events) can promote colloidal release and downward transport in both saturated and unsaturated soils,^{13,23–26} as can prolonged inundation.⁹ This may be due to physical dislodging (e.g., shearing, sloughing) by high flow rates and/or desorption induced by changes in water chemistry, which can subsequently lead to changes in colloid composition and rate (e.g., surface charges, ligand concen-


Special Issue: Environmental Redox Processes and Contaminant and Nutrient Dynamics

Received: September 2, 2023

Revised: January 20, 2024

Accepted: February 7, 2024

Published: March 22, 2024






Figure 1. Size range of solutes and environmental particles often considered colloids when between 0.5 nm and 1 μm . Commonly used aqueous sample filter sizes are shown for comparison.

trations).^{9,25,27} Notably, redox shifts during water table level fluctuations are particularly conducive to the generation/release of colloids.^{9,15,28–30} In anoxic environments, the abundance of colloids may also increase by processes such as reductive dissolution of redox-sensitive (oxyhydr)oxides, leading to the release of other minerals and/or organics as colloids.^{9,15,31} This may occur either directly by dissolution/transformation of the redox-sensitive mineral host phases or indirectly by changing the pH or ionic strength, which triggers dissolution/disassembly and stimulates particle dispersion.^{32,33} Secondary processes may also contribute to colloid formation or release; for example, reductively dissolved Fe(II) can react with sulfide to form FeS colloids,^{34–36} or organics released through reductive mineral dissolution can form chelating complexes with solid-phase, surface-associated elements/minerals to generate suspended organo-mineral colloids.^{37–39} Conversely, the addition of organic matter to the aqueous phase in anoxic environments has also been shown to result in a gradual decrease in colloid abundance by inducing flocculation (colloid aggregation and settling out of suspension).⁴⁰

When conditions switch to oxidizing, dissolved Fe(II), among other redox-sensitive metal(loid)s can oxidize quickly and form colloidal Fe(III) phases that may remain suspended in the aqueous phase because of their small size and/or the presence of organic chelating agents.^{40–44} However, oxidation tends to increase the partitioning of (previously) suspended materials toward the solid-phase and thereby lead to a more or less gradual (i.e., kinetically controlled) decrease in colloid abundance.⁴⁵

Depending on the biogeochemical conditions (e.g., nature of colloids, aqueous-phase composition, and solid-phase properties), colloids have been found to move faster,^{46,47} at the same rate,^{26,47} or slower¹⁰ than conservative tracers, highlighting the need to develop a better understanding of biogeochemical parameters that influence this transport in porous media. Additionally, the impact of redox conditions on colloid transport will also largely depend on the chemical and physical characteristics of these colloids. With all these dynamic processes, and codependency in colloidal generation and

behavior in response to redox-dynamic environmental systems, a useful modeling framework that includes redox-generated colloids and their transport into biogeochemical modeling is critical, but yet to be fully developed. This lack of a modeling tool currently leads to the underestimation of the impact of redox processes on colloid-facilitated transport in the environment, which leads to experimental designs that lack colloidal consideration, feeding back to dissuade researchers from collecting enough empirical data to develop such models in the first place.

Several studies have reviewed the characterization and behavior of colloids in natural systems.^{1,3,48–53} This review specifically focuses on critically evaluating how colloids in natural systems influence and are influenced by redox cycling. With S, Fe, Mn, and organic phases together composing the most ubiquitous redox-sensitive phases in soil, sediment, and water ecosystems, and Mn concentrations being typically lower, the redox transformation of S, Fe, and organic phases and subsequent colloid generation are the most studied colloidal phases in redox-affected ecosystems. As a consequence, several mechanisms underlined in this review that impact the generation, stability, and transport of colloids in redox-affected environments are mainly illustrated using examples focusing on S, Fe, and organic phases. We primarily focus on the definition (Section 2), the mechanisms of colloid generation in redox-dynamic environments (Section 3), the chemical controls over colloidal behavior in redox-dynamic environments (Section 4), the challenges of sampling in redox-dynamic field sites and associated experimental needs in predicting colloidal transport in redox-dynamic environments (Section 5), and the state of colloid transport modeling and associated challenges in incorporating redox chemistry (Section 6). Finally, we highlight the need to consider colloids in regular experimental and measurement plans and in biogeochemical modeling for a better understanding of the role and extent colloids have in elemental cycling and contaminant/nutrient transport. We also make recommendations and suggestions for promising approaches in future research of colloid biogeochemistry.

2. WHAT IS A COLLOID? THE DEFINITION CONUNDRUM

Much debate surrounds the definition of a colloid in environmental systems. Historically, particles have been termed “colloids” based on three general metrics: (i) they are within a certain particle size range, (ii) they remain stable in the aqueous phase for extended periods, and (iii) therefore, they exhibit some potential for mobility/transport. Here, we summarize the existing literature to provide background on the concept of colloids and provide criteria for when and how to use the term “colloid”. For this manuscript, we use the term “aqueous phase” to define a solution that contains dissolved species and suspended particles.

2.1. Size Cutoff and Beyond. The most common characteristic used to distinguish a colloid from dissolved or suspended particulate materials is its size—a solid particle with at least one dimension between 1 nm and 1 μm is often considered a colloid (Figure 1).^{1,30,54} Although convenient for methodological standardization purposes,⁵⁵ this operational definition is problematic from a biogeochemical point of view, as the functional differences between colloids and dissolved species are not considered. Furthermore, biases may arise due to the use of a specific filter pore size or material,⁵⁶ which can introduce filtration artifacts due to the inclusion/exclusion of specific colloidal particles (as discussed in later in Section 6.1).^{57,58} Within different research disciplines, additional terminology has emerged for colloid subclasses based on functional behavior, but still defined by size, for example, aqueous clusters (0.5–2 nm),^{31,59,60} nanoparticles (1–100 nm),^{61–63} and microbes/biocolloids (100 nm to 1 μm).^{64,65}

Aqueous clusters are polynuclear complexes in the aqueous phase that precede the formation of solid nanoparticles.⁶⁶ During particle precipitation at the nucleation stage, an aqueous cluster grows in size, accumulating additional molecular groups in a specific structure. Once the aqueous cluster reaches a critical radius and mass, the structure condensates to form a solid nanoparticle.⁶⁶ Thus, the structure of the aqueous cluster serves as the basis of the structure of the nanoparticle.⁶⁷ For example, FeS aqueous cluster stoichiometry ranges from Fe_2S_2 to $\text{Fe}_{150}\text{S}_{150}$, before the first condensed nanoparticle appears.⁶⁷ While aqueous clusters and nanoparticles can overlap in dimensions, particularly during the initial condensation phase, they have distinct physicochemical properties due to their different surface area to volume ratio. Importantly, despite their obvious metastable nature, some aqueous clusters have high stability constants^{68,69} and are resistant to redox transformation and dissociation,^{69,70} which explains their persistence and transport in surface waters.⁷¹ Due to the ability for these clusters to stay suspended in the aqueous phase for extended periods of time (>30 days; as discussed in Section 4.3.2) and similarity in size, chemical composition, and structure between aqueous cluster and nanoparticles, authors often simply refer to these aqueous clusters as colloids.^{72,73} However, because clusters with minimal stoichiometry can be smaller than 1 nm (up to 0.5 nm),⁶⁷ using a strict size definition for colloids is difficult. For the purposes of this manuscript, we consider “aqueous clusters” to be distinct from “nanoparticles”, but both can still be subcategorized within the broader “colloids” term.

Within the colloidal subclass of inorganic nanoparticles, studies have shown that the particle size has a significant impact on their chemical properties and structure, such as the

total free surface energy⁷⁴ and structural strain.^{75–78} The percentage of atoms localized at the surface increases compared to the bulk as the particle diameter decreases, leading to excess surface energy.⁷⁹ Strain-induced lattice contraction is caused by structural stress (the stress applied at the surface of the particle as a result of size reduction) and can significantly affect the particle reactivity.^{80–82} On the other hand, many organic compounds are poorly soluble or even insoluble in water, leading to the formation of organic nanoparticles^{83–85} such as (bio)polymers like lipoproteins, polysaccharides, and polyuronides.^{3,86,87} Overall, size and surface properties (including surface functional groups) of organic colloids drive their unique reactivity and suspension stability in the aqueous phase. Organic nanoparticles are negatively charged colloids, mainly due to the dissociation of hydroxyl and phenolic ($\sim\text{OH}$) groups, as well as carboxylic ($-\text{COOH}$) groups. These colloids are mostly pH-dependent because the Cation Exchange Capacity (CEC) process depends on the replacement of hydrogen; thus, CEC normally increases at higher pH values, becoming highly negative in neutral to alkaline soils.⁸⁸ Because of highly negative charges available on the surface, the organic nanoparticles sorb/complex different positively charged organic and inorganic constituents, thus being able to store essential nutrients and a high rate of contaminants.^{3,87} Additionally, organic nanoparticle colloids can associate with positively charged inorganic nanoparticles, promoting formation of colloids, thus combining the characteristics of organic and inorganic constituents at the nanoscale.⁸⁹

Finally, biocolloids are microbial cells either associated with colloidal assemblages⁹⁰ or present as individual entities, i.e., live organisms (bacteria, archaea, viruses, protozoa), that exhibit transportability in the aqueous phase.⁵³ Biocolloids may form by sloughing or detachment of organisms from biofilms or consist of planktonic organisms that exhibit a “nomadic lifestyle”.^{24,91} Notably, as they are living cells, biocolloids do not dissolve/reform (even if they can partially transform (e.g., rupture of cell wall, osmosis) in response to aqueous-phase chemistry such as pH and ionic strength) or even move like abiotic colloids in response to changing conditions. Although biocolloids typically respond to the same parameters as abiotic colloids (e.g., flow rate, ionic strength, pH), they also exhibit specific “biological strategies”, including sporulation (in suboptimal environmental conditions), multiplication (in beneficial environmental conditions), and/or have extracellular structures that allow them to actively attach to specific colloidal/solid-phase matrix surfaces (e.g., by lipopolysaccharides, curli, pili) or move to/from beneficial/detrimental conditions (e.g., by flagella, cilia).^{53,65,91}

2.2. Colloidal Stability in the Aqueous Phase. Colloids are thermodynamically metastable and, with time, transform to more stable phases. Thus, clusters will have a tendency to condense into solid nanoparticles, and nanoparticles to aggregates. The ability of a colloid to stay suspended in the aqueous phase, termed colloid stability, is thus related to resistance against condensation, aggregation, and chemical reactivity. Atomic and electronic structures at surface and near-surface of colloids vary with their intrinsic physicochemical properties (e.g., size, shape, morphology, surface properties), influencing resistance against dissolution, chemical transformation, and electrostatic and electrosteric repulsive forces that alter aggregation kinetics.⁹² As an example, a modified version of the Kelvin equation predicts solubility dependence

on size, stating that as particles get smaller through the nano range of sizes, their solubilities increase exponentially.⁹³ However, certain minerals are known to become less soluble as they get smaller in size.⁹⁴ Thus, intrinsic physicochemical properties, such as the size, directly influence the colloidal stability of nanoparticles. Among inorganic nanoparticles, some minerals only exist in the nano size range, such as iron (e.g., ferrihydrite) and manganese (oxyhydr)oxides, that have been defined as nanominerals.⁹² Unlike nanominerals, mineral nanoparticles are minerals that can also exist in larger sizes.⁹² With the mineral nature driving the nanoscale size range of a nanoparticle, one can anticipate a change in the colloidal stability depending on the mineral nature of the nanoparticle.⁹² Finally, coprecipitation of a cluster and nanoparticle, composing colloids, in the presence of trace metal, organic compounds, and nutrients, can influence the intrinsic physicochemical properties of colloids, such as solubility product (as discussed in later in Sections 4.2.1 and 4.3.1).

In addition to intrinsic physicochemical properties, the ability of a particle to stay suspended in the aqueous phase (i.e., colloid stability) is strongly dependent on aqueous-phase chemistry, including pH, ionic strength, ionic composition, and solid/solution ratio (Figure 2). These aqueous-phase compo-




Figure 2. Life cycle of natural colloids in the environment. (I) Colloids can be generated either by release from the solid-phase matrix or by *de novo* precipitation in aqueous phase. (II) Once formed, colloids can undergo dissolution, aggregation, or further stabilization. (III) Colloids, depending on their physicochemical properties, can either remain mobile in aqueous phase or become immobile and attached to or part of the solid-phase matrix.

sitions affect the magnitude and nature of the colloid's effective surface charges and thus the interparticle electrostatic repulsion/attraction between suspended colloids/particles, solutes, and the nondispersible solid-phase matrix.^{3,27,32,95–97} Electrostatic and/or steric forces at the colloid surface (and/or the solid-phase matrix) can repel particles from one another and keep colloids in suspension.³ In contrast, van der Waals and magnetic dipolar interactions can enhance interparticle attraction (or to the solid-phase matrix),³ promoting particle aggregation, flocculation, sorption, and/or physical filtration that can remove colloids from suspension.

Consequently, both the physical and chemical characteristics of colloids are highly sensitive to the conditions of the surrounding aqueous-phase and bulk solid-phase matrix (Figure 2). For example, mineral colloidal suspensions are

generally favored at pH values that are far away from the point of zero charge (PZC) of the particles,^{98–100} whereas rapid coagulation is promoted in the vicinity of the PZC.^{3,101} Additionally, aggregation becomes progressively less reversible with time spent at a aqueous-phase pH somewhat near the colloid's pH_{PZC}.¹⁰² Colloid dispersion is also favored at low ionic strength because increasing ionic strength can cause colloids to reach the minimum concentration of counterions to induce coagulation of colloids, referred to as the critical coagulation concentration (CCC).^{103,104} This process also applies to biocolloids, as observed by Jewett et al.,¹⁰⁵ who found that decreasing the ionic strength caused the bacterial attachment to decrease in column experiments. Changes in ionic strength, pH, and organic matter concentration/composition can trigger colloid aggregation or dissolution, chemical adsorption or desorption, physical attachment (filtering) or detachment (sloughing), and/or chemical (including redox) transformations.^{25,40,97,98,106–109} These geochemical factors that influence colloidal stability during redox cycling are further discussed in Section 4.

2.3. Colloid Mobility and Transport. Transport of colloids in a specific soil/sediment/aquatic environment at a given time point is dependent on a complex combination of factors, including the colloid chemical composition,^{3,110,111} the solid-phase matrix composition (e.g., texture, mineralogy),^{9,25,35,96,108,109,112} the aqueous-phase chemistry,^{10,27,104,113} the hydrological flow regime,^{25,108,112} the organic matter concentration and composition,^{40,96,114–119} and the macro-/microecological communities.¹¹⁰ Depending on the aforementioned factors, the time for which particles stay suspended in the aqueous phase as colloids can vary from seconds or hours to months or years.^{31,99} Thus, colloidal abundance is properly conceptualized as dynamic over both space and time—especially in redox-dynamic environments. These are further discussed in Section 4.

2.4. Guidelines for Using the Term “Colloids”. We propose the following guidelines that will facilitate future comparisons between colloid-related studies, while recognizing the methodological constraints involved. (1) The size range (or other criteria) used to define a colloid should be flexible, but clearly stated; a wider size range (0.5 nm to 1000 nm) is preferable, or one that is tailored to the substance(s)/element(s) of interest, i.e., the “function” should take precedence over operational partitioning to the extent possible. (2) The time frame at which colloid stability/mobility is monitored/considered should be duly noted, in order to assess the environmental persistence and potential influence of the colloids.

3. MECHANISMS OF COLLOID GENERATION IN REDOX-DYNAMIC ENVIRONMENTS

Redox processes, such as oxidative dissolution of sulfide minerals and Fe redox transformation,^{28,101,120–123} induce changes in aqueous-phase parameters (e.g., pH, ionic strength, and ionic composition) and chemical, organic, and mineral transformation of the solid-phase matrix that can generate (in)organic colloids, mobilizing nutrients and contaminants. NOM complexation/coating can also play a vital role in the liberation of metal(loid)s from the solid-phase matrix and their stabilization as colloids in suspension.^{3,96,97,124} While the generation of Fe-(oxyhydr)oxide colloids under oxic conditions has been extensively studied,^(e.g.,37,41,45,125) significantly less focus has been placed on colloid generation in redox

dynamic environments. Additionally, though recent studies have reported colloid formation in anoxic environments,^{31,63} including soil and sediments, the exact mechanism, kinetics, and fate are still elusive.

3.1. Formation of Colloids in Anoxic Environments.

Under anoxic conditions, mineral precipitation, change in chemical speciation, and organic complexation are widely understood to transform and remove nutrients (e.g., N, S, and P)^{15,126,127} and/or contaminants (e.g., divalent heavy metals,^{128–130} Cr,¹³¹ As,^{132–134} Sb,^{135,136} Se,¹³⁷ U,^{138,139} V, Tc,¹⁴⁰ and halogenated organic compounds^{141,142}) from pore- and groundwater. Paradoxically, colloids may readily transport nutrients and contaminants traditionally thought to be immobilized under anoxic conditions. In this section, we examine the mechanisms behind this seemingly paradoxical behavior and highlight its importance in redox-dynamic environments.

3.1.1. Colloid In Situ Release Due to Reductive Dissolution of the Solid-Phase Matrix. In general, reductive dissolution can promote colloid dispersion by dissolving the solid-phase matrix that holds the aggregated particles together.¹⁴³ Fe minerals, especially nanoparticulate and short-range-ordered (SRO) Fe phases,¹⁴⁴ are commonly cited as important binding agents for larger soil aggregates.¹⁴⁵ In natural systems, (bio)reduction of these SRO Fe phases may lead to *in situ* release of colloids.¹⁴⁶ Henderson et al.¹⁵ and Buettner et al.³⁹ showed that reductive dissolution of Fe(III)-cements could release P-bearing and C-bearing colloids, and Tadanier et al.¹⁴³ demonstrated that bioreduction of Fe-(oxyhydr)oxide aggregates can trigger the formation of nanometer-sized As-bearing Fe(III)-colloids. In all of these cases, the transportability of P, NOM, and As was increased as a result of colloidal *in situ* release generated from reductive dissolution.

Reductive dissolution of Fe(III)-(oxyhydr)oxides often triggers an increase in pH (via OH⁻ release) that can independently drive colloid *in situ* release due to (i) the solid-phase matrix dissolution and (ii) electrostatic repulsion between like-charged particles due to change in surface charge intensity by virtue of changes in pH.^{28,147} Several studies have attempted to untangle the effect of reductive dissolution of Fe(III)-(oxyhydr)oxides from any associated pH increase, but the literature shows contradictory conclusions. For example, De-Campos et al.¹⁴⁸ showed that Fe reduction decreased soil aggregate stability to a greater extent than changes in aqueous-phase pH alone, while Grybos et al.^{147,149} in a series of studies observed a greater *in situ* release of NOM and metal(loid)s from clay minerals due to aqueous-phase pH change than from the reduction of Mn and Fe-(oxyhydr)oxides alone in wetland soil.

3.1.2. Generation of Sulfurized Metal(loid) Colloids. Under sulfate-reducing conditions, a metal–aqueous sulfide interaction first generates small aqueous complexes, such as aqueous clusters,⁶⁶ preceding the formation of nanoparticles¹⁵⁰ (see Section 2.1). The thermodynamic and kinetic stability of these aqueous clusters against condensation into nanoparticles suggests their colloidal transport can remain relevant in sulfidic environments for extended periods.⁶⁶ Recently, Noël et al.³¹ showed reductive dissolution of ferrihydrite (poorly crystallized Fe(III)-(oxyhydr)oxides) by aqueous sulfide generates FeS colloidal aqueous clusters. These colloidal clusters, defined operationally in terms of their voltammetric characteristics, have been previously detected in many natural anoxic aqueous and sedimentary environments, including anoxic lake waters

and porewaters^{151,152} and sulfidic waters of bays, marshes, hydrothermal vents, and sedimentary porewaters.^{153–155} Rickard and Morse¹⁵⁰ suggested that rapidly forming metal sulfide colloidal aqueous clusters also exist for Cu, Zn, Ag, and Pb and can make up a substantial fraction of the sulfide and metal budget in anoxic environments. Additionally, the authors claim that colloidal aqueous clusters of FeS, ZnS, and CuS could constitute a major fraction of the dissolved metal load in anoxic oceanic, sedimentary, freshwater, and deep ocean vent environments, supporting the idea that divalent metals do not necessarily always settle under sulfidic conditions but rather can be significantly transported in anoxic environmental systems.

Condensation of these metal sulfide clusters can also produce nanoparticulate colloids by homogeneous precipitation¹⁵⁶ or by being templated on bacterial membranes.¹⁵⁷ For example, Kumar et al.¹⁵⁸ reported that As was transported under sulfate reducing conditions by FeS colloidal nanoparticles. Additionally, Weber et al.⁶³ demonstrated that sulfate reduction resulted in the release of Cd and Pb from a contaminated soil as either <50 nm Cu-rich sulfide nanoparticulate colloids or as associated with bacterial membrane biocolloids, which can be transported with advective flow.¹⁵⁹ Further, finely dispersed sulfide colloids, generally negatively charged at pH > 4,¹⁶⁰ may resist aggregation and deposition,¹⁶¹ as observed for dispersed Cu_xS and CdS colloids in flooded agricultural soils.¹⁶² Such stable metal sulfide colloids may serve as a mobile carrier phase enhancing the transport of chalcophile nutrients and contaminants by colloid-facilitated transport.

The extent to which the metal-sulfide colloids are formed is controlled by the solubility of the corresponding metal sulfides and limited by the availability of dissolved sulfide.^{17,66} The rates of microbial sulfate reduction and sulfate concentration are thus key parameters controlling the generation rate and the composition of metal-sulfide colloids. For instance, Xu et al.¹⁶³ observed that flooded soil samples with higher sulfate concentrations had greater and more rapid generation of metal colloids with the fraction of colloidal Cu_xS vs Cu(0) increasing with higher sulfate content. Similarly, Noël et al.³¹ showed that the rate of sulfidation (directly driven by the S(–II)/Fe(III) ratio) of ferrihydrite aggregates led to a mixture of ferrihydrite/FeS colloids at a very low sulfidation (S(–II)/Fe(III) ratio of 0.1), while complete conversion to FeS colloids was observed at a higher sulfidation (S(–II)/Fe(III) ratio of 0.5).

3.1.3. Anoxia-Induced In Situ Release of Colloidal Natural Organic Matter. Specifically, under anoxic conditions, there are three main proposed mechanisms for the *in situ* release of colloidal NOM (Figure 2). First, minerals undergo reductive dissolution, which subsequently releases any surface-bound NOM.⁴⁰ More specifically, as soils become anoxic, Mn- or Fe-(oxyhydr)oxide minerals, which often contain adsorbed or coprecipitated NOM^{164,165} or which are present as binding agents or cements, undergo reductive dissolution and release NOM,^{29,39,166,167} often still in colloidal co-association with relatively more reduction-resistant minerals, such as clay minerals^{168–170} and silicates.²⁹ Second, anoxia-induced changes in aqueous-phase pH and ionic strength can alter mineral surface charge and ionic composition such that NOM-associated colloids are desorbed or dispersed.^{32,39,147} Third, microbial respiration products have also been shown to produce colloidal NOM under anoxic conditions.^{171,172} Once

released, this reduced colloidal NOM can reduce redox-active metal(loid)s³⁸ and/or form stable metal(loid)-NOM complexes,^{173,174} thus potentially enhancing the mobility of these contaminants in subsurface environments.

3.2. Formation of Colloids in Oxidizing Environments. When conditions change back from anoxic to oxic, reduced forms can oxidize, releasing metal(loid)s, organics, and nutrients to pore, ground, and surface waters which may also (co)precipitate as/with (oxyhydr)oxide colloids.⁴¹

3.2.1. Generation of Colloids via Iron Oxidation. In pore, ground, and surface water, both the transport of dissolved Fe(II)- and Fe(II)-colloids into more oxic environments and the introduction of O₂ into anoxic zones can trigger the oxidation of Fe(II) to Fe(III), resulting in the formation of Fe(III)-rich colloids.^{37,125} From dissolved Fe(II), these Fe(III) phases begin as small aqueous clusters of octahedral Fe(O, OH, OH₂)₆ units and grow into larger clusters over time, eventually reaching the size of nanoparticles (i.e., >1 nm). The driving force for crystal growth and/or aggregation of these nanoparticles is the decrease in surface energy.¹⁷⁵ However, for ferrihydrite, and possibly other Fe phases, the surface energy appears to be low enough that these nanoparticles are more stable in the nano size range and do not grow further^{176,177} and are defined as nanominerals.⁹² Because ferrihydrite nanominerals consistently remain in the nano size range and thus in suspension,¹⁷⁸ they are among the most ubiquitous natural inorganic colloids.¹⁷⁹ Further discussion on the conversion of metastable ferrihydrite colloids into more stable Fe(III)-(hydroxy)oxide colloids can be found later in Section 4.3.1). During the oxidation processes, nutrients and contaminant metal(loid)s may associate with the Fe(III)-(oxyhydr)oxide colloids via (re)/coprecipitation or surface complexation.^{180–183} In particular, mobile ferrihydrite colloids commonly serve as geochemical vectors, facilitating the transport of a range of contaminants, such as As,^{184,185} Pb,¹⁸³ and nutrients such as phosphate^{186–189} within aqueous environments at anoxic–oxic interfaces.

Overall, oxidation increases the nucleation of (oxyhydr)oxide colloids, and even elements inert to redox changes, such as Al, can associate with (oxyhydr)oxide nanosized colloids to form larger colloids.¹⁶⁹ The extent to which the metal-(oxyhydr)oxide colloids are formed is controlled by the solubility of the corresponding metal-(oxyhydr)oxides and limited by the availability of the metal cation. As an example, Liang et al.⁴¹ studied the dynamics of dissolved, colloidal, and deposited Fe phases using a forced-gradient field experiment with the injection of oxygenated water with a high NOM concentration into a sandy aquifer. Under the increased dissolved oxygen (DO; 2 mg L⁻¹) conditions, Fe(II) oxygenation was rapid, resulting in the formation of Fe(III)-(oxyhydr)oxide colloids. However, when DO was low (0.2 mg L⁻¹), oxidation was much slower, promoting the formation of Fe-organic dissolved complexes (up to 80% of Fe(III) was in the dissolved phase) and subsequently limiting the Fe(III)-(oxyhydr)oxide colloid formation. These dissolved complexes could significantly limit the availability of metals to form metal-(oxyhydr)oxide colloids. However, in natural conditions, Fe(III)-organic dissolved complexes can sorb at the surface of (oxyhydr)oxide colloids, forming Fe(III)-organic-(oxyhydr)oxide colloids.

3.2.2. Generation of Colloidal Natural Organic Matter in Oxidic Systems. In pore, ground, and surface waters, colloids provide large surface areas to which NOM can sorb. Overall,

because NOM has a strong affinity for freshly precipitating Fe(III) relative to Fe(II),¹⁹⁰ associations of Fe(III)-colloids with NOM are abundant.^{169,191,192} Indeed, at typical natural pH values of 4–7, a fraction of NOM is negatively charged, while oxidized Fe(III)-(oxyhydr)oxide colloid surfaces, such as ferrihydrite, carry a net positive charge, creating favorable conditions for their electrostatic interaction.^{193–195} As an example, Riedel et al.¹⁶⁴ observed the precipitation of Fe-(oxyhydr)oxides in peatland samples at the anoxic/oxic interface, which subsequently coagulated with plant-derived and pyrogenic NOM. By extension, Lyvén et al.¹⁸¹ found that Fe(III)-(oxyhydr)oxide colloids in river waters are present as small nanosized colloids closely associated with humic-type NOM. The formation of these Fe(III)-(oxyhydr)oxide-NOM colloids enhances available surface area and, hence, available sorption capacity, potentially promoting the adsorption of nutrients and contaminants.¹⁹⁶ NOM interactions may also promote the dissolution of the Fe(III) colloids into Fe(III)-NOM dissolved complexes (usually defined in the literature to be <1 nm).^{197,198} Ultimately, there is a fine line between dissolved Fe-NOM and colloidal Fe-NOM, further emphasizing the challenges in identifying a colloid's potential to react and transport. Because Fe-NOM interactions are common in aqueous systems,¹⁸³ they are an important potential transport vector for nutrients and contaminants such as heavy metals.

3.3. In Situ Release of Biocolloids. Within redox transition zones of subsurface environments, redox-sensitive elements mediate electron-transfer reactions within living cells and subsequently undergo redox transformations linked to microbial energy generation through its utilization as a source of chemical energy or as an electron acceptor for anaerobic respiration.¹⁹⁹ Thus, redox-sensitive nutrients and contaminants are intimately linked to biocolloids (i.e., viruses, bacteria, spores, etc.), promoting their transports. For example, bacteria dispersed in the anoxic porewater can transport metal(loid)s and induce biomineralization. Xia et al.²⁰⁰ observed that the dynamics of Cu in flooded agricultural soils were dominated by microbe-associated colloids and consisted mostly of Cu(0) biomineralization and subsequent sulfidation. This recent study corroborates observations from Weber et al.,¹⁷ which suggest Cu(0)-bacteria colloids forming by disproportionation of Cu(I) are released by Cu-stressed bacteria to maintain Cu homeostasis.^{201,202} Under oxic conditions, more favorable for microbial growth, biocolloids can release organic carbon that can alter mineral surface properties and colloid aggregation.²⁰³

Notably, most of the literature regarding biocolloids in the terrestrial subsurface is focused on pathogens that enter the environment from sewage, landfills, and other wastewater/fecal contamination pathways,^{53,65} or organisms that are intentionally introduced for bioremediation purposes.^{26,53} Thus, there is limited information available on natural biocolloids in redox-dynamic environments.²⁴

4. GEOCHEMICAL CONTROLS OVER COLLOIDAL COMPOSITION, STABILITY AGAINST AGGREGATION, AND BEHAVIOR ACROSS REDOX-DYNAMIC ZONES

Colloidal suspensions are inherently metastable systems because the stability of colloids is primarily dependent on their ability to resist aggregation and chemical transformation, which are controlled by physical and chemical interactions that readily change.^{104,113,114,204–206} During aggregation, colloids

dispersed in the aqueous phase assemble together, leading to settling and decreased transport.²⁰⁷

Redox-dynamics and aqueous-phase chemical composition (e.g., ionic strength and composition, pH, presence of NOM or other reactive ions) influence intrinsic physicochemical properties (e.g., size, shape, morphology, strain, aging, surface area, chemistry, superparamagnetism, and surface coatings) of redox-generated colloids, which, retrospectively, affect aggregation.^{35,97,209,104,113–118,208} In addition to the resistance to aggregation, persistence and chemical stabilization of redox-sensitive colloids and sparingly soluble contaminants can occur, even when exposed to variable redox conditions.^{66,71,173,192}

4.1. Influence of Aqueous Phase Chemistry on Colloidal Stability against Aggregation in Redox-Dynamic Environments. The amount of colloid dispersion is strongly dependent on aqueous-phase pH, ionic strength, and ionic composition because these parameters affect the magnitude and nature of colloid and solid matrix surface charges and, as a result, interparticle attractive and repulsive forces. As discussed in Section 3, the redox processes directly impacting the structure/composition of colloids influence the stability of colloids against aggregation driven by these aqueous-phase chemical parameters. Retrospectively, the aqueous-phase chemical parameters are also affected by redox mechanisms, such as oxidative dissolution of sulfide minerals and Fe mineral redox transformations.^{28,120–123} Thus, redox processes can influence the stability of colloids against aggregation by directly impacting their structure/composition or indirectly by changing the aqueous-phase chemistry (e.g., change in pH and ion composition).^{32,95,96} Additionally, redox-driven biotic or abiotic transfer of electrons may impact surface properties of colloids,^{106,210} likely influencing their surface charges and binding environment. However, as this phenomenon is mainly described for engineered nanoparticles (e.g., Ag(0),²¹¹ CuO,²¹² zerovalent Fe,²¹³ and graphene²¹⁴), the influence of redox processes on the stability of natural colloids is largely overlooked.¹⁰⁶

4.1.1. Influence of Aqueous Phase pH on Redox-Generated Colloids. As discussed in Section 2.2, mineral colloidal suspensions are favored at pH values far away from their PZC. In redox-dynamic environments, due to protonation–deprotonation reactions, pH can change *in situ* that can influence the colloidal stability.²⁸ Under sustained reducing conditions mineralogy can also change, influencing the stability and reactivity of colloids that are generated. For instance, the low pH_{PZC} values of FeS colloids suggests colloid surfaces are negatively charged at pH values typical of naturally reduced zones (pH values of 5–9), resulting in higher interparticle repulsion and, consequently, lower colloid aggregation.^{31,215,216} In contrast, the pH of most natural pore, ground, and surface waters is in the same range as the pH_{PZC} values of ferrihydrite colloids (~ 7.5 – 7.9),²¹⁷ thus resulting in rapid coagulation of ferrihydrite-based colloids in natural systems. Thus, redox dynamics can periodically affect structure/composition of colloids (i.e., intrinsic physicochemical properties) and consequently impact their stability against pH-driven aggregation.

4.1.2. Influence of Ionic Strength and Composition on Redox-Generated Colloids. As discussed in Section 2.2, colloid dispersion is favored at ionic strengths below the colloid's CCC.^{103,104} Similarly to oxic conditions, in reducing environments, recent studies have shown that higher ionic

strength inhibits the generation of FeS colloids during ferrihydrite sulfidation.³¹ Similarly, column studies on the dispersion of Hg sulfide from mine tailings under anoxic conditions imply that sulfide colloids are only mobile at low ionic strengths.^{218,219} Additionally, the biotic or abiotic transfer of electrons driven by redox processes can affect localized ionic strength and pH in the immediate and confined surroundings (not necessarily representative of the bulk) that may lead to colloidal behavior different than that initially expected.

In addition to ionic strength, the ionic composition also impacts colloid stability in suspension.^{3,10,196} The newly formed or released colloids in response to redox fluctuations (see Section 3) are directly influenced by aqueous-phase ionic composition. Thus, nutrient (e.g., NOM, Si, S, P) and/or trace metal(loid) incorporation into the mineral structure or binding to the surface of colloids may change the intrinsic physicochemical properties, stabilizing their suspension and resistance to chemical (e.g., redox) transformations.²²⁰

4.1.3. Influence of Redox Reactions on Surface Charge. In anoxic conditions, reductive dissolution of ferrihydrite and the ensuing reaction products change surface properties of colloidal ferrihydrite.^{28,34} For example, under sulfate reducing conditions, the partial reductive dissolution of colloidal ferrihydrite promotes the formation and sorption of negatively charged S(0) colloids on the surfaces of the remaining positively charged ferrihydrite, which intensifies the electrostatic attractions between oppositely charged regions on adjacent ferrihydrite surfaces, thus decreasing their CCC, and overall zeta potential of ferrihydrite colloids.¹⁰⁶ This process accelerates as sulfide concentration increases and further accelerates ferrihydrite aggregation and/or retention in solid matrices. Thus, Liao et al.³⁴ concluded that mildly sulfidic anoxic porewater (7.8–46.9 μM sulfide) can significantly decrease the transport capacity of ferrihydrite colloids. However, aggregation of ferrihydrite colloids in mildly sulfidic anoxic porewater is dependent on the physical properties of the ferrihydrite colloids. Indeed, He et al.¹⁰⁶ showed that aggregation rates could inversely decrease (increasing the colloidal stability) with increasing dissolved sulfide concentration (from 156.3 to 312.5 μM) for the ferrihydrite colloids with higher hydrodynamic diameter. Finally, to date, stability between reduced colloids, such as metal-sulfide colloids, upon mild oxidative diffusion has not yet been thoroughly investigated to our knowledge.¹⁰⁶

4.2. Influence of Natural Organic Matter on Metal(loid) Colloidal Chemical Composition and Stability against Aggregation. NOM coatings on inorganic redox-generated colloids can strongly influence intrinsic physicochemical properties of colloids (e.g., size, shape, morphology, surface properties) and particle surface charge characteristics, inducing electrostatic and electrosteric repulsive forces that alter coagulation kinetics.^{4,124,221–228} Instead of aggregating/settling, redox-generated inorganic colloids can remain mobile due to interactions with NOM. Additionally, redox conditions affect the oxidation states of the redox-active elements, impacting their affinity toward the solid-phase matrix. Complexation with NOM can stabilize metal(loid)s in their less soluble oxidation state as aqueous solutes (usually defined by the community to be <1 nm) or colloids (usually defined by the community to be >1 nm; e.g., Fe(III)-NOM colloids) inhibiting their retention in a solid-phase matrix via adsorption or precipitation.

4.2.1. Impact on Strain and Size of Redox-Generated Colloids. Strain (mineral structure change resulting in response to a stress) has been shown to be affected by surface chemistry and, in particular, by the strength of surface–ligand interactions.²²⁹ As an example, Le Bars et al.²³⁰ showed that thiol functional groups can bind strongly to the nano-ZnS surface, reducing the internal strain regardless of particle size. The authors suggested that this strain release could result from a mechanism similar to that reported by Zhang et al.,²³¹ that is, a decrease in internal energy and an increase in crystallinity induced by the interaction between water and the ZnS surface. This alteration of strain may explain the transformation of Fe-(oxyhydr)oxides into more²³² or less crystalline phases²³³ upon exposure to redox fluctuations.

Based on nucleation and growth principles,²³⁴ organic ligand binding to the surface atoms during colloid formation can inhibit growth by blocking growth sites and, consequently, favor the formation of smaller colloids or aqueous complexes compared to organic-ligand-free systems. Also, if the surface energy decreases as a consequence of organic ligand binding, the system is thermodynamically more favorable for smaller crystal size colloids (reduced critical nucleus size) than an organic-ligand-free system. For example, the nano-ZnS synthesized in the presence of organic molecules containing a thiol group, such as cysteine, exhibits both smaller average crystallite domain sizes and higher strain than the nano-ZnS synthesized under the same conditions in the absence of cysteine.²³⁰ However, no significant effect on crystallite domain size was observed for laboratory-synthesized nano-ZnS formed in the presence of serine, acetate, and histidine, which highlights that the impact on the colloid crystal growth is dependent on the binding energy of specific organic ligands on the colloid surface.²³⁵ Thus, the impact of binding organics on redox-generated colloid strain and size is directly dependent on the organic ligand functionality.^{230,231,235}

4.2.2. Interplay between NOM and Metal(loid)s in Redox-Dynamic Environments. NOM (including organic ligands) typically has strong affinity for various metal(loid)s and redox-sensitive elements that contributes toward colloidal stability (due to negative surface charge exerted by NOM complexation), for example, of Zn sulfide,^{236–240} Cd sulfide,²⁴¹ Hg sulfide,^{242,243} Cu sulfide,¹⁶¹ and Fe sulfide and (oxyhydr)oxide^{31,39} in aquatic environments.

As discussed in Section 3, NOM and Fe are strongly linked. Acidic functional groups (carboxylic, phenolic, carbonyl) on NOM generally favor the formation of surface complexes on the Fe–OH sites of Fe-(oxyhydr)oxides via chemisorption²⁴⁴ and promote the stability of Fe(III)-(oxyhydr)oxide colloids.^{117,226,245,246} In particular, Fe-NOM colloids are ubiquitous at anoxic–oxic interfaces and have been extensively studied.^{39,41,164,165,247,248} In this coassociation, the ratio of NOM to Fe can impact the size and stability of these Fe-NOM colloids. Liao et al.³⁷ observed that as the C/Fe molar ratio increased, the truly dissolved Fe(II) (as defined by the authors to be <1 nm) decreased 7-fold and the colloidal Fe(II) (1–200 nm) increased accordingly. The C/Fe ratio can also influence colloidal behavior; the authors also observed the aggregation of HA-Fe colloids under both anoxic and oxic conditions decreased with increasing C/Fe molar ratio from 1.6 to 23.3 and the deposition kinetics of Fe(II)-HA colloids under anoxic conditions were slower than those of Fe(III)-HA colloids under oxic conditions.³⁷ Liao et al. also observed that the transport of ferrihydrite colloids was substantially lower in the

presence of reduced HA than in the presence of oxidized HA, which the authors attributed to enhanced electrostatic and steric stabilization.²⁴⁹ Finally, coating of magnetite with HA and phosphatidylcholine was also reported to affect interparticle electrostatic interactions and the colloidal behavior of nanosized magnetite colloids.²⁵⁰

The formation of stable colloidal Cr(III)-NOM (10 kDa to 220 nm) upon Cr(VI) reduction by reduced NOM under organic-rich anoxic conditions has been reported.³⁸ These colloids are relatively stable in pore, ground, and surface waters of low electrolyte concentrations and are readily transported through porous media.²⁵¹ Colloidal Fe-NOM can also influence Cr reduction. For example, Buerge and Hug²⁵² observed the formation of aqueous Cr(III)-NOM-Fe(III) complexes (<450 nm) after Cr(VI) reduction by Fe(II) in organic-rich waters. Pan et al.²⁵³ observed the formation of stable colloids (1–200 nm) composed of Cr(III), NOM, and Fe(III) during Cr(VI) removal via electrocoagulation at pH 8.0, but below pH 6.0 they aggregated and settled out of the aqueous phase. Upon reaction of Cr(VI) with Fe(II)-HA colloids under anoxic conditions, Cr was completely reduced and formed highly stable Cr(III)-HA-Fe colloids that persisted for at least 20 days, without substantial changes in particle size.^{254,255}

Mn is considered to be predominantly present as soluble Mn(II) in anoxic environments and in particulate form (Mn(III/IV)) in oxic environments, but several studies have also observed Mn(III) colloids in anoxic environments.^{173,256–258} Due to the strong affinity of Mn(III) with NOM, up to 90% of the Mn(III) in anoxic systems can be in colloidal form, and this fraction can increase with increasing molar C/Mn ratios.¹⁷³

4.3. Persistence, Chemical Preservation, and Transport of Redox-Sensitive Colloids. Pore, ground, and surface water mixing at oxic–anoxic interfaces, or seasonal hydrologic fluctuations, or colloidal transport can expose colloids to redox conditions different from those of their formation. Once exposed to different redox environments, redox-sensitive colloids could play an important role in biogeochemical processes of the surrounding environment. For example, sulfidic colloids exposed to oxidizing environments and oxidized colloids exposed to reducing environments promote electron shuttling that influences geochemical processes and microbial activities.^{67,172,259} Kumar et al.²⁶⁰ concluded that organic colloid transport from reducing conditions could influence the spatial extent of biotic and abiotic sulfidic conditions downstream in otherwise oxic aquifers, acting as a vector for biogeochemical reactivity. Persistence and chemical stabilization of redox-sensitive colloids, however, inhibit their redox-driven reactivity with the surrounding environment. Surprisingly, studies have observed the persistence of sulfidic colloids in oxidizing environments^{66,261} and oxidized colloids in reducing environments.^{192,262,263} For these (oxyhydr)oxide and sulfide colloids to exist and transport across oxic and anoxic zones and interfaces, continuous formation and/or dissolution must take place, or they need to be chemically/kinetically stabilized against redox-induced transformation. Thus, redox cycling needs to be considered within the overall theoretical framework for the stability of colloids and their environmental impacts. Some of these factors are discussed below.

4.3.1. Colloid Chemical Stabilization Driven by Mineral Transformations. At the nanoscale, three factors compete to

stabilize a given polymorph: enthalpy of polymorphic transition, surface enthalpy, and enthalpy of hydration. In general, the metastable polymorphs of coarse particles have lower surface energies, leading to crossovers in phase stability as the particle size decreases.⁹² This provides a thermodynamic explanation of why nanoparticulate oxides often crystallize as one polymorph, whereas a different polymorph is exhibited in coarser-grained material.⁹² Because a different polymorph exhibits different particle sizes, as well as different crystal structure and specific surface area, conversion from a thermodynamically metastable polymorph colloid to a more stable one can significantly modify colloidal stability.^{250,264,265} For instance, thermodynamically metastable ferrihydrite colloids could transform into crystalline Fe colloids (i.e., goethite, magnetite, and hematite) that are more thermodynamically stable against redox changes. While this transformation is mainly dependent on pH, temperature, Eh, and other dissolved ions, ferrihydrite is highly insoluble at circumneutral pH and the conversion to more thermodynamically stable Fe-(hydroxy)oxide colloids in reducing environments can only proceed in the presence of a catalyst such as Fe(II)_{aq}.^{266–270} Because Fe(II)_{aq} is stable and accumulates in high concentration under anoxic nonsulfidic conditions, this transformation pathway can play an important role in the formation of more thermodynamically stable Fe-(hydroxy)-oxide colloids at redox transition zones. Thompson et al.²³² showed that redox oscillations resulted in the transformation of SRO Fe-(oxyhydr)oxides (e.g., ferrihydrite and nanogoethite) into more crystalline Fe-(oxyhydr)oxides (goethite and hematite) in the bulk soil. If this is also true for colloids, the release of SRO Fe colloids and their eventual transformation might result in a different crystal structure and specific surface area of resultant colloids, hence, different behavior and transport potential. Goethite and hematite colloids are expected to resist reductive dissolution longer than ferrihydrite colloids based on reactions of flocculated phases, but this requires further exploration.²⁷¹

In addition to mineral transformation (dissolution/precipitation), clusters and nanoparticles composing newly generated colloids can coprecipitate in the presence of organic, nutrient, and trace metal, impacting the thermodynamic metastability of these colloids. As an example, in the presence of dissolved PO₄ and As, Fe-colloids coprecipitate as Fe[(OH)₃PO₄]_nH₂O and Fe[(OH)₃AsO₄PO₄]_nH₂O, respectively. These nanophases, similar to mixed, metastable nanophases found in certain natural sedimentary environments, do not have bulk counterparts or known solubilities. They are distinct from ferrihydrite or lepidocrocite with sorbed PO₄ and AsO₄ and by definition will exhibit different metastability than ferrihydrite or lepidocrocite colloids, which would precipitate in the absence of PO₄ and As impurities.²⁷²

4.3.2. Persistence of Sulfidic Colloids in Oxidizing Environments. In sulfidic environments, the generation of sulfide colloids has been proposed to bind and mobilize nutrients and contaminants, particularly chalcophile metals (e.g., Cu, Zn, Cd, Hg, Pb, Sb).^{63,72,161,240,273–275} Thermodynamic and kinetic considerations suggest that metal sulfide colloidal clusters exhibit high stability in oxic aqueous environments,⁶⁶ are remarkably resistant to oxidation, and have been found to contribute to nutrient and contaminant transport in rivers.⁷¹ Similarly, submicrometer-sized ZnS colloids²⁶¹ have previously been identified in oxic river water. For example, freshly prepared Zn sulfide clusters were

found to have a half-life of more than 30 days in the laboratory (Luther et al., Rozan et al., and Vazquez et al.),^{69–71} whereas, in water samples the natural Zn and Cu sulfide clusters (of unknown age) were found to persist after collection for an additional 30 and 42 days, respectively.²⁷⁶

In field samples, total metal data obtained from selected acidification or separation experiments,^{71,277–279} pseudovol-tammetry measured metal concentrations,^{280–282} and electro-chemical or gas chromatographic methods^{278,283–285} suggest that aqueous metal sulfide clusters can complex up to 90–100% of metals in sewage treatment plant waters^{71,286} and oxic waters of rivers, lakes, and the ocean. Consequently, these aqueous clusters can potentially account for a significant fraction of total metal load in oxic pore, ground, and surface waters.⁶⁶

4.3.3. Persistence of Oxidized Colloids in Sulfidic Waters. As a fingerprint of the resistance of colloids against redox-driven dissolution, Brendel and Luther²⁸⁷ detected Fe(III) colloids in anoxic sediment porewaters. In a recent study, Engel et al.¹⁹² found that despite anoxic conditions (dissolved oxygen <0.08 mg L⁻¹) and the presence of dissolved Fe(II), a significant portion of the colloidal Fe in a redox-dynamic floodplain remained in its oxidized form as ferrihydrite. Indeed, complexation or coprecipitation of ferrihydrite colloids with OM coatings has been shown to prolong the persistence of ferrihydrite colloids in an anoxic aqueous phase.^{247,288–292} Engel et al.¹⁹² reported that factors favoring the delay/inhibition of redox transformation likely include (1) high OM/Si loading and resulting coverage of the Fe-(oxyhydr)oxide surface,^{220,293–295} (2) the presence of OM-Fe(II) complexes in the aqueous phase, which can bond and stabilize Fe-(oxyhydr)oxide,²⁹⁶ and (3) surface organic functional groups—particularly the extent and type of associated ligands.^{288,297–299} Inversely, complexation of organic C on the surface of high surface charge colloids can potentially limit bioavailability of organic C, while being transported through a watershed.²⁹

4.3.4. Colloid-Facilitated Transport of Insoluble Redox-Sensitive Nutrients/Contaminants. Nutrients and contaminants that are relatively insoluble in their reduced forms (e.g., U, Cr, V, Se, Pu, Tc)³⁰⁰ are regarded as immobile under anoxic conditions. However, insoluble products can be mobile in the aquatic environment if they are present as colloids. For example, U(IV)-silica colloids have been observed at circum-neutral pH,³⁰¹ while U(IV) is relatively insoluble. More specifically, colloids precipitating from relatively insoluble reduced nutrients/contaminants could be small enough to transport in reduced aqueous environments. Indeed, Suzuki et al.³⁰² showed that uraninite (UO₂) colloids formed from bacterial reduction of U in sediments are typically of less than 2 nm diameter. Thus, precipitation of U as insoluble uraninite does not necessarily lead to complete immobilization. Hence, to predict the likely environmental fate of UO₂ nanosized colloids, it will be necessary to determine the transport physics of isolated nanosized colloids and of those sorbed to larger colloidal particles and organics, and to define models that link crystallization with crystal-growth kinetics, aggregation kinetics, and fluid flow.³⁰² Further, NOM and Fe-NOM has been shown to complex with sparingly soluble Cr(III) under anoxic conditions (see section 4.2.2) (e.g., refs 38 and 303–305).

5. EXPERIMENTAL CHALLENGES AND NEEDS FOR MEASURING AND CHARACTERIZING COLLOIDS AND COLLOIDAL TRANSPORT IN REDOX-DYNAMIC ENVIRONMENTS

Although redox influences on the generation (see Section 3), behavior (chemical transformation and aggregation; see Section 4), and subsequent transport of natural colloids have been clearly demonstrated, the colloidal aspects of redox dynamics are difficult to characterize in natural environments and field samples. Most of the existing research on colloid transport affected by redox environments has been observational, with the key model parameters (e.g., kinetic rate of generation, dynamic surface area) still poorly constrained and unquantified. This lack of knowledge is compounded by challenges associated with the detection and characterization of colloids in natural redox environments, as well as the lack of systematic experimental setups to obtain necessary data to parametrize and validate models to account for colloids.³⁰⁶

5.1. Operational Challenges Due to Lack of Systematic Size Definition of a Colloid. As discussed in Section 2.1, the lack of a unified size definition for colloids makes comparison across studies difficult. Most common sampling approaches ignore the presence of colloids, their chemical transformation/aggregation, and their transport. Many studies consider certain colloidal nutrients and contaminants in the aqueous phase to be “dissolved”, based on the utilized filtration membrane pore size.¹⁶⁹ Even when colloids are considered, their size range changes from study to study and specific ranges of colloidal sizes may still be excluded depending on the study and methodological approach taken and the research question targeted. As noted by Bao et al.,¹⁶⁹ many studies use a 450 nm filter to separate “solid particles” from “colloids”,^{163,173,307} whereas other studies use the same filter size to separate “colloidal” (<450 nm) from “dissolved” (<450 nm) species.³⁰⁸ Further, considering that “*in situ*” sampling equipment typically used to collect porewaters have orifices smaller than 1000 nm (150 and 600 nm rhizons being the most commonly used for *in situ* sampling in soils and sediments)¹⁹² and larger colloids are often disregarded. Finally, while Bao et al.¹⁶⁹ and numerous other studies define truly dissolved fractions as all material below 3 kDa, this excludes sulfide aqueous clusters observed by Luther and Rickard⁶⁶ (and discussed in detail earlier in Section 2.1). These operationally defined size limits for “colloids”^{163,173,307} are arbitrary from a functional perspective and can significantly impact our understanding of the transport and bioavailability of nutrients and contaminants that may be associated with colloids.

5.2. Locating Redox Changes and Their Temporal Variability in Natural Conditions. The majority of studies on colloids have been performed using either laboratory-synthesized colloids^{173,249,254} or lab-simulated experiments using colloids extracted from soils/sediments.^{29,200} These lab experiments allow researchers to constrain the system and focus on a limited number of parameters. However, they are not necessarily representative of pore, ground, and surface water colloids, which are expected to exhibit pronounced compositional complexity.¹⁹² Thus, understanding and simulating colloid transport affected by redox-dynamic environments requires, retrospectively, data comparison of lab-simulated experiments with natural conditions in order to validate the lab-simulation results and constrain future lab and numerical simulation experiments.

However, like most soil properties and processes, redox is spatiotemporally heterogeneous.³⁰⁹ Important questions still remain regarding where exactly redox-generated colloids are promoted in soils and sediments, and how best to track the potential stability of these colloids to chemical transformation and aggregation during colloidal transport. Because redox spatiotemporal heterogeneity has not yet been incorporated into mainstream conceptualizations of soil biogeochemistry,³⁰⁹ these questions remain unresolved for most environmental systems, affecting our ability to *in situ* track redox-generated colloids, monitor their behavior and transport, and fully understand the impacts on nutrients and contaminants associated with colloids in redox-dynamic environments. To address this knowledge gap, Lacroix et al. recently published a review where authors describe past and current approaches for detecting, quantifying, and characterizing redox heterogeneities in soils as a function of time.³¹⁰ The recent development of 2D imaging techniques, such as planar optode measurements, spatially resolved microbial techniques, and X-ray fluorescence imaging combined with spectroscopy, are promising and provide reliable forensic evidence for future research that integrates the influence of redox heterogeneities in generation, stability, and transport of colloids in soils and sediments.

The temporal dynamics of redox conditions are mainly driven by hydrologic regimes (e.g., snowmelt, precipitation, drought) that in turn induce changes in colloidal physicochemical composition. Thus, for the same location, the generation mechanisms of colloids, their physicochemical transformations and stability against aggregation, should also be tracked as a function of time/season and the changing of surrounding physicochemical parameters. However, the kinetics of redox changes are still poorly understood in natural environments. Thus, the time window needed between sampling events to observe changes in the nature and physicochemical composition of colloids remains unspecified and will likely differ for different sites.

5.3. Preservation of Redox Integrity. In addition to the lack of systematic and consistent colloid consideration/definition, the physicochemical properties of colloids in a natural aqueous phase can easily change during sampling, storage, and analyses, due to aging, changes in pH, ionic strength, redox conditions, and light exposure.^{311,312} Furthermore, associated nutrients and contaminants are also affected by sorption processes, complexation, and redox precipitations that can occur during transport and storage.

Characterizing suboxic/anoxic environmental colloids requires careful protocols for both sampling and analysis to preserve the native redox status of targeted elements. Unintentional introduction of O₂ can lead to the oxidation of pore, ground, or surface water samples; for example, oxidation of dissolved Fe(II) can form nanosized Fe-(oxyhydr)oxide colloids, which may not be naturally occurring in the sampled aquifer.¹²⁵ Such oxidation artifacts not only affect the physicochemical properties of colloids but could also impact the speciation of associated elements, such as phosphorus that may readily adsorb to nanosized Fe-(oxyhydr)oxide colloids³¹³ (as also discussed in Section 3.2.1). Consequently, sampling potentially anoxic solutions requires systematic collection and analysis methods in order to maintain redox and chemical integrity throughout the sampling-analysis continuum. For example, a study by Dai et al.³¹⁴ performed groundwater sampling and cross-flow ultrafiltration analysis with deliberate redox control measures and

found no significant association of Pu with colloids in groundwater at the U.S. Department of Energy Savannah River Site, which is in contrast to a similar study at the same field site with no redox control measures, where investigators found a significant amount of Pu association with colloids.³¹⁵ However, sampling anoxic waters requires proper training on elaborate techniques that are costly, difficult to obtain, and rarely tested for accuracy.

Preserving the redox state in collected samples is also a complicated task, particularly in the field. For example, even if oxygen is eliminated, the reduction reaction, if still active, could produce new colloids or change physicochemical properties of existing colloids after sampling, even if collected carefully. Most preservation approaches consist of freezing samples, whether by slowly cooling down to freezing temperatures³¹⁶ or flash-freezing (L-N₂) the samples immediately.^{317,318} However, the freezing process itself can facilitate irreversible precipitation of salts,³¹⁹ the formation of coatings on colloids,³²⁰ and/or colloid flocculation,³¹¹ altering the colloid concentration and requiring further consideration during analysis. Although, temperatures just above the freezing point (e.g., commonly used 2–5 °C) are believed to minimize reaction kinetics, these have not yet been systematically tested for their efficiency in halting the biogeochemical reactivities completely, particularly in anoxic environments. The lack of systematic and consistent redox preservation approaches could impact the accuracy of the biogeochemical parameter libraries used for simulating colloid transport.

Nonetheless, some general practices have been pointed out as being helpful to consistently apply while preserving natural colloid samples, for instance, minimizing exposure to light, maintaining low temperatures, and minimizing physical disturbance once collected, choosing materials (tubing, vials, etc.) that are known for restricting O₂ penetration and have lower affinity sorbent walls. If dilutions are absolutely necessary, maintaining pH, ionic strength, and ionic composition is critical.

6. CURRENT STATE OF COLLOIDAL TRANSPORT MODELING AND ASSOCIATED GAPS ACCOUNTING FOR REDOX

Based on the preceding sections, it becomes clear that effective colloid modeling requires the integration of several key processes: the generation of colloids in redox-dynamic environments, the stability of colloids as influenced by various chemical factors, and the transport mechanisms of colloids within different environmental systems. Additionally, colloid transport processes are inherently dynamic, varying both temporally and spatially. Specifically, generation/detachment (i.e., source) and/or retention/size exclusion (i.e., sink) processes are heavily influenced by spatial and temporal variability of unsaturated/saturated and redox conditions (see Sections 3 and 4),^{321–327} which are highly heterogeneous in soils and sediments.^{51,309} Contrasting conditions found in saturated and unsaturated porous media necessitate distinct approaches, which present additional challenges for modeling colloid transport.⁵ Similarly, redox variability can significantly change chemical conditions along a flow-path and/or over time (e.g., during wetting/drying or in relation to microbial metabolic activity).

In this context, addressing these challenges requires a thorough understanding of how redox fluctuations occur in the subsurface and the factors contributing to these variations.

This comprehensive understanding is pivotal in guiding the design of experiments for data collection, an essential step for developing robust reactive transport models (RTMs) that accurately and effectively represent environmental systems. The following sections present a concise overview of colloid transport continuum-based numerical models, aiming to highlight the capabilities and limitations in RTMs in effectively incorporating redox dynamics.

6.1. Modeling Colloid Transport in Porous Media under Dynamic Redox Conditions. Colloid Filtration Theory (CFT) in porous media, the process of removing suspended particles from a fluid by passing it through a porous medium such as a filter or a bed of particles,^{206,328} has played a significant role in the development of these colloid transport models and approaches for understanding and predicting colloid behavior in porous media. To describe the partitioning of colloids between the aqueous-phase and solid-phase matrix in variably saturated porous media, a general advective dispersion equation (ADE) for colloid transport is used (eq 1):

$$\frac{\partial(\theta C_c)}{\partial t} + \rho_B \frac{\partial S_c}{\partial t} = \frac{\partial}{\partial z} \left(D_c \theta \frac{\partial C_c}{\partial z} \right) - \frac{\partial \theta v_c C_c}{\partial z} + R \quad (1)$$

The left-hand side terms account for the temporal variations of both immobile (S_c [M M⁻¹]) and mobile (C_c [M L⁻³]) colloids on the solid surface of a porous solid-phase matrix; θ [L³ L⁻³] is the volumetric water content, ρ_B [M L⁻³] is the bulk density, and t [T] is time. The right-hand side terms represent the flow of colloids through the pores of the porous solid-phase matrix, where v_c [L T⁻¹] and D_c [L² T⁻¹] are the advective (i.e., Darcy or pore-water) velocity and the dispersion coefficient for colloidal particles, respectively; R describes any additional source or sink terms of colloids which are not integrated in the rest of the equation.

The presence of uncertain terms, such as R , within eq 1 presents a valuable opportunity for advancing the integration of redox dynamics into colloid transport by explicitly incorporating them into the equation. For instance, source and sink terms account for colloid generation and retention. However, distinguishing whether these processes are primarily driven by attachment or size exclusion mechanisms within the existing modeling framework is often challenging. The following section will explore these concepts in more detail and identify potential avenues for integrating redox dynamics with colloid transport.

6.1.1. Modeling Colloid Attachment in Porous Media. Commonly, quasi-first-order rate coefficients, or composites of rate coefficients, are used to model the partitioning of colloids between the surfaces of porous solid-phase matrix (stationary) and aqueous phases, via the attachment mechanism described by CFT.^{329,330} The rate of change in the concentration of immobilized colloids due to attachment can be modeled using first-order kinetics as follows (eq 2):

$$\rho_B \frac{\partial S_c}{\partial t} = k_{\text{att}} \theta C_c \quad (2)$$

The attachment rate constant (k_{att} [T⁻¹]), representing the rate at which colloidal particles deposit onto a porous media or soil/sediment solid-phase matrix, is directly influenced by two factors (eq 3): (i) the frequency of contact between colloids and the grain surface and (ii) the affinity of the colloids to adhere to the porous medium.^{206,328} The attachment rate constant is typically calculated as a function of the median

radius of the grain of porous media (a_c [L]), Darcy velocity (v_c [$L T^{-1}$]), porosity of the porous media (ϵ [-]), and collision (η [-]) and sticking (α [-]) efficiencies. These factors account for the diffusion, attachment, and gravitational sedimentation of colloidal particles.

$$k_{\text{att}} = \frac{3(1-\epsilon)\alpha\eta v_c}{4a_c} \quad (3)$$

The attachment of colloidal particles is contingent on the presence of available attachment sites, which are influenced by factors like characteristics (e.g., surface area, charge, roughness) of the solid surface of the solid-phase matrix and colloid properties (e.g., functional groups). The Derjaguin–Landau–Verwey–Overbeek (DLVO) theory^{331,332} proposes that an interplay occurs between attractive van der Waals and electrostatic forces (either attractive or repulsive depending upon the charges between the colloids and surfaces), and this interaction is dependent on the distance between colloids and the surface. The DLVO theory not only considers the impact of van der Waals forces and electrostatic repulsion/attraction but also takes into account additional factors, such as Brownian motion (the random motion of particles suspended in a fluid) and the ionic strength of the surrounding aqueous phase. Redox reactions significantly impact surface charges (see Section 4.1.3) and thus the prevalence of both favorable and unfavorable conditions conducive to colloidal attachment (i.e., the attachment efficiency, α). Furthermore, redox processes can also shift the parameters (pH, ionic strength, and ionic composition; see Section 4.1) and complexing agents (see Section 4.2) of the aqueous phase that govern colloid attachment to the solid surfaces of a porous solid-phase matrix. For instance, the generation or release of colloids with a high adsorption capacity (such as Fe-(oxyhydr)oxide, clays) can trap dissolved and particulate NOM in suspension (see Section 4.2) and reduce the attachment efficiency of NOM to a porous solid-phase matrix. Fully encapsulating all of the effect of these chemical processes into a single attachment efficiency term, α , is a challenge for current RTMs; thus, we recommend utilizing α terms that properly reflect changing particle and solution conditions.

6.1.2. Modeling Colloid Size Exclusion in Porous Media.

While CFT can reasonably estimate the collision [η] under conditions favorable for attachment, CFT does not work well under unfavorable conditions.^{327,333,334} Nonetheless, under unfavorable conditions, size exclusion has been suggested to account for filtration through porous media, including straining, wedging, and entrapment of colloids.^{335–339} Although these mechanisms remove colloids from the aqueous phase, they differ fundamentally from attachment mechanisms as they are primarily governed by geometrical constraints rather than electrostatic forces. However, there is a clear connection between the biogeochemical mechanisms, including redox and shifts in ionic strength, pH, and chemical composition, that physically change the shape and size of colloids (see Section 4.1), the collision efficiency, and size-exclusion effects.

While attachment removes colloids through collision, straining involves the filtration of colloids based on their size or shape. Similarly, wedging or entrapment involves trapping colloids between two surfaces enclosed within a larger solid aggregate or solid-phase matrix in porous media. A few studies have shown colloidal removal using both attachment and

straining mechanisms.^{337,339,340} In particular, Bradford and Bettahar³³⁵ and Xu et al.³³⁷ used a dual-site approach to explain colloidal retention mechanisms. One site was dedicated to attachment, while the other focused on straining. They suggested that attachment is reversible, but straining is irreversible. They further suggested modifications to eq 1 to include two sites for colloid retention in numerical representations:

$$\begin{aligned} \frac{\partial(\theta C_c)}{\partial t} + \rho_B \frac{\partial S_{c1}}{\partial t} + \rho_B \frac{\partial S_{c2}}{\partial t} \\ = \frac{\partial}{\partial z} \left(D_c \theta \frac{\partial C_c}{\partial z} \right) - \frac{\partial \theta v_c C_c}{\partial z} + R \end{aligned} \quad (4)$$

In the given context, the variables S_{c1} and S_{c2} represent the respective masses of colloids that have become attached to the solid-phase matrix and the masses of colloids that have been strained out of the aqueous phase. Furthermore, these authors aimed to model straining as a first-order kinetics process by determining the straining rate coefficient based on the distance calculated from the entrance of the porous solid-phase matrix.

Although the dual-site model approach is reasonable for simulating field scale colloidal transport processes, additional partitioning between attachment/detachment and straining fractions remain to be experimentally determined. Consequently, the dual-site approach has primarily been utilized in modeling studies where breakthrough curves can be calibrated. Future research is needed to calibrate colloidal behavior in broader applications, particularly accounting for dynamic redox conditions and associated physicochemical properties of colloids, aqueous-phase, and solid-phase matrices.

As discussed above, although different mechanisms (e.g., attachment, straining, entrapment) remove colloids from the system, they may yield different outcomes under prevalent redox conditions. Hence, understanding the specific role of redox on attachment, straining, wedging, or entrapment becomes crucial. Furthermore, redox conditions can alter the soil matrix through mineral dissolution (see Section 3.1), significantly affecting wedging or entrapment. This presents a valuable opportunity to expand and incorporate these mechanisms within RTMs while considering the influence of redox conditions.

6.1.3. Modeling Colloidal Detachment in Porous Media.

Detachment (as opposed to attachment) results in the separation of colloids from one another and/or from the porous solid-phase matrix. The detachment of colloids from the porous solid-phase matrix is the consequence of several physical factors, including hydraulic transients (such as infiltration and wet/dry cycles),^{327,341} colloid–colloid collisions,⁴⁸ an increase in shear stress,^{342–344} and movement of the air–water interface.³⁴⁵ On the other hand, chemical perturbations such as redox changes and subsequent shifts in aqueous-phase chemistry (e.g., ionic strength, pH, chemical composition; see Section 3), surface properties, and electrostatic interactions, may further promote detachment due to the weakening of bonds between colloids and the porous solid-phase matrix.^{5,339,346–351}

$$\rho_B \frac{\partial S_{c1}}{\partial t} = k_{\text{att}} \theta C_c - k_{\text{det}} \rho_B S_{c1} \quad (5)$$

Moisture content^{327,341} is a key parameter in modeling colloidal detachment, as demonstrated in eq 5, where S_{c1} [$M M^{-1}$] and C_c [$M L^{-3}$] are the concentrations of colloids in the

porous solid-phase matrix and mobile phases, respectively.^{327,341} Colloid detachment (k_{det} [T^{-1}]) is determined by porewater velocity above a critical moisture content. Various studies^{1,170,352–358} have found that DLVO forces, which balance van der Waals attractive forces with electrostatic repulsive forces, control colloidal detachment. Thus, we highly recommend improving DLVO models by incorporating the biogeochemical perturbations affecting surface properties and electrostatic interactions, such as changes in redox conditions.

6.1.4. Modeling Colloidal Generation in Porous Media. Research indicates that erosion processes have the potential to generate colloids. These newly generated colloids may not have originally existed in the dissolved phase but were removed from the porous solid-phase matrix.^{343,359} Researchers have categorized attached colloids into two distinct groups: those that are irreversibly attached and those that are attached but capable of being released. This categorization conceptually parallels eq 5, where attachment and straining are modeled separately. Additional terms representing colloid generation through physical detachment can be added to the equation as

$$\begin{aligned} \frac{\partial(\theta C_c)}{\partial t} + \rho_B \frac{\partial S_{\text{cir}}}{\partial t} + \rho_B \frac{\partial S_{\text{cim}}}{\partial t} \\ = \frac{\partial}{\partial z} \left(D_c \theta \frac{\partial C_c}{\partial z} \right) - \frac{\partial \theta v_c C_c}{\partial z} + R \end{aligned} \quad (6)$$

where S_{cir} and S_{cim} are irreversibly attached colloids and immobile colloids available for release, respectively.

In addition to erosion, colloids can be generated due to various other factors, including fluctuations in redox conditions and resulting changes in aqueous-phase chemistry, such as pH and ionic strength (see Section 4.1). However, colloid generation through redox variation is currently not well parametrized in models, primarily because more quantitative empirical evidence is needed. Most of the existing research on *in situ* release and new precipitation has been qualitative observational studies, and quantitative models that incorporate kinetics, dynamic surface area, and thermodynamic constraints are still lacking. The balance between *in situ* release and new precipitation is challenging to determine, especially for changing redox conditions, necessitating new experimental designs.

6.1.5. Retention of Colloids via Attachment and Straining at the Air–Water Interface. In variably saturated porous media, colloids experience a wide range of spatially and temporally varying physical and chemical conditions.^{105,360} These factors significantly influence colloid transport, making it more intricate in unsaturated media compared to saturated media.³²¹ Moreover, several processes occur in variably saturated porous media that do not occur in saturated conditions, such as discontinuous capillary fringes that promote colloid deposition or wetting–drying cycles that promote colloid transport.^{322,323} Overall, unsaturated conditions lead to a higher rate of colloid removal than saturated conditions.^{324–327} In addition, variations in subsurface redox states can alter solid-phase properties,^{327,341} surface charge, and microbial activity, ultimately impacting colloid behavior during these hydrological processes.

Moreover, redox states have the capacity to modulate the air–water interface by altering the local surface potential, consequently affecting the behavior of colloids at this interface.³⁶¹ For example, under unsaturated conditions, colloids can attach to the air–water interface and undergo

straining. This attachment is governed by electrostatic, hydrophobic, and steric interactions, and the resulting colloidal monolayers or multilayers³⁶² can significantly impede colloid transport. Wan and Wilson³⁶² found that colloids preferentially deposit at the air–water interface. This interface acts as a potent sorbing phase that significantly retards colloid movement in porous media. Hence, it is crucial to incorporate this phase into the modeling framework, for example, by introducing a Γ [L^{-1}] term into eq 1, to represent the interfacial surface area (eq 7)

$$\frac{\partial(\theta C_c)}{\partial t} + \rho_B \frac{\partial S_c}{\partial t} + \Gamma \frac{\partial A_c}{\partial t} = \frac{\partial}{\partial z} \left(D_c \theta \frac{\partial C_c}{\partial z} \right) - \frac{\partial \theta v_c C_c}{\partial z} + R \quad (7)$$

where A_c [M L^{-2}] is colloid concentration at the air–water interface.

Thin water films, referred to as film straining, pose another physical limitation on colloid transport in porous media under unsaturated conditions. This phenomenon has been extensively studied by Wan and Tokunaga,³⁴⁵ who suggested that colloid transport in unsaturated porous media is a function of the ratio between the size of colloids and the thickness of the water film. In addition, the air/water/solid-phase matrix contact line, which is formed by water molecules adsorbed onto the surfaces of colloids, has been observed to exert a noteworthy influence on the retention and release of colloids.^{327,344,363,364} Based on the previously established eq 8, Corapcioglu and Choi³³⁸ developed a numerical model to include colloid transport at the air/water/solid-phase matrix contact line into the modeling framework (eq 8). Here, F_c [ML^{-3}] represents the concentration of colloids trapped by film straining.

$$\begin{aligned} \frac{\partial(\theta C_c)}{\partial t} + \rho_B \frac{\partial S_c}{\partial t} + \Gamma \frac{\partial A_c}{\partial t} + \frac{\partial F_c}{\partial t} \\ = \frac{\partial}{\partial z} \left(D_c \theta \frac{\partial C_c}{\partial z} \right) - \frac{\partial \theta v_c C_c}{\partial z} + R \end{aligned} \quad (8)$$

Importantly, redox states can influence colloid behavior at the air–water interface through various mechanisms. It is crucial to understand the processes at this interface, including its modulation, which can be shaped by local surface properties influenced by redox conditions (see Section 4). These alterations can have implications for attachment, straining, and related phenomena. For example, attachment at the air–water interface may be governed by electrostatic conditions and redox states. Likewise, as discussed earlier in Section 4.2.1, straining is a unique process that can impact colloid transport. These processes should be quantified under varying redox conditions to enhance our comprehension, offering the possibility of a transformative shift in colloid transport modeling.

6.2. Parametrization of Colloid Generation and Behavior. Colloid generation, *in situ* release, and newly precipitated colloids (see Section 3) have not been parametrized enough in numerical models and lack empirical data (see Section 5). Furthermore, the distinction between *in situ* release and new precipitation is difficult, especially under changing redox conditions. The development of lab-simulation experiments, using a stepwise process from simple batch experiments to more complex column experiments that are capable of mimicking distinct colloid-generating redox processes and allow transport considerations, is crucial for

capturing the key parameters needed to model redox-generated colloids in porous media.

Once colloids are generated, their transport is dictated by their ability to interact with other colloids and the surrounding porous media, which is dependent on surface properties and electrostatic interactions. These surface- to pore-scale processes are poorly constrained for colloids. Thus, transport should be evaluated in column experiments with both well-constrained pore structure and material in order to identify the controlling mechanisms and kinetics.^{365–367} The transport of each well-characterized, monodisperse redox-generated colloid should be tested against a conservative tracer to calibrate transport properties. However, this approach complicates identifying distinct key parameters that are needed for modeling size exclusion (see Section 6.1.2) and attachment/detachment in porous media (see Sections 6.1.1 and 6.1.3). Specifically, thanks to the development of new imaging technologies, such as microfluidics and μ -X-ray fluorescence mapping, we suggest that future studies focus on understanding the influence of biogeochemical processes on the partitioning between size exclusion as a consequence of aggregation mechanism and attachment conditions.

6.3. Simulating the Impact of Colloidal Transport in Redox-Dynamic Environments. The challenges previously described in detecting and characterizing colloids associated with redox-dynamic environments in natural conditions go hand in hand with major challenges in integrating necessary data to parametrize and validate models to account for colloids at field scale. Thus, additional knowledge is needed about the relevance of colloidal transport and exported reactivity for ground and surface water quality and for RTMs to include colloidal transport at the field scale. As an example, Babey et al.¹⁷² showed, through RTM simulations based on data from a series of dual-domain column experiments,^{259,260,368,369} the necessity to consider transport of NOM colloids from anoxic environments as the key driver to explain the observed biogeochemistry downstream in an initially oxic surrounding environment. Indeed, the NOM colloidal transport from anoxic lenses to the oxic surrounding sediment drives the development of proximal secondary reduction zones (“halos”), characterized by high microbial activity (e.g., sulfate reduction) and accumulation of reduced reaction products (e.g., iron sulfide).^{22,370} This study corroborates that neglecting colloidal transport within and from redox-dynamic environments creates major uncertainties in model simulations of pore, ground, and surface water quality. Future research should focus first on developing RTMs to test if colloidal transport could explain some paradoxical/unexpected field-scale nutrient and contaminant cycling in redox-dynamic environments.

7. CONCLUSIONS AND FUTURE RECOMMENDATIONS

This review emphasizes the presence and role of colloids in the transport of nutrients, contaminants, and organic matter through environmental systems. As consistently highlighted in this review, colloid generation and transport mechanisms are not yet clearly understood and even less quantified/parametrized, partly due to methodological challenges in sampling and analyses of colloids in natural systems, especially redox-dynamic ones. However, it is only logical to anticipate that generation, behavior, and transport of colloids under redox dynamic environments can be different than those under stable oxic environments. Therefore, colloids in redox-dynamic

environments have yet to be part of routine sampling protocols or general predictive geochemical modeling approaches.

A major challenge in colloid research is the lack of a universal definition of a “colloid”, which leads to inconsistencies in identification and characterization of colloids in field- and lab-scale studies. Here, we propose a pragmatic approach toward nomenclature of colloids that could bring more uniformity, clarity, and consistency in colloidal research. We propose broadening the size range to cover various environmental colloids and inclusion of functionality in the definition of colloids. Being aware of biogeochemical conditions of the system that are conducive to colloidal generation will allow researchers to adopt a tailored sampling plan (frequency as well as key parameters) during an experiment. Additionally, attention is needed to address the lack of consistent experimental approaches for analyzing colloids in samples from potentially anoxic environments. We propose greater clarity in communicating how various methodological approaches, from sampling to analyses, may impact the results of colloid studies in redox dynamic environments, including the implications on what is considered a colloid.

While similar solution parameters (pH, ionic strength and composition, solid/solution ratio) control colloidal generation and stability in oxidized or reduced environments, the complexity is accentuated in redox-dynamic environments, where redox reactions can *in situ* catalyze sudden changes in solution chemistry that influence colloidal behavior. Consequently, colloidal transport-behavior relationships are challenging to predict, necessitating further model development to gain a deeper mechanistic understanding. To be able to study these reactions, sampling plans need to be spatially and temporally elaborate. This has implications for the mass balance of elements (trace metals, nutrients) and colloid-facilitated transport of contaminants. In addition, in anoxic environments, where results are more easily compounded by artifacts due to the materials used in sample collection (tubing, filter material, etc.), preservation methods and the ability of analytical techniques to maintain redox integrity during analysis are pertinent. In sum, there is need for tighter coupling between field, lab, and modeling studies to increase the efficiency and accuracy of data interpretation and knowledge on the environmental role of colloids, particularly in redox-dynamic systems.

In this review, we have examined existing literature on natural colloids and their environmental importance, with a focus toward redox-dynamic systems. We have identified and discussed challenges and opportunities related to the consideration of colloids in routine analyses, focused studies, and biogeochemical models. Our intent and hope is that this work will lead to broader and more open discussions that will address these issues, including workshops and hands-on training on sampling, analytical, and experimental approaches, as well as modeling tools that will help usher the research and understanding of colloids in environmental systems.

■ AUTHOR INFORMATION

Corresponding Author

Naresh Kumar – Soil Chemistry, Wageningen University and Research, Wageningen 6708 PB, The Netherlands;
orcid.org/0000-0002-8593-5758; Phone: +31 317 481797; Email: naresh.kumar@wur.nl

Authors

Eleanor Spielman-Sun – Environmental Geochemistry Group at SLAC, Stanford Synchrotron Radiation Lightsource (SSRL), SLAC National Accelerator Laboratory, Menlo Park, California 94025, United States; orcid.org/0000-0001-9626-2969

Kristin Boye – Environmental Geochemistry Group at SLAC, Stanford Synchrotron Radiation Lightsource (SSRL), SLAC National Accelerator Laboratory, Menlo Park, California 94025, United States; orcid.org/0000-0003-2087-607X

Dipankar Dwivedi – Earth and Environmental Sciences Area, Lawrence Berkeley National Laboratory, Berkeley, California 94720, United States

Maya Engel – Department of Soil and Water Sciences, Faculty of Agriculture, Food, and Environment, The Hebrew University of Jerusalem, Rehovot 7610001, Israel

Aaron Thompson – Department of Crop and Soil Sciences, University of Georgia, Athens, Georgia 30602, United States; orcid.org/0000-0001-6301-7377

Vincent Noël – Environmental Geochemistry Group at SLAC, Stanford Synchrotron Radiation Lightsource (SSRL), SLAC National Accelerator Laboratory, Menlo Park, California 94025, United States; orcid.org/0000-0002-5387-8664

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acsearthspacechem.3c00255>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Funding was provided by the U.S. Department of Energy (DOE) office of Biological and Environmental Research (BER), Environmental System Sciences Division, through its support of the SLAC Floodplain Hydro- Biogeochemistry Science Focus Area (SFA) under Contract No. DE- AC02-76SF00515.

REFERENCES

- (1) Kretzschmar, R.; Borkovec, M.; Grolimund, D.; Elimelech, M. Mobile Subsurface Colloids and Their Role in Contaminant Transport. *Adv. Agron.* **1999**, *66* (C), 121–193.
- (2) Stumm, W. Chemical Interaction in Particle Separation. *Environ. Sci. Technol.* **1977**, *11* (12), 1066–1070.
- (3) Goldberg, S. R.; Lebron, I.; Seaman, J. C.; Suarez, D. L. Soil Colloidal Behavior. In *Handbook of Soil Sciences Properties and Processes*; Huang, P. M., Li, Y., Sumner, M. E., Eds.; CRC Press: 2012; pp B195–B240.
- (4) Kretzschmar, R.; Sticher, H. Colloid Transport in Natural Porous Media: Influence of Surface Chemistry and Flow Velocity. *Phys. Chem. Earth* **1998**, *23* (2), 133–139.
- (5) Flury, M.; Qiu, H. Modeling Colloid-Facilitated Contaminant Transport in the Vadose Zone. *Vadose Zo. J.* **2008**, *7* (2), 682–697.
- (6) Means, J. C.; Wuayaratne, R. Role of Natural Colloids in the Transport of Hydrophobic Pollutants. *Science* (80-). **1982**, *215* (4535), 968–970.
- (7) Pédrot, M.; Dia, A.; Davranche, M.; Bouhnik-Le Coz, M.; Henin, O.; Gruau, G. Insights into Colloid-Mediated Trace Element Release at the Soil/Water Interface. *J. Colloid Interface Sci.* **2008**, *325* (1), 187–197.
- (8) Dia, A.; Gruau, G.; Olivé-Lauquet, G.; Riou, C.; Molénat, J.; Curmi, P. The Distribution of Rare Earth Elements in Groundwaters: Assessing the Role of Source-Rock Composition, Redox Changes and Colloidal Particles. *Geochim. Cosmochim. Acta* **2000**, *64* (24), 4131–4151.
- (9) Said-Pullicino, D.; Giannetta, B.; Demeglio, B.; Missong, A.; Gottselig, N.; Romani, M.; Bol, R.; Klumpp, E.; Celi, L. Redox-Driven Changes in Water-Dispersible Colloids and Their Role in Carbon Cycling in Hydromorphic Soils. *Geoderma* **2021**, *385*, 114894.
- (10) Pontoni, L.; van Hullebusch, E.; Pechaud, Y.; Fabbicino, M.; Esposito, G.; Pirozzi, F. Colloidal Mobilization and Fate of Trace Heavy Metals in Semi-Saturated Artificial Soil (OECD) Irrigated with Treated Wastewater. *Sustainability* **2016**, *8* (12), 1257.
- (11) Kim, J. I.; Zeh, P.; Delakowitz, B. Chemical Interactions of Actinide Ions with Groundwater Colloids in Gorleben Aquifer Systems. *Radiochim. Acta* **1992**, *58–59* (1), 147–154.
- (12) Flury, M.; Mathison, J. B.; Harsh, J. B. In Situ Mobilization of Colloids and Transport of Cesium in Hanford Sediments. *Environ. Sci. Technol.* **2002**, *36* (24), 5335–5341.
- (13) Lov, A.; Cornelis, G.; Larsbo, M.; Persson, I.; Sjøstedt, C.; Gustafsson, J. P.; Boye, K.; Kleja, D. B. Particle- and Colloid-Facilitated Pb Transport in Four Historically Contaminated Soils - Speciation and Effect of Irrigation Intensity. *Appl. Geochem.* **2018**, *96*, 327–338.
- (14) Deb, D.; Chakma, S. *Colloid and Colloid-Facilitated Contaminant Transport in Subsurface Ecosystem—a Concise Review*; Springer Berlin Heidelberg: 2022.
- (15) Henderson, R.; Kabengi, N.; Mantripragada, N.; Cabrera, M.; Hassan, S.; Thompson, A. Anoxia-Induced Release of Colloid- and Nanoparticle-Bound Phosphorus in Grassland Soils. *Environ. Sci. Technol.* **2012**, *46* (21), 11727–11734.
- (16) Missong, A.; Bol, R.; Nischwitz, V.; Krüger, J.; Lang, F.; Siemens, J.; Klumpp, E. Phosphorus in Water Dispersible-Colloids of Forest Soil Profiles. *Plant Soil* **2018**, *427* (1–2), 71–86.
- (17) Weber, F. A.; Voegelin, A.; Kretzschmar, R. Multi-Metal Contaminant Dynamics in Temporarily Flooded Soil under Sulfate Limitation. *Geochim. Cosmochim. Acta* **2009**, *73* (19), 5513–5527.
- (18) Lair, G. J.; Zehetner, F.; Fiebig, M.; Gerzabek, M. H.; van Gestel, C. A. M.; Hein, T.; Hohensinner, S.; Hsu, P.; Jones, K. C.; Jordan, G.; Koelmans, A. A.; Poot, A.; Slijkerman, D. M. E.; Totsche, K. U.; Bondar-Kunze, E.; Barth, J. A. C. How Do Long-Term Development and Periodical Changes of River-Floodplain Systems Affect the Fate of Contaminants? Results from European Rivers. *Environ. Pollut.* **2009**, *157* (12), 3336–3346.
- (19) Zachara, J. M.; Long, P. E.; Bargar, J.; Davis, J. A.; Fox, P.; Fredrickson, J. K.; Freshley, M. D.; Konopka, A. E.; Liu, C.; McKinley, J. P.; Rockhold, M. L.; Williams, K. H.; Yabusaki, S. B. Persistence of Uranium Groundwater Plumes: Contrasting Mechanisms at Two DOE Sites in the Groundwater-River Interaction Zone. *J. Contam. Hydrol.* **2013**, *147*, 45–72.
- (20) Lynch, S.; Batty, L.; Byrne, P. Environmental Risk of Metal Mining Contaminated River Bank Sediment at Redox-Transitional Zones. *Minerals* **2014**, *4* (1), 52–73.
- (21) Schulz-Zunkel, C.; Rinklebe, J.; Bork, H.-R. Trace Element Release Patterns from Three Floodplain Soils under Simulated Oxidized-Reduced Cycles. *Ecol. Eng.* **2015**, *83*, 485–495.
- (22) Yabusaki, S. B.; Wilkins, M. J.; Fang, Y.; Williams, K. H.; Arora, B.; Bargar, J.; Beller, H. R.; Bouskill, N. J.; Brodie, E. L.; Christensen, J. N.; Conrad, M. E.; Danczak, R. E.; King, E.; Soltanian, M. R.; Spycher, N. F.; Steefel, C. I.; Tokunaga, T. K.; Versteeg, R.; Waichler, S. R.; Wainwright, H. M. Water Table Dynamics and Biogeochemical Cycling in a Shallow, Variably-Saturated Floodplain. *Environ. Sci. Technol.* **2017**, *51* (6), 3307–3317.
- (23) Lehmann, K.; Lehmann, R.; Totsche, K. U. Event-Driven Dynamics of the Total Mobile Inventory in Undisturbed Soil Account for Significant Fluxes of Particulate Organic Carbon. *Sci. Total Environ.* **2021**, *756*, 143774.
- (24) Dibbern, D.; Schmalwasser, A.; Lueders, T.; Totsche, K. U. Selective Transport of Plant Root-Associated Bacterial Populations in Agricultural Soils upon Snowmelt. *Soil Biol. Biochem.* **2014**, *69*, 187–196.
- (25) Kjaergaard, C.; Moldrup, P.; de Jonge, L. W.; Jacobsen, O. H. Colloid Mobilization and Transport in Undisturbed Soil Columns. II.

- The Role of Colloid Dispersibility and Preferential Flow. *Vadose Zo. J.* **2004**, *3* (2), 424–433.
- (26) Natsch, A.; Keel, C.; Troxler, J.; Zala, M.; Von Albertini, N.; Defago, G. Importance of Preferential Flow and Soil Management in Vertical Transport of a Biocontrol Strain of *Pseudomonas fluorescens* in Structured Field Soil. *Appl. Environ. Microbiol.* **1996**, *62* (1), 33–40.
- (27) Dixit, S. P. Surface Charge Characteristics of Soil Colloids of Some Different Soil Types. *Zeitschrift für Pflanzenernährung und Bodenkd.* **1980**, *143* (4), 422–431.
- (28) Thompson, A.; Chadwick, O. A.; Boman, S.; Chorover, J. Colloid Mobilization during Soil Iron Redox Oscillations. *Environ. Sci. Technol.* **2006**, *40* (18), 5743–5749.
- (29) Rod, K. A.; Patel, K. F.; Kumar, S.; Cantando, E.; Leng, W.; Kukkadapu, R. K.; Qafoku, O.; Bowden, M.; Kaplan, D. I.; Kemner, K. M. Dispersible Colloid Facilitated Release of Organic Carbon From Two Contrasting Riparian Sediments. *Front. Water* **2020**, *2*, 1–8.
- (30) Afsar, M. Z.; Goodwin, C.; Beebe, T. P.; Jaisi, D. P.; Jin, Y. Quantification and Molecular Characterization of Organo-Mineral Associations as Influenced by Redox Oscillations. *Sci. Total Environ.* **2020**, *704*, 135454.
- (31) Noël, V.; Kumar, N.; Boye, K.; Barragan, L.; Lezama-Pacheco, J. S.; Chu, R.; Tolic, N.; Brown, G. E.; Bargar, J. R. FeS Colloids-Formation and Mobilization Pathways in Natural Waters. *Environ. Sci. Nano* **2020**, *7* (7), 2102–2116.
- (32) Bunn, R. A.; Magelky, R. D.; Ryan, J. N.; Elimelech, M. Mobilization of Natural Colloids from an Iron Oxide-Coated Sand Aquifer: Effect of PH and Ionic Strength. *Environ. Sci. Technol.* **2002**, *36* (3), 314–322.
- (33) Abgottsporn, F.; Bigalke, M.; Wilcke, W. Fast Colloidal and Dissolved Release of Trace Elements in a Carbonatic Soil after Experimental Flooding. *Geoderma* **2015**, *259–260*, 156–163.
- (34) Liao, P.; Yuan, S.; Wang, D. Impact of Redox Reactions on Colloid Transport in Saturated Porous Media: An Example of Ferrihydrite Colloids Transport in the Presence of Sulfide. *Environ. Sci. Technol.* **2016**, *50* (20), 10968–10977.
- (35) He, Y. T.; Wan, J.; Tokunaga, T. Kinetic Stability of Hematite Nanoparticles: The Effect of Particle Sizes. *J. Nanoparticle Res.* **2008**, *10* (2), 321–332.
- (36) Wan, M.; Schröder, C.; Peiffer, S. Fe(III):S(-II) Concentration Ratio Controls the Pathway and the Kinetics of Pyrite Formation during Sulfidation of Ferric Hydroxides. *Geochim. Cosmochim. Acta* **2017**, *217*, 334–348.
- (37) Liao, P.; Li, W.; Jiang, Y.; Wu, J.; Yuan, S.; Fortner, J. D.; Giammar, D. E. Formation, Aggregation, and Deposition Dynamics of NOM-Iron Colloids at Anoxic-Oxic Interfaces. *Environ. Sci. Technol.* **2017**, *51* (21), 12235–12245.
- (38) Li, B.; Liao, P.; Xie, L.; Li, Q.; Pan, C.; Ning, Z.; Liu, C. Reduced NOM Triggered Rapid Cr(VI) Reduction and Formation of NOM-Cr(III) Colloids in Anoxic Environments. *Water Res.* **2020**, *181*, 115923.
- (39) Buettner, S. W.; Kramer, M. G.; Chadwick, O. A.; Thompson, A. Mobilization of Colloidal Carbon during Iron Reduction in Basaltic Soils. *Geoderma* **2014**, *221–222*, 139–145.
- (40) Yan, J.; Lazouskaya, V.; Jin, Y. Soil Colloid Release Affected by Dissolved Organic Matter and Redox Conditions. *Vadose Zo. J.* **2016**, *15* (3), 1–10.
- (41) Liang, L.; McCarthy, J. F.; Jolley, L. W.; McNabb, J. A.; Mehlhorn, T. L. Iron Dynamics: Transformation of Fe(II)/Fe(III) during Injection of Natural Organic Matter in a Sandy Aquifer. *Geochim. Cosmochim. Acta* **1993**, *57* (9), 1987–1999.
- (42) Sharma, V. K.; Filip, J.; Zboril, R.; Varma, R. S. Natural Inorganic Nanoparticles - Formation, Fate, and Toxicity in the Environment. *Chem. Soc. Rev.* **2015**, *44* (23), 8410–8423.
- (43) Stolze, L.; Zhang, D.; Guo, H.; Rolle, M. Surface Complexation Modeling of Arsenic Mobilization from Goethite: Interpretation of an in-Situ Experiment. *Geochim. Cosmochim. Acta* **2019**, *248*, 274–288.
- (44) Stolze, L.; Battistel, M.; Rolle, M. Oxidative Dissolution of Arsenic-Bearing Sulfide Minerals in Groundwater: Impact of Hydrochemical and Hydrodynamic Conditions on Arsenic Release and Surface Evolution. *Environ. Sci. Technol.* **2022**, *56* (8), 5049–5061.
- (45) Pullin, M. J.; Cabaniss, S. E. The Effects of PH, Ionic Strength, and Iron-Fulvic Acid Interactions on the Kinetics of Non-Photochemical Iron Transformations. I. Iron(II) Oxidation and Iron(III) Colloid Formation. *Geochim. Cosmochim. Acta* **2003**, *67* (21), 4067–4077.
- (46) Ranville, J. F.; Chittleborough, D. J.; Beckett, R. Particle-Size and Element Distributions of Soil Colloids. *Soil Sci. Soc. Am. J.* **2005**, *69* (4), 1173–1184.
- (47) Puls, R. W.; Clark, D. A.; Paul, C. J. Surface Chemical Effects on Colloid Stability and Transport through Natural Porous Media. In *Colloids in the Aquatic Environment*; Elsevier: 1993; pp 287–300.
- (48) Ryan, J. N.; Elimelech, M. Colloid Mobilization and Transport in Groundwater. *Colloids Surfaces A Physicochem. Eng. Asp.* **1996**, *107*, 1–56.
- (49) Molnar, I. L.; Johnson, W. P.; Gerhard, J. I.; Willson, C. S.; O'Carroll, D. M. Predicting Colloid Transport through Saturated Porous Media: A Critical Review. *Water Resour. Res.* **2015**, *51* (9), 6804–6845.
- (50) Ouyang, Y.; Shinde, D.; Mansell, R. S.; Harris, W. Colloid-enhanced Transport of Chemicals in Subsurface Environments: A Review. *Crit. Rev. Environ. Sci. Technol.* **1996**, *26* (2), 189–204.
- (51) Zhang, Z.; Furman, A. Soil Redox Dynamics under Dynamic Hydrologic Regimes - A Review. *Sci. Total Environ.* **2021**, *763*, 143026.
- (52) Kanti Sen, T.; Khilar, K. C. Review on Subsurface Colloids and Colloid-Associated Contaminant Transport in Saturated Porous Media. *Adv. Colloid Interface Sci.* **2006**, *119* (2–3), 71–96.
- (53) Sen, T. K. Processes in Pathogenic Biocolloidal Contaminants Transport in Saturated and Unsaturated Porous Media: A Review. *Water, Air, Soil Pollut.* **2011**, *216* (1–4), 239–256.
- (54) Hunter, R. J.; Ekdawi, N. The Sedimentation Behaviour of Coagulated Particles of Small Radius. *Colloids Surf.* **1986**, *18* (2–4), 325–340.
- (55) Horowitz, A. J. A Review of Selected Inorganic Surface Water Quality-Monitoring Practices: Are We Really Measuring What We Think, and If So, Are We Doing It Right? *Environ. Sci. Technol.* **2013**, *47* (6), 2471–2486.
- (56) Morrison, M. A.; Benoit, G. Filtration Artifacts Caused by Overloading Membrane Filters. *Environ. Sci. Technol.* **2001**, *35* (18), 3774–3779.
- (57) Horowitz, A. J. Some Thoughts on Problems Associated With Various Sampling Media Used for Environmental Monitoring†. *Analyst* **1997**, *122* (11), 1193–1200.
- (58) Zirkler, D.; Lang, F.; Kaupenjohann, M. Lost in Filtration”—The Separation of Soil Colloids from Larger Particles. *Colloids Surfaces A Physicochem. Eng. Asp.* **2012**, *399*, 35–40.
- (59) Rickard, D. Sulfidic Sediments and Sedimentary Rocks. In *Developments in Sedimentology*; Elsevier: 2012.
- (60) Combes, J. M.; Manceau, A.; Calas, G.; Bottero, J. Y. Formation of Ferric Oxides from Aqueous Solutions: A Polyhedral Approach by X-Ray Absorption Spectroscopy: I. Hydrolysis and Formation of Ferric Gels. *Geochim. Cosmochim. Acta* **1989**, *53* (3), 583–594.
- (61) Hotze, E. M.; Phenrat, T.; Lowry, G. V. Nanoparticle Aggregation: Challenges to Understanding Transport and Reactivity in the Environment. *J. Environ. Qual.* **2010**, *39* (6), 1909–1924.
- (62) Mansor, M.; Winkler, C.; Hochella, M. F.; Xu, J. Nanoparticulate Nickel-Hosting Phases in Sulfidic Environments: Effects of Ferrous Iron and Bacterial Presence on Mineral Formation Mechanism and Solid-Phase Nickel Distribution. *Front. Earth Sci.* **2019**, *7* (June), 1–15.
- (63) Weber, F. A.; Voegelin, A.; Kaegi, R.; Kretzschmar, R. Contaminant Mobilization by Metallic Copper and Metal Sulphide Colloids in Flooded Soil. *Nat. Geosci.* **2009**, *2* (4), 267–271.
- (64) Bosch, J.; Heister, K.; Hofmann, T.; Meckenstock, R. U. Nanosized Iron Oxide Colloids Strongly Enhance Microbial Iron Reduction. *Appl. Environ. Microbiol.* **2010**, *76* (1), 184–189.

- (65) Bradford, S. A.; Morales, V. L.; Zhang, W.; Harvey, R. W.; Packman, A. I.; Mohanram, A.; Welty, C. Transport and Fate of Microbial Pathogens in Agricultural Settings. *Crit. Rev. Environ. Sci. Technol.* **2013**, *43* (8), 775–893.
- (66) Luther, G. W.; Rickard, D. T. Metal Sulfide Cluster Complexes and Their Biogeochemical Importance in the Environment. *J. Nanoparticle Res.* **2005**, *7* (4–5), 389–407.
- (67) Rickard, D.; Luther, G. W. Chemistry of Iron Sulfides. *Chem. Rev.* **2007**, *107* (2), 514–562.
- (68) Luther, G. W.; Rickard, D. T.; Theberge, S.; Olroyd, A. Determination of Metal (Bi)Sulfide Stability Constants of Mn 2+, Fe 2+, Co 2+, Ni 2+, Cu 2+, and Zn 2+ by Voltammetric Methods. *Environ. Sci. Technol.* **1996**, *30* (2), 671–679.
- (69) Luther, G. W.; Theberge, S. M.; Rickard, D. T. Evidence for Aqueous Clusters as Intermediates during Zinc Sulfide Formation. *Geochim. Cosmochim. Acta* **1999**, *63* (19–20), 3159–3169.
- (70) Vazquez G, F.; Zhang, J.; Millero, F. J. Effect of Metals on the Rate of the Oxidation of H₂S in Seawater. *Geophys. Res. Lett.* **1989**, *16* (12), 1363–1366.
- (71) Rozan, T. F.; Lassman, M. E.; Ridge, D. P.; Luther, G. W. Evidence for Iron, Copper and Zinc Complexation as Multinuclear Sulphide Clusters in Oxidizing Rivers. *Nature* **2000**, *406* (6798), 879–882.
- (72) Gammons, C. H.; Fradsen, A. K. Fate and Transport of Metals in H₂S-Rich Waters at a Treatment Wetland. *Geochem. Trans.* **2001**, *2*, 1–15.
- (73) Sukola, K.; Wang, F.; Tessier, A. Metal-Sulfide Species in Oxidizing Waters. *Anal. Chim. Acta* **2005**, *528* (2), 183–195.
- (74) Zhang, H.; Chen, B.; Banfield, J. F. Particle Size and PH Effects on Nanoparticle Dissolution. *J. Phys. Chem. C* **2010**, *114* (35), 14876–14884.
- (75) Gilbert, B.; Huang, F.; Zhang, H.; Waychunas, G. A.; Banfield, J. F. Nanoparticles: Strained and Stiff. *Science* (80-). **2004**, *305* (5684), 651–654.
- (76) Jiang, Q.; Liang, L. H.; Zhao, D. S. Lattice Contraction and Surface Stress of Fcc Nanocrystals. *J. Phys. Chem. B* **2001**, *105* (27), 6275–6277.
- (77) Lu, S. H.; Chen, T. F.; Wang, A. J.; Wu, Z. L.; Wang, Y. S. Lattice and Optical Property Evolution of Ultra-Small ZnS Quantum Dots Grown from a Single-Source Precursor. *Appl. Surf. Sci.* **2014**, *299*, 116–122.
- (78) Masadeh, A. S.; Božin, E. S.; Farrow, C. L.; Paglia, G.; Juhas, P.; Billinge, S. J. L.; Karkamkar, A.; Kanatzidis, M. G. Quantitative Size-Dependent Structure and Strain Determination of CdSe Nanoparticles Using Atomic Pair Distribution Function Analysis. *Phys. Rev. B* **2007**, *76* (11), 115413.
- (79) Auffan, M.; Rose, J.; Bottero, J.-Y.; Lowry, G. V.; Jolivet, J.-P.; Wiesner, M. R. Towards a Definition of Inorganic Nanoparticles from an Environmental, Health and Safety Perspective. *Nat. Nanotechnol.* **2009**, *4* (10), 634–641.
- (80) Mavrikakis, M.; Hammer, B.; Nørskov, J. K. Effect of Strain on the Reactivity of Metal Surfaces. *Phys. Rev. Lett.* **1998**, *81* (13), 2819–2822.
- (81) Strasser, P.; Koh, S.; Annyev, T.; Greeley, J.; More, K.; Yu, C.; Liu, Z.; Kaya, S.; Nordlund, D.; Ogasawara, H.; Toney, M. F.; Nilsson, A. Lattice-Strain Control of the Activity in Dealloyed Core-Shell Fuel Cell Catalysts. *Nat. Chem.* **2010**, *2* (6), 454–460.
- (82) Pratt, A.; Lari, L.; Hovorka, O.; Shah, A.; Woffinden, C.; Tear, S. P.; Binns, C.; Kröger, R. Enhanced Oxidation of Nanoparticles through Strain-Mediated Ionic Transport. *Nat. Mater.* **2014**, *13* (1), 26–30.
- (83) Horn, D.; Rieger, J. Organic Nanoparticles in the Aqueous Phase—Theory, Experiment, and Use. *Angew. Chemie Int. Ed.* **2001**, *40* (23), 4330.
- (84) Distler, D. *Polymeric Dispersions: Principles and Applications*; Asua, J. M., Ed.; Springer Netherlands: 1997.
- (85) Brinker, C. J.; Scherer, G. W. *Sol-Gel Science*; Academic Press: 1990.
- (86) Brady, N. C.; Weil, R. R. *The Nature and Properties of Soil*; Pearson: 2008.
- (87) Stevenson, F. J.; Olsen, R. A. A Simplified Representation of the Chemical Nature and Reactions of Soil Humus. *J. Agron. Educ.* **1989**, *18* (2), 84–88.
- (88) Elias, E. *Soils of the Ethiopian Highlands*; CASCAPE Project: 2016.
- (89) Bourgeat-Lami, E. Organic/Inorganic Nanocomposite Colloids. *Encyclopedia of Nanoscience and Nanotechnology* **2004**, 305–332.
- (90) Li, S.; Zhang, W.; Zhang, D.; Xiu, W.; Wu, S.; Chai, J.; Ma, J.; Jat Baloch, M. Y.; Sun, S.; Yang, Y. Migration Risk of Escherichia Coli O157:H7 in Unsaturated Porous Media in Response to Different Colloid Types and Compositions. *Environ. Pollut.* **2023**, *323*, 121282.
- (91) Steenhuis, T. S.; Morales, V. L.; Cakmak, M. E.; Salvucci, A. E.; Zhang, W. Biocolloids: Transport and Retention in Soils. In *Encyclopedia of Earth Sciences Series*; Gliński, J.; Horabik, J.; Lipiec, J., Eds.; Springer: 2011; pp 66–70.
- (92) Hochella, M. F.; Lower, S. K.; Maurice, P. A.; Penn, R. L.; Sahai, N.; Sparks, D. L.; Twining, B. S. Nanominerals, Mineral Nanoparticles, and Earth Systems. *Science* (80-). **2008**, *319* (5870), 1631–1635.
- (93) Gilbert, B.; Banfield, J. F. Molecular Geomicrobiology. In *Reviews in Mineralogy and Geochemistry*; Banfield, J. F., Cervini-Silva, J., Nealon, K. H., Eds.; Mineralogical Society of America: 2005; pp 109–155.
- (94) Tang, R.; Wang, L.; Nancollas, G. H. Size-Effects in the Dissolution of Hydroxyapatite: An Understanding of Biological Demineralization. *J. Mater. Chem.* **2004**, *14* (14), 2341.
- (95) Suarez, D. L.; Rhoades, J. D.; Lavado, R.; Grieve, C. M. Effect of PH on Saturated Hydraulic Conductivity and Soil Dispersion. *Soil Sci. Soc. Am. J.* **1984**, *48* (1), 50–55.
- (96) Heil, D.; Sposito, G. Organic Matter Role in Illitic Soil Colloids Flocculation: I. Counter Ions and PH. *Soil Sci. Soc. Am. J.* **1993**, *57* (5), 1241–1246.
- (97) Li, D.; Chang, H.-S.; Seaman, J. C.; Kaplan, D. I. Effects of Matrix Heterogeneity and Aqueous Humic Acid on Transport and Deposition of Mineral Colloids in Sandy Sediments. *J. Environ. Chem. Eng.* **2013**, *1* (4), 875–883.
- (98) Grolimund, D.; Elimelech, M.; Borkovec, M. Aggregation and Deposition Kinetics of Mobile Colloidal Particles in Natural Porous Media. *Colloids Surfaces A Physicochem. Eng. Asp.* **2001**, *191* (1–2), 179–188.
- (99) Su, Y.; Jassby, D.; Zhang, Y.; Keller, A. A.; Adeleye, A. S. Comparison of the Colloidal Stability, Mobility, and Performance of Nanoscale Zerovalent Iron and Sulfidated Derivatives. *J. Hazard. Mater.* **2020**, *396*, 122691.
- (100) Kretzschmar, R.; Sticher, H. Transport of Humic-Coated Iron Oxide Colloids in a Sandy Soil: Influence of Ca 2+ and Trace Metals. *Environ. Sci. Technol.* **1997**, *31* (12), 3497–3504.
- (101) Sposito, G. *The Surface Chemistry of Natural Particles*; Oxford University Press: 2004.
- (102) Yuwono, V. M.; Burrows, N. D.; Soltis, J. A.; Anh Do, T.; Lee Penn, R. Aggregation of Ferrihydrite Nanoparticles in Aqueous Systems. *Faraday Discuss.* **2012**, *159*, 235.
- (103) Hsu, J.-P.; Kuo, Y.-C. The Critical Coagulation Concentration of Counterions: Spherical Particles in Asymmetric Electrolyte Solutions. *J. Colloid Interface Sci.* **1997**, *185* (2), 530–537.
- (104) Legg, B. A.; Zhu, M.; Comolli, L. R.; Gilbert, B.; Banfield, J. F. Impacts of Ionic Strength on Three-Dimensional Nanoparticle Aggregate Structure and Consequences for Environmental Transport and Deposition. *Environ. Sci. Technol.* **2014**, *48* (23), 13703–13710.
- (105) Jewett, D. G.; Hilbert, T. A.; Logan, B. E.; Arnold, R. G.; Bales, R. C. Bacterial Transport in Laboratory Columns and Filters: Influence of Ionic Strength and PH on Collision Efficiency. *Water Res.* **1995**, *29* (7), 1673–1680.
- (106) He, L.; Xie, L.; Wang, D.; Li, W.; Fortner, J. D.; Li, Q.; Duan, Y.; Shi, Z.; Liao, P.; Liu, C. Elucidating the Role of Sulfide on the Stability of Ferrihydrite Colloids under Anoxic Conditions. *Environ. Sci. Technol.* **2019**, *53* (8), 4173–4184.

- (107) Chin, Y.; Gschwend, P. M. The Abundance, Distribution, and Configuration of Porewater Organic Colloids in Recent Sediments. *Geochim. Cosmochim. Acta* **1991**, *55* (5), 1309–1317.
- (108) Kjaergaard, C.; Poulsen, T. G.; Moldrup, P.; de Jonge, L. W. Colloid Mobilization and Transport in Undisturbed Soil Columns. I. Pore Structure Characterization and Tritium Transport. *Vadose Zo. J.* **2004**, *3* (2), 413–423.
- (109) Kjaergaard, C.; de Jonge, L. W.; Moldrup, P.; Schjønning, P. Water-Dispersible Colloids: Effects of Measurement Method, Clay Content, Initial Soil Matrix Potential, and Wetting Rate. *Vadose Zo. J.* **2004**, *3* (2), 403–412.
- (110) Gu, S.; Lian, F.; Han, Y.; Wang, Z.; Xing, B. Effect of Root Exudates on the Release, Surface Property, Colloidal Stability, and Phytotoxicity of Dissolved Black Carbon. *Ecotoxicol. Environ. Saf.* **2022**, *239*, 113687.
- (111) Seta, A. K.; Karathanasis, A. D. Atrazine Adsorption by Soil Colloids and Co-Transport through Subsurface Environments. *Soil Sci. Soc. Am. J.* **1997**, *61* (2), 612.
- (112) Poulsen, T. G.; Moldrup, P.; de Jonge, L. W.; Komatsu, T. Colloid and Bromide Transport in Undisturbed Soil Columns: Application of Two-Region Model. *Vadose Zo. J.* **2006**, *5* (2), 649–656.
- (113) Dale, J. G.; Stegemeier, J. P.; Kim, C. S. Aggregation of Nanoscale Iron Oxyhydroxides and Corresponding Effects on Metal Uptake, Retention, and Speciation: I. Ionic-Strength and PH. *Geochim. Cosmochim. Acta* **2015**, *148*, 100–112.
- (114) Amal, R.; Raper, J.; Waite, T. Effect of Fulvic Acid Adsorption on the Aggregation Kinetics and Structure of Hematite Particles. *J. Colloid Interface Sci.* **1992**, *151* (1), 244–257.
- (115) Mylon, S. E.; Chen, K. L.; Elimelech, M. Influence of Natural Organic Matter and Ionic Composition on the Kinetics and Structure of Hematite Colloid Aggregation: Implications to Iron Depletion in Estuaries. *Langmuir* **2004**, *20* (21), 9000–9006.
- (116) Chen, K. L.; Mylon, S. E.; Elimelech, M. Aggregation Kinetics of Alginate-Coated Hematite Nanoparticles in Monovalent and Divalent Electrolytes. *Environ. Sci. Technol.* **2006**, *40* (5), 1516–1523.
- (117) Baalousha, M. Aggregation and Disaggregation of Iron Oxide Nanoparticles: Influence of Particle Concentration, PH and Natural Organic Matter. *Sci. Total Environ.* **2009**, *407* (6), 2093–2101.
- (118) Baalousha, M.; Manciuola, A.; Cumberland, S.; Kendall, K.; Lead, J. R. Aggregation and Surface Properties of Iron Oxide Nanoparticles: Influence of PH and Natural Organic Matter. *Environ. Toxicol. Chem.* **2008**, *27* (9), 1875.
- (119) Engel, M.; Lezama Pacheco, J. S.; Noël, V.; Boye, K.; Fendorf, S. Organic Compounds Alter the Preference and Rates of Heavy Metal Adsorption on Ferrihydrite. *Sci. Total Environ.* **2021**, *750*, 141485.
- (120) Ponnampertuma, F. N.; Martinez, E.; Loy, T. Influence of Redox Potential and Partial Pressure of Carbon Dioxide on PH Values and Suspension Effect of Flooded Soils. *Soil Sci.* **1966**, *101* (6), 421–431.
- (121) Gillespie, L. J. Reduction Potentials of Bacterial Cultures and of Water-Logged Soils. *Soil Sci.* **1920**, *9* (4), 199–216.
- (122) Joffe, J. S. Soil Profile Studies: VII. The Gleit Process. *Soil Sci.* **1935**, *39* (5), 391–402.
- (123) Vepraskas, M. J.; Faulkner, S. P. Redox Chemistry of Hydric Soils. In *Wetland Soils: Genesis, Hydrology, Landscapes, and Classification*; Richardson, J. L., Vepraskas, M., Eds.; CRC Press: 2001; pp 85–106.
- (124) Tiller, C. L.; O'Melia, C. R. Natural Organic Matter and Colloidal Stability: Models and Measurements. *Colloids Surfaces A Physicochem. Eng. Asp.* **1993**, *73*, 89–102.
- (125) Wolthoorn, A.; Temminghoff, E. J. M.; Weng, L.; van Riemsdijk, W. H. Colloid Formation in Groundwater: Effect of Phosphate, Manganese, Silicate and Dissolved Organic Matter on the Dynamic Heterogeneous Oxidation of Ferrous Iron. *Appl. Geochem.* **2004**, *19* (4), 611–622.
- (126) Whitmore, S. L.; Hamilton, S. K. Rapid Removal of Nitrate and Sulfate in Freshwater Wetland Sediments. *J. Environ. Qual.* **2005**, *34* (6), 2062–2071.
- (127) Noël, V.; Boye, K.; Kukkadapu, R. K.; Bone, S.; Lezama Pacheco, J. S.; Cardarelli, E.; Janot, N.; Fendorf, S.; Williams, K. H.; Bargar, J. R. Understanding Controls on Redox Processes in Floodplain Sediments of the Upper Colorado River Basin. *Sci. Total Environ.* **2017**, *603–604*, 663–675.
- (128) Morse, J. W.; Arakaki, T. Adsorption and Coprecipitation of Divalent Metals with Mackinawite (FeS). *Geochim. Cosmochim. Acta* **1993**, *57* (15), 3635–3640.
- (129) Jeong, H. Y.; Klaue, B.; Blum, J. D.; Hayes, K. F. Sorption of Mercuric Ion by Synthetic Nanocrystalline Mackinawite (FeS). *Environ. Sci. Technol.* **2007**, *41* (22), 7699–7705.
- (130) Lewis, A. E. Review of Metal Sulphide Precipitation. *Hydrometallurgy* **2010**, *104* (2), 222–234.
- (131) Patterson, R. R.; Fendorf, S.; Fendorf, M. Reduction of Hexavalent Chromium by Amorphous Iron Sulfide. *Environ. Sci. Technol.* **1997**, *31* (7), 2039–2044.
- (132) Gallegos, T. J.; Hyun, S. P.; Hayes, K. F. Spectroscopic Investigation of the Uptake of Arsenite from Solution by Synthetic Mackinawite. *Environ. Sci. Technol.* **2007**, *41* (22), 7781–7786.
- (133) Jeong, H. Y.; Han, Y.-S.; Park, S. W.; Hayes, K. F. Aerobic Oxidation of Mackinawite (FeS) and Its Environmental Implication for Arsenic Mobilization. *Geochim. Cosmochim. Acta* **2010**, *74* (11), 3182–3198.
- (134) Han, Y.-S.; Demond, A. H.; Gallegos, T. J.; Hayes, K. F. Dependence of Particle Concentration Effect on PH and Redox for Arsenic Removal by FeS-Coated Sand under Anoxic Conditions. *Chemosphere* **2015**, *134*, 499–503.
- (135) Besold, J.; Eberle, A.; Noël, V.; Kujala, K.; Kumar, N.; Scheinost, A. C.; Pacheco, J. L.; Fendorf, S.; Planer-Friedrich, B. Antimonite Binding to Natural Organic Matter: Spectroscopic Evidence from a Mine Water Impacted Peatland. *Environ. Sci. Technol.* **2019**, *53* (18), 10792–10802.
- (136) Besold, J.; Kumar, N.; Scheinost, A. C.; Lezama Pacheco, J.; Fendorf, S.; Planer-Friedrich, B. Antimonite Complexation with Thiol and Carboxyl/Phenol Groups of Peat Organic Matter. *Environ. Sci. Technol.* **2019**, *53* (9), 5005–5015.
- (137) Han, D. S.; Batchelor, B.; Abdel-Wahab, A. Sorption of Selenium(IV) and Selenium(VI) to Mackinawite (FeS): Effect of Contact Time, Extent of Removal, Sorption Envelopes. *J. Hazard. Mater.* **2011**, *186* (1), 451–457.
- (138) Hyun, S. P.; Davis, J. A.; Sun, K.; Hayes, K. F. Uranium(VI) Reduction by Iron(II) Monosulfide Mackinawite. *Environ. Sci. Technol.* **2012**, *46* (6), 3369–3376.
- (139) Veeramani, H.; Scheinost, A. C.; Monsegue, N.; Qafoku, N. P.; Kukkadapu, R.; Newville, M.; Lanzirrotti, A.; Pruden, A.; Murayama, M.; Hochella, M. F. Abiotic Reductive Immobilization of U(VI) by Biogenic Mackinawite. *Environ. Sci. Technol.* **2013**, *47* (5), 2361–2369.
- (140) Livens, F. R.; Jones, M. J.; Hynes, A. J.; Charnock, J. M.; Mosselmans, J. F. W.; Hennig, C.; Steele, H.; Collison, D.; Vaughan, D. J.; Patrick, R. A. D.; Reed, W. A.; Moyes, L. N. X-Ray Absorption Spectroscopy Studies of Reactions of Technetium, Uranium and Neptunium with Mackinawite. *J. Environ. Radioact.* **2004**, *74* (1–3), 211–219.
- (141) Butler, E. C.; Hayes, K. F. Kinetics of the Transformation of Halogenated Aliphatic Compounds by Iron Sulfide. *Environ. Sci. Technol.* **2000**, *34* (3), 422–429.
- (142) Kyung, D.; Sihm, Y.; Kim, S.; Bae, S.; Amin, M. T.; Alazba, A. A.; Lee, W. Synergistic Effect of Nano-Sized Mackinawite with Cyano-Cobalamin in Cement Slurries for Reductive Dechlorination of Tetrachloroethylene. *J. Hazard. Mater.* **2016**, *311*, 1–10.
- (143) Tadanier, C. J.; Schreiber, M. E.; Roller, J. W. Arsenic Mobilization through Microbially Mediated Deflocculation of Ferrihydrite. *Environ. Sci. Technol.* **2005**, *39* (9), 3061–3068.

- (144) Bonneville, S.; Van Cappellen, P.; Behrends, T. Microbial Reduction of Iron(III) Oxyhydroxides: Effects of Mineral Solubility and Availability. *Chem. Geol.* **2004**, *212* (3–4), 255–268.
- (145) Arduino, E.; Barberis, E.; Boero, V. Iron Oxides and Particle Aggregation in B Horizons of Some Italian Soils. *Geoderma* **1989**, *45* (3–4), 319–329.
- (146) Ryan, J. N.; Gschwend, P. M. Colloid Mobilization in Two Atlantic Coastal Plain Aquifers: Field Studies. *Water Resour. Res.* **1990**, *26* (2), 307–322.
- (147) Grybos, M.; Davranche, M.; Gruau, G.; Petitjean, P.; Pédrot, M. Increasing PH Drives Organic Matter Solubilization from Wetland Soils under Reducing Conditions. *Geoderma* **2009**, *154* (1–2), 13–19.
- (148) De-Campos, A. B.; Mamedov, A. I.; Huang, C. Short-Term Reducing Conditions Decrease Soil Aggregation. *Soil Sci. Soc. Am. J.* **2009**, *73* (2), 550–559.
- (149) Grybos, M.; Davranche, M.; Gruau, G.; Petitjean, P. Is Trace Metal Release in Wetland Soils Controlled by Organic Matter Mobility or Fe-Oxyhydroxides Reduction? *J. Colloid Interface Sci.* **2007**, *314* (2), 490–501.
- (150) Rickard, D.; Morse, J. W. Acid Volatile Sulfide (AVS). *Mar. Chem.* **2005**, *97* (3–4), 141–197.
- (151) Buffle, J.; Vitre, R. R. de; Perret, D.; Leppard, G. G. Combining Field Measurements for Speciation in Non Perturbable Waters. In *Metal Speciation: Theory, Analysis and Application*; Kramer, J. R., Allen, H. E., Eds.; Lewis Publishers: 1988; pp 99–124.
- (152) Luther, G. W., III; Glazer, B.; Ma, S.; Trouwborst, R.; Shultz, B. R.; Druschel, G.; Kraiya, C. Iron and Sulfur Chemistry in a Stratified Lake: Evidence for Iron-Rich Sulfide Complexes. *Aquat. Geochemistry* **2003**, *9* (2), 87–110.
- (153) Theberge, S. M.; Luther, G. W., III Determination of the Electrochemical Properties of a Soluble Aqueous FeS Species Present in Sulfidic Solutions. *Aquat. Geochemistry* **1997**, *3*, 191–211.
- (154) Luther, G. W.; Rozan, T. F.; Taillefert, M.; Nuzzio, D. B.; Di Meo, C.; Shank, T. M.; Lutz, R. A.; Cary, S. C. Chemical Speciation Drives Hydrothermal Vent Ecology. *Nature* **2001**, *410* (6830), 813–816.
- (155) Taillefert, M.; Hover, V. C.; Rozan, T. F.; Theberge, S. M.; Luther, G. W. The Influence of Sulfides on Soluble Organic-Fe(III) in Anoxic Sediment Porewaters. *Estuaries* **2002**, *25* (6), 1088–1096.
- (156) Luther, G. W.; Theberge, S. M.; Rozan, T. F.; Rickard, D.; Rowlands, C. C.; Oldroyd, A. Aqueous Copper Sulfide Clusters as Intermediates during Copper Sulfide Formation. *Environ. Sci. Technol.* **2002**, *36* (3), 394–402.
- (157) Ferris, F. G.; Fyfe, W. S.; Beveridge, T. J. Bacteria as Nucleation Sites for Authigenic Minerals in a Metal-Contaminated Lake Sediment. *Chem. Geol.* **1987**, *63* (3–4), 225–232.
- (158) Kumar, N.; Couture, R.-M.; Millot, R.; Battaglia-Brunet, F.; Rose, J. Microbial Sulfate Reduction Enhances Arsenic Mobility Downstream of Zerovalent-Iron-Based Permeable Reactive Barrier. *Environ. Sci. Technol.* **2016**, *50* (14), 7610–7617.
- (159) Tufenkji, N. Modeling Microbial Transport in Porous Media: Traditional Approaches and Recent Developments. *Adv. Water Resour.* **2007**, *30* (6–7), 1455–1469.
- (160) Bebie, J.; Schoonen, M. A. A.; Fuhrmann, M.; Strongin, D. R. Surface Charge Development on Transition Metal Sulfides: An Electrokinetic Study. *Geochim. Cosmochim. Acta* **1998**, *62* (4), 633–642.
- (161) Horzempa, L. M.; Helz, G. R. Controls on the Stability of Sulfide Sols: Colloidal Covellite as an Example. *Geochim. Cosmochim. Acta* **1979**, *43* (10), 1645–1650.
- (162) Xu, H.; Xia, B.; He, E.; Qiu, R.; Peijnenburg, W. J. G. M.; Qiu, H.; Zhao, L.; Xu, X.; Cao, X. Dynamic Release and Transformation of Metallic Copper Colloids in Flooded Paddy Soil: Role of Soil Reducible Sulfate and Temperature. *J. Hazard. Mater.* **2021**, *402*, 123462.
- (163) Xu, H.; Ji, L.; Kong, M.; Jiang, H.; Chen, J. Molecular Weight-Dependent Adsorption Fractionation of Natural Organic Matter on Ferrihydrite Colloids in Aquatic Environment. *Chem. Eng. J.* **2019**, *363*, 356–364.
- (164) Riedel, T.; Zak, D.; Biester, H.; Dittmar, T. Iron Traps Terrestrially Derived Dissolved Organic Matter at Redox Interfaces. *Proc. Natl. Acad. Sci. U. S. A.* **2013**, *110* (25), 10101–10105.
- (165) Lalonde, K.; Mucci, A.; Ouellet, A.; Gélinas, Y. Preservation of Organic Matter in Sediments Promoted by Iron. *Nature* **2012**, *483* (7388), 198–200.
- (166) Adhikari, D.; Zhao, Q.; Das, K.; Mejia, J.; Huang, R.; Wang, X.; Poulson, S. R.; Tang, Y.; Roden, E. E.; Yang, Y. Dynamics of Ferrihydrite-Bound Organic Carbon during Microbial Fe Reduction. *Geochim. Cosmochim. Acta* **2017**, *212*, 221–233.
- (167) Han, L.; Sun, K.; Keiluweit, M.; Yang, Y.; Yang, Y.; Jin, J.; Sun, H.; Wu, F.; Xing, B. Mobilization of Ferrihydrite-Associated Organic Carbon during Fe Reduction: Adsorption versus Coprecipitation. *Chem. Geol.* **2019**, *503*, 61–68.
- (168) Giannetta, B.; Oliveira de Souza, D.; Aquilanti, G.; Celi, L.; Said-Pullicino, D. Redox-Driven Changes in Organic C Stabilization and Fe Mineral Transformations in Temperate Hydromorphic Soils. *Geoderma* **2022**, *406*, 115532.
- (169) Bao, T.; Wang, P.; Hu, B.; Wang, X.; Qian, J. Mobilization of Colloids during Sediment Resuspension and Its Effect on the Release of Heavy Metals and Dissolved Organic Matter. *Sci. Total Environ.* **2023**, *861*, 160678.
- (170) Ryan, J. N.; Gschwend, P. M. Effect of Solution Chemistry on Clay Colloid Release from an Iron Oxide-Coated Aquifer Sand. *Environ. Sci. Technol.* **1994**, *28* (9), 1717–1726.
- (171) Klüpfel, L.; Piepenbrock, A.; Kappler, A.; Sander, M. Humic Substances as Fully Regenerable Electron Acceptors in Recurrently Anoxic Environments. *Nat. Geosci.* **2014**, *7* (3), 195–200.
- (172) Babey, T.; Boye, K.; Tolar, B.; Engel, M.; Noël, V.; Perzan, Z.; Kumar, N.; Francis, C. A.; Bargar, J. R.; Maher, K. Simulation of Anoxic Lenses as Exporters of Reactivity in Alluvial Aquifer Sediments. *Geochim. Cosmochim. Acta* **2022**, *334*, 119–134.
- (173) Li, Q.; Xie, L.; Jiang, Y.; Fortner, J. D.; Yu, K.; Liao, P.; Liu, C. Formation and Stability of NOM-Mn(III) Colloids in Aquatic Environments. *Water Res.* **2019**, *149*, 190–201.
- (174) Tang, Y.; Zhang, M.; Sun, G.; Pan, G. Impact of Eutrophication on Arsenic Cycling in Freshwaters. *Water Res.* **2019**, *150*, 191–199.
- (175) Sun, Y.; Zuo, X.; Sankaranarayanan, S. K. R. S.; Peng, S.; Narayanan, B.; Kamath, G. Quantitative 3D Evolution of Colloidal Nanoparticle Oxidation in Solution. *Science* (80-.). **2017**, *356* (6335), 303–307.
- (176) Navrotsky, A. Thermochemistry of Nanomaterials. *Rev. Mineral. Geochemistry* **2001**, *44* (1), 73–103.
- (177) Waychunas, G. A.; Kim, C. S.; Banfield, J. F. Nanoparticulate Iron Oxide Minerals in Soils and Sediments: Unique Properties and Contaminant Scavenging Mechanisms. *J. Nanoparticle Res.* **2005**, *7* (4–5), 409–433.
- (178) Wigginton, N. S.; Haus, K. L.; Hochella, M. F., Jr Aquatic Environmental Nanoparticles. *J. Environ. Monit.* **2007**, *9* (12), 1306.
- (179) Filella, M. Colloidal Properties of Submicron Particles in Natural Waters. In *Environmental Colloids and Particles*; Wiley: 2007; pp 17–93.
- (180) Gunnars, A.; Blomqvist, S.; Johansson, P.; Andersson, C. Formation of Fe(III) Oxyhydroxide Colloids in Freshwater and Brackish Seawater, with Incorporation of Phosphate and Calcium. *Geochim. Cosmochim. Acta* **2002**, *66* (5), 745–758.
- (181) Lyvén, B.; Hasselöv, M.; Turner, D. R.; Haraldsson, C.; Andersson, K. Competition between Iron- and Carbon-Based Colloidal Carriers for Trace Metals in a Freshwater Assessed Using Flow Field-Flow Fractionation Coupled to ICPMS. *Geochim. Cosmochim. Acta* **2003**, *67* (20), 3791–3802.
- (182) Hochella, M. F. Environmentally Important, Poorly Crystalline Fe/Mn Hydrous Oxides: Ferrihydrite and a Possibly New Vernadite-like Mineral from the Clark Fork River Superfund Complex. *Am. Mineral.* **2005**, *90* (4), 718–724.

- (183) Hasselov, M.; von der Kammer, F. Iron Oxides as Geochemical Nanovectors for Metal Transport in Soil-River Systems. *Elements* **2008**, *4* (6), 401–406.
- (184) Fritzsche, A.; Rennert, T.; Totsche, K. U. Arsenic Strongly Associates with Ferrihydrite Colloids Formed in a Soil Effluent. *Environ. Pollut.* **2011**, *159* (5), 1398–1405.
- (185) Puls, R. W.; Powell, R. M. Transport of Inorganic Colloids through Natural Aquifer Material: Implications for Contaminant Transport. *Environ. Sci. Technol.* **1992**, *26* (3), 614–621.
- (186) Gunnars, A.; Blomqvist, S. Phosphate Exchange across the Sediment-Water Interface When Shifting from Anoxic to Oxidic Conditions: An Experimental Comparison of Freshwater and Brackish-Marine Systems. *Biogeochemistry* **1997**, *37*, 203–226.
- (187) Ren, J.; Packman, A. I. Coupled Stream-Subsurface Exchange of Colloidal Hematite and Dissolved Zinc, Copper, and Phosphate. *Environ. Sci. Technol.* **2005**, *39* (17), 6387–6394.
- (188) Baken, S.; Regelink, I. C.; Comans, R. N. J.; Smolders, E.; Koopmans, G. F. Iron-Rich Colloids as Carriers of Phosphorus in Streams: A Field-Flow Fractionation Study. *Water Res.* **2016**, *99*, 83–90.
- (189) Huang, S.; Chen, L.; Li, J.; Xu, J.; Xie, W.; Zhang, C. The Effects of Colloidal Fe and Mn on P Distribution in Groundwater System of Jiangnan Plain, China. *Sci. Total Environ.* **2023**, *854*, 158739.
- (190) Nierop, K. G. J.; Jansen, B.; Verstraten, J. M. Dissolved Organic Matter, Aluminium and Iron Interactions: Precipitation Induced by Metal/Carbon Ratio, PH and Competition. *Sci. Total Environ.* **2002**, *300* (1–3), 201–211.
- (191) Orem, W. H.; Gaudette, H. E. Organic Matter in Anoxic Marine Pore Water: Oxidation Effects. *Org. Geochem.* **1984**, *5* (4), 175–181.
- (192) Engel, M.; Noël, V.; Pierce, S.; Kovarik, L.; Kukkadapu, R. K.; Pacheco, J. S. L.; Qafoku, O.; Runyon, J. R.; Chorover, J.; Zhou, W.; Cliff, J.; Boye, K.; Bargar, J. R. Structure and Composition of Natural Ferrihydrite Nano-Colloids in Anoxic Groundwater. *Water Res.* **2023**, *238*, 119990.
- (193) Fu, D.; Keech, P. G.; Sun, X.; Wren, J. C. Iron Oxyhydroxide Nanoparticles Formed by Forced Hydrolysis: Dependence of Phase Composition on Solution Concentration. *Phys. Chem. Chem. Phys.* **2011**, *13* (41), 18523.
- (194) Angelico, R.; Ceglie, A.; He, J.-Z.; Liu, Y.-R.; Palumbo, G.; Colombo, C. Particle Size, Charge and Colloidal Stability of Humic Acids Coprecipitated with Ferrihydrite. *Chemosphere* **2014**, *99*, 239–247.
- (195) Li, W.; Liu, D.; Wu, J.; Kim, C.; Fortner, J. D. Aqueous Aggregation and Surface Deposition Processes of Engineered Superparamagnetic Iron Oxide Nanoparticles for Environmental Applications. *Environ. Sci. Technol.* **2014**, *48* (20), 11892–11900.
- (196) Wang, Y.; Fruttschi, M.; Suvorova, E.; Phrommavanh, V.; Descostes, M.; Osman, A. A. A.; Geipel, G.; Bernier-Latmani, R. Mobile Uranium(IV)-Bearing Colloids in a Mining-Impacted Wetland. *Nat. Commun.* **2013**, *4* (May), 1–9.
- (197) Liang, L.; Hofmann, A.; Gu, B. Ligand-Induced Dissolution and Release of Ferrihydrite Colloids. *Geochim. Cosmochim. Acta* **2000**, *64* (12), 2027–2037.
- (198) Hofmann, A.; Liang, L. Mobilization of Colloidal Ferrihydrite Particles in Porous Media—An Inner-Sphere Complexation Approach. *Geochim. Cosmochim. Acta* **2007**, *71* (24), 5847–5861.
- (199) Blöthe, M.; Roden, E. E. Microbial Iron Redox Cycling in a Circumneutral-PH Groundwater Seep. *Appl. Environ. Microbiol.* **2009**, *75* (2), 468–473.
- (200) Xia, B.; Qiu, H.; Knorr, K.-H. H.; Blodau, C.; Qiu, R. Occurrence and Fate of Colloids and Colloid-Associated Metals in a Mining-Impacted Agricultural Soil upon Prolonged Flooding. *J. Hazard. Mater.* **2018**, *348*, 56–66.
- (201) Rensing, C.; Grass, G. Escherichia Coli Mechanisms of Copper Homeostasis in a Changing Environment. *FEMS Microbiol. Rev.* **2003**, *27* (2–3), 197–213.
- (202) Outten, F. W.; Huffman, D. L.; Hale, J. A.; O'Halloran, T. V. The Independent Cue and CusSystems Confer Copper Tolerance during Aerobic and Anaerobic Growth InEscherichia Coli. *J. Biol. Chem.* **2001**, *276* (33), 30670–30677.
- (203) Kretzschmar, R.; Schafer, T. Metal Retention and Transport on Colloidal Particles in the Environment. *Elements* **2005**, *1* (4), 205–210.
- (204) Tosco, T.; Bosch, J.; Meckenstock, R. U.; Sethi, R. Transport of Ferrihydrite Nanoparticles in Saturated Porous Media: Role of Ionic Strength and Flow Rate. *Environ. Sci. Technol.* **2012**, *46* (7), 4008–4015.
- (205) Burrows, N. D.; Hale, C. R. H.; Penn, R. L. Effect of Ionic Strength on the Kinetics of Crystal Growth by Oriented Aggregation. *Cryst. Growth Des.* **2012**, *12* (10), 4787–4797.
- (206) Elimelech, M.; Gregory, J.; Jia, X. *Particle Deposition and Aggregation: Measurement, Modelling and Simulation*; Williams, R. A. F., Ed.; Butterworth-Heinemann: 2013.
- (207) Allain, C.; Cloitre, M.; Parris, F. Settling by Cluster Deposition in Aggregating Colloidal Suspensions. *J. Colloid Interface Sci.* **1996**, *178* (2), 411–416.
- (208) Sheng, A.; Liu, F.; Shi, L.; Liu, J. Aggregation Kinetics of Hematite Particles in the Presence of Outer Membrane Cytochrome OmcA of Shewanella Oneidensis MR-1. *Environ. Sci. Technol.* **2016**, *50* (20), 11016–11024.
- (209) Chekli, L.; Phuntsho, S.; Roy, M.; Lombi, E.; Donner, E.; Shon, H. K. Assessing the Aggregation Behaviour of Iron Oxide Nanoparticles under Relevant Environmental Conditions Using a Multi-Method Approach. *Water Res.* **2013**, *47* (13), 4585–4599.
- (210) Amstaetter, K.; Borch, T.; Kappler, A. Influence of Humic Acid Imposed Changes of Ferrihydrite Aggregation on Microbial Fe(III) Reduction. *Geochim. Cosmochim. Acta* **2012**, *85*, 326–341.
- (211) Milne, C. J.; Lapworth, D. J.; Gooddy, D. C.; Elgy, C. N.; Valsami-Jones, E. Role of Humic Acid in the Stability of Ag Nanoparticles in Suboxic Conditions. *Environ. Sci. Technol.* **2017**, *51* (11), 6063–6070.
- (212) Ma, R.; Stegemeier, J. P.; Levard, C.; Dale, J. G.; Noack, C. W.; Yang, T.; Brown, G. E.; Lowry, G. V. Sulfidation of Copper Oxide Nanoparticles and Properties of Resulting Copper Sulfide. *Environ. Sci. Nano* **2014**, *1* (4), 347–357.
- (213) Cao, Z.; Liu, X.; Xu, J.; Zhang, J.; Yang, Y.; Zhou, J.; Xu, X.; Lowry, G. V. Removal of Antibiotic Florfenicol by Sulfide-Modified Nanoscale Zero-Valent Iron. *Environ. Sci. Technol.* **2017**, *51* (19), 11269–11277.
- (214) Fu, H.; Qu, X.; Chen, W.; Zhu, D. Transformation and Destabilization of Graphene Oxide in Reducing Aqueous Solutions Containing Sulfide. *Environ. Toxicol. Chem.* **2014**, *33* (12), 2647–2653.
- (215) Wu, L.; Druschel, G.; Findlay, A.; Beard, B. L.; Johnson, C. M. Experimental Determination of Iron Isotope Fractionations among Feaq2+-FeSaq-Mackinawite at Low Temperatures: Implications for the Rock Record. *Geochim. Cosmochim. Acta* **2012**, *89*, 46–61.
- (216) Matamoros-Veloz, A.; Stawski, T. M.; Benning, L. G. Nanoparticle Assembly Leads to Mackinawite Formation. *Cryst. Growth Des.* **2018**, *18* (11), 6757–6764.
- (217) Schwertmann, U.; Fechter, H. The Point of Zero Charge of Natural and Synthetic Ferrihydrites and Its Relation to Adsorbed Silicate. *Clay Miner.* **1982**, *17* (4), 471–476.
- (218) Lowry, G. V.; Shaw, S.; Kim, C. S.; Rytuba, J. J.; Brown, G. E. Macroscopic and Microscopic Observations of Particle-Facilitated Mercury Transport from New Idria and Sulphur Bank Mercury Mine Tailings. *Environ. Sci. Technol.* **2004**, *38* (19), 5101–5111.
- (219) Anderson, M. S. The Influence of Substituted Cations on the Properties of Soil Colloids. *J. Agric. Res.* **1929**, *38* (10), 565–584.
- (220) Schulz, K.; Thomasarigo, L. K.; Kaegi, R.; Kretzschmar, R. Stabilization of Ferrihydrite and Lepidocrocite by Silicate during Fe(II)-Catalyzed Mineral Transformation: Impact on Particle Morphology and Silicate Distribution. *Environ. Sci. Technol.* **2022**, *56* (9), 5929–5938.

- (221) Liang, L.; Morgan, J. J. Chemical Aspects of Iron Oxide Coagulation in Water: Laboratory Studies and Implications for Natural Systems. *Aquat. Sci.* **1990**, *52* (1), 32–55.
- (222) Au, K.-K.; Penisson, A. C.; Yang, S.; O'Melia, C. R. Natural Organic Matter at Oxide/Water Interfaces: Complexation and Conformation. *Geochim. Cosmochim. Acta* **1999**, *63* (19–20), 2903–2917.
- (223) Kretzschmar, R.; Robarge, W. P.; Weed, S. B. Flocculation of Kaolinitic Soil Clays: Effects of Humic Substances and Iron Oxides. *Soil Sci. Soc. Am. J.* **1993**, *57* (5), 1277–1283.
- (224) Chorover, J.; Sposito, G. Colloid Chemistry of Kaolinitic Tropical Soils. *Soil Sci. Soc. Am. J.* **1995**, *59* (6), 1558–1564.
- (225) Vilg -Ritter, A.; Rose, J.; Masion, A.; Bottero, J.-Y.; Lain , J.-M. Chemistry and Structure of Aggregates Formed with Fe-Salts and Natural Organic Matter. *Colloids Surfaces A Physicochem. Eng. Asp.* **1999**, *147* (3), 297–308.
- (226) Sander, S.; Mosley, L. M.; Hunter, K. A. Investigation of Interparticle Forces in Natural Waters: Effects of Adsorbed Humic Acids on Iron Oxide and Alumina Surface Properties. *Environ. Sci. Technol.* **2004**, *38* (18), 4791–4796.
- (227) Tomb cz, E.; Libor, Z.; Ill s, E.; Majzik, A.; Klumpp, E. The Role of Reactive Surface Sites and Complexation by Humic Acids in the Interaction of Clay Mineral and Iron Oxide Particles. *Org. Geochem.* **2004**, *35* (3), 257–267.
- (228) Hajd , A.; Ill s, E.; Tomb cz, E.; Borb th, I. Surface Charging, Polyanionic Coating and Colloid Stability of Magnetite Nanoparticles. *Colloids Surfaces A Physicochem. Eng. Asp.* **2009**, *347* (1–3), 104–108.
- (229) Gilbert, B.; Huang, F.; Lin, Z.; Goodell, C.; Zhang, H.; Banfield, J. F. Surface Chemistry Controls Crystallinity of ZnS Nanoparticles. *Nano Lett.* **2006**, *6* (4), 605–610.
- (230) Le Bars, M.; Levard, C.; Legros, S.; Vidal, V.; Fernandez-Martinez, A.; Michel, F. M.; Thill, A.; Prelot, B.; Dublet-Adli, G.; Borschneck, D.; Rose, J.; Doelsch, E. Size and Strain of Zinc Sulfide Nanoparticles Altered by Interaction with Organic Molecules. *Environ. Sci. Technol.* **2022**, *56* (23), 16831–16837.
- (231) Zhang, H.; Gilbert, B.; Huang, F.; Banfield, J. F. Water-Driven Structure Transformation in Nanoparticles at Room Temperature. *Nature* **2003**, *424* (6952), 1025–1029.
- (232) Thompson, A.; Chadwick, O. A.; Rancourt, D. G.; Chorover, J. Iron-Oxide Crystallinity Increases during Soil Redox Oscillations. *Geochim. Cosmochim. Acta* **2006**, *70* (7), 1710–1727.
- (233) Winkler, P.; Kaiser, K.; Thompson, A.; Kalbitz, K.; Fiedler, S.; Jahn, R. Contrasting Evolution of Iron Phase Composition in Soils Exposed to Redox Fluctuations. *Geochim. Cosmochim. Acta* **2018**, *235*, 89–102.
- (234) De Yoreo, J. J. Principles of Crystal Nucleation and Growth. *Rev. Mineral. Geochemistry* **2003**, *54* (1), 57–93.
- (235) Nawrocki, G.; Cieplak, M. Interactions of Aqueous Amino Acids and Proteins with the (110) Surface of ZnS in Molecular Dynamics Simulations. *J. Chem. Phys.* **2014**, *140* (9), 095101.
- (236) Moreau, J. W.; Weber, P. K.; Martin, M. C.; Gilbert, B.; Hutcheon, I. D.; Banfield, J. F. Extracellular Proteins Limit the Dispersal of Biogenic Nanoparticles. *Science* (80-). **2007**, *316* (5831), 1600–1603.
- (237) Deonaraine, A.; Lau, B. L. T.; Aiken, G. R.; Ryan, J. N.; Hsu-Kim, H. Effects of Humic Substances on Precipitation and Aggregation of Zinc Sulfide Nanoparticles. *Environ. Sci. Technol.* **2011**, *45* (8), 3217–3223.
- (238) Gondikas, A. P.; Jang, E. K.; Hsu-Kim, H. Influence of Amino Acids Cysteine and Serine on Aggregation Kinetics of Zinc and Mercury Sulfide Colloids. *J. Colloid Interface Sci.* **2010**, *347* (2), 167–171.
- (239) Gondikas, A. P.; Masion, A.; Auffan, M.; Lau, B. L. T.; Hsu-Kim, H. Early-Stage Precipitation Kinetics of Zinc Sulfide Nanoclusters Forming in the Presence of Cysteine. *Chem. Geol.* **2012**, *329*, 10–17.
- (240) Lau, B. L. T.; Hsu-Kim, H. Precipitation and Growth of Zinc Sulfide Nanoparticles in the Presence of Thiol-Containing Natural Organic Ligands. *Environ. Sci. Technol.* **2008**, *42* (19), 7236–7241.
- (241) Hoffmann, K.; Christl, I.; Kaegi, R.; Kretzschmar, R. Effects of Natural Organic Matter (NOM), Metal-to-Sulfide Ratio and Mn²⁺ on Cadmium Sulfide Nanoparticle Growth and Colloidal Stability. *Environ. Sci. Nano* **2020**, *7* (11), 3385–3404.
- (242) Deonaraine, A.; Hsu-Kim, H. Precipitation of Mercuric Sulfide Nanoparticles in NOM-Containing Water: Implications for the Natural Environment. *Environ. Sci. Technol.* **2009**, *43* (7), 2368–2373.
- (243) Ravichandran, M.; Aiken, G. R.; Ryan, J. N.; Reddy, M. M. Inhibition of Precipitation and Aggregation of Metacinnabar (Mercuric Sulfide) by Dissolved Organic Matter Isolated from the Florida Everglades. *Environ. Sci. Technol.* **1999**, *33* (9), 1418–1423.
- (244) Vindedahl, A. M.; Stemig, M. S.; Arnold, W. A.; Penn, R. L. Character of Humic Substances as a Predictor for Goethite Nanoparticle Reactivity and Aggregation. *Environ. Sci. Technol.* **2016**, *50* (3), 1200–1208.
- (245) Ill s, E.; Tomb cz, E. The Effect of Humic Acid Adsorption on PH-Dependent Surface Charging and Aggregation of Magnetite Nanoparticles. *J. Colloid Interface Sci.* **2006**, *295* (1), 115–123.
- (246) Tomb cz, E.; T th, I. Y.; Nesztor, D.; Ill s, E.; Hajd , A.; Szekeres, M.; V k s, L. Adsorption of Organic Acids on Magnetite Nanoparticles, PH-Dependent Colloidal Stability and Salt Tolerance. *Colloids Surfaces A Physicochem. Eng. Asp.* **2013**, *435*, 91–96.
- (247) Lapworth, D. J.; Stolpe, B.; Williams, P. J.; Gooddy, D. C.; Lead, J. R. Characterization of Suboxic Groundwater Colloids Using a Multi-Method Approach. *Environ. Sci. Technol.* **2013**, *47* (6), 2554–2561.
- (248) Daugherty, E. E.; Gilbert, B.; Nico, P. S.; Borch, T. Complexation and Redox Buffering of Iron(II) by Dissolved Organic Matter. *Environ. Sci. Technol.* **2017**, *51* (19), 11096–11104.
- (249) Liao, P.; Li, W.; Wang, D.; Jiang, Y.; Pan, C.; Fortner, J. D.; Yuan, S. Effect of Reduced Humic Acid on the Transport of Ferrihydrite Nanoparticles under Anoxic Conditions. *Water Res.* **2017**, *109*, 347–357.
- (250) Demangeat, E.; P drot, M.; Dia, A.; Bouhnik-le-Coz, M.; Grasset, F.; Hanna, K.; Kamagate, M.; Cabello-Hurtado, F. Colloidal and Chemical Stabilities of Iron Oxide Nanoparticles in Aqueous Solutions: The Interplay of Structural, Chemical and Environmental Drivers. *Environ. Sci. Nano* **2018**, *5* (4), 992–1001.
- (251) Li, B.; Liao, P.; Liu, P.; Wang, D.; Ye, Z.; Wang, J.; Chen, J.; Ning, Z.; Jiang, Y.; Liu, C. Formation, Aggregation, and Transport of NOM-Cr(III) Colloids in Aquatic Environments. *Environ. Sci. Nano* **2022**, *9* (3), 1133–1145.
- (252) Buerge, I. J.; Hug, S. J. Influence of Organic Ligands on Chromium(VI) Reduction by Iron(II). *Environ. Sci. Technol.* **1998**, *32* (14), 2092–2099.
- (253) Pan, C.; Troyer, L. D.; Liao, P.; Catalano, J. G.; Li, W.; Giammar, D. E. Effect of Humic Acid on the Removal of Chromium(VI) and the Production of Solids in Iron Electrocoagulation. *Environ. Sci. Technol.* **2017**, *51* (11), 6308–6318.
- (254) Liao, P.; Pan, C.; Ding, W.; Li, W.; Yuan, S.; Fortner, J. D.; Giammar, D. E. Formation and Transport of Cr(III)-NOM-Fe Colloids upon Reaction of Cr(VI) with NOM-Fe(II) Colloids at Anoxic-Oxic Interfaces. *Environ. Sci. Technol.* **2020**, *54* (7), 4256–4266.
- (255) Li, B.; Zhang, S.; Liao, P.; Liu, P.; Ye, Z.; Liu, C. NOM-Induced Dissolution of Cr x Fe 1- x (OH) 3 Precipitates and Formation of Cr(III)-NOM-Fe Colloids under Oxic and Anoxic Conditions. *ACS Earth Sp. Chem.* **2022**, *6* (12), 2995–3006.
- (256) Tebo, B. M.; Neelson, K. H.; Emerson, S.; Jacobs, L. Microbialmediation of Mn(II) and Co(II) Precipitation at Oxic/Anoxic Interfaces Inthe Water Columns of Saanich Inlet, British Columbia and Framvaren Fjord, Norway. *Limnol. Ocean.* **1984**, *29*, 1247–1258.
- (257) Jones, M. E.; Nico, P. S.; Ying, S.; Regier, T.; Thieme, J.; Keiluweit, M. Manganese-Driven Carbon Oxidation at Oxic-Anoxic Interfaces. *Environ. Sci. Technol.* **2018**, *52* (21), 12349–12357.
- (258) Spencer, D. W.; Brewer, P. G. Vertical Advection Diffusion and Redox Potentials as Controls on the Distribution of Manganese

- and Other Trace Metals Dissolved in Waters of the Black Sea. *J. Geophys. Res.* **1971**, *76* (24), 5877–5892.
- (259) Aeppli, M.; Babey, T.; Engel, M.; Lacroix, E. M.; Tolar, B. B.; Fendorf, S.; Bargar, J. R.; Boye, K. Export of Organic Carbon from Reduced Fine-Grained Zones Governs Biogeochemical Reactivity in a Simulated Aquifer. *Environ. Sci. Technol.* **2022**, *56* (4), 2738–2746.
- (260) Kumar, N.; Noël, V.; Planer-Friedrich, B.; Besold, J.; Lezama-Pacheco, J. S.; Bargar, J. R.; Brown, G. E.; Fendorf, S.; Boye, K. Redox Heterogeneities Promote Thioarsenate Formation and Release into Groundwater from Low Arsenic Sediments. *Environ. Sci. Technol.* **2020**, *54* (6), 3237–3244.
- (261) Priadi, C.; Le Pape, P.; Morin, G.; Ayrault, S.; Maillot, F.; Juillot, F.; Hochreutener, R.; Llorens, I.; Testemale, D.; Proux, O.; Brown, G. E. X-Ray Absorption Fine Structure Evidence for Amorphous Zinc Sulfide as a Major Zinc Species in Suspended Matter from the Seine River Downstream of Paris, Ile-de-France, France. *Environ. Sci. Technol.* **2012**, *46* (7), 3712–3720.
- (262) Homoky, W. B.; Conway, T. M.; John, S. G.; König, D.; Deng, F.; Tagliabue, A.; Mills, R. A. Iron Colloids Dominate Sedimentary Supply to the Ocean Interior. *Proc. Natl. Acad. Sci. U. S. A.* **2021**, *118* (13), e2016078118.
- (263) Gomez-Gonzalez, M. A.; Voegelin, A.; Garcia-Guinea, J.; Bolea, E.; Laborda, F.; Garrido, F. Colloidal Mobilization of Arsenic from Mining-Affected Soils by Surface Runoff. *Chemosphere* **2016**, *144*, 1123–1131.
- (264) Nedkov, I.; Merodiiska, T.; Slavov, L.; Vandenberghe, R. E.; Kusano, Y.; Takada, J. Surface Oxidation, Size and Shape of Nano-Sized Magnetite Obtained by Co-Precipitation. *J. Magn. Magn. Mater.* **2006**, *300* (2), 358–367.
- (265) Shokrollahi, H. A Review of the Magnetic Properties, Synthesis Methods and Applications of Maghemite. *J. Magn. Magn. Mater.* **2017**, *426*, 74–81.
- (266) Tronc, E.; Belleville, P.; Jolivet, J. P.; Livage, J. Transformation of Ferric Hydroxide into Spinel by Iron(II) Adsorption. *Langmuir* **1992**, *8* (1), 313–319.
- (267) Benner, S. G.; Hansel, C. M.; Wielinga, B. W.; Barber, T. M.; Fendorf, S. Reductive Dissolution and Biomineralization of Iron Hydroxide under Dynamic Flow Conditions. *Environ. Sci. Technol.* **2002**, *36* (8), 1705–1711.
- (268) Fredrickson, J. K.; Zachara, J. M.; Kennedy, D. W.; Dong, H.; Onstott, T. C.; Hinman, N. W.; Li, S. Biogenic Iron Mineralization Accompanying the Dissimilatory Reduction of Hydrated Ferric Oxide by a Groundwater Bacterium. *Geochim. Cosmochim. Acta* **1998**, *62* (19–20), 3239–3257.
- (269) Glasauer, S.; Weidler, P. G.; Langley, S.; Beveridge, T. J. Controls on Fe Reduction and Mineral Formation by a Subsurface Bacterium. *Geochim. Cosmochim. Acta* **2003**, *67* (7), 1277–1288.
- (270) Hansel, C. M.; Benner, S. G.; Neiss, J.; Dohnalkova, A.; Kukkadapu, R. K.; Fendorf, S. Secondary Mineralization Pathways Induced by Dissimilatory Iron Reduction of Ferrihydrite under Advective Flow. *Geochim. Cosmochim. Acta* **2003**, *67* (16), 2977–2992.
- (271) Kumar, N.; Lezama-Pacheco, J. S.; Noël, V.; Dublet, G.; Brown, G. E. Sulfidation Mechanisms of Fe(III)-(Oxyhydr)Oxide Nanoparticles: A Spectroscopic Study. *Environ. Sci. Nano* **2018**, *5* (4), 1012–1026.
- (272) Sahai, N.; Lee, Y. J.; Xu, H.; Ciardelli, M.; Gaillard, J.-F. Role of Fe(II) and Phosphate in Arsenic Uptake by Coprecipitation. *Geochim. Cosmochim. Acta* **2007**, *71* (13), 3193–3210.
- (273) Hofacker, A. F.; Voegelin, A.; Kaegi, R.; Weber, F. A.; Kretzschmar, R. Temperature-Dependent Formation of Metallic Copper and Metal Sulfide Nanoparticles during Flooding of a Contaminated Soil. *Geochim. Cosmochim. Acta* **2013**, *103*, 316–332.
- (274) Hockmann, K.; Planer-Friedrich, B.; Johnston, S. G.; Peiffer, S.; Burton, E. D. Antimony Mobility in Sulfidic Systems: Coupling with Sulfide-Induced Iron Oxide Transformations. *Geochim. Cosmochim. Acta* **2020**, *282*, 276–296.
- (275) Jacobs, L.; Emerson, S.; Skei, J. Partitioning and Transport of Metals across the Interface in a Permanently Anoxic Basin: Framvaren Fjord, Norway. *Geochim. Cosmochim. Acta* **1985**, *49* (6), 1433–1444.
- (276) Rozan, T. F.; Benoit, G. Geochemical Factors Controlling Free Cu Ion Concentrations in River Water. *Geochim. Cosmochim. Acta* **1999**, *63* (19–20), 3311–3319.
- (277) Landing, W. M.; Lewis, B. L. Thermodynamic Modeling of Trace Metal Speciation in the Black Sea. In *Black Sea Oceanography*; Springer Netherlands: 1991; pp 125–160. DOI: 10.1007/978-94-011-2608-3_8.
- (278) Radford-Knery, J.; Cutter, G. A. Biogeochemistry of Dissolved Hydrogen Sulfide Species and Carbonyl Sulfide in the Western North Atlantic Ocean. *Geochim. Cosmochim. Acta* **1994**, *58* (24), 5421–5431.
- (279) Rozan, T. F.; Benoit, G.; Luther, G. W. Measuring Metal Sulfide Complexes in Oxidic River Waters with Square Wave Voltammetry. *Environ. Sci. Technol.* **1999**, *33* (17), 3021–3026.
- (280) Lewis, B. L.; Luther, G. W.; Lane, H.; Church, T. M. Determination of Metal-Organic Complexation in Natural Waters by SWASV with Pseudopolarograms. *Electroanalysis* **1995**, *7* (2), 166–177.
- (281) Croot, P. L.; Moffett, J. W.; Luther, G. W. Polarographic Determination of Half-Wave Potentials for Copper-Organic Complexes in Seawater. *Mar. Chem.* **1999**, *67* (3–4), 219–232.
- (282) Rozan, T. F.; Luther, G. W.; Ridge, D.; Robinson, S. Determination of Pb Complexation in Oxidic and Sulfidic Waters Using Pseudovoltammetry. *Environ. Sci. Technol.* **2003**, *37* (17), 3845–3852.
- (283) Luther, G. W.; Tsamakis, E. Concentration and Form of Dissolved Sulfide in the Oxidic Water Column of the Ocean. *Mar. Chem.* **1989**, *27* (3–4), 165–177.
- (284) Kuwabara, J. S.; Luther, G. W. Dissolved Sulfides in the Oxidic Water Column of San Francisco Bay, California. *Estuaries* **1993**, *16* (3), 567.
- (285) Luther, G. W.; Ferdeman, T. G. Voltammetric Characterization of Iron(II) Sulfide Complexes in Laboratory Solutions and in Marine Waters and Porewaters. *Environ. Sci. Technol.* **1993**, *27* (6), 1154–1163.
- (286) Rozan, T. F.; Luther, G. W. Voltammetric Evidence Suggesting Ag Speciation Is Dominated by Sulfide Complexation in River Water. *ACS Symp. Ser.* **2002**, *811*, 371–387.
- (287) Brendel, P. J.; Luther, G. W. I. Development of a Gold Amalgam Voltammetric Microelectrode for the Determination of Dissolved Fe, Mn, O₂, and S(-II) in Porewaters of Marine and Freshwater Sediments. *Environ. Sci. Technol.* **1995**, *29* (3), 751–761.
- (288) Thomasarrigo, L. K.; Bouchet, S.; Kaegi, R.; Kretzschmar, R. Organic Matter Influences Transformation Products of Ferrihydrite Exposed to Sulfide. *Environ. Sci. Nano* **2020**, *7* (11), 3405–3418.
- (289) Sharma, P.; Ofner, J.; Kappler, A. Formation of Binary and Ternary Colloids and Dissolved Complexes of Organic Matter, Fe and As. *Environ. Sci. Technol.* **2010**, *44* (12), 4479–4485.
- (290) Kaegi, R.; Voegelin, A.; Folini, D.; Hug, S. J. Effect of Phosphate, Silicate, and Ca on the Morphology, Structure and Elemental Composition of Fe(III)-Precipitates Formed in Aerated Fe(II) and As(III) Containing Water. *Geochim. Cosmochim. Acta* **2010**, *74* (20), 5798–5816.
- (291) Moens, C.; Montalvo, D.; Smolders, E. The Concentration and Size Distribution of Iron-rich Colloids in Pore Waters Are Related to Soil Organic Matter Content and Pore Water Calcium Concentration. *Eur. J. Soil Sci.* **2021**, *72* (5), 2199–2214.
- (292) Stolpe, B.; Guo, L.; Shiller, A. M. Binding and Transport of Rare Earth Elements by Organic and Iron-Rich Nanocolloids in Alaskan Rivers, as Revealed by Field-Flow Fractionation and ICP-MS. *Geochim. Cosmochim. Acta* **2013**, *106*, 446–462.
- (293) Giannetta, B.; Balint, R.; Said-Pullicino, D.; Plaza, C.; Martin, M.; Zaccone, C. Fe(II)-Catalyzed Transformation of Fe (Oxyhydr)-Oxides across Organic Matter Fractions in Organically Amended Soils. *Sci. Total Environ.* **2020**, *748*, 141125.
- (294) Cismasu, A. C.; Levard, C.; Michel, F. M.; Brown, G. E. Properties of Impurity-Bearing Ferrihydrite II: Insights into the

Surface Structure and Composition of Pure, Al- and Si-Bearing Ferrihydrite from Zn(II) Sorption Experiments and Zn K-Edge X-Ray Absorption Spectroscopy. *Geochim. Cosmochim. Acta* **2013**, *119*, 46–60.

(295) Thomasarrigo, L. K.; Byrne, J. M.; Kappler, A.; Kretzschmar, R. Impact of Organic Matter on Iron(II)-Catalyzed Mineral Transformations in Ferrihydrite-Organic Matter Coprecipitates. *Environ. Sci. Technol.* **2018**, *52* (21), 12316–12326.

(296) Pédrot, M.; Boudec, A. Le; Davranche, M.; Dia, A.; Henin, O. How Does Organic Matter Constrain the Nature, Size and Availability of Fe Nanoparticles for Biological Reduction? *J. Colloid Interface Sci.* **2011**, *359* (1), 75–85.

(297) Zhou, Z.; Latta, D. E.; Noor, N.; Thompson, A.; Borch, T.; Scherer, M. M. Fe(II)-Catalyzed Transformation of Organic Matter-Ferrihydrite Coprecipitates: A Closer Look Using Fe Isotopes. *Environ. Sci. Technol.* **2018**, *52* (19), 11142–11150.

(298) Thomasarrigo, L. K.; Kaegi, R.; Kretzschmar, R. Ferrihydrite Growth and Transformation in the Presence of Ferrous Iron and Model Organic Ligands. *Environ. Sci. Technol.* **2019**, *53* (23), 13636–13647.

(299) ThomasArrigo, L. K.; Kretzschmar, R. Iron Speciation Changes and Mobilization of Colloids during Redox Cycling in Fe-Rich, Icelandic Peat Soils. *Geoderma* **2022**, *428*, 116217.

(300) Borch, T.; Kretzschmar, R.; Skappler, A.; Van Cappellen, P.; Ginder-Vogel, M.; Voegelin, A.; Campbell, K. Biogeochemical Redox Processes and Their Impact on Contaminant Dynamics. *Environ. Sci. Technol.* **2010**, *44* (1), 15–23.

(301) Dreissig, I.; Weiss, S.; Hennig, C.; Bernhard, G.; Zänker, H. Formation of Uranium(IV)-Silica Colloids at near-Neutral PH. *Geochim. Cosmochim. Acta* **2011**, *75* (2), 352–367.

(302) Suzuki, Y.; Kelly, S. D.; Kemner, K. M.; Banfield, J. F. Nanometre-Size Products of Uranium Bioreduction. *Nature* **2002**, *419* (6903), 134–134.

(303) Gustafsson, J. P.; Persson, I.; Oromieh, A. G.; van Schaik, J. W. J.; Sjöstedt, C.; Kleja, D. B. Chromium(III) Complexation to Natural Organic Matter: Mechanisms and Modeling. *Environ. Sci. Technol.* **2014**, *48* (3), 1753–1761.

(304) James, B. R.; Bartlett, R. J. Behavior of Chromium in Soils: V. Fate of Organically Complexed Cr(III) Added to Soil. *J. Environ. Qual.* **1983**, *12* (2), 169–172.

(305) James, B. R.; Bartlett, R. J. Behavior of Chromium in Soils. VI. Interactions Between Oxidation-Reduction and Organic Complexation. *J. Environ. Qual.* **1983**, *12* (2), 173–176.

(306) Blankinship, J. C.; Berhe, A. A.; Crow, S. E.; Druhan, J. L.; Heckman, K. A.; Keiluweit, M.; Lawrence, C. R.; Marin-Spiotta, E.; Plante, A. F.; Rasmussen, C.; Schädel, C.; Schimel, J. P.; Sierra, C. A.; Thompson, A.; Wagai, R.; Wieder, W. R. Improving Understanding of Soil Organic Matter Dynamics by Triangulating Theories, Measurements, and Models. *Biogeochemistry* **2018**, *140* (1), 1–13.

(307) Cuss, C. W.; Glover, C. N.; Javed, M. B.; Nagel, A.; Shoty, W. Geochemical and Biological Controls on the Ecological Relevance of Total, Dissolved, and Colloidal Forms of Trace Elements in Large Boreal Rivers: Review and Case Studies. *Environ. Rev.* **2020**, *28* (2), 138–163.

(308) Wu, J.; Boyle, E.; Sunda, W.; Wen, L.-S. Soluble and Colloidal Iron in the Oligotrophic North Atlantic and North Pacific. *Science* (80-.). **2001**, *293* (5531), 847–849.

(309) Lacroix, E. M.; Aeppli, M.; Boye, K.; Brodie, E.; Fendorf, S.; Keiluweit, M.; Naughton, H.; Noël, V.; Sihi, D. Consider the Anoxic Microsite: Acknowledging and Appreciating Spatiotemporal Redox Heterogeneity in Soils and Sediments. *ACS Earth Sp. Chem.* **2023**, *7*, 1592.

(310) Lacroix, E. M.; Aeppli, M.; Boye, K.; Brodie, E.; Fendorf, S.; Keiluweit, M.; Naughton, H. R.; Noël, V.; Sihi, D. Consider the Anoxic Microsite: Acknowledging and Appreciating Spatiotemporal Redox Heterogeneity in Soils and Sediments. *ACS Earth Sp. Chem.* **2023**, *7* (9), 1592–1609.

(311) Chen, Y.-W.; Buffle, J. Physicochemical and Microbial Preservation of Colloid Characteristics of Natural Water Samples. I: Experimental Conditions. *Water Res.* **1996**, *30* (9), 2178–2184.

(312) Chen, Y.-W.; Buffle, J. Physicochemical and Microbial Preservation of Colloid Characteristics of Natural Water Samples. II: Physicochemical and Microbial Evolution. *Water Res.* **1996**, *30* (9), 2185–2192.

(313) Hassellöv, M.; Buesseler, K. O.; Pike, S. M.; Dai, M. Application of Cross-Flow Ultrafiltration for the Determination of Colloidal Abundances in Suboxic Ferrous-Rich Ground Waters. *Sci. Total Environ.* **2007**, *372* (2–3), 636–644.

(314) Dai, M.; Kelley, J. M.; Buesseler, K. O. Sources and Migration of Plutonium in Groundwater at the Savannah River Site. *Environ. Sci. Technol.* **2002**, *36* (17), 3690–3699.

(315) Kaplan, D. I.; Bertsch, P. M.; Adriano, D. C.; Orlandini, K. A. Actinide Association with Groundwater Colloids in a Coastal Plain Aquifer. *Radiochim. Acta* **1994**, *66–67* (s1), 181–188.

(316) Deville, S. *Freezing Colloids: Observations, Principles, Control, and Use*; Springer International: 2017; Engineering Materials and Processes.

(317) Hofmann, A.; Pelletier, M.; Michot, L.; Stradner, A.; Schurtenberger, P.; Kretzschmar, R. Characterization of the Pores in Hydrous Ferric Oxide Aggregates Formed by Freezing and Thawing. *J. Colloid Interface Sci.* **2004**, *271* (1), 163–173.

(318) Mikutta, C.; Mikutta, R.; Bonneville, S.; Wagner, F.; Voegelin, A.; Christl, I.; Kretzschmar, R. Synthetic Coprecipitates of Exopolysaccharides and Ferrihydrite. Part I: Characterization. *Geochim. Cosmochim. Acta* **2008**, *72* (4), 1111–1127.

(319) Murase, N.; Franks, F. Salt Precipitation during the Freeze-Concentration of Phosphate Buffer Solutions. *Biophys. Chem.* **1989**, *34* (3), 293–300.

(320) Ilett, M.; Matar, O.; Bamiduro, F.; Sanchez-Segado, S.; Brydson, R.; Brown, A.; Hondow, N. Nanoparticle Corona Artefacts Derived from Specimen Preparation of Particle Suspensions. *Sci. Rep.* **2020**, *10* (1), 5278.

(321) Strathmann, M.; Leon-Morales, C. F.; Flemming, H.-C. Influence of Biofilms on Colloid Mobility in the Subsurface. In *Colloidal Transport in Porous Media*; Springer Berlin Heidelberg: 2007; pp 143–173. DOI: 10.1007/978-3-540-71339-5_6.

(322) Powelson, D. K.; Gerba, C. P. Virus Removal from Sewage Effluents during Saturated and Unsaturated Flow through Soil Columns. *Water Res.* **1994**, *28* (10), 2175–2181.

(323) McCarthy, J. F.; McKay, L. D. Colloid Transport in the Subsurface: Past, Present, and Future Challenges. *Vadose Zo. J.* **2004**, *3* (2), 326–337.

(324) Keller, A. A.; Sirivithayapakorn, S. Transport of Colloids in Unsaturated Porous Media: Explaining Large-Scale Behavior Based on Pore-Scale Mechanisms. *Water Resour. Res.* **2004**, *40* (12), 1.

(325) Elimelech, M.; Chen, J. Y.; Kuznar, Z. A. Particle Deposition onto Solid Surfaces with Micropatterned Charge Heterogeneity: The “Hydrodynamic Bump” Effect. *Langmuir* **2003**, *19* (17), 6594–6597.

(326) Elimelech, M.; Nagai, M.; Ko, C.-H.; Ryan, J. N. Relative Insignificance of Mineral Grain Zeta Potential to Colloid Transport in Geochemically Heterogeneous Porous Media. *Environ. Sci. Technol.* **2000**, *34* (11), 2143–2148.

(327) Lenhart, J. J.; Saiers, J. E. Colloid Mobilization in Water-Saturated Porous Media under Transient Chemical Conditions. *Environ. Sci. Technol.* **2003**, *37* (12), 2780–2787.

(328) Yao, K.-M.; Habibian, M. T.; O’Melia, C. R. Water and Waste Water Filtration. Concepts and Applications. *Environ. Sci. Technol.* **1971**, *5* (11), 1105–1112.

(329) Harvey, R. W. Parameters involved in Modeling Movement of Bacteria in Groundwater. In *Modeling the Environmental Fate of Microorganisms*; American Society for Microbiology, 1991; 89–114.

(330) Johnson, P. R.; Sun, N.; Elimelech, M. Colloid Transport in Geochemically Heterogeneous Porous Media: Modeling and Measurements. *Environ. Sci. Technol.* **1996**, *30* (11), 3284–3293.

(331) Verwey, E. J. W. Theory of the Stability of Lyophobic Colloids. *J. Phys. Colloid Chem.* **1947**, *51* (3), 631–636.

- (332) Derjaguin, B.; Landau, L. D. Theory of the Stability of Strongly Charged Lyophobic Sols and of the Adhesion of Strongly Charged Particles in Solutions of Electrolytes. *Acta Physicochim. U.R.S.S.* **1941**, *14*, 633–662.
- (333) Cornelis, G. Fate Descriptors for Engineered Nanoparticles: The Good, the Bad, and the Ugly. *Environ. Sci. Nano* **2015**, *2* (1), 19–26.
- (334) Tufenkji, N.; Elimelech, M. Correlation Equation for Predicting Single-Collector Efficiency in Physicochemical Filtration in Saturated Porous Media. *Environ. Sci. Technol.* **2004**, *38* (2), 529–536.
- (335) Bradford, S. A.; Bettahar, M. Concentration Dependent Transport of Colloids in Saturated Porous Media. *J. Contam. Hydrol.* **2006**, *82* (1–2), 99–117.
- (336) Bradford, S. A.; Bettahar, M. Straining, Attachment, and Detachment of Cryptosporidium Oocysts in Saturated Porous Media. *J. Environ. Qual.* **2005**, *34* (2), 469–478.
- (337) Xu, S.; Gao, B.; Saiers, J. E. Straining of Colloidal Particles in Saturated Porous Media. *Water Resour. Res.* **2006**, *42* (12), 1.
- (338) Corapcioglu, M. Y.; Choi, H. Modeling Colloid Transport in Unsaturated Porous Media and Validation with Laboratory Column Data. *Water Resour. Res.* **1996**, *32* (12), 3437–3449.
- (339) Bradford, S. A.; Simunek, J.; Bettahar, M.; Tadassa, Y. F.; van Genuchten, M. T.; Yates, S. R. Straining of Colloids at Textural Interfaces. *Water Resour. Res.* **2005**, *41* (10), 1.
- (340) Gargiulo, G.; Bradford, S. A.; Simunek, J.; Ustohal, P.; Vereecken, H.; Klumpp, E. Bacteria Transport and Deposition under Unsaturated Flow Conditions: The Role of Water Content and Bacteria Surface Hydrophobicity. *Vadose Zo. J.* **2008**, *7* (2), 406–419.
- (341) Saiers, J. E.; Lenhart, J. J. Ionic-Strength Effects on Colloid Transport and Interfacial Reactions in Partially Saturated Porous Media. *Water Resour. Res.* **2003**, *39* (9), 1.
- (342) Sen, T. K.; Nalwaya, N.; Khilar, K. C. Colloid-Associated Contaminant Transport in Porous Media: 2. Mathematical Modeling. *AIChE J.* **2002**, *48* (10), 2375–2385.
- (343) Sen, T. K.; Shanbhag, S.; Khilar, K. C. Subsurface Colloids in Groundwater Contamination: A Mathematical Model. *Colloids Surfaces A Physicochem. Eng. Asp.* **2004**, *232* (1), 29–38.
- (344) Shang, J.; Flury, M.; Chen, G.; Zhuang, J. Impact of Flow Rate, Water Content, and Capillary Forces on in Situ Colloid Mobilization during Infiltration in Unsaturated Sediments. *Water Resour. Res.* **2008**, *44* (6), 1.
- (345) Wan, J.; Tokunaga, T. K. Film Straining of Colloids in Unsaturated Porous Media: Conceptual Model and Experimental Testing. *Environ. Sci. Technol.* **1997**, *31* (8), 2413–2420.
- (346) Dąbroś, T.; Van De Ven, T. G. Kinetics of Coating by Colloidal Particles. *J. Colloid Interface Sci.* **1982**, *89* (1), 232–244.
- (347) Corapcioglu, M. Y.; Jiang, S. Colloid-Facilitated Groundwater Contaminant Transport. *Water Resour. Res.* **1993**, *29* (7), 2215–2226.
- (348) van de Weerd, H.; Leijnse, A.; van Riemsdijk, W. H. Transport of Reactive Colloids and Contaminants in Groundwater: Effect of Nonlinear Kinetic Interactions. *J. Contam. Hydrol.* **1998**, *32* (3–4), 313–331.
- (349) Schijven, J. F.; Hassanizadeh, S. M.; de Bruin, R. H. A. M. Two-Site Kinetic Modeling of Bacteriophages Transport through Columns of Saturated Dune Sand. *J. Contam. Hydrol.* **2002**, *57* (3–4), 259–279.
- (350) Dwivedi, D.; Mohanty, B. P.; Lesikar, B. J. Impact of the Linked Surface Water-Soil Water-Groundwater System on Transport of E. Coli in the Subsurface. *Water, Air, Soil Pollut.* **2016**, *227* (9), 351.
- (351) Knabe, D.; Dwivedi, D.; Wang, H.; Griebler, C.; Engelhardt, I. Numerical Investigations to Identify Environmental Factors for Field-Scale Reactive Transport of Pathogens at Riverbank Filtration Sites. *Adv. Water Resour.* **2023**, *173*, 104389.
- (352) Ryan, J. N.; Elimelech, M. Colloid Mobilization and Transport in Groundwater. *Colloids Surfaces A Physicochem. Eng. Asp.* **1996**, *107*, 1–56.
- (353) Roy, S. B.; Dzombak, D. A. Chemical Factors Influencing Colloid-Facilitated Transport of Contaminants in Porous Media. *Environ. Sci. Technol.* **1997**, *31* (3), 656–664.
- (354) Johnson, P. R.; Elimelech, M. Dynamics of Colloid Deposition in Porous Media: Blocking Based on Random Sequential Adsorption. *Langmuir* **1995**, *11* (3), 801–812.
- (355) Roy, S. B.; Dzombak, D. A. Colloid Release and Transport Processes in Natural and Model Porous Media. *Colloids Surfaces A Physicochem. Eng. Asp.* **1996**, *107*, 245–262.
- (356) Boström, M.; Williams, D. R. M.; Ninham, B. W. Specific Ion Effects: Why DLVO Theory Fails for Biology and Colloid Systems. *Phys. Rev. Lett.* **2001**, *87* (16), 168103.
- (357) Hermansson, M. The DLVO Theory in Microbial Adhesion. *Colloids Surfaces B Biointerfaces* **1999**, *14* (1–4), 105–119.
- (358) Ginn, T. R. On the Distribution of Multicomponent Mixtures over Generalized Exposure Time in Subsurface Flow and Reactive Transport: Theory and Formulations for Residence-Time-Dependent Sorption/Desorption with Memory. *Water Resour. Res.* **2000**, *36* (10), 2885–2893.
- (359) Massoudieh, A.; Ginn, T. R. Colloid-Facilitated Contaminant Transport in Unsaturated Porous Media. *Modelling of Pollutants in Complex Environmental Systems* **2010**, *II*, 259–287.
- (360) Powelson, D. K.; Mills, A. L. Transport of Escherichia Coli in Sand Columns with Constant and Changing Water Contents. *J. Environ. Qual.* **2001**, *30* (1), 238–245.
- (361) Palacin, S.; Barraud, A. Chemical Reactivity at the Air–Water Interface: Redox Properties of the Tetrapyrridino Porphyrinium Ring. *Colloids Surf.* **1991**, *52*, 123–147.
- (362) Wan, J.; Wilson, J. L. Colloid Transport in Unsaturated Porous Media. *Water Resour. Res.* **1994**, *30* (4), 857–864.
- (363) Crist, J. T.; McCarthy, J. F.; Zevi, Y.; Baveye, P.; Throop, J. A.; Steenhuis, T. S. Pore-Scale Visualization of Colloid Transport and Retention in Partly Saturated Porous Media. *Vadose Zo. J.* **2004**, *3* (2), 444–450.
- (364) Crist, J. T.; Zevi, Y.; McCarthy, J. F.; Throop, J. A.; Steenhuis, T. S. Transport and Retention Mechanisms of Colloids in Partially Saturated Porous Media. *Vadose Zo. J.* **2005**, *4* (1), 184.
- (365) Arah, J. R. M.; Vinten, A. J. A. Simplified Models of Anoxia and Denitrification in Aggregated and Simple-Structured Soils. *Eur. J. Soil Sci.* **1995**, *46* (4), 507–517.
- (366) Riley, K. W.; Matson, P. A. NLOSS: A Mechanistic Model of Denitrified N₂O and N₂ Evolution from Soil. *Soil Sci.* **2000**, *165* (3), 237–249.
- (367) Davidson, E. A. Pulses of Nitric Oxide and Nitrous Oxide Flux Following Wetting of Dry Soil: An Assessment of Probable Sources and Importance Relative to Annual Fluxes. *Ecol. Bull.* **1992**, *42*, 149–155.
- (368) Engel, M.; Boye, K.; Noël, V.; Babey, T.; Bargar, J. R.; Fendorf, S. Simulated Aquifer Heterogeneity Leads to Enhanced Attenuation and Multiple Retention Processes of Zinc. *Environ. Sci. Technol.* **2021**, *55* (5), 2939–2948.
- (369) Engel, M.; Noël, V.; Kukkadapu, R. K.; Boye, K.; Bargar, J. R.; Fendorf, S. Nitrate Controls on the Extent and Type of Metal Retention in Fine-Grained Sediments of a Simulated Aquifer. *Environ. Sci. Technol.* **2022**, *56* (20), 14452–14461.
- (370) Schulze, E.-D.; Chapin, F. S.; Gebauer, G. Nitrogen Nutrition and Isotope Differences among Life Forms at the Northern Treeline of Alaska. *Oecologia* **1994**, *100* (4), 406–412.