

Diary of a wimpy fault

- [Roland Bürgmann](#)

Nature Geoscience **volume8**, pages331–332 (2015) | [Download Citation](#)

Subduction zone faults can slip slowly, generating tremor. The varying correlation between tidal stresses and tremor occurring deep in the Cascadia subduction zone suggests that the fault is inherently weak, and gets weaker as it slips.


The discovery of tectonic tremor in the deep roots of plate-boundary subduction zones in Japan and North America was a great surprise. These unusual and low-amplitude seismic signals emanate from a depth in the Earth where rocks should be too hot and too viscous to produce seismic events. The tremors were found to be caused by episodic slow-slip events on the deep sections of plate-boundary faults, well below the parts of the fault that produce great earthquakes. Tremors can be triggered and modulated by the small stresses associated with body and ocean tides, implying that the faults that generate them must be very weak. Writing in *Nature Geoscience*, Houston¹ analyses tremor events in the Cascadia subduction zone between 2007 and 2012. She shows that not only are the deep parts of the fault very weak, but that the fault becomes weaker during individual slip events.

Tremors are the weak cousins of regular earthquakes. While it took decades of painstaking work to document the modest effect of tides on regular earthquakes, tremors were found to be easily triggered by these transient stresses². This extreme sensitivity to very small stress changes indicates that the tremor-producing fault is frictionally weak, in large part due to fluids in the fault zone being at near-lithostatic pressure. That is, the overburden of rock is counteracted by fluid pressures of comparable magnitude in the fault. The high fluid pressure leads to a very low effective frictional strength of the fault, and slip and tremor can readily respond even to tiny tidal shear stresses of less than 100 Pa (ref. [3](#)).

Houston¹ explores in detail the occurrence of tens of thousands of tremors created by six slow-slip events in the Cascadia subduction zone between 2007 and 2012 ([Fig. 1](#)). Each slow-slip event took several weeks to complete, with the

slip front propagating at rates of several kilometres per day along a nearly 200-km-long stretch of the fault. At any given point on the fault, tremors and slip continue for several days but accumulate only two to three centimetres of total offset during each slip event⁴. Houston finds that not only is tremor (and thus fault slip) modulated by tides, but the strength of the correlation between the tides and tremors increases as slow slip progresses. This implies that the fault becomes increasingly sensitive to tides and progressively weakens as slip reaches a total of a few centimetres. So, not only are tremor-producing faults easily pushed around by external forces, they continue to weaken as they go.

Figure 1: Recurring slip events deep on the Cascadia subduction zone are illuminated by tremors.



The tremor locations (circles) are coloured by date of occurrence during a month-long tremor-and-slip episode in 2010. Houston₁ finds that the correlation between tremor and tidal stress at a point on the

fault strengthens during a slip event, implying that the fault weakens as slip evolves. At depths of less than about 20 km (purple shaded region), the megathrust appears to be locked and therefore builds up stress towards the next great subduction earthquake. Figure courtesy of Aaron Wech, US Geological Survey.

[Full size image](#)

Tremors have a story to tell about how deep faults work. The complex distribution of tremors in space and time and their response to external stresses illuminate the mechanical properties of the deep fault⁵⁶. Some frictional models of slow-slip events feature slip-weakening distances that are large compared with those found in laboratory experiments (10^{-5} to 10^{-6} m); however, there are a number of alternative ways in which models can be parameterized to produce such events⁶⁷. The observation of a slowly increasing correlation of tides and tremors during the Cascadia slow-slip events indicates that the fault may indeed need to slip a relatively large distance, on the scale of a centimetre, before it weakens to its frictional sliding strength.

Houston suggests that the observed drop in fault strength over the course of a slow-slip event could involve the breakage of mineral precipitates in the fault zone. The minerals rapidly regrow in the approximately 14-month-long recurrence interval between slow-slip events, during which time the fault strength recovers and stress accumulates⁸. Interestingly, a transition from tremor modulated by the rate of peak tidal shear stress to tremor modulated by the amplitude of peak tidal shear stress during a large slow slip event has recently been documented in Cascadia⁹. This observation adds another twist to the relationship of tides and tremors and should further illuminate the underlying physics of the slow-slip process.

Tremor-producing faults are weak, but it was not known if the weakness stems only from very high fluid pressures or if the fault-zone rocks also have a very low friction coefficient. Taking advantage of differences in the temporal variations of tidal fault-perpendicular stress and mean stress (pressure), Houston is able to estimate the intrinsic, fluid-pressure-independent friction coefficient for the fault *in situ*. She finds values for the average friction coefficient well below 0.2 — much lower than most known rock types, with friction values between 0.6 and 0.8. Such low values require unusually weak fault zone materials, such as talc, graphite and saponite, to effectively lubricate

the fault zone. Saponite breaks down at temperatures well below those found in the tremor zone, but talc and graphite can be stable to very high temperatures and pressures¹⁰, so could exist in the deep fault zone beneath the Cascades.

Above the weak tremor zone in Cascadia, the seismogenic part of the fault is strongly coupled and capable of producing great megathrust ruptures. The last such event in 1700 was a not-so-wimpy magnitude 9 earthquake. Given that the Cascades are home to large cities such as Seattle, Portland and Vancouver, we must continue to investigate potential links between slip on the deep subduction zone and megathrust earthquakes on locked parts of the fault. In the process, we are bound to learn more about the physics of deep plate-boundary faulting with improved and targeted seismologic, geodetic, gravity, and electro-magnetic observations, including those from space-borne systems, borehole sensors, and seafloor instrumentation. As large earthquakes often initiate close to the tremor zone, insights gained from these observations may enable improved characterization of time-dependent earthquake hazard and of precursory processes that appear to precede some large earthquake ruptures.

References

1. 1.
Houston, H. *Nature Geosci.* **8**, 409–415 (2015).
o Show context for reference 1
o [Google Scholar](#)
2. 2.
Rubinstein, J. L., La Rocca, M., Vidale, J. E., Creager, K. C. & Wech, A. G. *Science* **319**, 186–189 (2008).
o Show context for reference 2
o [CAS](#)
o [PubMed](#)
o [Article](#)

- [Google Scholar](#)

3. 3.

Thomas, A. M., Nadeau, R. M. & Bürgmann, R. *Nature* **462**, 1048–1051 (2009).

o Show context for reference 3

o

- [CAS](#)
- [PubMed](#)
- [Article](#)
- [Google Scholar](#)

4. 4.

Wech, A. G., Creager, K. C. & Melbourne, T. I. *J. Geophys. Res.* **114**, B10316 (2009).

o Show context for reference 4

o

- [Article](#)
- [Google Scholar](#)

5. 5.

Beeler, N. M., Thomas, A., Bürgmann, R. & Shelly, D. *J. Geophys. Res.* **118**, 5976–5990 (2013).

o Show context for reference 5

o

- [Article](#)
- [Google Scholar](#)

6. 6.

Hawthorne, J. C. & Rubin, A. *J. Geophys. R.* **118**, 1216–1239 (2013).

o Show context for reference 6

o

- [Article](#)
- [Google Scholar](#)

7. 7.

Liu, Y. & Rice, J. R. *J. Geophys. Res.* **112**, 1978–2012 (2007).

o Show contextfor reference 7

o

▪ [Google Scholar](#)

8. 8.

Audet, P. & Bürgmann, R. *Nature* **510**, 389–392 (2014).

o Show contextfor reference 8

o

▪ [CAS](#)

▪ [PubMed](#)

▪ [Article](#)

▪ [Google Scholar](#)

9. 9.

Royer, A. A., Thomas, A. M. & Bostock, M. G. *J. Geophys. R.* **120**, 384–405 (2015).

o Show contextfor reference 9

o

▪ [Article](#)

▪ [Google Scholar](#)

10. 10.

Moore, D. E. & Rymer, M. J. *Nature* **448**, 795–797 (2007).

o Show contextfor reference 10

o

▪ [CAS](#)

▪ [PubMed](#)

▪ [Article](#)

▪ [Google Scholar](#)

[Download references](#)

Author information

Affiliations

1. *Roland Bürgmann is in the Department of Earth and Planetary Science, University of California, Berkeley, California 94720–4767, USA*

- Roland Bürgmann

Corresponding author

Correspondence to [Roland Bürgmann](#).