Lawrence Berkeley National Laboratory

Recent Work

Title

TERPENES FROM PITTOSPORACEAE

Permalink

<https://escholarship.org/uc/item/449932k8>

Authors

Nemethy, E.K.
Calvin, M.

Publication Date

1982-03-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

CHEMICAL BIODYNAMICS DIVISION

Submitted to Phytochemistry

TERPENES FROM PITTOSPORACEAE

Esther K. Nemethy and Melvin Calvin

March 1982

RECEIVED
LAWRENCE
BERKELEY LABORATORY

APR 5 1982

LIBRARY AND
DOCUMENTS SECTION

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782*

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

TERPENES FROM PITTOSPORACEAE

Esther K. Nemethy and Melvin Calvin

Lawrence Berkeley Laboratory & Department of Chemistry

University of California, Berkeley, California 94720

Key word index: Pittosporum resiniferum, Pittosporum undulatum, essential oil, monoterpenes, α -pinene; myrcene, limonene.

Abstract: The monoterpene constituents of the fruits of Pittosporum resiniferum and Pittosporum undulatum have been identified. The essential oil of Pittosporum resiniferum contains myrcene and α -pinene in equal quantities, whereas the major terpene constituent of Pittosporum undulatum oil is limonene.

This work was supported by the Assistant Secretary for Conservation and Solar Energy, Office of Solar Applications for Industry of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

The fruits of the tree Pittosporum resiniferum (family Pittosporaceae) have been reported to contain the unusual natural product n-heptane (Fernandez, E., personal communication). It is reported that even the green fresh fruit will burn brilliantly when a match is applied to it and that the expressed oil of the fruit is flammable. This species is native to the Philippines where the nuts are sometimes used as torch light [1]. Pittosporum fruit, therefore, shows promise as a potential alternate liquid energy source, and we have examined the chemical composition of this oil. The two alkanes n-heptane and n-nonane are minor components of this oil; the major components are two volatile monoterpenes, α -pinene and myrcene.

Another species of this family, Pittosporum undulatum, is widespread in California, and it produces small fragrant orange fruits in abundance. P. undulatum fruits were collected on the University of California Berkeley Campus during November and December 1981. The n-alkanes could not be detected in the oil of P. undulatum. Limonene is the only major monoterpene present in this oil, accounting for 99% of the monoterpene fraction; α -pinene is the minor component. No oxygenated monoterpenes were detected in either of these oils, but both contained minor quantities of sesquiterpene hydrocarbons.

A few members of the family Pittosporaceae have been subject to phytochemical investigation. Phytosterols have been isolated from the bark of P. floribundum [2], P. colensoi, and P. eugenoides [3]. Polyacetylenic compounds have been isolated from roots of P. buechanani [4], P. crassifolium, P. tobira and P. undulatum [5]. However, there are no prior reports on the essential oils of any Pittosporum species.

Experimental:

Capillary GC analyses were obtained using either a 40 m SP2250 SCOT, an 80 m Carbowax 20M column, temp. programmed from 40⁰ to 200⁰ at 4⁰/min. Where indicated, percentages refer to computer-calculated peak areas without correction to response. Preparatory GC was carried out on a 10' 4 mm ID 10% SP2250 glass column at 110⁰. Coupled GC/MS were carried out on the SP2250 column interfaced to a Finnigan 4000 mass spectrometer at 70 eV. ¹H NMR spectra were recorded at 250 MHz in CDCl₃ with TMS as internal standard.

Analysis of *P. resiniferum* essential oil. Ripe fruits were homogenized in a Waring blender, then extracted with CH₂Cl₂ by stirring under N₂ at ambient temperature for 24 hrs. After slow distillation of the CH₂Cl₂ a light orange oil, 8-10% of the fr.wt. was obtained. GC/MS analysis of the oil indicated the presence of n-heptane (5%)*, n-nonane (7%), three isomeric monoterpenes (85%) and six minor sesquiterpenes (6%). The two major monoterpenes were isolated by preparative GC and identified as α-pinene (38%) and myrcene (40%) by ¹H NMR, MS and coelution with standards on the SP2250 and Carbowax 20M capillary columns. Catalytic hydrogenation of the oil (Pd on C, 1 atm) gave pinane and 2,6-dimethyl-octane, identified by MS and GC retention time.

α-pinene: ¹H NMR (250 MHz) δ 5.17 (1H,s), 2.33 (1H,m), 2.2 (2H, br.s), 2.18(1H,br.s), 1.94 (1H,m), 1.65 (3H,s), 1.26 (3H,s), 1.5 (2H,m), 0.83 (3H,s).

MS: m/z: 136, 121, 107, 105, 93 (b.p.), 79, 77.

Myrcene: ¹H NMR (250 MHz) δ 6.37 (1H,m; J = 10.7, 17.6 Hz), 5.4-4.9 (5H), 2.2 (4H,m), 1.7 (3H,s), 1.6 (3H,s).

MS: m/z: 136, 131, 107, 94, 93 (b.p.), 69, 41.

Analysis of P. undulatum essential oil. Fresh fruits were extracted in the same manner as before, but with petroleum ether. A light orange oil, 4-5% of the fresh fruit weight, was obtained. Chromatography on silica gel, eluting with petroleum ether, gave the hydrocarbon fraction which was 50% of the oil. GC and GC/MS analyses showed the presence of two monoterpenes in a ratio of 10:1, which coeluted with limonene and α -pinene, respectively. Limonene was purified by preparative GC and identified by ^1H NMR. Catalytic hydrogenation of the oil (Pd on C, 1 atm) gave 1-isopropyl-4-methylcyclohexane and pinane, identified by MS and GC retention times.

Limonene: ^1H NMR (250 MHz) δ 5.7 (1H,s), 4.7 (2H,s), 2.2-1.67 (5H,m), 1.67 (3H,s), 1.59 (3H,s), 1.47-1.4 (2H,m).

MS: m/z: 136, 121, 107, 93, 79, 68 (b.p.).

Acknowledgment: We thank R. Hurov for the supply of Pittosporum resiniferum nuts. This work was supported by the Assistant Secretary for Conservation and Solar Energy, Office of Solar Applications for Industry, Department of Energy under Contract No. DE-AC03-76SF00098.

References:

- [1] Noble, B. (1978), Canopy, 4, 4.
- [2] Desai, H. K., Gawad, D. H., Joshi, B. S., Parthasarathy, P. D., Ravindranath, K. R., Saindane, M. T., Sidhaye, A. K., and Viswanathan, N. (1976), Ind. J. Chem. 15B, 291.
- [3] Cambrie, R., and Parnell, J. C. (1969), N. Z. J. Sci. 12, 453.
- [4] Bohlman, F., and Rode, K. (1968). Chem. Ber. 101, 1889.
- [5] Bohlman, F., and Zdero, C. (1975), Chem. Ber. 108, 2541.

* The % n-heptane may be too low, since some may have evaporated in transit.

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720