

Lawrence Berkeley National Laboratory

Recent Work

Title

THE BETA-DELAYED PROTON DECAY OF ^{61}Ge

Permalink

<https://escholarship.org/uc/item/44w5f41g>

Author

Hotchkis, M.A.C.

Publication Date

1986-08-01



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

RECEIVED
LAWRENCE
BERKELEY LABORATORY

OCT 2 1986

LIBRARY AND
DOCUMENTS SECTION

Submitted to Physical Review C


THE BETA-DELAYED PROTON DECAY OF ^{61}Ge

M.A.C. Hotchkis, J.E. Reiff, D.J. Vieira,
F. Blonnigen, T.F. Lang, D.M. Moltz, X. Xu,
and J. Cerny

August 1986

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.*



LBL-22006
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THE BETA-DELAYED PROTON DECAY OF ^{61}Ge

M. A. C. Hotchkis, J. E. Reiff, D. J. Vieira*), F. Blonnigen*),
T. F. Lang, D. M. Moltz, X. Xu and Joseph Cerny

Department of Chemistry and
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

August 1986

This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

THE BETA-DELAYED PROTON DECAY OF ^{61}Ge

M. A. C. Hotchkis, J. E. Reiff, D. J. Vieira^{*}), F. Blonnigen⁺),
T. F. Lang, D. M. Moltz, X. Xu and Joseph Cerny

Department of Chemistry and
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

August 1986

Abstract:

The $T_z = -3/2$, $A = 4n + 1$ nuclide ^{61}Ge has been produced via the $^{40}\text{Ca}(^{24}\text{Mg}, 3n)$ reaction and identified by its beta-delayed proton emission. A single proton group at 3.11 ± 0.03 MeV with $t_{1/2} = 40 \pm 15$ ms has been observed, corresponding to decay of the $T = 3/2$ isobaric analog state in ^{61}Ga to the ground state of ^{60}Zn . Searches for the beta-delayed proton decay of the next member of this series, ^{65}Se , to be produced by the $^{40}\text{Ca}(^{28}\text{Si}, 3n)$ reaction, have proven unsuccessful.

Keyword Abstract

RADIOACTIVITY ^{61}Ge from $^{40}\text{Ca}(^{24}\text{Mg}, 3n)$ and $^{40}\text{Ca}(^{28}\text{Si}, \alpha 3n)$; measured E_p , $t_{1/2}$; deduced partial decay scheme; Si surface barrier detectors.

This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

INTRODUCTION

The decays of many proton rich light nuclei have been identified by their beta-delayed proton emission. This work is facilitated by the fact that the isobaric analog state, which is well populated by the superallowed beta transition, is unbound to (isospin forbidden) proton emission. In such cases the overall delayed proton branching ratio is large, and the proton spectra are dominated by transitions from the isobaric analog state to low-lying levels in the proton daughter. All members of the $T_z = -3/2$, $A = 4n + 1$ series of nuclei from ^{17}Ne to ^{57}Zn , and all known beta-decaying $T_z = -2$ nuclei, exhibit this decay pattern. In the present work we have extended the $T_z = -3/2$, $A = 4n + 1$ series by observing the delayed proton decay of ^{61}Ge . Additionally, we have searched for the next member of this series, ^{65}Se .

Observation of the decays of these nuclei in this mass region provides a means of probing the proton drip line, whose location we have predicted by applying the Kelson-Garvey mass relation¹. This charge symmetric mass relation requires as input the masses of the $T_z = +1/2$ and $-1/2$ nuclei and the mass of the mirror of the nucleus being calculated. For $A \geq 61$, the masses of the $T_z = -1/2$ series are not known, so they have been calculated from their mirror nuclei using a formula for Coulomb displacement energies derived from a recent fit to data². These calculations indicate that the lightest particle stable isotope for each element in this region is expected to be ^{60}Ga , ^{60}Ge , ^{66}As , ^{65}Se , ^{70}Br and ^{69}Kr . Lack of mass measurements for the $T_z = +1/2$ series with $A > 70$ prevents application of this technique for higher Z elements. The energies of beta-delayed protons corresponding to the decay of the isobaric analog state in the daughter

nuclide can also be estimated in each case, by again applying the Coulomb displacement energy formula, with the calculated mass of the $T_z = -3/2$ nucleus and the known mass of the proton daughter; hence such measurements provide indirect tests of these mass predictions.

These mass calculations predict that ^{65}As , the beta daughter of ^{65}Se , should be unbound to proton emission by 570 (± 300) keV, the uncertainty being dominated by the experimental uncertainty in the mass of ^{64}Ge ; hence, all beta decays of ^{65}Se should lead to proton emission. It is also possible that ^{65}As is a ground state proton emitter with a measurable lifetime. However, its observation would be experimentally difficult since its half-life may be very short, and the proton energies would be too low for conventional $\Delta E-E$ silicon detector telescope techniques. ^{65}As is also of considerable interest for its role in the astrophysical rp process³.

EXPERIMENTAL METHOD

Beta-delayed proton emitters have been observed among the products of the $^{24}\text{Mg} + \text{Ca}$ reaction at energies from 77 to 120 MeV and the $^{28}\text{Si} + \text{Ca}$ reaction from 73 to 128 MeV. The beams were provided by the Lawrence Berkeley Laboratory 88-Inch Cyclotron. Two different methods were used to observe beta-delayed protons. One set-up employed a pulsed beam technique observing radioactive recoils stopped in the target, while the other used a helium-jet transport system. In both cases silicon detector telescopes were used to detect the activities.

In the pulsed beam experiment, the detectors were protected from radiation damage by a rotating tantalum wheel located between the target and the

detectors. The slotted wheel controlled the pulsing of the beam, and permitted the detectors to view the target through a slot in the wheel while the beam was off. The target, biased to +2.5 kV to suppress electron emission, was positioned at an angle of 20° to the beam direction. A detector telescope, which consisted of a $15 \mu\text{m}$ ΔE detector and a $250 \mu\text{m}$ E detector, was located perpendicular to the beam and detected protons emitted by products retained in the target, with an effective solid angle of 0.24 sr. A typical timing cycle consisted of an irradiation period of 85 ms with an inspection period of 15 ms (to establish beam cessation) followed by a counting period of 120 ms which was divided into eight 15 ms channels. This set-up, identical to that used to discover the beta-delayed proton emission of ^{53}Ni and ^{57}Zn ⁴, is shown in Fig. 1. This method was used with 78 and 85 MeV ^{24}Mg beams incident upon a 1.5 mg cm^{-2} natural calcium target. The beam was produced in a Penning Ion Gauge source and gave beam currents of 20 to 50 particle nA. Known peaks in the delayed proton spectra of ^{25}Si and ^{41}Ti , produced in $^3\text{He} + \text{Mg}$ and $^3\text{He} + \text{Ca}$ reactions respectively, were used to calibrate the detector telescope.

The other experiments employed a helium-jet transport system which is shown schematically in Fig. 2. Nuclei recoiling from a 1.9 mg cm^{-2} natural calcium target normal to the beam were thermalized in 1.3 atm of helium. These reaction products were transported on NaCl aerosols through a 70 cm long capillary (1.27 mm i.d.) to a shielded detector box. There they were deposited on the surface of a 15 cm diameter rotating disc, directly in front of a three-element silicon detector telescope consisting of $20 \mu\text{m}$ ΔE , $272 \mu\text{m}$ E, and a $300 \mu\text{m}$ reject detector. The ΔE -E combination permitted the identification of protons with energies ranging from 1.2 to 6.5

MeV. Calibration of the telescope was achieved using delayed protons from ^{21}Mg , ^{25}Si , ^{29}S , ^{37}Ca , ^{41}Ti and ^{45}Cr produced in the $^{14}\text{N} + \text{Ca}$ reaction at 165 MeV (all listed bombarding energies are "on-target" values). The rotating disc served a dual purpose: 1) it moved long-lived activities away from the detectors, and 2) by using two different disc speeds, qualitative information on the half-lives of the activities was obtained. The overall efficiency of this system is approximately twice that of the target counting method. The beams used in these experiments were produced in an Electron Cyclotron Resonance (ECR) source coupled to the cyclotron; typical beam currents were 100 particle nA for the ^{24}Mg beams and 100-200 particle nA for the ^{28}Si beams.

RESULTS

$^{24}\text{Mg} + \text{Ca}$

The beta-delayed proton spectrum obtained at 85 MeV from the pulsed-beam target observation technique is shown in Fig. 3. This spectrum is the result of 25 mC of integrated beam (10^+) on target. A peak is observed at $E_{\text{lab}} = 3.10 \pm 0.07$ MeV, which can be attributed to the decay of ^{61}Ge (see below). At 78 MeV this peak is also evident, with a yield two-thirds of that at 85 MeV. Its half-life was deduced to be 40 ± 15 ms using data from both bombardments; the decay curve is shown in Fig. 4. Peaks at 1.78 and 2.08 MeV, which had long half-lives, are identified as being due to the delayed proton decay of ^{59}Zn ($t_{1/2} = 210$ ms)⁵.

Figure 5 shows the delayed proton spectrum obtained with the helium-jet transport system in the 120 MeV bombardment. In addition to a peak at 3.11 MeV, delayed protons from ^{53}Co , ^{57}Zn , ^{59}Zn and ^{25}Si are evident.

The yields of the proton groups at this energy have been compared to those at 77 and 90 MeV, also measured with this system. These observations have been compared to calculations made with the statistical model fusion-evaporation code ALICE⁶. The yields of ^{53m}Co , ^{57}Zn and ^{59}Zn were found to be consistent with these calculations. ^{25}Si is believed to be produced in transfer reactions on the Havar target chamber entrance foils where the beam energy is sufficiently high (146 MeV) for such reactions to occur. The 3.11 MeV peak is attributed to the decay of ^{61}Ge , since its yield is a maximum at 90 MeV, as expected from the ALICE calculations. This peak is identified as the transition from the isobaric analog state in ^{61}Ga , fed by the superallowed beta decay of ^{61}Ge , to the ground state of ^{60}Zn . The predicted proton energy for this transition obtained from the Kelson-Garvey mass relation and the Coulomb displacement energy calculations previously mentioned is 3.21 MeV.

Since the energy of the 3.11 MeV peak observed here coincides with that of the dominant group in the delayed proton spectrum of ^{37}Ca , it has been necessary to evaluate the possibility that ^{37}Ca , produced in reactions on the small oxygen contamination in the target, could be the source of this activity. The effective threshold for the $^{16}\text{O}(^{24}\text{Mg},3n)^{37}\text{Ca}$ reaction predicted by the ALICE code is at 95 MeV; ^{37}Ca may in principle contribute to the spectrum at 120 MeV, but cannot account for the yield of the 3.11 MeV peak being a maximum at 90 MeV and being readily observed at lower energies. Also, the observed half-life of the 3.11 MeV peak is too short for ^{37}Ca ($t_{1/2} = 175$ ms). Further evidence against ^{37}Ca is provided by the data obtained with the ^{28}Si beam, discussed below, where the peak was again observed while the maximum beam energy was below threshold for the

$^{16}_0(^{28}\text{Si}, \alpha 3n)^{37}\text{Ca}$ reaction.

$^{28}\text{Si} + \text{Ca}$

The $^{28}\text{Si} + \text{Ca}$ reaction was studied at several energies to search for the delayed proton decay of ^{65}Se , the next nuclide in the $T_z = -3/2$, $A = 4n + 1$ series; these experiments also served to confirm the identity of the 3.11 MeV delayed proton peak seen in the $^{24}\text{Mg} + \text{Ca}$ data. Figure 6 shows two spectra that were obtained with the 128 MeV ^{28}Si beam. Only the speed of the rotating disc was changed between obtaining these two spectra. The figure shows how the delayed proton peak yields from $^{53\text{m}}\text{Co}$, ^{61}Ge and ^{41}Ti vary with rotation speed. The observed yield of $^{53\text{m}}\text{Co}$ ($t_{1/2} = 247$ ms) is clearly suppressed at the faster speed while those of ^{61}Ge ($t_{1/2} = 40$ ms), and ^{41}Ti ($t_{1/2} = 80$ ms) are not appreciably affected. At the high disc speed, much of the $^{53\text{m}}\text{Co}$ is removed before decaying while most of the ^{41}Ti and ^{61}Ge nuclei decay while under the detectors. ^{65}Ge , ($t_{1/2} = 31$ s) produced in the high yield 2pn evaporation channel, emits a continuum of delayed protons ranging from 1.1 to 2.3 MeV⁷; however, the high reaction cross section is offset by its very small proton branch (0.013%). ^{41}Ti is produced in reactions on the oxygen contaminant in the target. The ^{61}Ge peak has also been observed in much reduced yield at 113 MeV bombarding energy, but is absent at lower energies, as expected for the 3n evaporation channel.

The spectra in Fig. 6 show no evidence for delayed protons arising from the decay of ^{65}Se , which has a predicted energy for the transition from the isobaric analog state to the ground state of 3.75 MeV. If the proton decay

were to feed either the first or second excited state of ^{64}Ge , a proton peak at 2.8 or 2.0 MeV would be expected. Since no data are available for ^{64}Ge , its level scheme has been assumed to be similar to that of ^{66}Zn , as expected according to the $N_p N_n$ scheme of Casten⁸.

Figure 7 shows the spectrum obtained when the calcium target was bombarded with 97 MeV ^{28}Si . This is below threshold for the production of ^{61}Ge . Present in the spectrum are delayed protons from $^{53\text{m}}\text{Co}$, ^{65}Ge , and ^{25}Si . The ^{25}Si is once again assumed to be produced in transfer reactions off the entrance windows of the target chamber. Again, there is no evidence of protons arising from the decay of ^{65}Se in this spectrum, or in the data taken at 113 MeV, where its yield is expected to be a maximum. An early attempt to observe ^{65}Se with the target counting method using the same reaction at 105 MeV was also unsuccessful⁹.

Figures 6 and 7 also show several small peaks superimposed on the ^{65}Ge continuum, as do data from experiments which used 73 and 81 MeV ^{28}Si beams. These peaks, at 1.41, 2.22 and 2.41 MeV, cannot be associated with ^{65}Se as 81 MeV is below threshold for its production in this reaction. These peaks are believed to arise from the weak beta-delayed proton decays of one or more unknown $T_z = -1/2$ and -1 nuclei also produced in this reaction. In particular, ^{60}Ga , ^{61}Ga , ^{62}Ge , ^{63}Ge , ^{66}Se and ^{67}Se may be contributing. Mass separation would be required to identify the origin of these weak proton groups.

CONCLUSION

In Fig. 8 a partial decay scheme for ^{61}Ge is shown, deduced from the results of this work. The beta branching ratio to the isobaric analog state

has been estimated assuming $\log ft = 3.3$ for the superallowed transition. The spins and parities of the ground states of ^{61}Ge and ^{61}Ga indicated in Fig. 8 are taken to be the same as in their mirror nuclei. The laboratory energy of the observed delayed proton peak due to ^{61}Ge is 3.11 ± 0.03 MeV, being the weighted mean of five values obtained in the present experiments; this is in good agreement with the predicted energy of 3.21 MeV for the transition, lending credence to the mass calculations. No evidence has been found in our data of proton decay feeding the first excited state of ^{60}Zn , which would give rise to a peak at $E_{\text{lab}} = 2.11$ MeV. An upper limit of 10% of the proton decay can be ascribed to this branch. The mass excess of the lowest $T = 3/2$ state in ^{61}Ga deduced from this work is -43.73 ± 0.03 MeV. Since the $T = 3/2$ analog state in ^{61}Zn is not known, it is not possible to apply the Isobaric Multiplet Mass Equation to predict the ground state mass of ^{61}Ge . Such calculations have been possible for all lighter $T = 3/2$, $A = 4n + 1$ multiplets (see, for example, ref. 4).

While the uncertainties in the mass predictions do increase farther from stability, our calculations indicate that ^{65}Se should decay by beta emission. It is predicted to be stable with respect to particle decay, with $S_p = 1.02$ MeV and $S_{2p} = 0.62$ MeV. Our failure to observe ^{65}Se may be due to an unexpectedly low cross section for the $^{40}\text{Ca}(^{28}\text{Si}, 3n)$ reaction. However, according to the ALICE code, the cross section of this reaction is predicted to be about two-thirds of the ^{61}Ge cross section produced in the $\alpha 3n$ channel at 128 MeV, so it should have been readily observable in these experiments. Alternatively, it is possible that selenium is not transported efficiently in the helium-jet with NaCl aerosols.

This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

References

*⁾Present address: Isotope and Nuclear Chemistry Division, Los Alamos National Laboratory, Los Alamos, NM 87544.

+⁾Permanent address: II Phys. Inst., University of Giessen, W. Germany.

1. I. Kelson and G. T. Garvey, Phys. Lett. 23, 689 (1966).
2. M. S. Antony, J. Britz and A. Pape, At. Data Nucl. Data Tables 34, 279 (1986).
3. The rp process is effectively terminated at ^{64}Ge if ^{65}As is unbound, as it almost certainly must be at the temperature considered for the process ($\sim 10^9\text{K}$). See S. E. Woosley, Proceedings of the Accelerated Radioactive Beams Workshop, Parksville, Canada, 5-7 September 1985, p. 4.
4. D. J. Vieira, D. F. Sherman, M. S. Zisman, R. A. Gough and J. Cerny, Phys. Lett. 60B, 261 (1976).
5. J. Honkanen, M. Kortelahti, K. Eskola and K. Vierinen, Nucl. Phys. A366, 109 (1981).

6. M. Blann and J. Birplinghoff, Lawrence Livermore National Laboratory report UCID-19614, unpublished (1982).
7. J. C. Hardy, J. A. Macdonald, H. Schmeing, T. Faestermann, H. R. Andrews, J. S. Geiger, R. L. Graham and K. P. Jackson, Phys. Lett. 63B, 27 (1976).
8. R. F. Casten, Phys. Rev. C 33, 1819 (1986).
9. D. J. Vieira, Ph.D. Thesis, Report No. LBL-7161, unpublished (1978).

Figure Captions

Fig. 1. Diagram of the pulsed-beam target observation system and associated electronics which control the pulsing of the cyclotron beam and initiation of the counting period.

Fig. 2. Schematic diagram of the helium-jet transport system.

Fig. 3. The delayed proton spectrum from the $^{24}\text{Mg} + \text{Ca}$ reaction at 85 MeV using the pulsed-beam target observation system.


Fig. 4. The decay curve for ^{61}Ge , incorporating data taken at 78 and 85 MeV.

Fig. 5. The delayed proton spectrum from the $^{24}\text{Mg} + \text{Ca}$ reaction at 120 MeV using the helium-jet transport system.

Fig. 6. Delayed proton spectra obtained in the $^{28}\text{Si} + \text{Ca}$ reaction at 128 MeV with two different disc speeds.


Fig. 7. Delayed proton spectrum from the $^{28}\text{Si} + \text{Ca}$ reaction at 97 MeV. This is below the threshold for the production of ^{61}Ge .

Fig. 8. The proposed decay scheme of ^{61}Ge . Energies are given in MeV relative to the ground state of ^{61}Ga . Predicted energies and assumed spins, parities and isospins are indicated in parentheses.



XBL 7710-2075

Fig. 1



XBL 867-11649

Fig. 2






Fig. 3




XBL 868-3044

Fig. 4




XBL 864-10764A

Fig. 5




XBL 864-10767A

Fig. 6



XBL 864-10766A

Fig. 7



XBL 867-11648

Fig. 8

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*