

UC Berkeley

UC Berkeley Previously Published Works

Title

Energy Flux, Lactate Shuttling, Mitochondrial Dynamics, and Hypoxia.

Permalink

<https://escholarship.org/uc/item/4633t3wn>

Journal

Advances in experimental medicine and biology, 903

ISSN

0065-2598

Author

Brooks, George A

Publication Date

2016

DOI

10.1007/978-1-4899-7678-9_29

Peer reviewed

47. Wolfel, E.E., B.M. Groves, G.A. Brooks, G.E. Butterfield, R.S. Mazzeo, L.G. Moore, J.R. Sutton, P.R. Bender, T.E. Dahms, R.E. McCullough, R.G. McCullough, S.-Y. Huang, S.-F. Sun, R.F. Grover, H.N. Hultgren, and J.T. Reeves. Oxygen transport during steady-state submaximal exercise in chronic hypoxia. *J. Appl. Physiol.* 70:1129-1136, 1991.
48. Yoon, H., A. Fanelli, E. F. Grollman, and N. J. Philp. Identification of a unique monocarboxylate transporter (MCT3) in retinal pigment epithelium. *Biochem. Biophys. Res. Comm.* 234: 90-94, 1997.
49. Zinker, B.A., R.D. Wilson, and D.H. Wasserman. Interaction of decreased arterial PO₂ and exercise on carbohydrate metabolism in the dog. *Am. J. Physiol.* 269:E409-417, 1995.

Chapter 16

Are Arterial, Muscle and Working Limb Lactate Exchange Data Obtained on Men at Altitude Consistent with the Hypothesis of an Intracellular Lactate Shuttle?

George A. Brooks

Exercise Physiology Laboratory, Department of Integrative Biology, University of California, Berkeley, CA 94720 USA

Key words: mitochondria, LDH, MCT1, hypoxia, cell redox, exercise, exertion

Abstract: The "Lactate Shuttle" Hypothesis posits that lactate removal requires exchange among producing and consuming cells. The "Intra-cellular Lactate Shuttle" hypothesis posits that lactate exchange occurs among compartments within cells, and that mitochondria are the major sites of cellular lactate disposal. Thus, cells with high mitochondrial densities (cardiocytes, myocytes, hepatocytes) are those which participate in lactate clearance. The model of an Intracellular Lactate Shuttle recognizes that the K_{eq} for LDH is $3.6 \times 10^4 M^{-1}$; thus, glycolysis results in cytosolic lactate production regardless of the intracellular PO₂. The model also requires presence of a mitochondrial monocarboxylate transporter (MCT) that allows uptake of lactate as well as pyruvate, and intra-mitochondrial LDH whose function is linked to the ETC, and which permits lactate → pyruvate conversion and oxidation. Recently, we have shown that liver, heart and muscle mitochondria readily oxidize lactate and contain LDH and MCT1. Accordingly, we have concluded that lactate is the predominant monocarboxylate oxidized by mitochondria *in vivo*. The model of an "Intra-cellular Lactate Shuttle" is consistent with many of the observations on men at sea level and altitude. The observations include: oxidation is the primary fate of lactate disposal during rest and exercise; lactate production and oxidation occur simultaneously within resting and working muscle; increasing [lactate]_i increases muscle lactate extraction, and that by increasing SaO₂ acclimatization reduces blood [lactate].

1. INTRODUCTION

Since the initial observations of Edwards (18), blood lactate responses to exercise at altitude have been observed to demonstrate several unexplained features. Not only has the term "lactate paradox" been used, but investigators have used different definitions for the "paradox" (24, 27, 34). However, since the initial observations and subsequent articulation of terms, our understanding of carbohydrate metabolism at altitude has grown immensely (5-9, 12, 27, 36, 37). Rather than dwell on any specific term whose utility may have passed, in this report the effort will be to use new knowledge in an attempt to explain blood metabolite responses to altitude.

Commencing with brief reviews and presentation of data necessary to illustrate presence of the "Cell-Cell" and "Intra-cellular Lactate Shuttles," an attempt will be made to evaluate whether those concepts are of use in understanding apparently disparate data observed on men at altitude.

2. PHENOMENA TO EXPLAIN:

- Why is arterial [lactate] elevated during exercise at a given power output at altitude compared to sea level?
- Is the elevation in arterial [lactate] at altitude attributable to increased production and appearance in the blood (Ra), or are lactate disposal (Rd) and clearance rates (MCR) inhibited at altitude?
- How can working human muscle display a "Stainsby Effect" at altitude wherein the start of contractions results in muscle net lactate release. Thereafter, net lactate release follows the fall of lactate [v-a] with net release declining to zero even as arterial [lactate] is elevated and constant?
- If there is a "Stainsby Effect" at altitude, what explains the elevation in arterial [lactate]?
- How can working muscle simultaneously release, consume and oxidize lactate at altitude?

3. WHAT IS THE CELL-CELL LACTATE SHUTTLE?

The original "Cell-Cell" lactate shuttle posited that shuttling of lactate through the interstitium and vasculature provides a significant carbon source for oxidation and gluconeogenesis during rest and exercise (4, 5). In Figure 1, Type IIB (fast-glycolytic) fibers are indicated to be lactate producing

31. Molé, P.A., P.A. VanHandel, and W.R. Sandel. Extra O₂ consumption attributable to NADH₂ during maximum lactate oxidation in the heart. *Biochem. Biophys. Res. Comm.* 85:1143-1149, 1978.
32. Philp, N., P. Chu, T-C. Pan, R.Z. Zhng, M-L. Chu, K. Stark, D. Boettinger, H. Yoon, and T. Kieber-Emmons. Developmental expression and molecular cloning of REMP, a novel membrane protein. *Exp. Cell Res.* 219: 64-73, 1995.
33. Price, N. T., V.N. Jackson, and A.P. Halestrap. Cloning and sequencing of four new mammalian monocarboxylate transporter (MCT) homologues confirms the existence of transporter family with an ancient past. *Biochem. J.* 329: 321-328, 1998.
34. Reeves. J.T., E.E. Wolfel, H.J. Green, R.S. Mazzeo, A.J. Young, J.R. Sutton, and G.A. Brooks. Oxygen transport during exercise at high altitude and the lactate paradox: lessons from Operation Everest II and Pikes Peak. In: *Exercise and Sports Sciences Reviews*. K.B. Pandolf (Ed.), Vol. 20, Williams and Wilkins, 1992, pp. 275-296.
35. Richter, E.A., B. Kiens, B. Saltin, N.J. Christensen, and G. Savard. Skeletal muscle glucose uptake during dynamic exercise in humans: role of muscle mass. *Am. J. Physiol.* 254:E555-E561, 1988.
36. Roberts, A.C., G.E. Butterfield, J.T. Reeves, A. Cymerman, E.E. Wolfel, and G.A. Brooks. Altitude and β -blockade augment glucose utilization during submaximal exercise. *J. Appl. Physiol.* 80:606-615, 1996.
37. Roberts, A.C., G.E. Butterfield, A. Cymerman, J.T. Reeves, E.E. Wolfel, and G.A. Brooks. Acclimatization to 4,300 m altitude decreases reliance on fat as a substrate. *J. Appl. Physiol.* 81:1762-1771, 1996.
38. Roth, D.A. and G.A. Brooks. Lactate and pyruvate transport is dominated using a pH gradient-sensitive carrier in rat skeletal muscle sarcolemmal vesicles. *Arch. of Biochem. Biophys.* 279:386-394, 1990.
39. Roth, D.A. and G.A. Brooks. Lactate transport is mediated by a membrane-borne carrier in rat skeletal muscle sarcolemmal vesicles. *Arch. of Biochem. Biophys.* 279:377-385, 1990.
40. Stainsby, W.N. and G.A. Brooks. Control of lactic acid metabolism in contracting muscles and during exercise. In: *Exercise and Sport Sciences Reviews*. K.B. Pandolf and J.O. Holloszy (Eds.). Baltimore: Williams and Wilkins, 1990, pp. 29-63.
41. Stanley, W.C., E.W. Gertz, J.A. Wisneski, D.L. Morris, R. Neese, and G.A. Brooks. Systemic lactate turnover during graded exercise in man. *Am. J. Physiol. (Endocrinol. Metab. 12)*. 249:E595-E602, 1985.
42. Stanley, W.C., E.W. Gertz, J.A. Wisneski, D.L. Morris, R. Neese, and G.A. Brooks. Lactate metabolism in exercising human skeletal muscle: Evidence for lactate extraction during net lactate release. *J. Appl. Physiol.* 60:1116-1120, 1986.
43. Stanley, W.C., J.A. Wisneski, E.W. Gertz, R.A. Neese, and G.A. Brooks. Glucose and lactate interrelations during moderate intensity exercise in man. *Metabolism* 37:850-858, 1988.
44. Wasserman, D.H., D.B. Lacey, D.R. Green, P.E. Williams, and A.D. Cherrington. Dynamics of hepatic lactate and glucose balances during prolonged exercise and recovery in the dog. *J. Appl. Physiol.* 63:2411-2417, 1987.
45. Welch, H.G. and W.N. Stainsby. Oxygen debt in contracting dog skeletal muscle in situ. *Resp. Physiol.* 3:229-242, 1967.
46. Wilson, M.C., V.N. Jackson, C. Hedde, N.T. Price, H. Pilegaard, C. Juel, A. Bonen, I. Montgomery, O.F. Hutter, and A.P. Halestrap. Lactic acid efflux from white skeletal muscle is catalyzed by the monocarboxylate transporter isoform MCT3. *J. Biol. Chem.* 273:15920-15926, 1998.

- steady-rate exercise at 4,300 m altitude. *Am. J. Physiol.* 275 (Regulatory Integrative and Comparative Physiol. 44): R1192-1201, 1998.
13. Brooks, G.A., E.E. Wolfel, B.M. Groves, P.R. Bender, G.E. Butterfield, A. Cymerman, R.S. Mazzeo, J.R. Sutton, R.R. Wolfe, and J.T. Reeves. Muscle accounts for glucose disposal but not blood lactate appearance during exercise after acclimatization to 4,300 m. *J. Appl. Physiol.* 72:2435-2445, 1992.
 14. Connett, R.J., T.E.J. Gayeski, and C.R. Honig. Lactate accumulation in fully aerobic, working, dog gracilis muscle. *Am. J. Physiol.* 246:H120-H128, 1984.
 15. Connett, R.J., C.R. Honig, T.E.J. Gayeski, and G.A. Brooks. Defining hypoxia: a systems view of VO₂, glycolysis, energetics and intracellular PO₂. *J. Appl. Physiol.* 68:833-842, 1990.
 16. Depocas, F., Y. Minaire, and J. Chatonnet. Rates of formation and oxidation of lactic acid in dogs at rest and during moderate exercise. *Can. J. Physiol. Pharmacol.* 47:603-610, 1969.
 17. Donovan, C.M. and G.A. Brooks. Endurance training affects lactate clearance, not lactate production. *Am. J. Physiol.* 244 (Endocrinol. Metab. 7) 244:E83-E92, 1983.
 18. Edwards, H.T. Lactic acid at rest and work at high altitudes. *Am. J. Physiol.* 116:367-375, 1936.
 19. Faintrenie, G. and A. Gélöën. Alpha-1 adrenergic regulation of lactate production in white adipocytes. *J. Pharmacol. and Experimental Therapeutics.* 277:235-238, 1995.
 20. Garcia, C. K., M.S. Brown, R.K. Pathak, and J.L. Goldstein. cDNA cloning of MCT2, a second monocarboxylate transporter expressed in different cells than MCT1. *J. Biol. Chem.* 270: 1843-1849, 1995.
 21. Garcia, C. K., J.L. Goldstein, R.K. Pathak, R.G. Anderson, and M.S. Brown. Molecular characterization of a membrane transporter for lactate, pyruvate and other monocarboxylates: implications for the Cori cycle. *Cell* 76: 865-873, 1994.
 22. Gertz, E.W., J.A. Wisneski, W.C. Stanley, and R.A. Neese. Myocardial substrate utilization during exercise in humans. Dual carbon-labeled carbohydrate isotope experiments. *J. Clin. Invest.* 82:2017-2025, 1988.
 23. Gladden, L.B., R.E. Crawford, and M.J. Webster. Effect of lactate concentration and metabolic rate on net lactate uptake by canine skeletal muscle. *Am. J. Physiol.* 266:R1095-101, 1994.
 24. Hochachka, P.W. The lactate paradox: analysis of underlying mechanisms. *Ann. Sports Med.* 4:184-188, 1989.
 25. Jöbsis F.F. and W. N. Stainsby. Oxidation of NADH during contractions of circulated skeletal muscle. *Resp. Physiol.* 4:292-300, 1968.
 26. Johnson, J.A. and R.M. Fusaro. The role of skin in carbohydrate metabolism. *Adv. Metab. Disord.* 6:1-55, 1972.
 27. Kayser, B. Lactate during exercise at high altitude. *Eur. J. Appl. Physiol.* 74: 195-205, 1996.
 28. Kim, C.M., J.L. Goldstein, and M.S. Brown. cDNA cloning of MEV, a mutant protein that facilitates cellular uptake of mevalonate, and identification of the point mutation responsible for its gain of function. *J. Biol. Chem.* 267: 23113-23121, 1992.
 29. Kopperschlager G. and Kirchnerberger J. Methods for the separation of lactate dehydrogenases and clinical significance of the enzyme. *J. Chromat. B, Biomed. App.* 684:25-49, 1996.
 30. Mazzeo, R.S., G.A. Brooks, D.A. Schoeller and T.F. Budinger. Disposal of [1-¹³C]lactate during rest and exercise. *J. Appl. Physiol.* 60:232-241, 1986.




Figure 1. Model of the "Cell-Cell" Lactate Shuttle. Lactate is formed in fast-twitch white (type IIb) skeletal muscle fibers, or transiently in populations of red fibers at the start of exercise. Lactate released into the interstitium from cellular sites of production can be taken up and oxidized by adjacent lactate consuming (type I) muscle fibers. Alternatively, during exercise some lactate released into the circulation reperfuses the active muscle bed within a fraction of a minute, where uptake and oxidation in red, highly oxidative fibers occurs. Some of the lactate released from an active muscle bed can be taken up by the heart and oxidized, or taken up by the liver and kidneys where lactate serves as a gluconeogenic precursor. Redrawn from (4).

cells, whereas Type I fibers are indicated as sites of lactate oxidation. This hypothesis was developed at a time when fiber heterogeneity in skeletal muscle was recognized (1, 3). Further, the hypothesis was based on results of rats and dogs made to exercise and infused with radioactive glucose and lactate tracers (10, 11, 16, 17). On those species it was shown that during exercise lactate turnover and oxidation exceed glucose turnover and oxidation. Subsequently, the same phenomena of lactate flux and oxidation were demonstrated in humans (30, 43). Data in Figure 2 shows the direct relationships between lactate turnover and metabolic rate and between lactate oxidation and metabolic rate (30). Lactate turnover is high at rest and increases as a direct function of exercise intensity and metabolic power output. Further, lactate disposal through oxidation (approximately 50% at rest), increases both absolutely and relatively during exercise such that oxidation accounts for 75% of lactate disposal during exercise.

While systemic lactate flux and whole-body oxidation data supported the concept of a lactate shuttle, the technology could not be used to demonstrate tissue lactate exchange. Therefore, studies were conducted to determine the lactate concentration differences across active (leg) and inactive (arm)




Figure 2. Lactate turnover determined with $[1-^{13}\text{C}]$ lactate tracer in men at sea level during rest, exercise at 50% $\text{VO}_{2\text{max}}$, and exercise at 75% $\text{VO}_{2\text{max}}$. Data from Mazzeo et al. (30).




Figure 3. [Lactate] in blood sampled simultaneously from the radial artery, brachial vein, and iliac vein in men during continual, progressive exercise protocol. The results illustrate tissue lactate exchange through the blood during exercise. From Stanley et al. (42).

muscle beds during progressive leg cycling exercise. The results (42) (Figure 3) demonstrate lactate release from active (leg) muscle beds, transit through the vasculature, and uptake by inactive muscle (arm) beds. In the same set of studies, measurements of arterial-coronary sinus concentration differences

10. CONCLUSION

The Cell-Cell and Intramuscular Lactate Shuttle Concepts appear to be of assistance in understanding several phenomena observed in men exercising at altitude. Further work is needed to evaluate the effects of hypoxia on expression of mitochondrial and extra-mitochondrial lactate transporters and lactate dehydrogenase isoforms.

ACKNOWLEDGEMENT

Supported by NIH grants DK19577 and AR 42906.

REFERENCES

- Baldwin, K.M., P.J. Campbell, and D.A. Cooke. Glycogen, lactate and alanine changes in muscle during graded exercise. *J. Appl. Physiol.* 43:288-291, 1977.
- Baldwin, K.M., A.M. Hooker, and R.E. Herrick. Lactate oxidative capacity in different types of muscle. *Biochem. Biophys. Res. Comm.* 83:151-157, 1978.
- Barnard, R., V.R. Edgerton, T. Furukawa, and J.B. Peter. Histochemical, biochemical and contractile properties of red, white, and intermediate fibers. *Am. J. Physiol.* 220:410-414, 1971.
- Brooks, G.A. Lactate: Glycolytic product and oxidative substrate during sustained exercise in mammals—the "lactate shuttle," In: *Comparative Physiology and Biochemistry - Current Topics and Trends, Volume A, Respiration - Metabolism - Circulation*, R. Gilles (Ed.), Berlin, Springer-Verlag, 1985, pp. 208-218.
- Brooks, G.A. Current concepts in lactate exchange. *Med. Sci. Sports Exerc.* 23:895-906, 1991.
- Brooks, G.A. Mammalian Fuel Utilization During Sustained Exercise. *Comp. Biochem. Physiol.* 120:89-107, 1998.
- Brooks, G.A., H. Dubouchaud, M. Brown, J. P. Sicurello, and C.E. Butz. Role of mitochondrial lactic dehydrogenase and lactate oxidation in the 'intra-cellular lactate shuttle.' *Proc. Natl. Acad. Sci. USA* 96:1129-1134, 1999.
- Brooks, G.A., G.E. Butterfield, R.R. Wolfe, B.M. Groves, R.S. Mazzeo, J.R. Sutton, E.E. Wolfel, and J.T. Reeves. Increased dependence on blood glucose after acclimatization to 4,300m. *J. Appl. Physiol.* 70:919-927, 1991.
- Brooks, G.A., G.E. Butterfield, R.R. Wolfe, B.M. Groves, R.S. Mazzeo, J.R. Sutton, E.E. Wolfel, and J.T. Reeves. Decreased reliance on lactate during exercise after acclimatization to 4,300 m. *J. Appl. Physiol.* 71:333-341, 1991.
- Brooks, G.A. and C.M. Donovan. Effect of training on glucose kinetics during exercise. *Am. J. Physiol.* 244 (Endocrinol. Metab. 7):E505-E512, 1983.
- Brooks, G.A. and G.A. Gaesser. End points of lactate and glucose metabolism after exhausting exercise. *J. Appl. Physiol.* 49:1057-1069, 1980.
- Brooks, G.A., E. E. Wolfel, G. E. Butterfield, A. Cymerman, A.C. Roberts, R.S. Mazzeo, and J.T. Reeves. Poor relationship between arterial [lactate] and leg net release during

extraction and oxidation by working muscle beds. The Lactate Shuttle Concepts accommodate all of these phenomena.

Arterial [lactate] is elevated at altitude because lactate Ra (production) is increased. However, it is clear working muscle is not the sole contributor to the circulating lactate load. Non-working muscle, skin, adipose and liver are all candidates as contributors to the circulating lactate load during exercise at altitude. Lactate production in non-contracting tissues appears to be under β -adrenergic control.

Lactate production is increased at altitude because glycolysis is stimulated. This stimulation occurs transiently in working muscle, but persists elsewhere in the body.

Working human muscle displays a "Stainsby Effect" during exercise at altitude. The start of contractions results in net lactate release followed the fall of lactate [v-a] and net release to zero. However, measurements of working limb (muscle) lactate concentration differences are inadequate to reveal active intramuscular production and oxidation.

Working skeletal muscle can consume and oxidize lactate at altitude. Further, working muscle lactate extraction and oxidation depend on the arterial lactate concentration. Because lactate extraction and oxidation are related to arterial [lactate], the greatest intramuscular lactate oxidation is observed during exercise upon acute altitude exposure.

It is now known that muscle possesses at least two MCT isoforms that facilitate sarcolemmal exchange of lactate; these are MCT1 and MCT4. The effects of chronic hypoxia on expression of these lactate transport proteins is unknown. However, the model (Figure 4) indicates a major role of sarcolemmal lactate transporters at altitude. Probably, MCT1 is present in mitochondria and sarcolemmal membranes whereas MCT4 is the constitutive muscle cell membrane isoform.

In addition to skeletal muscle, numerous other cells and tissues (erythrocytes, heart, liver, kidneys, brain, testes) express lactate transporter isoforms. Again, the effects of hypoxia are unknown.

Intra-myocyte, -hepatocyte, and -cardiocyte lactate oxidation depends on the presence of mitochondrial lactate-pyruvate transporter. When cytosolic [lactate] is high, lactate enters mitochondria where it is oxidized. Apparently, lactate oxidation in muscle is facilitated by high mitochondrial density and oxygen consumption rates.

Intra-myocyte, -hepatocyte, and -cardiocyte lactate oxidation depends on the presence of mitochondrial LDH. In mitochondria, LDH probably exists in the inter-membrane space as well as on the inner membrane. Mitochondrial LDH is essential for the oxidation of lactate.

and coronary blood flow indicated that exogenous lactate becomes the major fuel for the heart during exercise (22). Further, in those studies the appearance of ^{13}C and ^{14}C in blood glucose following tracer-lactate infusion indicated participation of the Cori Cycle in lactate removal (41, 43). Thus, skeletal as well as cardiac muscle and liver participate in lactate exchange and metabolism during exercise.

With regard to the issue of oxygen insufficiency as the cause of lactate production in muscle, the results obtained on resting, as well as contracting muscle indicate that lactate production occurs in well perfused and fully oxygenated tissues (47). Not only is lactate always present in resting muscle, but lactate release and oxidation occur simultaneously (6) while cardiac output and limb blood flow rise so that oxygen transport meets demand, and significant O_2 remains in femoral venous blood. Data on intact functioning humans as well as animal muscle preparations indicate that lactate production occurs under fully aerobic conditions (14, 15, 25).

4. WHAT IS THE INTRACELLULAR LACTATE SHUTTLE?

To resolve the dilemma of tissue lactate production under fully aerobic conditions, the "Intracellular Lactate Shuttle" was posited. While there was good evidence to support activity of the "Cell-Cell Lactate Shuttle" during exercise, at rest recruitment of Type IIB fibers could not be invoked. A better model that allowed for lactate production in all fiber types during rest and exercise was necessary (6). The "Intracellular" Shuttle (Figure 4) allows that glycolysis in the cytosol results in lactate production, and lactate shuttles to mitochondria within the cell of production for oxidative removal (7).

That glycolysis inevitably results in cytosolic lactate production relates to the energetics of the terminal enzyme, lactate dehydrogenase (LDH). The K_{eq} for LDH is $3.6 \times 10^4 \text{ M}^{-1}$, and the free energy change (ΔG°) approximates -6 kcal/mol . This means that glycolysis inevitably proceeds to lactate production, and that its reversal in the cytosol is unlikely *in vivo*. How then is lactate oxidized?

To evaluate the role of mitochondria in balancing lactate production and oxidation as part of an "Intra-cellular Lactate Shuttle," isolated rat cardiac, skeletal muscle and liver mitochondria were respired with lactate and pyruvate in presence or absence of known inhibitors of metabolism. As well, efforts were made to detect the presence of LDH isoforms in mitochondria by electrophoresis and electron microscopy. Results support the conclusion of a mitochondrial role in cellular lactate oxidation (6, 7), and provide an

explanation for the high correlation between lactate clearance during exercise and muscle mitochondrial respiratory capacity (2, 31).




Figure 4. Depiction of the functional relationship between mLDH and mMCT in operation of the "Intra-cellular Lactate Shuttle." The predominant monocarboxylate entering the mitochondrial inter-membrane space is lactate. Entry of lactate and pyruvate into the mitochondrial matrix is facilitated by mMCT. Thus, lactate enters mitochondria; lactate is oxidized to pyruvate via mLDH when mitochondrial Redox decreases, and pyruvate is oxidized via the TCA Cycle and ETC. From Brooks et al. (7).

In one experiment, mitochondria were isolated and respired with pyruvate-malate and lactate-malate as substrates. Respiration was stimulated by the presence of ADP. Figure 5 shows that maximal, ADP-stimulated respiration rates were obtained with both lactate and pyruvate as substrates (7). Further, the figure shows that addition of the known LDH inhibitor oxamate, blocks mitochondrial lactate oxidation. Thus, LDH is necessary for mitochondrial lactate oxidation.

To determine if LDH existed in mitochondria, organelles from rat liver, heart and skeletal muscle were isolated. Agarose gel electrophoresis of LDH isoenzyme patterns in cytosolic fractions of different tissues (Figure 6) are consistent with results of previous investigations showing tissue specificity (e.g., 29). However, mitochondrial fractions also revealed presence of LDH isoenzymes. Further, LDH isoenzyme patterns differed among tissues, and between mitochondria and the surrounding cytosol in each tissue. Heart and




Figure 12. Leg blood lactate extraction as a function of arterial lactate concentration during rest (closed symbols) and leg cycling exercise (open symbols) at sea level (squares), upon acute exposure to 4,300 m altitude (circles), and after 3-wk acclimatization (triangles). Each point $n = 6-7$, mean \pm SEM. Lactate extraction is highly correlated with arterial concentration. From Brooks et al. (13).

Although the results portrayed in Figure 12 are unique to men working at altitude, they are not unique in nature. With canine muscle preparations made to contract *in situ*, Gladden and associates (23) showed that by raising the arterial [lactate], working muscle switches from net release to uptake. Similarly, by adding arm to leg exercise (arm exercise raises the arterial [lactate]), Richter et al. (35) showed that leg muscle switches from net release to uptake. Thus, based on either chemical balance (Figures 3 and 9), or isotope tracers (Figure 12), muscle is capable of lactate exchange and oxidation.

9. RESOLUTION OF QUESTIONS:

The Cell-Cell and Intramuscular Lactate Shuttle Concepts may be of assistance in understanding several phenomena observed in men exercising at altitude. Such phenomena of interest include: elevated arterial lactate levels, response of circulating lactate concentration to β -blockade, temporal variation in lactate [v-a] and net release, and simultaneous lactate release,




Figure 11. Mean (\pm SEM) leg net lactate venous-arterial concentration differences at rest and over time during exercise at sea level, upon acute exposure to 4,300 m altitude, and after a 3-wk acclimatization period: (A) control ($n = 5$), and (B) β -blocked subjects ($n = 6$). Corresponding mean (\pm SEM) net lactate release from the two legs over time during rest and exercise time in the same: (C) control, and (D) β -blocked subjects. From Brooks et al. (6).

Figure 12 portrays lactate extraction by leg muscles of men during rest and altitude at sea level, upon acute exposure to 4,300 m, and after a 3-week acclimatization period (9). The values were calculated from the decrease in tracer ($[3-^{13}\text{C}]$ lactate) mass across limbs. Not shown is that during exercise extraction equaled oxidation (i.e., the rise in $^{13}\text{CO}_2$ equaled the decrease in ^{13}C -lactate). As shown in Figure 12, lactate extraction and oxidation are directly related to arterial lactate concentration. Thus, greatest lactate extraction and oxidation rates were observed upon acute altitude exposure when the arterial lactate concentration was highest.

muscle mitochondria were noted by prevalence of both LDH-1 (H4) and LDH-5 (M4), while liver mitochondria were distinguished by presence of LDH-5 (M4) (7).




Figure 5. Depiction of a Clarke- O_2 electrode tracing showing that oxamate blocks lactate, but not pyruvate oxidation. Such experiments indicate that the ability of mitochondria isolated from muscle, liver and heart to oxidize lactate requires LDH for conversion to pyruvate within mitochondria. Data from Brooks et al. (7).

Even though studies with oxamate indicated functionality of mitochondrial LDH (Figure 5), and electrophoresis showed mitochondrial LDH (Figure 6), to further exclude the possibility of isolation artifacts, antibodies to LDH isoforms were used to detect presence of mitochondrial LDH, *in situ*. The electron micrograph in Figure 7 clearly shows a mitochondrial location of LDH (7).

The model of an "Intracellular Lactate Shuttle" (6, 7) depends heavily on presence of a mitochondrial lactate-pyruvate, or monocarboxylate transporter (MCT) (38, 39). To demonstrate the effects of inhibiting the mitochondrial lactate-pyruvate transporter, mitochondria were respired in the presence or absence of the known MCT inhibitor, cinnamate (CINN); results are shown in Figure 8. CINN blocked respiration of both substrates. However, in the presence of CINN, State 3 respiration was restored with




Figure 6. Agarose gel electrophoresis of LDH in mitochondria from rat liver and heart. LDH isoenzyme patterns differ between cytosol and mitochondria in both tissues. From Brooks et al. (7).

addition of either succinate or glutamate. Restoration of respiration with succinate or glutamate indicates that CINN does not block mitochondrial respiratory Complex I (glutamate, an NADH linked substrate), or Complex II (succinate, an FADH₂-linked substrate). Upstream inhibition of mitochondrial lactate and pyruvate oxidation, such as a transport limitation, by CINN is indicated. Thus, consistent with the concept of an "Intracellular Lactate Shuttle" (Figure 4), results as in Figures 5 and 8 show that mitochondrial lactate oxidation proceeds very rapidly requiring a transporter and LDH.

To date, preliminary studies have been conducted to identify the mitochondrial lactate-pyruvate transporter. Several possibilities exist. The cDNA of the first sarcolemmal lactate transporter from the hamster ovary cells was isolated by expression cloning (28). The protein was expressed in several tissues including heart, red skeletal muscle and erythrocytes (21). The cDNA was subsequently used for the screening of hamster liver cDNA library and resulted in isolation of the second transporter isoform which was expressed mainly in liver (20). These isoforms shared 60% identity with each other. The third MCT isoform was initially isolated as a protein exclusively expressed in the chicken retinal epithelium (32). All three isoforms stimulate proton-coupled lactate and pyruvate uptake when expressed in a heterologous system and show high affinities for propionate and ketone bodies (20, 21), and were named monocarboxylate transporters (MCT1, MCT2 and MCT3). More recently, four new mammalian monocarboxylate transporter (MCT) homologues were cloned and sequenced (33), and other candidates exist (Aivazachvili and Brooks, unpublished). Although scattered and partial homology exists among the

working human limbs (12) (Figure 11). Under all non- β -blocked conditions studied, resting limbs released lactate on a net basis. Then, when contractions started, there was a burst of lactate release, an occurrence most exaggerated upon acute altitude exposure (Figure 11A). However, even then, \dot{V}_O_2 declined to zero as exercise continued, all the while arterial [lactate] remained elevated and constant. Clearly, working muscle contributes to the circulating lactate load when exercise starts. However, the persistence of an elevation in circulating lactate (Figure 9), despite rapid turnover (Figure 10), must be attributable to tissue sites other than working muscle.

At present, the extra-muscular sites of lactate production and release during exercise at altitude are undetermined. In dogs made to exercise under normoxic (44) and hypoxic conditions (45), the liver is a site of net lactate release. However, similar data are not available on humans. Other possibilities include adipose and skin, both of which respond to adrenergic stimulation by increasing glycolysis and lactate release (19, 26).

At altitude, β -adrenergic blockade had some perplexing effects on muscle lactate exchange (Figure 11B). On acute exposure under β -blockade, there was a negative lactate [v-a], indicating uptake. This result was different from that observed under all blocked and unblocked conditions. The start of exercise upon acute exposure with β -blockade resulted in increased net lactate release, but release was blunted. Interestingly, net lactate release from the working limb was greatest after acclimatization, when the circulating [lactate] had returned almost to sea level values (Figure 9). As discussed previously (12), muscle and femoral venous lactate contents were minimally effected by β -blockade at altitude. Therefore, the lactate [v-a] was exaggerated due to the lesser circulatory lactate load.

8. LACTATE EXTRACTION AND OXIDATION DURING NET RELEASE:

Previously, Stanley et al. (42) demonstrated that working human muscle is capable of simultaneous lactate extraction and release with almost all of the extraction attributable to oxidation in working muscle. Leg extraction and oxidation can be calculated in two ways, which usually yield good agreement in the steady state. One method is to compare the amount of tracer lactate leaving the limb with that entering. The difference (i.e., the decrease in tracer mass leaving the leg) is quantitatively related to the increase in labeled CO₂ release. The alternative method is to compare the mass of tracer CO₂ entering and leaving the leg.




Figure 10. Blood glucose disposal rate (Rd), determined with $[6,6-^2\text{H}]$ glucose, and lactate appearance rate (Ra), determined with $[3-^{13}\text{C}]$ lactate, in men during rest (top) and exercise (bottom), at sea level (left) and acute exposure to 4,300 m altitude (right). During acute altitude exposure, lactate flux exceeds glucose flux during exercise. From Brooks et al. (13).

7. ILLUSTRATION OF THE "STAINSBY EFFECT" BY WORKING HUMAN MUSCLE AT ALTITUDE

To evaluate the role of working muscle in contributing to the circulating lactate load during exercise at high altitude, femoral arterial and venous catheterizations were performed and limb blood flow rates as well as $[v-a]$ lactate differences were determined. In this way, net lactate release rates (\dot{L}) were determined during rest and exercise under each environmental condition (12). That canine muscle preparations made to contract *in situ* release lactate on a net basis when exercise starts, and then switch to zero release or net consumption as contractions progress was first made by Stainsby and associates (40, 45). A similar 'Stainsby Effect' is observable in

MCT isoforms, it is unlikely that all isoforms are lactate-pyruvate transporters. Most probably, MCT1 and MCT4 are the lactate transporters present in muscle (46).




Figure 7. Electron micrograph of high pressure frozen rat liver showing mitochondria, rough ER, and cytosol. Immunolocalization of anti-LDH-5 (M4) antibodies is indicated by the 15 nm gold particles. Magnification = 54,100 X and scale bar = 400 nm. Note presence of LDH-5 in mitochondria and surrounding matrix and organelles. From Brooks et al. (7).




Figure 8. Cinnamate blocks oxidation of lactate and pyruvate. Because muscle, liver and heart mitochondria can respire both glutamate (donates electrons at Complex I) and succinate (donates electrons at Complex II), the blockage of cinnamate is upstream of PDH, GDH or SDH. The essential role of a mitochondrial MCT (lactate/pyruvate) transporter is illustrated. Data from Brooks et al. (7).

Because there was a good relationship between the oxidative capacity of muscle and the abundance of MCT1, we raised an antibody to the C-terminus of rat MCT1. Western blots of cardiac and skeletal muscle mitochondria prepared from rats show presence of MCT1 in striated muscle mitochondria. Further, electron microscopy and immunolocalization techniques show MCT1 on inner mitochondrial membranes (Brooks, et al., unpublished). Thus, MCT1 is a candidate mitochondrial lactate/pyruvate transporter.

5. STUDIES ON MEN EXERCISING AT ALTITUDE

Having now introduced the Cell-Cell and Intracellular Lactate Shuttle Concepts, the pertinent data on men exercising at altitude will be reviewed and presented with an eye toward evaluating whether the concepts are of assistance in evaluating the data.

When men are acutely exposed to moderate altitude (e.g., 4,300 m on Pikes Peak), the circulating [lactate] is elevated during exercise compared to during the same, sub-maximal power output as at sea level. In each environmental condition portrayed (sea level, acute and chronic altitude exposure), the arterial [lactate] is elevated, and stable during exercise (Figure 9A). On acute altitude exposure, the greatest lactate concentration response is observed. With acclimatization, the circulating [lactate] declines even though whole body and working limb VO_2 are unchanged at altitude compared to sea level. Thus, there can be no possibility of a change in lactate production due to O_2 lack.

Figure 9B shows the sympathetic effect on circulating lactate level. β -adrenergic blockade with the non-specific β -antagonist propranolol decreases arterial [lactate] by approximately 60%, a result most obvious on acute altitude exposure. At altitude, β -blockade attenuates, but does not completely block the blood lactate response to exercise. What is the source of circulating lactate during exercise at altitude?

6. LACTATE TURNOVER (PRODUCTION) AT ALTITUDE

In the 1988 Pikes Peak study, fluxes of glucose and lactate were determined simultaneously using primed-continuous infusions of [6,6-




Figure 9. Arterial lactate concentrations (mean \pm SEM) over time in: (A) control (n = 5), and (B) β -blocked subjects (n = 6) at sea level (SL), upon acute exposure and chronic (3-wk) exposure to 4,300 m altitude; symbols: * different from SL, ‡ different from A1, § different from rest, † different from control, $p < 0.05$. From Brooks et al. (6).

^2H]glucose and [3- ^{13}C]lactate. The results for sea level and acute exposure to 4,300 m are shown in Figure 10. During rest at sea level, glucose disposal (Rd) exceeds lactate appearance (Ra). Upon altitude exposure, glucose Rd rises significantly, but lactate Ra rises relatively more. During exercise at 50% VO_2max at sea level, glucose flux rises compared to rest, and lactate flux rises to approximate glucose flux. Then, during exercise upon acute altitude exposure, lactate flux exceeds glucose flux. Thus, it is clear that altitude exposure causes a shift to carbohydrate metabolism, with both glucose and lactate fluxes exaggerated compared to sea level. Most importantly, blood lactate flux, a key measure of tissue lactate production, is increased significantly at altitude.