

DERIVED TRACES OF SOERGEL CATEGORIES

EUGENE GORSKY, MATTHEW HOGANCAMP, AND PAUL WEDRICH

ABSTRACT. We study two kinds of categorical traces of (monoidal) dg categories, with particular interest in categories of Soergel bimodules. First, we explicitly compute the usual Hochschild homology, or derived vertical trace, of the category of Soergel bimodules in arbitrary types. We show that this dg algebra is formal, and calculate its homology explicitly, for all Coxeter groups. Secondly, we introduce the notion of derived horizontal trace of a monoidal dg category and identify the derived horizontal trace of Soergel bimodules in type A with the homotopy category of perfect dg modules of an explicit algebra. As an application we obtain a derived annular Khovanov–Rozansky link invariant with an action of full twist insertion, and thus a categorification of the HOMFLY-PT skein module of the solid torus.

CONTENTS

1. Introduction	1
Acknowledgements	9
Funding	9
2. Facts from homological algebra	9
3. Standard dg categorical constructions	14
4. Homotopy idempotents and the Karoubi envelope	17
5. The bar complex for dg categories	28
6. The dg monoidal center and trace	37
7. Traces of the Soergel category	51
8. The Soergel category in type A	56
References	67

1. INTRODUCTION

1.1. **Traces.** Traces are ubiquitous in mathematics. If A is an algebra over a field \mathbb{k} , its trace (or cocenter) is defined as the quotient $\mathrm{HH}_0(A) = A/[A, A]$. Given any finite-dimensional A -module M , the trace $\mathrm{tr}_M: A \rightarrow \mathbb{k}$ of the A -action on M satisfies $\mathrm{tr}_M(xy) = \mathrm{tr}_M(yx)$ and hence factors through $A/[A, A]$. The projection $A \rightarrow A/[A, A]$ can thus be considered as a universal trace on A .

We are interested in traces for categories. If \mathcal{C} is a \mathbb{k} -linear category, then its trace is a vector space over \mathbb{k} . If \mathcal{C} is a monoidal category (or a 2-category) then its trace is a 1-category. Abstractly, the trace of an n -category satisfying certain assumptions is an $(n - 1)$ -category related to the factorization homology of the circle (see e.g.

[AF19] for an introduction), but we will not employ this point of view in the present paper. Categorical traces have been studied in various settings, see for example Ocneanu [Ocn94], Evans–Kawahigashi [EK95], Walker [Wal], Ben-Zvi–Nadler [BN09], Ponto–Shulman [PS13], Beliakova–Lauda–Habiro–Živković [BHLv17], Hoyois–Schroetzke–Sibilla [HSS17], Beliakova–Putyra–Wehrli [BPW19].

In this paper we construct and study the derived traces of monoidal dg categories, with a view towards applications in higher representation theory and link homology.

Given a \mathbb{k} -linear dg category \mathcal{C} , one can define its Hochschild homology $\mathrm{HH}(\mathcal{C})$ which is a vector space over \mathbb{k} . It is defined as homology of the explicitly defined *cyclic bar complex* which we review in Section 5.5. Keller proved [Kel06] that $\mathrm{HH}(\mathcal{C})$ is a derived invariant of \mathcal{C} , for example, if \mathcal{C} is the category of perfect complexes over an algebra A then $\mathrm{HH}(\mathcal{C})$ is isomorphic to the usual Hochschild homology of A . Note that $\mathrm{HH}_0(A) = A/[A, A]$ suggesting the interpretation of Hochschild homology as a *derived (vertical) trace*.

If \mathcal{C} is a monoidal category (or a bicategory), then there is a richer notion of *horizontal trace* $\mathrm{Tr}_0(\mathcal{C})$ which is well-studied in various levels of generality. This is a category equipped with the “trace functor” $\mathrm{Tr}_0: \mathcal{C} \rightarrow \mathrm{Tr}_0(\mathcal{C})$. If \mathcal{C} has left duals, then Tr_0 is initial among all trace-like functors¹ $F: \mathcal{C} \rightarrow \mathcal{D}$, i.e. functors equipped with a natural transformation $F(X \otimes Y) \rightarrow F(Y \otimes X)$ respecting the tensor product in \mathcal{C} (if \mathcal{C} also has right duals, then the components of these natural transformations are necessarily isomorphisms). The horizontal trace is indeed a richer notion than the vertical trace, since the endomorphism algebra of $\mathrm{Tr}_0(\mathbb{1})$ in the horizontal trace naturally agrees with the vertical trace $\mathrm{HH}_0(\mathcal{C})$.

Example 1.1. The horizontal trace of the bicategory of tangles (where objects are finite sets of points in I^2 , 1-morphisms are tangles in I^3 , and 2-morphisms are tangle cobordisms in I^4 up to isotopy rel boundary) is the category of links in the thickened annulus (objects are links in $I^2 \times S^1$, morphisms are link cobordisms in $I^3 \times S^1$ up to isotopy rel boundary).

In this paper we define a derived version of the horizontal trace and prove the following:

Theorem 1.2. *There is a natural dg functor $\mathcal{C} \rightarrow \mathrm{Tr}(\mathcal{C})$, which is homotopy trace-like i.e. it is equipped with transformations $\mathrm{Tr}(X \otimes Y) \rightarrow \mathrm{Tr}(Y \otimes X)$ that are natural in Y and natural up to coherent homotopy in X . The endomorphism algebra of $\mathrm{Tr}(\mathbb{1})$ is naturally isomorphic to $\mathrm{HH}(\mathcal{C})$.*

If \mathcal{C} has left duals we expect that Tr is initial among all homotopy trace-like dg functors out of \mathcal{C} .

We also define the notion of dg Drinfeld center of \mathcal{C} and prove that it acts on $\mathrm{Tr}(\mathcal{C})$; see section 6.6.

¹Variations of trace-like functors are known under the names *shadows* [PS13], *commutator functors* [BFO09], *categorical traces* [HPT16], *trace functors* [Zhu18].

It is desirable to consider the closure of $\mathrm{Tr}(\mathcal{C})$ with respect to mapping cones and homotopy direct summands, which we denote by:

$$(1) \quad \widetilde{\mathrm{Tr}}(\mathcal{C}) := \mathrm{Pretr}(\mathrm{Kar}^{dg}(\mathrm{Tr}(\mathcal{C}))),$$

where Kar^{dg} and $\mathrm{Pretr}(-)$ respectively denote the homotopy idempotent completion and pretriangulated hull; see sections 3 and 4 for details.

1.2. Traces of Soergel bimodules. Next, we apply all the machinery of derived traces to categories of Soergel bimodules, starting with a computation of the derived vertical trace. Let W be a Coxeter group with simple reflections $S \subset W$ and a realization V over \mathbb{C} , and $\mathrm{SBim}(W)$ the associated monoidal category of Soergel bimodules [Soe92], which is a categorification of the Hecke algebra associated to the Coxeter system (W, S) . We set $R := \mathrm{Sym}^\bullet(V^*)$, graded by placing V^* in bidegree $(2, 0)$ and $\Lambda := \Lambda^\bullet(V^*)$, graded by placing V^* in bidegree $(2, -1)$.

Theorem 1.3. *We have an isomorphism of associative bigraded algebras*

$$\mathrm{HH}(\mathrm{SBim}(W)) \cong \mathrm{HH}(R) \rtimes \mathbb{C}[W].$$

Furthermore, $\mathrm{HH}(\mathrm{SBim}(W))$ is formal as dg algebra, so higher Massey products on Hochschild homology vanish.

Remark 1.4. The Hochschild homology of the polynomial ring R is canonically isomorphic to $R \otimes \Lambda$. After choosing a basis of V we can identify $\mathrm{HH}(\mathrm{SBim}(W))$ with the algebra $\mathbb{C}[x_1, \dots, x_r, \theta_1, \dots, \theta_r] \rtimes \mathbb{C}[W]$ in which x_i are even variables of degree $(2, 0)$ and the θ_i are odd variables of degree $(2, -1)$, and $r = \dim(V)$.

Remark 1.5. In Theorem 1.3 the generators of the wreath product algebra on the right hand side are identified with the Hochschild cycles on the left hand side as follows. The generators of R correspond to cycles $x \in \mathrm{Hom}(\mathbb{1}, \mathbb{1})$, the generators of Λ correspond to the cycles $x \parallel \mathrm{Id} - \mathrm{Id} \parallel x \in \mathrm{Hom}(\mathbb{1}, \mathbb{1}) \otimes \mathrm{Hom}(\mathbb{1}, \mathbb{1})$ (see section 5 for notation conventions), and the elements $w \in W$ correspond to $(-1)^{\ell(w)} \mathrm{Id}_{\Delta_w} \in \mathrm{Hom}(\Delta_w, \Delta_w)$ where Δ_w is the Rouquier complex corresponding to positive braid lift of w , and $\ell(w)$ is the length of w . See Theorem 7.7 for details.

Remark 1.6. In [EL16] the isomorphism $\mathrm{HH}_0(\mathrm{SBim}(W)) \cong R \rtimes \mathbb{C}[W]$ was proved by completely different methods, using cellularity of $\mathrm{SBim}(W)$.

After proving Theorem 1.2 we specialise to Soergel bimodules SBim_n for the symmetric group S_n , which feature in triply-graded Khovanov–Rozansky link homology [Kho07]. In [GW19] the first and third authors studied the category of annular webs and foams, which can be regarded as Karoubi completion of the horizontal trace $\mathrm{Tr}_0(\mathrm{SBim}_n)$. In particular, they proved that this Karoubi completion is generated by the direct summands of $\mathrm{Tr}_0(\mathbb{1}_n)$. Here we generalise this result to the derived horizontal trace using slightly different methods.

Theorem 1.7. *The dg functor $\mathrm{Hom}_{\mathrm{Tr}(\mathrm{SBim}_n)}(\mathrm{Tr}(\mathbb{1}_n), -)$ induces a quasi-equivalence relating $\widetilde{\mathrm{Tr}}(\mathrm{SBim}_n)$ to the category of perfect right dg-modules over $\mathrm{End}(\mathrm{Tr}(\mathbb{1}_n))$. In other words, we have a quasi-equivalence*

$$\begin{aligned} \widetilde{\mathrm{Tr}}(\mathrm{SBim}_n) &\cong \mathrm{Perf}(\mathrm{HH}(R) \rtimes \mathbb{C}[S_n]) \\ &\cong \mathrm{Perf}(\mathbb{C}[x_1, \dots, x_n, \theta_1, \dots, \theta_n] \rtimes \mathbb{C}[S_n]), \end{aligned}$$

where $\deg(x_i) = (2, 0)$ and $\deg(\theta_i) = (2, -1)$.

Note that projective modules over the algebra $\mathrm{HH}(R) \rtimes \mathbb{C}[S_n]$ are naturally indexed up to isomorphism by partitions of n , and under the equivalence of categories above every object of $\widetilde{\mathrm{Tr}}(\mathrm{SBim}_n)$ can be expressed as a twisted complex whose terms are (direct sums of shifts of) these projective modules. In other words, if e_λ is a projector in $\mathbb{C}[S_n]$ onto an irreducible representation V_λ , then we define the *Schur object*

$$(2) \quad \mathbb{S}^\lambda := e_\lambda \mathrm{Tr}(\mathbb{1}_n)$$

in $\widetilde{\mathrm{Tr}}(\mathrm{SBim}_n)$. A perfect dg module over $\mathrm{HH}(R) \rtimes \mathbb{C}[S_n]$ is then a twisted complex built out of \mathbb{S}^λ .

Remark 1.8. The type A Soergel bimodule categories taken together form the monoidal bicategory $\bigoplus_{n \geq 0} \mathrm{SBim}_n$, with the (new) tensor product \boxtimes provided by induction $\mathrm{SBim}_n \times \mathrm{SBim}_m \rightarrow \mathrm{SBim}_{m+n}$. Its trace inherits the monoidal structure and is expected to admit a braiding that induces the S_n action on $\mathrm{Tr}(\mathbb{1}_1)^{\boxtimes n} = \mathrm{Tr}(\mathbb{1}_n)$. The Schur object \mathbb{S}^λ is designed to be the evaluation of the λ -Schur functor on $\mathrm{Tr}(\mathbb{1}_1)$.

1.3. Derived annular link invariants and categorification of the skein module of the solid torus. The main motivation for this paper is to develop a framework for the categorification of the HOMFLY-PT skein module of the solid torus that is compatible with expectations from topological field theory, while at the same time allowing for explicit computations of the associated link invariant.

To describe this skein module, recall that the type A_{n-1} Hecke algebra H_n can be described as the linear span of braids on n strands modulo *skein relations* and isotopies. The multiplication in H_n is inherited from stacking braids, and the unit is represented by identity braid.

Similarly, the (positive half of the) skein module of the annulus $\mathrm{Sk}_n^+(\mathbb{A})$ is defined as the linear span of annular braid closures modulo skein relations and isotopies. It is easy to see from the definition that $\mathrm{Sk}_n^+(\mathbb{A})$ is isomorphic to the cocenter of the Hecke algebra:

$$\mathrm{Sk}_n^+(\mathbb{A}) \simeq \frac{H_n}{[H_n, H_n]}.$$

Any trace function f on H_n , i.e. a linear function satisfying $f(xy) = f(yx)$, naturally factors through the cocenter, and hence can be viewed as a function on $\mathrm{Sk}_n^+(\mathbb{A})$.

Let Λ_q denote the ring of symmetric functions in infinitely many variables over $\mathbb{C}(q)$, and let $\Lambda_q^{(n)}$ denote the subspace of degree n symmetric functions. The skein module of the annulus enjoys the following properties:

- (a) $\mathrm{Sk}_n^+(\mathbb{A})$ is isomorphic to $\Lambda_q^{(n)}$. It has a basis of Schur functions s_λ labeled by partitions λ with n boxes.
- (b) The HOMFLY-PT invariant of links yields a trace on $H_n \rightarrow \mathbb{C}(q)[a, a^{-1}]$ (called the Jones-Ocneanu trace), and can be computed by projecting H_n to $\mathrm{Sk}_n^+(\mathbb{A}) \cong \Lambda_q^{(n)}$ and applying a certain algebra map $\Lambda_q \rightarrow \mathbb{C}(q)[a, a^{-1}]$.
- (c) The center of H_n naturally acts on its cocenter. On the level of annular link diagrams this corresponds to cutting open the annular link diagram and inserting a central element before closing it again. In particular, the full twist is central in the braid group and hence acts on $\mathrm{Sk}_n^+(\mathbb{A})$.

Links in $S^1 \times D^2$ may be studied by means of their diagrams in \mathbb{A} , after choosing a homeomorphism $S^1 \times D^2 \cong \mathbb{A} \times I$. Such a homeomorphism will be referred to as an *I-bundle structure* on $S^1 \times D^2$. A *framing* is a choice of *I*-bundle structure up to isotopy. A choice of *I*-bundle structure gives us a well-defined link diagram associated to generic links L , whereas a choice of framing determines a diagram only up to Reidemeister moves. Two different framings are related by some number of twists, which on the level of link diagrams corresponds to the insertion of some power of the full-twist braid, as in (c) above.

For this reason, if one is interested in (say, the positive half) of the skein module of a 3-manifold Y which is homeomorphic to $S^1 \times D^2$ (but with no preferred homeomorphism) then it is necessary to understand not just the skein module $\mathrm{Sk}_n^+(\mathbb{A})$, but also the automorphism of full twist insertion.

The categorification of the skein module proceeds in several steps. First, the Hecke algebra H_n is categorified by the monoidal category SBim_n of Soergel bimodules in type A_{n-1} , or by a closely related monoidal category of webs and foams defined by Queffelec–Rose [QR16, Remark 3.24], see [MV10], [Wed19, Remark 3.3], and [RW16, Section 4.4] for the connection. In the second step one must categorify the cocenter of H_n . Traditionally (see Queffelec–Rose [QR18], Beliakova–Putyra–Wehrli [BPW19], and Queffelec–Rose–Sartori [QRS18]) this is done using the underived horizontal trace Tr_0 . This underived horizontal trace is satisfactory for many purposes. For instance Queffelec–Rose–Sartori proved in [QRS18] that the triply-graded Khovanov–Rozansky homology KhR [KR08, Kho07] factors through the underived horizontal trace, which gives a categorification of (b) above. Additionally, in [GW19] the first and third authors connected the annular Khovanov–Rozansky invariant of Queffelec–Rose [QR18] to the underived horizontal trace of type A Soergel bimodules $\mathrm{Tr}_0(\mathrm{SBim}_n)$ and showed that a categorification of (a) holds upon Karoubi completion.

Note that $\mathrm{Tr}_0(\mathrm{SBim}_n)$ indeed categorifies $\Lambda_q^{(n)}$, as it is generated by objects \mathbb{S}^λ as in (2), which correspond to Schur functions s_λ . So the Grothendieck group of $\mathrm{Tr}_0(\mathrm{SBim}_n)$ is naturally isomorphic to $\Lambda_q^{(n)}$.

For a categorification of the skein module of the solid torus, we also need automorphisms of the target category which realise changes in *I*-bundle structure,

as in property (c) above. The following example shows that the ordinary annular Khovanov–Rozansky link invariant, which is constructed using the underived horizontal trace, does not enjoy this property.

Example 1.9. The annular Khovanov–Rozansky invariant of a 2-component unlink decomposes into two non-trivial direct summands $\text{AKhR}(\mathbb{1}_2) \cong S^2 \oplus \wedge^2$. Here S^2 and \wedge^2 denote the Schur objects $\mathbb{S}^{(2)}$ and $\mathbb{S}^{(1,1)}$ as in (2). Twisting the I -bundle structure turns the unlink into an annular Hopf link—the braid closure of the full twist on two strands—whose invariant is a chain complex

$$\text{AKhR}(\text{FT}_2) \simeq \underline{\wedge^2(-2)} \oplus \wedge^2 \oplus \left(\underline{0} \longrightarrow \wedge^2 \xrightarrow{x_1 - x_2} S^2(2) \right).$$

which decomposes into three non-trivial direct summands. Changes in I -bundle structure do not induce isomorphisms on annular Khovanov–Rozansky invariant. Here we have used the version of AKhR defined in [GW19], but the same argument applies to all other constructions employing the horizontal trace.

To remedy this issue, we use the derived horizontal trace.

Definition 1.10. We define the *derived annular Khovanov–Rozansky invariant* of a braid word $\underline{\beta}$ on n strands, denoted $\text{AKhR}_{\text{dg}}(\underline{\beta})$, to be the derived horizontal trace class of the Rouquier complex of $\underline{\beta}$ in $\text{Tr}(\text{SBim}_n)$.

By Theorem 1.7, this invariant can be considered as taking values in perfect dg modules over the dg algebra $\mathbb{C}[x_1, \dots, x_n, \theta_1, \dots, \theta_n] \rtimes \mathbb{C}[S_n]$, where S_n is supported in cohomological degree zero and the variables x_i and θ_i have cohomological degree 0 and -1 respectively. Unlike for AKhR , changes in I -bundle structure induce automorphisms on AKhR_{dg} . These arise naturally through the action of the derived central Rouquier complex of the full twist braid on the derived horizontal trace.

Example 1.11. The derived annular Khovanov–Rozansky invariant of the full twist is a twisted complex

$$\text{AKhR}_{\text{dg}}(\text{FT}_2) \simeq \underline{\wedge^2(-2)} \oplus \left(\underline{\wedge^2} \begin{array}{c} \xrightarrow{x_1 - x_2} \\ \xrightarrow{\theta_1 - \theta_2} \end{array} \wedge^2 \xrightarrow{x_1 - x_2} S^2(2) \right).$$

Like $\text{AKhR}_{\text{dg}}(\mathbb{1}_2) \cong S^2 \oplus \wedge^2$, this has two indecomposable direct summands. The action of the derived central full twist on AKhR_{dg} sends \wedge^2 to $\wedge^2(-2)$ and S^2 to the twisted complex shown as the second direct summand above.

Remark 1.12. The indecomposable summands of $\text{AKhR}_{\text{dg}}(\text{FT}_n)$ are nothing but the images of indecomposable summands of $\text{AKhR}_{\text{dg}}(\mathbb{1}_n)$ (that is, \mathbb{S}^λ for partitions λ of n) under the action of the full twist. Following the conjectures of the first author, Neguț and Rasmussen [GNR16], we expect the action of the full twist to be closely related to the action of Bergeron–Garsia operator ∇ originating in theory of Macdonald polynomials [BGHT99].

In future work, we will use $\mathrm{AKhR}_{\mathrm{dg}}$ to study *cabling operations* for Khovanov–Rozansky link homologies. We also anticipate that the technology of derived traces will be useful in the program to categorify skein algebras, see e.g. [QW18], and for explicit computations of the vector space-valued 4-manifold invariants derived from Khovanov–Rozansky link homology [MWW19].

1.4. Comparison with character sheaves. In this section we briefly compare our results to the theory of character sheaves. Let G be a semi-simple split algebraic group with a Borel subgroup B and Weyl group W . Recall that a geometric categorification of the Hecke algebra for W is given by the category of B -equivariant constructible sheaves on the flag variety G/B , or, equivalently, B -biequivariant sheaves on G . Other, very similar versions of the Hecke category include B -equivariant D -modules on G/B , or Harish-Chandra bimodules. By the work of Soergel the geometric Hecke category is closely related to the category of Soergel bimodules. The polynomial ring \mathbb{R} corresponds to the B -equivariant cohomology of a point. For more details, see [Soe90].

In [BZFN10, BN09] both the trace and the center of the geometric Hecke category were identified with the category of Lusztig’s character sheaves [Lus84]. In particular, the object $\mathrm{Tr}(\mathbb{1})$ which plays a prominent role in the paper corresponds to the so-called Springer sheaf. The derived endomorphism algebra of the Springer sheaf is known (in particular, it is isomorphic to $\mathbb{C}[W]$ in degree zero), and the formality result similar to Theorem 1.3 was proved by Rider [Rid13]. It is important to mention that the results of [Rid13] hold in the category of *mixed perverse sheaves* which is equipped with an additional grading. See also [PVdB19, eq. 0.0.4] for a related formality result.

Finally, it is known [Lus84, RR16] that in type A the summands of the Springer sheaf generate the category of character sheaves, while this is not the case in other types.

Note that we do not claim any results about the Drinfeld center of SBim_n or the corresponding category of complexes, but plan to compute it in the future work.

1.5. Comparison with Hilbert scheme of points. In [GNR16] the first author, Neguț and Rasmussen proposed a set of conjectures relating the category of Soergel bimodules to the Hilbert scheme of points on the plane $\mathrm{Hilb}^n(\mathbb{C}^2)$. In particular, they conjectured that both the trace and the center of $\mathcal{K}^b(\mathrm{SBim}_n)$ are closely related to the derived category of coherent sheaves on $\mathrm{Hilb}^n(\mathbb{C}^2)$. We plan to work out the precise connection between this work and [GNR16] in the future, and only comment on one remarkable formal similarity.

Haiman constructed in [Hai01] a rank $n!$ vector bundle \mathcal{P} on $\mathrm{Hilb}^n(\mathbb{C}^2)$ called *the Procesi bundle*. Its endomorphism algebra has the form

$$(3) \quad \mathrm{Hom}(\mathcal{P}, \mathcal{P}) = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] \rtimes \mathbb{C}[S_n], \quad \mathrm{Ext}^i(\mathcal{P}, \mathcal{P}) = 0, \quad i > 0.$$

It is known that the direct summands of \mathcal{P} generate the derived category of $\mathrm{Hilb}^n(\mathbb{C}^2)$, and hence the functor

$$\mathrm{RHom}(\mathcal{P}, -): D^b(\mathrm{Hilb}^n(\mathbb{C}^2)) \rightarrow D^b(\mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] \rtimes \mathbb{C}[S_n]\text{-mod})$$

is an equivalence [BKR01]. The equation (3) is very similar to the endomorphism algebra of $\mathrm{Tr}(\mathbb{1}_n)$ appearing in Theorem 1.3, but the odd variables θ_i are replaced by the even variables y_i . It is likely that the dg enhancement of $D^b(\mathrm{Hilb}^n(\mathbb{C}^2))$ is related to the horizontal trace of SBim_n by some kind of Koszul duality.

1.6. Organization of the paper. In section 2 we set up notation and conventions for differential graded (dg) categories and functors between them. Throughout the paper we chose to avoid any discussion of A_∞ categories and A_∞ functors, so we use the formalism of quasi-functors instead (see subsection 2.3). We also review the notion of formality for dg algebras and its relation to Massey products, see subsection 2.5.

In section 3 and 4 we discuss various notions of completion of dg categories with respect to direct sums, cones and homotopy idempotents. In particular, we define the pretriangulated hull and the dg Karoubi completion for an abstract dg category. This material is quite standard, and can be found, for example, in Seidel’s book [Sei08], which we more or less follow; however, we decided to present it as concretely as possible for the readers’ convenience. In particular, we avoid Yoneda embeddings altogether and explicitly construct A_∞ lifts of homotopy idempotents (Proposition 4.15) which allows us to give a dg model for the Karoubi completion.

In section 5 we define and study the 2-sided bar complex of a dg category, its cyclic version and Hochschild homology of a dg category. In subsection 5.5 we compare the full Hochschild homology of the dg category with its vertical trace. The main result of this section is Theorem 5.19 where we prove that if a dg category admits a semiorthogonal decomposition then its cyclic bar complex retracts onto the direct sum of cyclic bar complexes for summands. This is a dg version of a result of Kuznetsov [Kuz09] on additivity of Hochschild homology in semiorthogonal decompositions.

The next section 6 is the technical core of the paper. We define and study the derived Drinfeld center and derived horizontal trace for monoidal dg categories. We prove Theorem 1.2 on the universal trace functor $\mathrm{Tr}: \mathcal{C} \rightarrow \mathrm{Tr}(\mathcal{C})$ and its properties, and also define an action of the derived center on the derived trace. Note that $\mathrm{Tr}(\mathcal{C})$ is usually not pretriangulated or idempotent complete, but the results of section 5 allow us to consider the corresponding completions.

In section 7 we apply this machinery to the monoidal category of Soergel bimodules and prove Theorem 1.3.

In section 8 we prove Theorem 1.7 and describe an explicit “annular simplification” algorithm which allows us to identify the trace of any type A Soergel bimodule with a homotopy summand in the direct sum of several copies of $\mathrm{Tr}(\mathbb{1})$. We also discuss the connections of the derived trace with annular Khovanov-Rozansky invariants and the work of the first and third author [GW19]. In particular, we construct a “forgetful functor” from the derived to the “underived” horizontal trace, and show

that Khovanov-Rozansky homology of a braid closure factors through it, see Theorem 8.24.

ACKNOWLEDGEMENTS

The authors would like to thank Anna Beliakova, Roman Bezrukavnikov, Tudor Dimofte, Ben Elias, Tina Kanstrup, Oscar Kivinen, Andrei Neguț, Alexei Oblomkov, Jacob Rasmussen, David Reutter, Raphaël Rouquier, Lev Rozansky, Sarah Scherotzke, Catharina Stroppel, Kostya Tolmachov and Geordie Williamson for many useful discussions.

This work also profited from discussions during the AIM workshop “Categorified Hecke algebras, link homology, and Hilbert schemes”, the ESI Workshop “Categorification in quantum topology and beyond”, the workshop “Hidden Algebraic Structures in Topology” at Caltech, the workshop “Hilbert schemes, categorification and combinatorics” at UC Davis, and the MSRI programs “Quantum Symmetries” and “Higher Categories and Categorification”. We thank the host institutions and sponsors of these activities for their support.

FUNDING

E.G. was partially supported by NSF grants DMS-1700814 and DMS-1760329. The work of E. G. in Section 8 was partially supported by the RSF grant 16-11-10018.

P.W. was partially supported by the Australian Research Council grants ‘Braid groups and higher representation theory’ DP140103821 and ‘Low dimensional categories’ DP160103479 while at the Australian National University during early stages of this project.

2. FACTS FROM HOMOLOGICAL ALGEBRA

This section serves to recall important notions from homological algebra and to fix notation and conventions.

2.1. Complexes. Let \mathbb{k} be a commutative ring. The category of complexes of \mathbb{k} -modules will be denoted $\text{Ch}(\mathbb{k})$. Its objects are complexes of \mathbb{k} -modules, also called *dg \mathbb{k} -modules*, and we will use the cohomological convention for differentials:

$$\dots \xrightarrow{d} X^k \xrightarrow{d} X^{k+1} \xrightarrow{d} \dots$$

In particular, the differentials are considered to be of cohomological degree 1. The morphism spaces between objects X and Y in $\text{Ch}(\mathbb{k})$ are the complexes with

$$\text{Hom}_{\text{Ch}(\mathbb{k})}^k(X, Y) = \prod_{i \in \mathbb{Z}} \text{Hom}_{\mathbb{k}}(X^i, Y^{i+k}), \quad d_{\text{Hom}_{\text{Ch}(\mathbb{k})}(X, Y)}(f) := d_Y \circ f - (-1)^{|f|} f \circ d_X$$

where $|f|$ denotes the cohomological degree of f . The full subcategory of bounded complexes will be denoted $\text{Ch}^b(\mathbb{k})$. The categories $\text{Ch}(\mathbb{k})$ and $\text{Ch}^b(\mathbb{k})$ are symmetric monoidal, with the tensor product defined on objects as

$$(X \otimes_{\mathbb{k}} Y)^k = \bigoplus_{i+j=k} X^i \otimes_{\mathbb{k}} Y^j, \quad d_{X \otimes_{\mathbb{k}} Y} = d_X \otimes \text{Id}_Y + \text{Id}_X \otimes d_Y$$

and on morphisms f, g by $(f \otimes g)(x \otimes y) = (-1)^{|x||g|} f(x) \otimes g(y)$, with braiding

$$\tau_{X,Y}: X \otimes_{\mathbb{k}} Y \rightarrow Y \otimes_{\mathbb{k}} X, \quad \tau_{X,Y}(x \otimes y) = (-1)^{|x||y|} y \otimes x.$$

Finally, for a complex X and $l \in \mathbb{Z}$, we denote by $\Sigma^l X$ the complex with

$$(\Sigma^l X)^k = X^{k+l}, \quad d_{\Sigma^l X} = (-1)^l d_X.$$

In particular, for $l > 0$, the translation Σ^l shifts the complex X to the left by l steps.

2.2. Differential graded categories. A differential \mathbb{Z} -graded \mathbb{k} -linear category \mathcal{C} , or short a *dg category* is a category enriched in $\text{Ch}(\mathbb{k})$. This means every morphism space $\text{Hom}_{\mathcal{C}}(X, Y)$ is an object in $\text{Ch}(\mathbb{k})$ and composition of morphisms forms chain maps

$$\text{Hom}_{\mathcal{C}}(Y, Z) \otimes_{\mathbb{k}} \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{C}}(X, Z)$$

which means that the differentials satisfy the Leibniz rule with respect to composition: $d_{\mathcal{C}}(f \circ g) = d_{\mathcal{C}}(f) \circ g + (-1)^{|f|} f \circ d_{\mathcal{C}}(g)$. Other abelian groups besides \mathbb{Z} can be used for gradings, and later we will consider $\mathbb{Z} \times \mathbb{Z}$ -graded complexes with differentials of degree $(0, 1)$. In any case, the cohomological degree of a homogenous morphism f will be denoted by $|f|$.

Example 2.1. Any \mathbb{k} -linear category can and will be regarded as a dg category with morphism complexes concentrated in cohomological degree zero, thus necessarily with zero differential.

Example 2.2. For any \mathbb{k} -linear category \mathcal{A} , the category $\text{Ch}(\mathcal{A})$ (resp. $\text{Ch}^b(\mathcal{A})$) of (bounded) complexes in \mathcal{A} (whose definition mimics the one of $\text{Ch}(\mathbb{k})$ (resp. $\text{Ch}^b(\mathbb{k})$)) is a dg category.

Example 2.3. Any (dg) \mathbb{k} -algebra A can and will be regarded as a dg category with one object $*$ and $\text{End}_A(*) = A$.

A morphism $f \in \text{Hom}_{\mathcal{C}}(X, Y)$ is said to be *closed* if $d_{\mathcal{C}}(f) = 0$ and *exact* or *null-homotopic* if $f = d_{\mathcal{C}}(h)$ for some $h \in \text{Hom}_{\mathcal{C}}(X, Y)$, which in this case is called a *null-homotopy* for f . For $f, g \in \text{Hom}_{\mathcal{C}}^k(X, Y)$, we write $f \simeq g$ and say f and g are *homotopic* if $f - g$ is null-homotopic.

The *cohomology category* of \mathcal{C} , denoted $H^0(\mathcal{C})$ is defined to be the additive category with the same objects as \mathcal{C} , and with

$$\text{Hom}_{H^0(\mathcal{C})}(X, Y) := \frac{\{f \in \text{Hom}_{\mathcal{C}}^0(X, Y) \mid d_{\mathcal{C}}(f) = 0\}}{d_{\mathcal{C}}(\text{Hom}_{\mathcal{C}}^{-1}(X, Y))}$$

By *isomorphism* in \mathcal{C} we mean degree zero closed invertible morphisms. If there exists an isomorphism in $\text{Hom}_{\mathcal{C}}(X, Y)$, we write $X \cong Y$. A degree zero closed morphism $f \in \text{Hom}_{\mathcal{C}}(X, Y)$ is said to be a *homotopy equivalence* if it induces an isomorphism in

$H^0(\mathcal{C})$, and in this case we write $X \simeq Y$ and say X and Y are homotopy equivalent. If $X \simeq 0$, then we say X is *contractible*.

For a dg category \mathcal{C} , we denote by \mathcal{C}^{op} the dg category with the same objects as \mathcal{C} and with $\text{Hom}_{\mathcal{C}^{\text{op}}}(X, Y) := \text{Hom}_{\mathcal{C}}(Y, X)$, where the composite of $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(X, Y)$ with $g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, Z)$, denoted $g \circ_{\mathcal{C}^{\text{op}}} f$, is given by $(-1)^{|f||g|} f \circ_{\mathcal{C}} g$.

For two dg categories \mathcal{C}, \mathcal{D} we denote by $\mathcal{C} \otimes_{\mathbb{k}} \mathcal{D}$ the category with objects given by pairs of objects (X, Y) for $X \in \mathcal{C}$ and $Y \in \mathcal{D}$ and morphisms given by complexes

$$\text{Hom}_{\mathcal{C} \otimes_{\mathbb{k}} \mathcal{D}} \left((X, Y), (X', Y') \right) := \text{Hom}_{\mathcal{C}}(X, X') \otimes_{\mathbb{k}} \text{Hom}_{\mathcal{D}}(Y, Y')$$

with composition

$$(f \otimes g) \circ (f' \otimes g') := (-1)^{|g||f'|} (f \circ f') \otimes (g \circ g').$$

Let \mathcal{C} be a dg category. A *subcategory* $\mathcal{J} \subset \mathcal{C}$ is a collection of objects $\text{Obj}(\mathcal{J}) \subset \text{Obj}(\mathcal{C})$ with hom spaces $\text{Hom}_{\mathcal{J}}(X, X')$ for $X, X' \in \text{Obj}(\mathcal{J})$ being subcomplexes $\text{Hom}_{\mathcal{C}}(X, X')$, which are closed under composition. The subcategory $\mathcal{J} \subset \mathcal{C}$ is *full* if $\text{Hom}_{\mathcal{J}}(X, X') = \text{Hom}_{\mathcal{C}}(X, X')$ for all $X, X' \in \text{Obj}(\mathcal{J})$. The subcategory $\mathcal{J} \subset \mathcal{C}$ is *unital* if $\text{Id}_X \in \mathcal{J}$ whenever $X \in \text{Obj}(\mathcal{J})$. Henceforth all subcategories are unital.

Example 2.4. We will denote by $\mathcal{J} \subset \mathcal{C}$ the subcategory spanned by the identity maps in \mathcal{C} . More generally, if $\mathcal{B} \subset \mathcal{C}$ is a full subcategory, then we have $\mathcal{J}_{\mathcal{B}} \subset \mathcal{C}$, the (unital, but not full) subcategory spanned by the identity morphisms in \mathcal{C} which are contained in \mathcal{B} .

2.3. DG functors. If \mathcal{B}, \mathcal{C} are dg categories, a dg functor $F: \mathcal{B} \rightarrow \mathcal{C}$ is a functor whose action on hom complexes $\text{Hom}_{\mathcal{B}}(X, Y) \rightarrow \text{Hom}_{\mathcal{C}}(F(X), F(Y))$ is a degree zero chain map. The collection of dg functors $\mathcal{B} \rightarrow \mathcal{C}$ itself forms a dg category. Objects of this functor category are functors, and morphisms are natural transformations, as defined next. If F, G are functors $\mathcal{B} \rightarrow \mathcal{C}$, a natural transformation $\alpha: F \rightarrow G$ of degree k is an assignment $X \mapsto \alpha_X \in \text{Hom}_{\mathcal{C}}^k(F(X), G(X))$ such that

$$G(f) \circ \alpha_X = (-1)^{k|f|} \alpha_Y \circ F(f)$$

for all morphisms $f \in \text{Hom}_{\mathcal{B}}(X, Y)$. The differential of α by definition sends $X \mapsto d_{\mathcal{C}}(\alpha_X)$ (the naturality of $d(\alpha_X)$ so defined follows from the Leibniz rule).

By an *isomorphism of dg functors* we mean a degree zero closed invertible natural transformation of dg functors. If $F, G: \mathcal{B} \rightarrow \mathcal{C}$ are isomorphic, we write $F \cong G$.

Any dg functor $F: \mathcal{B} \rightarrow \mathcal{C}$ naturally induces a functor between the corresponding homotopy categories $H^0(F): H^0(\mathcal{B}) \rightarrow H^0(\mathcal{C})$.

A dg functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is an *equivalence* of dg categories if there is a dg functor $G: \mathcal{D} \rightarrow \mathcal{C}$ such that $F \circ G \cong \text{Id}_{\mathcal{D}}$ and $G \circ F \cong \text{Id}_{\mathcal{C}}$.

A dg functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is *quasi-fully faithful*, if restricts to quasi-isomorphisms on hom complexes, i.e. for every pair of objects X, Y in \mathcal{C} , the induced map $F^*: H^*(\text{Hom}_{\mathcal{C}}(X, Y)) \rightarrow H^*(\text{Hom}_{\mathcal{D}}(F(X), F(Y)))$ is an isomorphism; it is *quasi-essentially*

surjective if the functor $H^0(F)$ between the respective homotopy categories is essentially surjective. If F is quasi-fully faithful and quasi-essentially surjective, then it is called a *quasi-equivalence*.

For many applications in this paper we will need a weaker notion of a functor between dg categories. If dgcats is the category where the objects are all (small) dg categories and the morphisms are all dg functors, one can define [Tab05, Toë07] the category \mathbf{Hqe} as a localization of dgcats with respect to quasi-equivalences. More abstractly, dgcats has a model category structure whose weak equivalences are the quasi-equivalences, and \mathbf{Hqe} is the corresponding localization.

Two dg categories \mathcal{C} and \mathcal{D} are called *quasi-equivalent* if there exist dg categories $\mathcal{B}_1, \dots, \mathcal{B}_n$ and a chain of quasi-equivalences

$$\mathcal{C} \leftarrow \mathcal{B}_1 \rightarrow \dots \leftarrow \mathcal{B}_n \rightarrow \mathcal{D}.$$

Then \mathcal{C} and \mathcal{D} are quasi-equivalent if and only if they are isomorphic in \mathbf{Hqe} .

A *quasi-functor* between two dg categories is a morphism in \mathbf{Hqe} . For example, if we have a dg functor $F: \mathcal{C} \rightarrow \mathcal{B}$ and a quasi-equivalence $G: \mathcal{D} \rightarrow \mathcal{B}$ then F induces a quasi-functor from \mathcal{C} to \mathcal{D} . Since quasi-equivalences induces equivalences of homotopy categories, a quasi-functor between dg categories \mathcal{C} and \mathcal{D} induces an honest functor between the homotopy categories $H^0(\mathcal{C}) \rightarrow H^0(\mathcal{D})$.

Remark 2.5. Instead of working with \mathbf{Hqe} and quasi-functors, one could choose to work with A_∞ functors between dg categories. Over a field \mathbb{k} , this is essentially an equivalent viewpoint since every quasi-equivalence admits an inverse, which is in general not a dg functor but an A_∞ functor [Sei08]. However, we decided to stay away from A_∞ functors in this paper.

2.4. Bimodules. If \mathcal{C} and \mathcal{D} are dg categories, a \mathcal{D}, \mathcal{C} -bimodule \mathcal{M} is the data of

- for each pair of objects $Y \in \mathcal{D}, X \in \mathcal{C}$, a dg \mathbb{k} -module $Y\mathcal{M}X$.
- for each quadruple of objects $Y, Y' \in \mathcal{D}$ and $X, X' \in \mathcal{C}$, action maps

$$\text{Hom}_{\mathcal{D}}(Y, Y') \otimes_{\mathbb{k}} Y\mathcal{M}X \otimes_{\mathbb{k}} \text{Hom}_{\mathcal{C}}(X', X) \rightarrow Y'\mathcal{M}X'$$

satisfying the usual associativity constraints.

The action maps are required to be chain maps of degree zero. This is equivalent to $|f \cdot m \cdot g| = |f| + |m| + |g|$ and the Leibniz rule

$$d_{\mathcal{M}}(f \cdot m \cdot g) = d_{\mathcal{D}}(f) \cdot m \cdot g + (-1)^{|f|} f \cdot d_{\mathcal{M}}(m) \cdot g + (-1)^{|f|+|m|} f \cdot m \cdot d_{\mathcal{C}}(g)$$

for all $f \in \text{Hom}_{\mathcal{D}}(Y, Y'), m \in Y\mathcal{M}X, g \in \text{Hom}_{\mathcal{C}}(X', X)$.

The notation ${}_{\mathcal{D}}\mathcal{M}_{\mathcal{C}}$ will be used to indicate that \mathcal{M} is a \mathcal{D}, \mathcal{C} -bimodule.

A left \mathcal{C} -module is the same as a \mathcal{C}, \mathbb{k} -bimodule, and a right \mathcal{C} -module is the same as a \mathbb{k}, \mathcal{C} -bimodule, by definition.

Example 2.6. If \mathcal{C} is a dg category and $X, Y \in \mathcal{C}$ are objects, then we denote

$$Y\mathcal{C}X := \text{Hom}_{\mathcal{C}}(X, Y), \quad \mathcal{C}X := \bigoplus_Y Y\mathcal{C}X, \quad Y\mathcal{C} := \bigoplus_X Y\mathcal{C}X.$$

The composition of morphisms in \mathcal{C} equips $\mathcal{C}X$ with the structure of a left \mathcal{C} -module, $Y\mathcal{C}$ with the structure of a right \mathcal{C} -module (called the *Yoneda modules*), and $\mathcal{C} = \bigoplus_{X,Y} Y\mathcal{C}X$ with the structure of a \mathcal{C} , \mathcal{C} -bimodule (the *regular bimodule*).

We will use the Yoneda modules and the regular bimodule to streamline notation in certain places.

Remark 2.7. For dg categories \mathcal{C} , \mathcal{D} , the following notions are equivalent:

- (1) \mathcal{D} , \mathcal{C} -bimodules,
- (2) left $\mathcal{D} \otimes_{\mathbb{k}} \mathcal{C}^{\text{op}}$ -modules,
- (3) right $\mathcal{C} \otimes_{\mathbb{k}} \mathcal{D}^{\text{op}}$ -modules,
- (4) functors $\mathcal{D} \otimes_{\mathbb{k}} \mathcal{C}^{\text{op}} \rightarrow \mathbb{k}\text{-dgmod}$.

However, such identifications necessarily involve choices and hidden signs; for this reason, we will typically not use them.

Given dg categories \mathcal{B} , \mathcal{C} , \mathcal{D} and bimodules ${}_{\mathcal{D}}\mathcal{M}_{\mathcal{C}}$, ${}_{\mathcal{C}}\mathcal{N}_{\mathcal{B}}$, their tensor product ${}_{\mathcal{D}}(\mathcal{M} \otimes_{\mathcal{C}} \mathcal{N})_{\mathcal{B}}$ is the bimodule with

$$Z(\mathcal{M} \otimes_{\mathcal{C}} \mathcal{N})X := \bigoplus_{Y \in \mathcal{C}} \left(ZMY \otimes YNX \right) / \sim,$$

where \sim is the equivalence relation $(m \cdot f) \otimes n \sim m \otimes (f \cdot n)$ for all $m \in ZMY$, $f \in Y\mathcal{C}Y'$, $n \in Y'NX$.

Remark 2.8. The category of \mathcal{C} , \mathcal{C} -bimodules is monoidal with tensor product as defined above, and monoidal identity given by the regular bimodule \mathcal{C} .

Remark 2.9. We often regard \mathcal{C} and \mathcal{D} as the (very big) non-unital algebras

$$\mathcal{C} = \bigoplus_{X', X \in \mathcal{C}} X'\mathcal{C}X, \quad \mathcal{D} = \bigoplus_{Y', Y \in \mathcal{D}} Y'\mathcal{D}Y,$$

and a \mathcal{D} , \mathcal{C} -bimodule \mathcal{M} as the (very big) dg bimodule

$$\mathcal{M} = \bigoplus_{Y \in \mathcal{D}, X \in \mathcal{C}} YMX.$$

In this language an object $X \in \mathcal{C}$ corresponds to the distinguished idempotent Id_X in the big algebra $\bigoplus_{X', X} X'\mathcal{C}X$. In this way, essentially all of one's intuition from the usual world of algebras and bimodules carries over into the world of dg categories and their bimodules.

2.5. A_{∞} algebras and deformation retracts. Recall, an A_{∞} algebra is a graded \mathbb{k} -module A equipped with maps $\mu_n : A^{\otimes n} \rightarrow A$ degree $2 - n$, $n \geq 1$, satisfying the following family of identities for $M \geq 1$

$$\sum_{M=r+s+t} (-1)^{r+st} \mu_{r+1+t}(\text{Id}^{\otimes r} \otimes \mu_s \otimes \text{Id}^{\otimes t}) = 0.$$

In particular a dg algebra is an A_∞ algebra in which μ_n vanish for $n \neq 1, 2$. In this case $\mu_1 : A \rightarrow A$ is the differential and $\mu_2 : A \otimes A \rightarrow A$ is honestly associative and satisfies the Leibniz rule

$$\mu_1 \circ \mu_2 - \mu_2 \circ (\mu_1 \otimes \text{Id}_A) - \mu_2 \circ (\text{Id}_A \otimes \mu_1) = 0$$

Suppose $X, Y \in \mathcal{C}$ are objects in a dg category. A deformation retract $X \rightarrow Y$ is the data of closed degree zero morphisms $\pi : X \rightarrow Y$, $\sigma : Y \rightarrow X$ and a degree -1 homotopy $h \in \text{End}^{-1}(X)$ such that

$$\pi \circ \sigma = \text{Id}_Y, \quad d(h) = \text{Id}_X - \sigma \circ \pi, \quad h \circ \sigma = 0 = \pi \circ h.$$

The following is well known [Kel01, Kad80, Mer99].

Theorem 2.10. *If A is a dg algebra and V is a dg \mathbb{k} -module, then any deformation retract $A \rightarrow V$ gives V the structure of an A_∞ algebra quasi-isomorphic to A .*

Proof. The construction follows [Mer99]. Let $m : A \otimes A \rightarrow A$ be the multiplication in A . We define recursively a sequence of maps $\lambda_k : V^{\otimes k} \rightarrow A$ by $\lambda_2 = m(\sigma \otimes \sigma)$ and

$$\lambda_n = -m(\sigma \otimes h\lambda_{n-1}) + \sum_{s=2}^{n-2} (-1)^{s+1} m(h\lambda_s \otimes h\lambda_{n-s}) + (-1)^{n+1} m(h\lambda_{n-1} \otimes \sigma), \quad n \geq 3$$

Then $\mu_n = \pi \circ \lambda_n$ defines the structure of a strictly unital A_∞ algebra on V . \square

Remark 2.11. Suppose that A and V have an additional grading which is preserved by the differential and the maps σ, π and h . Then the A_∞ structure maps on V can be chosen to preserve this grading as well.

Remark 2.12. If $R \subset A$ is a commutative dg subalgebra and V admits the structure of a dg R -module, and the data of the deformation retract (that is σ, π and h) can be chosen to be R -linear, then the A_∞ structure maps on V can be chosen to be R -linear, in the sense that they descend to the quotient

$$\mu_n : V \otimes_R V \otimes_R \cdots \otimes_R V \rightarrow V.$$

This elementary fact is often very useful.

If \mathbb{k} is a field, then A deformation retracts onto its homology $H(A)$ (regarded as a dg \mathbb{k} -module with zero differential), and so $H(A)$ inherits the structure of an A_∞ algebra with $\mu_1 = 0$. A differential graded algebra A is called *formal* if it is quasi-isomorphic to its homology $H(A)$. The above discussion shows that a dg algebra A over a field \mathbb{k} is formal if and only if the A_∞ structure on $H(A)$ is trivial, that is, $\mu_k = 0$ for $k > 2$.

3. STANDARD DG CATEGORICAL CONSTRUCTIONS

In this and the following section we describe the processes of adjoining finite direct sums, suspension, and twists to a dg category.

A functor $F : \mathcal{C} \rightarrow \mathbb{k}\text{-dgm}$ is said to be *representable* if there is an object $X \in \mathcal{C}$ such that F is isomorphic to $\text{Hom}_{\mathcal{C}}(X, -)$ (or $\text{Hom}_{\mathcal{C}}(-, X)$ if F is contravariant).

The dg category \mathcal{C} is *additive* if for each finite collection of objects $X_i \in \mathcal{C}$ the functor $Y \mapsto \bigoplus_i Y \mathcal{C} X_i$ is representable. This means \mathcal{C} has finite coproducts, for which we use the symbol \oplus .

The dg category \mathcal{C} is *suspended* (or *has suspension*) if for each $X \in \mathcal{C}$ and $l \in \mathbb{Z}$ the functor $Y \mapsto \Sigma^l(Y \mathcal{C} X)$ is representable. This is equivalent to the existence of an object $\Sigma^l X$, an *l-translate*, for every object X of \mathcal{C} , together with a given closed degree l invertible morphism $\Sigma^l X \rightarrow X$.

Let $\alpha \in \text{End}_{\mathcal{C}}^1(X)$ be an endomorphism in \mathcal{C} satisfying the *Maurer–Cartan equation* $d_{\mathcal{C}}(\alpha) + \alpha \circ \alpha = 0$. Then we have a functor $\phi_{\alpha}: \mathcal{C} \rightarrow \mathbb{k}\text{-dgm}$ sending an object Y to the complex $(Y \mathcal{C} X, d_{\alpha})$ with *twisted differential* $d_{\alpha}(f) := d(f) - (-1)^{|f|} f \circ \alpha$.

The category $\text{dg } \mathcal{C}$ is said to *have twists* if for each Maurer–Cartan element α in \mathcal{C} the functors ϕ_{α} is representable.

3.1. Additive suspended envelope. If \mathcal{C} is a dg category, the *additive suspended envelope* $\Sigma \mathcal{C}$ of \mathcal{C} is the dg category whose objects are collections $\{\Sigma^{a_i} X^i\}_{i \in I}$ where $I \subset \mathbb{Z}$ is a finite set, $X^i \in \mathcal{C}$ and $a_i \in \mathbb{Z}$. Morphism complexes in $\Sigma \mathcal{C}$ are by definition

$$\text{Hom}_{\Sigma \mathcal{C}}^l(\{\Sigma^{a_i} X^i\}_{i \in I}, \{\Sigma^{b_j} Y^j\}_{j \in J}) = \prod_{i \in I} \bigoplus_{j \in J} \text{Hom}_{\mathcal{C}}^{l+b_j-a_i}(X^i, Y^j)$$

with differential

$$(4) \quad d_{\Sigma \mathcal{C}}((f_{ji})_{(j,i) \in J \times I}) = ((-1)^{b_j} d_{\mathcal{C}}(f_{ji}))_{(j,i) \in J \times I}.$$

An element of this hom space can be thought of as a $J \times I$ matrix (f_{ji}) of morphisms $f_{ji} \in \text{Hom}_{\mathcal{C}}(X^i, Y^j)$. Composition of morphisms is given by usual matrix multiplication and composition in \mathcal{C} .

There is a canonical fully faithful dg functor $\mathcal{C} \rightarrow \Sigma \mathcal{C}$ defined object-wise by $X \mapsto \{X\}$ (with indexing set I a singleton), and we may identify \mathcal{C} with its image in $\Sigma \mathcal{C}$. It is straightforward to verify that $\Sigma \mathcal{C}$ is additive and suspended (with Σ^l indicating l -translates), and we will henceforth abuse notation by writing

$$\bigoplus_{i \in I} \Sigma^{a_i} X^i := \{\Sigma^{a_i} X^i\}_{i \in I} \in \Sigma \mathcal{C}.$$

We also write 0 for the empty direct sum, corresponding to the case $I = \emptyset$.

It is not hard to check that $\Sigma \Sigma \mathcal{C} \cong \Sigma \mathcal{C}$ and this idempotent property of the assignment $\mathcal{C} \mapsto \Sigma \mathcal{C}$ justifies the name additive suspended envelope.

Remark 3.1. We also define an additive suspended envelope $\Sigma^{\mathbb{I}} \mathcal{C}$ with countable direct products, where the finiteness assumption on the indexing sets $I \subset \mathbb{Z}$ is removed. Homogeneous morphisms in $\Sigma^{\mathbb{I}} \mathcal{C}$ are by definition matrices of morphisms in \mathcal{C} , each row of which has only finitely many nonzero entries.

3.2. Twisted envelope and pretriangulated hull. The *twisted envelope* of \mathcal{C} can be constructed explicitly as follows. Let $\text{Tw}(\mathcal{C})$ be the category with objects $\text{tw}_{\alpha}(X)$

where $X \in \mathcal{C}$ and $\alpha \in \text{End}_{\mathcal{C}}^1(X)$ satisfying $d_{\mathcal{C}}(\alpha) + \alpha^2 = 0$ as above. The morphism complexes in $\text{Tw}(\mathcal{C})$ are by definition

$$\text{Hom}_{\text{Tw}(\mathcal{C})}(\text{tw}_{\alpha}(X), \text{tw}_{\beta}(Y)) := \text{Hom}_{\mathcal{C}}(X, Y)$$

with differential

$$d_{\text{Tw}(\mathcal{C})}(f) = d_{\mathcal{C}}(f) + \beta \circ f - (-1)^{|f|} f \circ \alpha.$$

We say that \mathcal{C} has *twists* if the obvious fully faithful dg functor $\mathcal{C} \rightarrow \text{Tw}(\mathcal{C})$ sending $X \mapsto \text{tw}_0(X)$ is an equivalence. The natural inclusion $\text{Tw}(\mathcal{C}) \rightarrow \text{Tw}(\text{Tw}(\mathcal{C}))$ sending $\text{tw}_{\alpha}(X) \mapsto \text{tw}_0(\text{tw}_{\alpha}(X))$ is an equivalence, with an inverse equivalence $\text{Tw}(\text{Tw}(\mathcal{C})) \rightarrow \text{Tw}(\mathcal{C})$ defined by $\text{tw}_{\beta}(\text{tw}_{\alpha}(X)) \mapsto \text{tw}_{\alpha+\beta}(X)$ (compare with the procedure of taking the total complex of a bicomplex). Thus, $\text{Tw}(\mathcal{C})$ has twists. This idempotent property of the assignment $\mathcal{C} \mapsto \text{Tw}(\mathcal{C})$ justifies our referring to $\text{Tw}(\mathcal{C})$ as the twisted envelope.

Example 3.2. Any additive category \mathcal{A} can be thought of as a dg category with zero differential and trivial grading (all morphisms are placed in degree zero). Then $\text{Tw}(\Sigma\mathcal{A})$ is equivalent to the usual category of bounded complexes $\text{Ch}^b(\mathcal{A})$.

If $f \in \text{Hom}_{\mathcal{C}}^1(X, Y)$ is a degree 1 closed morphism, then the cone of f is the object $\text{tw}_{\alpha}(X \oplus Y)$ with $\alpha = \begin{bmatrix} 0 & 0 \\ f & 0 \end{bmatrix}$ inside $\text{Tw}(\mathcal{C})$. If instead $f: X \rightarrow Y$ is a degree zero closed morphism and \mathcal{C} has suspension, then we first replace f by a degree 1 closed morphism $\Sigma^1 X \rightarrow Y$ and apply the previous construction.

Definition 3.3. We say a dg category \mathcal{C} is *pretriangulated* if it is suspended and closed under taking cones.

The *pretriangulated hull* $\text{Pretr}(\mathcal{C})$ of a dg category \mathcal{C} is the full subcategory of $\text{Tw}(\Sigma\mathcal{C})$ generated by $\Sigma\mathcal{C}$ under taking mapping cones.

It follows from the discussion above that $\text{Pretr}(\mathcal{C})$ is pretriangulated, and \mathcal{C} itself is pretriangulated if and only if the natural embedding $\mathcal{C} \rightarrow \text{Pretr}(\mathcal{C})$ is an equivalence.

Objects in the pretriangulated hull can be expressed as iterated mapping cones of objects in $\Sigma\mathcal{C}$, also known as *one-sided twisted complexes*.

Example 3.4. For $X = \text{tw}_{\alpha}(\bigoplus_{j \in J} \Sigma^{a_j} X^j)$ we can collect terms with equal shifts, i.e. set $Y^i := \bigoplus_{j: a_j = -i} X^j$ for $i \in \mathbb{Z}$, and write $\alpha_{j,i} \in \text{Hom}^{1+i-j}(Y^i, Y^j)$ for the components of the twist. Then X can be illustrated as:

$$\begin{array}{ccccccc}
 & & \alpha_{i+1,i-2} & & \alpha_{i+2,i-1} & & \\
 & & \curvearrowright & & \curvearrowright & & \\
 & & \alpha_{i,i-2} & & \alpha_{i+1,i-1} & & \alpha_{i+2,i} \\
 & & \curvearrowright & & \curvearrowright & & \curvearrowright \\
 \dots & \xrightarrow{\alpha_{i-1,i-2}} & \Sigma^{1-i} Y^{i-1} & \xrightarrow{\alpha_{i,i-1}} & \Sigma^{-i} Y^i & \xrightarrow{\alpha_{i+1,i}} & \Sigma^{-1-i} Y^{i+1} & \xrightarrow{\alpha_{i+2,i+1}} & \dots
 \end{array}$$

The Maurer–Cartan equation for α now is $(-1)^j d_{\mathcal{C}}(\alpha_{j,i}) + \sum_{j>k>i} \alpha_{j,k} \circ \alpha_{k,i} = 0$.

Remark 3.5. Let A be a dg algebra supported in non-positive degrees, considered as a dg category with one object. Then $\text{Pretr}(A) = \text{Tw}(\Sigma A)$ since every twisted complex is one-sided, i.e. an iterated cone.

4. HOMOTOPY IDEMPOTENTS AND THE KAROUBI ENVELOPE

4.1. Homotopy idempotents. Let \mathcal{C} be a dg category. A *homotopy idempotent* in \mathcal{C} is an endomorphism $e \in \text{End}_{\mathcal{C}}^0(X)$ such that $e^2 \simeq e$ (i.e. an idempotent in $H^0(\mathcal{C})$). We say that Y is an *image* of e if there exist closed degree zero morphisms $\sigma: Y \rightarrow X$, $\pi: X \rightarrow Y$ such that

$$\pi \circ \sigma \simeq \text{Id}_Y, \quad \sigma \circ \pi \simeq e.$$

Lemma 4.1. *Images of homotopy idempotents satisfy the following basic properties:*

- (1) Suppose $e_1, e_2 \in \text{End}_{\mathcal{C}}(X)$ are homotopy idempotents with $e_1 \simeq e_2$. If Y_i is an image of e_i ($i = 1, 2$) then $Y_1 \simeq Y_2$. In particular the image of a homotopy idempotent is unique up to homotopy equivalence.
- (2) If Y is an image of a homotopy idempotent $e \in \text{End}_{\mathcal{C}}(X)$ then any homotopy idempotent $e' \in \text{End}_{\mathcal{C}}(Y)$ determines a homotopy idempotent $e'' \in \text{End}_{\mathcal{C}}(X)$ with the property that Z is an image of e' if and only if it is an image of e'' .
- (3) If Y_0, Y_1 are the images of homotopy idempotents $e_i \in \text{End}_{\mathcal{C}}(X_i)$ ($i = 0, 1$) then

$$\text{Hom}_{H^0(\mathcal{C})}(Y_0, Y_1) \cong e_1 \text{Hom}_{H^0(\mathcal{C})}(X_0, X_1)e_0.$$

Proof. Exercise. □

Definition 4.2. We say that \mathcal{C} is *homotopy idempotent complete* if $H^0(\mathcal{C})$ is idempotent complete, i.e. each homotopy idempotent in \mathcal{C} has an image.

Our goal in this section is to construct the *homotopy Karoubi envelope* $\text{Kar}^{dg}(\mathcal{C})$ and prove the following.

Theorem 4.3. *Every dg category \mathcal{C} admits an embedding $\mathcal{C} \hookrightarrow \text{Kar}^{dg}(\mathcal{C})$ into a homotopy idempotent complete dg category characterized up to quasi-equivalence by the following universal property: if \mathcal{D} is a homotopy idempotent complete dg category equipped with a dg functor $\mathcal{C} \rightarrow \mathcal{D}$, then there is unique morphism (quasi-functor) $\text{Kar}^{dg}(\mathcal{C}) \rightarrow \mathcal{D}$ in \mathbf{Hqe} such that the following diagram commutes:*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\quad} & \mathcal{D} \\ \downarrow & \nearrow \text{---} & \\ \text{Kar}^{dg}(\mathcal{C}) & & \end{array}$$

Furthermore, \mathcal{C} is idempotent complete if and only if the canonical functor $\mathcal{C} \rightarrow \text{Kar}^{dg}(\mathcal{C})$ is a quasi-equivalence.

4.2. A_∞ idempotents. Note that if $e_0 \in \text{End}_{\mathcal{C}}^0(X)$ is a homotopy idempotent and $h_0 \in \text{End}_{\mathcal{C}}^{-1}(X)$ satisfies $d_{\mathcal{C}}(h_0) = e_0 \circ (\text{Id}_X - e_0)$, then $e_0 \circ h_0 - h_0 \circ e_0$ is automatically closed. This morphism obstructs certain constructions, and it is natural to require it to be null-homotopic (as we will see below, one can choose a homotopy h_0 such that this holds, but not every homotopy satisfies this condition), via an endomorphism we will denote e_1 . There is a higher family of obstructions which is natural to require to be trivial, via homotopies e_k, h_k for $k \in \mathbb{Z}_{\geq 1}$. This results in the notion of an A_∞ idempotent or idempotent up to coherent homotopy, which we describe next.

Definition 4.4. An A_∞ idempotent in \mathcal{C} is a triple $(X, \underline{e}, \underline{h})$, consisting of an object X , and a collection of endomorphisms $\underline{e} = \{e_k \in \text{End}_{\mathcal{C}}^{-2k}(X)\}_{k=0}^\infty$, $\underline{h} = \{h_k \in \text{End}_{\mathcal{C}}^{-1-2k}(X)\}_{k=0}^\infty$ satisfying

$$(5a) \quad d(e_k) = \sum_{i+j=k-1} (e_i h_j - h_i e_j)$$

$$(5b) \quad d(h_k) = e_k - \sum_{i+j=k} e_i e_j - \sum_{i+j=k-1} h_i h_j$$

The *complement* of the A_∞ idempotent $(X, \underline{e}, \underline{h})$ is the A_∞ idempotent $(X, \underline{e}^\perp, \underline{h}^\perp)^\perp := (X, \underline{e}^\perp, \underline{h}^\perp)$ where

$$e_0^\perp = \text{Id}_X - e_0, \quad e_k^\perp = -e_k \quad (k \geq 1), \quad h_k^\perp = h_k \quad (k \geq 0).$$


Verification that this defines an A_∞ idempotent is left to the reader.

Definition 4.5. If $(X, \underline{e}, \underline{h})$ is an A_∞ idempotent, then let $Z(X, \underline{e}, \underline{h}) \in \text{Tw}(\Sigma^{\mathbb{I}}\mathcal{C})$ denote the twisted complex of the form $\text{tw}_\delta(\prod_{k \geq 0} \Sigma^{-k} X)$ and differential δ given in terms of components by

$$\delta_{ji} = \begin{cases} -e_k^\perp & \text{for } i \text{ even, } j = i + 1 + 2k \\ h_k & \text{for } i \text{ even, } j = i + 2k + 2 \\ e_k & \text{for } i \text{ odd, } j = i + 1 + 2k \\ -h_k & \text{for } i \text{ odd, } j = i + 2k + 2 \end{cases}$$

(recall that $-e_k^\perp = e_k$ for $k \geq 1$).

This twisted complex can be visualized as



with length > 3 components of the differential not pictured.

Example 4.6. The identity of X gives an A_∞ idempotent with $e_0 = \text{Id}_X$ and $e_{k+1} = 0 = h_k$ for $k \geq 0$. The resulting twisted complex

$$Z(X, \text{Id}_X, 0) = \underline{X} \xrightarrow{0} \Sigma^{-1}X \xrightarrow{\text{Id}_X} \Sigma^{-2}X \xrightarrow{0} \Sigma^{-3}X \xrightarrow{\text{Id}_X} \dots$$

is homotopy equivalent to X after the cancellation of contractible summands.

We wish to prove that $Z(X, \underline{e}, \underline{h})$ is a well-defined twisted complex, and is an image of e_0 . To prove this requires a considerable amount of bookkeeping, for which it is useful to consider the generating functions $e(z) = \sum_{k \geq 0} e_k z^k$ and $h(z) = \sum_{k \geq 0} h_k z^k$ where z is a formal indeterminate of degree 2. Below we work an abstract dg category which formalizes the relations satisfied by these generating functions.

4.3. Abstract A_∞ -idempotents and their images.

Definition 4.7. Let z be a formal indeterminate of cohomological degree 2. Let $\mathcal{R}_{\text{idem}}$ be the $\mathbb{k}[[z]]$ -linear dg category with one object \mathbf{X} whose endomorphism complex is freely generated by endomorphisms $e, h \in \text{End}_{\mathcal{R}_{\text{idem}}}(\mathbf{X})$ satisfying

$$\deg(e) = 0, \quad \deg(h) = -1,$$

$$(6a) \quad d(e) = z(eh - he)$$

$$(6b) \quad d(h) = e - e^2 - zh^2,$$

extended to arbitrary morphisms by the Leibniz rule.

To check that $\mathcal{R}_{\text{idem}}$ is a dg category, one must check that $d^2 = 0$ on all morphisms. It suffices to check on the generating morphisms, which is straightforward. For instance to verify that $d^2(e) = 0$ it suffices to verify that $d(eh) = d(he)$, which follows from the computations

$$d(eh) = d(e)h + ed(h) = z(eh - he)h + e(e - e^2 - zh^2) = e^2 - e^3 - zheh$$

and

$$d(he) = d(h)e - hd(e) = (e - e^2 - zh^2)e - zh(eh - he) = e^2 - e^3 - zheh.$$

The proof that $d^2(h) = 0$ is equally straightforward.

The notion of an A_∞ idempotent in \mathcal{C} can now be described as follows. Consider the dg category $\mathcal{C}[[z]]$ with the same objects as \mathcal{C} , and morphism complexes

$$\text{Hom}_{\mathcal{C}[[z]]}(X, Y) := \text{Hom}_{\mathcal{C}}(X, Y) \otimes_{\mathbb{k}} \mathbb{k}[[z]].$$

Then an A_∞ idempotent in \mathcal{C} is equivalent to a dg functor $\mathcal{R}_{\text{idem}} \rightarrow \mathcal{C}[[z]]$. The image of \mathbf{X} in \mathcal{C} is an object $X \in \mathcal{C}$, and the images of e, h are formal series of morphisms $h(z)$,

$$e(z) = \sum_{k \geq 0} e_k z^k \quad (e_k \in \text{End}_{\mathcal{C}}^{-2k}(X)),$$

$$h(z) = \sum_{k \geq 0} h_k z^k \quad (h_k \in \text{End}_{\mathcal{C}}^{-1-2k}(X)),$$

satisfying the identities (6a), (6b). In terms of components, this yields (5a),(5b).

The following is responsible for the notion of complementary A_∞ idempotents.

Lemma 4.8. *There is an automorphism of \mathcal{R}_{idem} which sends $e \mapsto 1 - e$ and fixes h . \square*

Now we have an analogue of Definition 4.5.

Definition 4.9. Let $Z(e, h)$ denote the twisted complex $\text{tw}_\alpha(\mathbf{X} \oplus \Sigma^{-1}\mathbf{X})$ where

$$\alpha = \begin{bmatrix} zh & ze \\ e - 1 & -zh \end{bmatrix}.$$

We also have an analogue of X , viewed through Example 4.6.

Definition 4.10. Let $Z(1, 0) \in \text{Tw}(\Sigma\mathcal{R}_{idem})$ denote the twisted complex $\text{tw}_\beta(\mathbf{X} \oplus \Sigma^{-1}\mathbf{X})$ where

$$\beta = \begin{bmatrix} 0 & z \\ 0 & 0 \end{bmatrix}.$$

Our goal is show that $Z(e, h)$ is the image of a homotopy idempotent acting on $Z(1, 0)$.

Lemma 4.11. *The object $Z(1, 0) = \text{tw}_\beta(\mathbf{X} \oplus \Sigma^{-1}\mathbf{X})$ is a well-defined twisted complex in $\text{Tw}(\Sigma^{\text{II}}\mathcal{R}_{idem})$, and the following defines a homotopy idempotent E acting on $Z(1, 0)$:*

$$E := \begin{bmatrix} e & 0 \\ he - eh & e \end{bmatrix}.$$

Proof. The first statement is clear since $d(\beta) = 0 = \beta^2$. For the second statement, let

$$H := \begin{bmatrix} h & 0 \\ h^2 & -h \end{bmatrix},$$

regarded as an endomorphism of $\mathbf{X} \oplus \Sigma^{-1}\mathbf{X}$. It is straightforward to check that

$$d(H) + \beta H + \beta H = E - E^2.$$

For this, one must keep in mind that a sign appears in the bottom row of the matrix representing $d(H)$, due to the signs involved in differentiating morphisms in the suspended envelope (4):

$$d\left(\begin{bmatrix} h & 0 \\ h^2 & -h \end{bmatrix}\right) = \begin{bmatrix} d(h) & 0 \\ -d(h^2) & d(h) \end{bmatrix}.$$

\square

Lemma 4.12. *The object $Z(e, h) = \text{tw}_\alpha(\mathbf{X} \oplus \Sigma^{-1}\mathbf{X})$ from Definition 4.9 is a well-defined twisted complex in $\text{Tw}(\Sigma^{\text{II}}\mathcal{R}_{idem})$; moreover $Z(e, h)$ is an image of the homotopy idempotent E acting on $Z(1, 0)$ (from Lemma 4.11).*

Proof. To show that $Z(e, h)$ is a well-defined twisted complex we must check that $\alpha \in \text{End}_{\Sigma \mathcal{R}_{\text{idem}}}(\mathbf{X} \oplus \Sigma^{-1}\mathbf{X})$ satisfies the Maurer-Cartan equation

$$\begin{bmatrix} zd(h) & zd(e) \\ -d(e) & zd(h) \end{bmatrix} + \begin{bmatrix} zh & ze \\ e-1 & -zh \end{bmatrix} \begin{bmatrix} zh & ze \\ e-1 & -zh \end{bmatrix} = 0.$$

This is easily verified. Now, define maps $\sigma : Z(e, h) \rightarrow Z(1, 0)$ and $\pi : Z(1, 0) \rightarrow Z(e, h)$ by the matrices

$$\sigma = \begin{bmatrix} 1 & 0 \\ h & e \end{bmatrix}, \quad \pi = \begin{bmatrix} e & 0 \\ -h & 1 \end{bmatrix}.$$

Observe that $\sigma \circ \pi = E$. To check σ is closed is a computation:

$$\begin{bmatrix} 0 & 0 \\ -d(h) & -d(e) \end{bmatrix} + \begin{bmatrix} 0 & z \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ h & e \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ h & e \end{bmatrix} \begin{bmatrix} zh & ze \\ e-1 & -zh \end{bmatrix} = 0,$$

and that π is closed is the computation

$$\begin{bmatrix} d(e) & 0 \\ d(h) & 0 \end{bmatrix} + \begin{bmatrix} zh & ze \\ e-1 & -zh \end{bmatrix} \begin{bmatrix} e & 0 \\ -h & 1 \end{bmatrix} - \begin{bmatrix} e & 0 \\ -h & 1 \end{bmatrix} \begin{bmatrix} 0 & z \\ 0 & 0 \end{bmatrix} = 0,$$

both of which are straightforward. It remains to show that $\pi \circ \sigma \simeq \text{Id}_{Z(e, h)}$. Observe that

$$\pi \circ \sigma = \begin{bmatrix} e & 0 \\ -h & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ h & e \end{bmatrix} = \begin{bmatrix} e & 0 \\ 0 & e \end{bmatrix}.$$

Now, let $K = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \in \text{End}_{\Sigma \mathcal{R}_{\text{idem}}}(\mathbf{X} \oplus \Sigma^{-1}\mathbf{X})$. The following computes $d(K) + \alpha K + K\alpha$:

$$d\left(\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}\right) + \begin{bmatrix} zh & ze \\ e-1 & -zh \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} zh & ze \\ e-1 & -zh \end{bmatrix} = \begin{bmatrix} e-1 & zh-zh \\ 0 & e-1 \end{bmatrix},$$

which shows that $\pi \circ \sigma \simeq \text{Id}_{Z(e, h)}$. This completes the proof that $Z(e, h)$ is an image of the idempotent E acting on $Z(1, 0)$. \square

The twisted complexes $Z(1, 0)$ and $Z(e, h)$ and the maps relating them can be pictured diagrammatically as

$$\begin{array}{ccc} Z(1, 0) & = & \begin{array}{ccc} \mathbf{X} & \xleftarrow{z} & \Sigma^{-1}\mathbf{X} \\ \downarrow e & \searrow -h & \downarrow 1 \\ \mathbf{X} & \xrightleftharpoons[ze]{e-1} & \Sigma^{-1}\mathbf{X} \\ \downarrow 1 & \searrow h & \downarrow e \\ \mathbf{X} & \xleftarrow{z} & \Sigma^{-1}\mathbf{X} \end{array} \\ \downarrow \pi & & \text{Circular arrows on } \mathbf{X} \text{ and } \Sigma^{-1}\mathbf{X} \text{ with labels } zh \text{ and } -zh \\ Z(e, h) & = & \\ \downarrow \sigma & & \\ Z(1, 0) & = & \end{array}$$

Suppose (X, e, h) is an A_∞ idempotent and $\Phi : \mathcal{R}_{\text{idem}} \rightarrow \mathcal{C}[[z]]$ the corresponding dg functor. We may regard $\mathcal{C}[[z]]$ as a (non-full) subcategory of $\Sigma^{\mathbb{I}}\mathcal{C}$ via the functor

sending $X \mapsto X[[z]] := \prod_{k \geq 0} \Sigma^{-2k} X$ with the formal endomorphism z given by the rightward shift on $X[[z]] = X \times \Sigma^{-2} X \times \Sigma^{-4} X \times \dots$.

Thus the images of the abstract twisted complexes $Z(1, 0)$ and $Z(e, h)$ under Φ can be viewed as twisted complexes in $\mathrm{Tw}(\Sigma^{\mathrm{II}}\mathcal{C})$. A moment's thought confirms that these twisted complexes are precisely $Z(X, \mathrm{Id}_X, 0)$ from Example 4.6 and $Z(X, \underline{e}, \underline{h})$ from Definition 4.5. To see this, note that half terms in $Z(X, \underline{e}, \underline{h})$ yield a copy of $X[[z]] = \prod_{k \geq 0} \Sigma^{-2k} X$, while the other half yield a copy of $\Sigma^{-1} X[[z]] = \prod_{k \geq 0} \Sigma^{-2k-1} X$.

Since $Z(X, \mathrm{Id}_X, 0) \simeq X$, Lemma 4.12 shows that $Z(X, \underline{e}, \underline{h})$ is the image of some homotopy idempotent acting on X . It is not hard to see that this homotopy idempotent is e_0 , thereby proving the following.

Proposition 4.13. *Let $(X, \underline{e}, \underline{h})$ be an A_∞ idempotent in \mathcal{C} . The object $Z(X, \underline{e}, \underline{h})$ is a well-defined twisted complex in $\mathrm{Tw}(\Sigma^{\mathrm{II}}\mathcal{C})$; moreover this twisted complex is an image of the homotopy idempotent e_0 acting on X . \square*

Remark 4.14. The complex $Z(X, \underline{e}, \underline{h})$ has a natural endomorphism z representing the 2-periodicity in this construction. This endomorphism is null-homotopic (by an explicit homotopy), so z^k is null homotopic for all $k \geq 1$ as well.

The cone of z^k is homotopy equivalent to a finite twisted complex (a truncated version of $Z(X, \underline{e}, \underline{h})$), which by the above represents the image of e_0 acting on $X \oplus \Sigma^{2k-1} X$.

4.4. From homotopy idempotents to A_∞ idempotents. Finally, we show that any homotopy idempotent e in \mathcal{C} can be given the structure of an A_∞ idempotent. This is well known to experts, but we will give an explicit construction of the higher homotopies following ideas of Seidel [Sei08, Lemma 4.2]. See [BN93, Propositions 3.2 and 3.4] for an alternative proof. We were not able to find explicit formulas for \underline{e} and \underline{h} in the literature.

Proposition 4.15. *Suppose that $e, h \in \mathrm{End}_{\mathcal{C}}(X)$ are such that e is degree zero and closed, and h is degree -1 and satisfies $d_{\mathcal{C}}(h) = e^2 - e$. Then there exist endomorphisms $h^{(k)} \in \mathrm{End}^{-k}(X)$ for $k \geq 1$ such that $h^{(0)} = 1 - e$, $h^{(1)} = h$ and*

$$d(h^{(k)}) = \sum_{i=0}^{k-1} (-1)^i h^{(i)} h^{(k-1-i)} + \begin{cases} -h^{(k-1)} & \text{if } k \text{ is odd} \\ 0 & \text{if } k \text{ is even.} \end{cases}$$

Corollary 4.16. *Suppose that $e, h \in \mathrm{End}_{\mathcal{C}}(X)$ are such that e is degree zero and closed, and h is degree -1 and satisfies $d_{\mathcal{C}}(h) = e - e^2$. Let $h^{(k)}$ denote the morphisms obtained from Proposition 4.15 starting at $(e, -h)$. We define $e_0 := e$ and $e_k = (-1)^{k+1} h^{(2k)}$ for $k \geq 1$, as well as $h_k := (-1)^{k+1} h^{(2k+1)}$ for $k \geq 0$. Then $(X, \underline{e}, \underline{h})$ is an A_∞ idempotent in \mathcal{C} .*

Proof. It is straightforward that these morphisms satisfy the equations from Definition 4.4. \square

Proof of Proposition 4.15. We will construct $h^{(n)}$ inductively. Recall that $h^{(0)} = 1 - e$. Suppose that we found $h^{(1)}, \dots, h^{(n-1)}, h_{\text{temp}}^{(n)}$ such that

$$(7) \quad d(h^{(k)}) = \sum_{i=0}^{k-1} (-1)^i h^{(i)} h^{(k-1-i)} + \begin{cases} -h^{(k-1)} & \text{if } k \text{ is odd} \\ 0 & \text{if } k \text{ is even.} \end{cases}, \quad 1 \leq k \leq n-1$$

$$(8) \quad d(h_{\text{temp}}^{(n)}) = \sum_{i=0}^{n-1} (-1)^i h^{(i)} h^{(n-1-i)} + \begin{cases} -h^{(n-1)} & \text{if } n \text{ is odd} \\ 0 & \text{if } n \text{ is even.} \end{cases}$$

For $n = 1$, this follows from the assumptions of the proposition if we set $h_{\text{temp}}^{(1)} = h$. Our goal is to find an $h_{\text{temp}}^{(n+1)}$ and a closed y_n , such that (8) will be satisfied for $n \mapsto n+1$ if we set $h^{(n)} = h_{\text{temp}}^{(n)} + y_n$. Moreover, (7) will then hold for $k = n$ since y_n is closed. To this end, we define

$$x_n = h^{(0)} h_{\text{temp}}^{(n)} + \sum_{i=1}^{n-1} (-1)^i h^{(i)} h^{(n-i)} + (-1)^n h_{\text{temp}}^{(n)} h^{(0)} + \begin{cases} -h_{\text{temp}}^{(n)} & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

$$q_n = -h^{(1)} h_{\text{temp}}^{(n)} + \sum_{i=2}^{n-1} (-1)^i h^{(i)} h^{(n+1-i)} + (-1)^n h_{\text{temp}}^{(n)} h^{(1)}$$

It is not hard to check that $d(x_n) = 0$. For $x \in \text{End}_{\mathbb{C}}(X)$ we define

$$A(x) = ex - xe, \quad B(x) = ex - x(1 - e).$$

We then set $y_n = B(x_n)$ if n is even, and $y_n = A(x_n)$ if n is odd. In either case we again have $d(y_n) = 0$. Further, one can check that $d(q_n) = A(x_n)$ if n is even, and $d(q_n) = B(x_n)$ if n is odd, which helps to verify:

$$B(y_n) = B(B(x_n)) = d(hx_n + 3x_n h + 2q_n e - q_n) + x_n \quad \text{if } n \text{ is even}$$

$$A(y_n) = A(A(x_n)) = d(hx_n - 3x_n h - 2q_n e + q_n) + x_n \quad \text{if } n \text{ is odd.}$$

Now we set $h^{(n)} = h_{\text{temp}}^{(n)} + y_n$ and $h_{\text{temp}}^{(n+1)} = -(hx_n + 3x_n h + 2q_n e - q_n)$ if n is even, and $h_{\text{temp}}^{(n+1)} = -(hx_n - 3x_n h - 2q_n e + q_n)$ if n is odd. Let us check that (8) is now satisfied for $n \mapsto n+1$. We only consider the case of odd $n+1$, as the other one is analogous.

$$\begin{aligned} d(h_{\text{temp}}^{(n+1)}) &= x_n - B(y_n) = h^{(0)} h_{\text{temp}}^{(n)} + \sum_{i=1}^{n-1} (-1)^i h^{(i)} h^{(n-i)} + h_{\text{temp}}^{(n)} h^{(0)} - h_{\text{temp}}^{(n)} \\ &\quad + h^{(0)} y_n + y_n h^{(0)} - y_n \\ &= h^{(0)} h^{(n)} + \sum_{i=1}^{n-1} (-1)^i h^{(i)} h^{(n-i)} + h^{(n)} h^{(0)} - h^{(n)} \end{aligned}$$

where we have used $-B(y_n) = -ey_n - y_n e + y_n = h^{(0)} y_n + y_n h^{(0)} - y_n$. \square

4.5. The Karoubi envelope.

Definition 4.17. For a dg category \mathcal{C} we define the dg Karoubi envelope $\text{Kar}^{dg}(\mathcal{C})$ as the full dg subcategory of $\text{Tw}(\Sigma^{\text{II}}\mathcal{C})$ with objects the twisted complexes homotopy equivalent to $Z(X, \underline{e}, \underline{h})$ for some A_∞ idempotent $(X, \underline{e}, \underline{h})$ in $\Sigma\mathcal{C}$.

Note that by Example 4.6, we have $\mathcal{C} \hookrightarrow \text{Kar}^{dg}(\mathcal{C})$.

Lemma 4.18. *The category $\text{Kar}^{dg}(\mathcal{C})$ is homotopy idempotent complete.*

Proof. If $e_0 \in \text{End}_{\mathcal{C}}(X)$ is a homotopy idempotent, then e_0 admits a lift to an A_∞ idempotent by Proposition 4.15, and Proposition 4.13 constructs an image of e_0 .

On the other hand, every object of $\text{Kar}^{dg}(\mathcal{C})$ is the image of some idempotent in $H^0(\mathcal{C})$ by construction, and conversely every idempotent in \mathcal{C} has an image in $\text{Kar}^{dg}(\mathcal{C})$. So if $Y \in \text{Kar}^{dg}(\mathcal{C})$ is the image of a homotopy idempotent $e_0 \in \text{End}_{\mathcal{C}}(X)$ then all images of all homotopy idempotents in $\text{End}_{\text{Kar}^{dg}(\mathcal{C})}(Y)$ can be constructed as images of some induced homotopy idempotents $e_0'' \in \text{End}_{\mathcal{C}}(X)$ by part (2) of Lemma 4.1. \square

Lemma 4.19. *There is an equivalence of additive categories $H^0(\text{Kar}^{dg}(\mathcal{C})) \simeq \text{Kar} H^0(\mathcal{C})$, where the latter denotes the usual idempotent completion of the additive category $H^0(\mathcal{C})$.*

Proof. Lemma 4.18 implies that $H^0(\text{Kar}^{dg}(\mathcal{C}))$ is idempotent complete in the usual sense, for additive categories. Now, the canonical functor $H^0(\mathcal{C}) \rightarrow H^0(\text{Kar}^{dg}(\mathcal{C}))$ induces a functor on the Karoubi envelope $\text{Kar}(H^0(\mathcal{C})) \rightarrow H^0(\text{Kar}^{dg}(\mathcal{C}))$ because the latter is Karoubian. This functor is essentially surjective because every object in $\text{Kar}^{dg}(\mathcal{C})$ is the image of some homotopy idempotent acting on some object of \mathcal{C} , and fully faithful by part (3) of Lemma 4.1. \square

Lemma 4.20. *A dg category \mathcal{D} is homotopy idempotent complete if and only if the natural functor $\mathcal{D} \rightarrow \text{Kar}^{dg}(\mathcal{D})$ is a quasi-equivalence.*

Proof. Assume that $H^0(\mathcal{D})$ is idempotent complete. Then $H^0(\Sigma\mathcal{D})$ is also idempotent complete. By Lemma 4.19 the natural functor $\Sigma\mathcal{D} \rightarrow \text{Kar}^{dg}(\Sigma\mathcal{D})$ induces an equivalence $H^0(\Sigma\mathcal{D}) \rightarrow H^0(\text{Kar}^{dg}(\Sigma\mathcal{D})) \simeq \text{Kar}(H^0(\Sigma\mathcal{D}))$, hence $\mathcal{D} \rightarrow \text{Kar}^{dg}(\mathcal{D})$ is a quasi-equivalence.

For the converse, suppose that the canonical functor $\mathcal{D} \rightarrow \text{Kar}^{dg}(\mathcal{D})$ is a quasi-equivalence. Then we have an equivalence of categories $H^0(\mathcal{D}) \rightarrow H^0(\text{Kar}^{dg}(\mathcal{D}))$. Since the latter category is idempotent complete, so is the former, i.e. \mathcal{D} is homotopy idempotent complete. \square

We are ready to check the universal property of $\text{Kar}^{dg}(\mathcal{C})$.

Proof of Theorem 4.3. Let \mathcal{D} be a homotopy idempotent complete dg category, and let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a dg functor. We extend F to a dg functor $\text{Tw}(\Sigma^{\text{II}}\mathcal{C}) \rightarrow \text{Tw}(\Sigma^{\text{II}}\mathcal{D})$; this restricts to a dg functor on the full subcategories $\text{Kar}^{dg}(\mathcal{C}) \rightarrow \text{Kar}^{dg}(\mathcal{D})$. Since \mathcal{D} is homotopy idempotent complete, the canonical dg functor $\mathcal{D} \rightarrow \text{Kar}^{dg}(\mathcal{D})$ is a

quasi-equivalence by Lemma 4.20, and composing with the inverse (in \mathbf{Hqe}) gives a quasi-functor $\tilde{F}: \text{Kar}^{dg}(\mathcal{C}) \rightarrow \mathcal{D}$ lifting F .

The uniqueness of this lift up to homotopy (again in \mathbf{Hqe}) follows because if $e \in \text{End}_{\mathcal{C}}(X)$ is a homotopy idempotent then $F(\text{im } e)$ is determined uniquely up to homotopy by $F(X)$ and $F(e)$. \square

Lemma 4.21. *Let I be a finite poset and suppose we have objects $X_i \in \mathcal{C}$ equipped with homotopy idempotents $e_i \in \text{End}_{\mathcal{C}}^0(X)$. Suppose $Y_i \simeq \text{im } e_i$ for $i \in I$. Then any one-sided twist $\text{tw}_{\alpha}(\bigoplus_i Y_i)$ is the image of some homotopy idempotent f acting on some one-sided twist $\text{tw}_{\beta}(\bigoplus_i X_i)$.*

Proof. It is sufficient to prove the statement for two-term twisted complexes, i.e. cones. Suppose that Y_0, Y_1 are homotopy summands of X_0, X_1 corresponding to homotopy idempotents e_0, e_1 and $g: Y_0 \rightarrow Y_1$ is a closed degree zero morphism. There are inclusions $\sigma_i: Y_i \rightarrow X_i$, projections $\pi_i: X_i \rightarrow Y_i$ and homotopies $h_i: Y_i \rightarrow Y_i$ such that $\pi_i \sigma_i = \text{Id}_{Y_i} + d(h_i)$. Define $g' := \sigma_1 g \pi_0$, and consider the following chain maps between $\text{Cone}(g)$ and $\text{Cone}(g')$:

$$p := \begin{array}{ccc} X_0 & \xrightarrow{g'} & X_1 \\ \downarrow \pi_0 & \searrow H & \downarrow \pi_1 \\ Y_0 & \xrightarrow{g} & Y_1 \end{array}, \quad s := \begin{array}{ccc} X_0 & \xrightarrow{g'} & X_1 \\ \sigma_0 \uparrow & \nearrow H' & \sigma_1 \uparrow \\ Y_0 & \xrightarrow{g} & Y_1 \end{array}$$

where $H = h_1 g \pi_0$ and $H' = \sigma_1 g h_0$. It is easy to see that $p \circ s$ is homotopic to identity, so $\text{Cone}(g)$ is a homotopy summand of $\text{Cone}(g')$. \square

Theorem 4.22. *If \mathcal{C} is pretriangulated, then so is $\text{Kar}^{dg}(\mathcal{C})$.*

Proof. Every object of $\text{Kar}^{dg}(\mathcal{C})$ is (isomorphic to) a homotopy summand of an object in \mathcal{C} . A one-sided twisted complex constructed from homotopy summands of $Y^i \in \mathcal{C}$ ($i \in I$) is also a homotopy summand of a twisted complex constructed from Y^i by Lemma 4.21; such objects live in $\text{Kar}^{dg}(\mathcal{C})$ since \mathcal{C} is pretriangulated and $\text{Kar}^{dg}(\mathcal{C})$ is idempotent complete. \square

4.6. Perfect complexes are Karoubian. Let A be a dg algebra. Throughout this section we will assume that A is supported in *non-positive* cohomological degrees.

Let $\langle A \rangle$ to be the category of finitely generated free A -modules, that is, direct sums of finitely many copies of A . We define $\text{Pretr}(A)$ to be the category of (finite) twisted complexes of free A -modules or, equivalently, the pretriangulated hull of $\langle A \rangle$. We define $\text{Kar}^{dg}\langle A \rangle$ to be the category of projective A -modules, that is, homotopy direct summands of objects in $\langle A \rangle$. Let $\text{Perf}(A)$ be the category of *perfect twisted complexes* over A , that is, bounded twisted complexes built out of objects in $\text{Kar}^{dg}\langle A \rangle$. In other words,

$$\text{Perf}(A) = \text{Pretr}(\text{Kar}^{dg}\langle A \rangle).$$

Finally, we define $\text{Ch}^b(A)$ (resp. $\text{Ch}^b(\text{Kar}^{dg}\langle A \rangle)$) to be the category of bounded complexes of free (resp. projective) A -modules.

By definition, an object of $\text{Pretr}(A)$ is a graded free dg module $M = \bigoplus_i \Sigma^{-i} M_i$ with M_i in $\langle A \rangle$, and a differential given by a matrix $d = (d_{ij})$ where $d_{ij}: M_j \rightarrow M_i$ and d_{ii} agrees with the internal differential on M_i . The differential d is homogeneous of cohomological degree 1, so d_{ij} has cohomological degree $1 + j - i$. Since A is non-positively graded, we have $d_{ij} = 0$ unless $i \geq j$, so the matrix $d = (d_{ij})$ is lower-triangular. Also, $d^2 = 0$ which means

$$(9) \quad \sum_{k: i \geq k \geq j} d_{ik} d_{kj} = 0 \text{ for all } i > j.$$

Similarly, if $M = (\bigoplus_i \Sigma^{-i} M_i, d)$ and $N = (\bigoplus_i \Sigma^{-i} N_i, d)$ then a morphism from M to N is given by an lower-triangular matrix of morphisms $f = (f_{ij})$ where f_{ij} is a morphism from M_j to N_i of cohomological degree $j - i$.

This immediately implies the following:

Proposition 4.23. *Suppose that the differential on A vanishes and let (M, d) be a twisted complex in $\text{Pretr}(A)$, and $d' = \sum_i d_{i+1,i}$. Then $(d')^2 = 0$, so (M, d') is a well-defined chain complex in $\text{Ch}^b(A)$.*

Remark 4.24. A similar construction works if the differential on A is non-trivial, but its component from A_{-1} to A_0 vanishes.

We will call (M, d') the underlying complex of the twisted complex (M, d) . This construction defines dg functors

$$\epsilon: \text{Pretr}(A) \rightarrow \text{Ch}^b(A), \quad \epsilon: \text{Perf}(A) \rightarrow \text{Ch}^b(\text{Kar}^{dg}\langle A \rangle).$$

We will denote both of these functors by ϵ since it will be clear from the context which one is used.

The following lemma is a standard application of perturbation theory.

Lemma 4.25. *Suppose that X is an object in $\text{Pretr}(A)$ or $\text{Perf}(A)$. If $\epsilon(X)$ is contractible then X is contractible. If $f: X \rightarrow Y$ is a morphism such that $\epsilon(f)$ is a homotopy equivalence, then f is a homotopy equivalence.*

Proof. For the first part see e.g [GH17, Lemma 2.19]. For the second part, observe that f is a homotopy equivalence if and only if the cone of f is contractible. \square

Theorem 4.26. *Assume that A is non-positively graded and the differential on A vanishes. Then the category $\text{Perf}(A)$ is homotopy idempotent complete, and*

$$\text{Kar}^{dg}(\text{Pretr}(A)) = \text{Perf}(A).$$

Remark 4.27. The proof is similar to [BV08, Lemma 1.5.6(iii)] and [GW19, Appendix], but we include it here for completeness and add more details for reader's convenience.

Proof. Let X be a twisted complex in $\text{Perf}(A)$, and let e be a homotopy idempotent endomorphism of X . Without loss of generality we can suppose that X is a (one-sided) twist of

$$\bigoplus_{0 < i < (2N-1)} \Sigma^{-i} X_i$$

for some sufficiently large N .

As in Section 4.4, we can lift e to an A_∞ idempotent $(X, \underline{e}, \underline{h})$ and consider a twisted complex Z built out of several copies of X connected by the maps from \underline{e} and \underline{h} with appropriate signs. As in Remark 4.14 we build a bounded Z from $2N$ of copies of X , which is homotopy equivalent to the image of e acting on $X \oplus \Sigma^{2N-1}X$ via an argument similar to Lemma 4.12. By our assumptions, X is a direct sum of copies of $\Sigma^{-i}A$ with $i > 0$, and $\Sigma^{2N-1}X$ is a direct sum of copies of $\Sigma^{-i}A$ with $i < 0$. There are no nonzero A -module maps from $\Sigma^{-j}A \rightarrow \Sigma^{-i}A$ unless $i > j$, so we get the following diagram:

$$\begin{array}{ccccc}
 \Sigma^{2N-1}X & & & & X \\
 \downarrow & \searrow & \searrow & & \downarrow \\
 Z_{<0} & \longrightarrow & Z_0 & \longrightarrow & Z_{>0} \\
 \downarrow & \searrow & \searrow & & \downarrow \\
 \Sigma^{2N-1}X & & & & X
 \end{array}$$

where the top half of the diagram represents the projection $\pi: X \oplus \Sigma^{2N-1}X \rightarrow Z$ and the bottom half represents the inclusion $i: Z \rightarrow X \oplus \Sigma^{2N-1}X$. We decomposed $Z = \bigoplus_a \Sigma^{i_a}A$ into its summands $Z_{>0}$, Z_0 , $Z_{<0}$ corresponding to those indices a for which $i_a > 0$, $i_a = 0$, or $i_a < 0$, respectively (recall that Σ is the suspension so in fact lowers cohomological degree).

Observe that $\pi \circ i$ is homotopic to identity on Z and vanishes on Z_0 . Since the differential on A vanishes, by Proposition 4.23 one can define the forgetful dg functor $\epsilon: \text{Perf}(A) \rightarrow \text{Ch}(\text{Kar}^{dg}(A))$. By applying it to Z we get a chain complex $\epsilon(Z)$ and a homotopy $h \in \text{End}(\epsilon(Z))$ such that $dh + hd|_{Z_0} = \text{Id}_{Z_0}$. Now

$$(dh)^2 = (dh + hd)dh = dh, \quad (hd)^2 = hd(dh + hd) = hd,$$

so dh and hd are two orthogonal idempotents on Z_0 . Since $\text{Kar}^{dg}(A)$ is Karoubian, we can split $Z_0 = Q \oplus Q'$.

For the final step, let T be the natural twisted complex built out of Q and $Z_{>0}$ (abstractly speaking, this is the cone of the composition $\Sigma^{-1}Q \hookrightarrow \Sigma^{-1}Z_0 \rightarrow Z_{>0}$). It is easy to see that T is a subcomplex of Z and we can restrict the maps i and π from the above diagram to $i': T \rightarrow X$ and $\pi': X \rightarrow T$. Then $i' \circ \pi' = i \circ \pi|_X \simeq e$ and $\epsilon(\pi' \circ i') \simeq \text{Id}_{\epsilon(T)}$. By Lemma 4.25 we get that $\pi' \circ i'$ is homotopic to Id_T . Therefore T represents the image of e , and we conclude that X is homotopy equivalent to a perfect A -module.

This shows that $\text{Perf}(A)$ is homotopy Karoubian. \square

Suppose that a dg algebra A retracts to its dg module V . Recall that in section 2.5 we defined A_∞ operations $\mu_k: V^{\otimes k} \rightarrow V$.

Lemma 4.28. *Assume that $\mu_k = 0$ for $k \geq 3$. Then the dg categories $\text{Perf}(A)$ and $\text{Perf}(V)$ are quasi-equivalent.*

Proof. Since $\mu_k = 0$ for $k > 2$, we can regard V as a dg algebra with differential μ_1 and multiplication μ_2 . Furthermore, we can consider A and V as dg categories $(*, A)$ and $(*, V)$ with one object with endomorphism algebra A or V , respectively. We claim that $(*, A)$ and $(*, V)$ are quasi-equivalent.

Indeed, there exist A_∞ algebra homomorphisms between A and V^2 (the A_∞ maps $\lambda_k : V^{\otimes k} \rightarrow A$ are defined in the proof of Theorem 2.10). By Theorem 2.10, these A_∞ algebra homomorphisms induce quasi-isomorphisms on homology, so $(*, A)$ and $(*, V)$ are quasi-equivalent as A_∞ categories. By [LH03, Kel01], homotopy category of A_∞ algebras (regarded as a model category) is equivalent to homotopy category of dg algebras, so $(*, A)$ and $(*, V)$ are quasi-equivalent as dg algebras.

The corresponding categories of free dg modules $\langle A \rangle$ and $\langle V \rangle$ are nothing but additive closures of $(*, A)$ and $(*, V)$, so these are quasi-equivalent. Therefore $\text{Kar}^{dg}\langle A \rangle$ and $\text{Kar}^{dg}\langle V \rangle$ are quasi-equivalent by Theorem 4.3 and their pretriangulated hulls are quasi-equivalent as well. \square

5. THE BAR COMPLEX FOR DG CATEGORIES

One can motivate the introduction of the (two-sided) bar complex of a dg category as the object which governs the notion of “naturality up to coherent homotopy”. To illustrate, let \mathcal{C} and \mathcal{D} be dg categories and $F, G : \mathcal{C} \rightarrow \mathcal{D}$ dg functors. We have already introduced the notion of a (say, degree zero) natural transformation, which is a choice of morphism $\alpha_X \in \text{Hom}_{\mathcal{D}}(F(X), G(X))$ for all $X \in \mathcal{C}$, natural with respect to morphisms in \mathcal{C} . This means for every morphism $f : X \rightarrow Y$ in \mathcal{C} we have

$$G(f) \circ \alpha_X - \alpha_Y \circ F(f) = 0.$$

Said differently, the pair of dg functors F, G determines a \mathcal{C}, \mathcal{C} -bimodule $\mathcal{B}(G, F)$, which for $X, Y \in \mathcal{C}$ has

$$Y\mathcal{B}(G, F)X := G(Y)\mathcal{D}F(X),$$

(in the notation of §2.4) and a natural transformation is simply a map of \mathcal{C}, \mathcal{C} -bimodules

$$\alpha : \mathcal{C} \rightarrow \mathcal{B}(G, F).$$

The image of $\text{Id}_X \in \mathcal{C}$ is the chosen morphism α_X , and naturality is equivalent to α being a \mathcal{C}, \mathcal{C} -bimodule map.

Now suppose that we are in a situation where naturality does not hold on the nose, but only up to homotopies $\alpha_f \in \text{Hom}_{\mathcal{D}}^{-1}(F(X), G(Y))$:

$$d_{\mathcal{D}}(\alpha_f) = G(f) \circ \alpha_X - \alpha_Y \circ F(f).$$

²A word of caution: there may be no dg algebra homomorphisms between A and V unless $\lambda_k = 0$ for $k > 2$

Let us consider another morphism $g: Y \rightarrow Z$ in \mathcal{C} and suppose that f and g are closed. It is straightforward to check that the expression

$$\alpha_{g \circ f} - G(g) \circ \alpha_f - \alpha_g \circ F(f)$$

is closed. It is often desirable to suppose this expression is also exact, i.e. there exists a homotopy $\alpha_{g,f} \in \text{Hom}_{\mathcal{D}}^{-2}(F(X), G(Z))$. Roughly speaking, this says that the assignment $f \mapsto \alpha_f$ satisfies a version of the Leibniz rule, up to homotopy.

Now, assuming the existence of such ‘‘higher homotopies’’ $\alpha_{f,g}$ for all composable morphisms f, g one can then define a degree -3 closed morphism associated to each triple of composable morphisms f_1, f_2, f_3 , the exactness of which would allow us to define a family of closed morphisms (obstructions) associated to each 4-tuple of composable morphisms, and so on.

If all such obstructions are exact (and a family of homotopies realizing this exactness is given) then then we say that the system $(\alpha_X, \alpha_f, \alpha_{g,f}, \dots)$ is a *homotopy coherent natural transformation*.

The two-sided bar complex $\mathbf{B}(\mathcal{C})$ of \mathcal{C} , which we describe explicitly below, can be considered as a free resolution of the trivial \mathcal{C}, \mathcal{C} -bimodule \mathcal{C} , and is spanned as a bimodule by sequences of composable morphisms of arbitrary finite length $r \geq 0$. The data of a homotopy coherent natural transformation $F \rightarrow G$ is then encoded as a map of \mathcal{C}, \mathcal{C} -modules

$$\mathbf{B}(\mathcal{C}) \rightarrow \mathcal{B}(G, F).$$

Various operations on natural transformations can be understood in terms of various structures on the two-sided bar complex; for instance the composition of natural transformations can be understood via a natural comultiplication on $\mathbf{B}(\mathcal{C})$.

After this motivation, we now give an explicit description of the two-sided bar complex.

5.1. The two-sided bar complex of a pair. Let \mathcal{C} be a dg category and $\mathcal{J} \subset \mathcal{C}$ a (unital) subcategory. We wish to define the two-sided bar complex associated to $(\mathcal{J}, \mathcal{C})$. First, consider \mathcal{C}, \mathcal{C} -bimodules of the form $\mathcal{C} \otimes_{\mathcal{J}} \cdots \otimes_{\mathcal{J}} \mathcal{C}$.

Example 5.1. In case $\mathcal{J} = \mathcal{J}_{\mathcal{C}}$ is the subcategory of identity maps from Example 2.4, $\mathcal{C} \otimes_{\mathcal{J}} \cdots \otimes_{\mathcal{J}} \mathcal{C}$ is spanned by composable morphisms in \mathcal{C} . More generally, if $\mathcal{B} \subset \mathcal{C}$ is a full subcategory and $\mathcal{J} = \mathcal{J}_{\mathcal{B}} \subset \mathcal{C}$ is the subcategory of identity morphisms in \mathcal{B} , then

$$\underbrace{\mathcal{C} \otimes_{\mathcal{J}} \otimes \cdots \otimes_{\mathcal{J}} \mathcal{C}}_{r+2 \text{ factors}}$$

is spanned as a bimodule by sequences of composable morphisms, which we denote by $f_0 \| f_1 \| \cdots \| f_{r+1}$, in which f_1, \dots, f_r are morphisms in the full subcategory $\mathcal{B} \subset \mathcal{C}$.

The *two-sided bar complex* associated to the pair $\mathcal{J} \subset \mathcal{C}$ is by definition the total complex $\mathbf{B}(\mathcal{C}; \mathcal{J})$ of the following bicomplex:

$$(10) \quad \cdots \longrightarrow \mathcal{C} \otimes_{\mathcal{J}} \mathcal{C} \otimes_{\mathcal{J}} \mathcal{C} \otimes_{\mathcal{J}} \mathcal{C} \longrightarrow \mathcal{C} \otimes_{\mathcal{J}} \mathcal{C} \otimes_{\mathcal{J}} \mathcal{C} \longrightarrow \mathcal{C} \otimes_{\mathcal{J}} \mathcal{C}$$

in which the horizontal arrows are

$$f_0 \parallel \cdots \parallel f_{r+1} \mapsto \sum_{i=0}^r (-1)^i f_0 \parallel \cdots \parallel f_i \circ f_{i+1} \parallel \cdots \parallel f_{r+1}.$$

More precisely,

$$\mathbf{B}(\mathcal{C}; \mathcal{J}) = \bigoplus_{r \geq 0} \bigoplus_{X_1, \dots, X_{r+1}} \Sigma^r \left(\mathcal{C}X_1 \otimes X_1 \mathcal{C}X_2 \otimes \cdots \otimes X_{r+1} \mathcal{C} \right),$$

where the direct sum is over objects X_1, \dots, X_{r+1} for which $\text{Id}_{X_i} \in \mathcal{J}$ for $i = 1, \dots, r+1$. The differential on the two-sided bar complex is $d_{\oplus} + d_{\text{bar}}$, where d_{\oplus} is the direct sum of differentials on the terms above (inherited from $d_{\mathcal{C}}$ by the usual tensor product rule, and a sign $(-1)^r$ coming from the r -translation), and d_{bar} is defined by

$$d_{\text{bar}}(f_0 \parallel \cdots \parallel f_{r+1}) = \sum_{i=0}^r (-1)^i f_0 \parallel \cdots \parallel f_i \circ f_{i+1} \parallel \cdots \parallel f_{r+1}.$$

(The notation is suggestive: the bar differential is the alternating sum of erasing bars.) The bimodule structure is given term-wise by

$$g \cdot (f_0 \parallel f_1 \parallel \cdots \parallel f_r \parallel f_{r+1}) \cdot g' := (-1)^{|g|r} (g \circ f_0) \parallel f_1 \parallel \cdots \parallel f_r \parallel (f_{r+1} \circ g').$$

Note that the left-action of bimodules obtains a sign-twist under translation.

By convention $\mathbf{B}(\mathcal{C})$ denotes $\mathbf{B}(\mathcal{C}, \mathcal{J})$ where $\mathcal{J} = \mathcal{J}_{\mathcal{C}}$ is the subcategory of identity maps in \mathcal{C} .

Remark 5.2. At the beginning of this section we have motivated the bar complex by claiming that \mathcal{C}, \mathcal{C} -bimodule maps $\alpha: \mathbf{B}(\mathcal{C}) \rightarrow \mathcal{B}(G, F)$ encode homotopy coherent natural transformations between dg functors $F, G: \mathcal{C} \rightarrow \mathcal{D}$. Indeed, the data of such a bimodule map is precisely the choice of closed degree zero morphisms $\alpha_X := \alpha(\text{Id}_X \parallel \text{Id}_X)$ from $F(X)$ to $G(X)$ in \mathcal{D} , together with morphisms $\alpha(\text{Id}_Y \parallel f \parallel \text{Id}_X)$ which realize naturality up to homotopy, and the higher homotopies $\alpha(\text{Id}_{X_0} \parallel f_1, \dots, f_r \parallel \text{Id}_{X_{r+1}})$ which provide the required higher naturality data.

Proposition 5.3. *The two-sided bar complex $\mathbf{B}(\mathcal{C}, \mathcal{J})$ is a coalgebra object in the category of \mathcal{C}, \mathcal{C} -bimodules: we have maps of bimodules $\mathbf{B}(\mathcal{C}, \mathcal{J}) \rightarrow \mathcal{C}$ and $\mathbf{B}(\mathcal{C}, \mathcal{J}) \rightarrow \mathbf{B}(\mathcal{C}, \mathcal{J}) \otimes_{\mathcal{C}} \mathbf{B}(\mathcal{C}, \mathcal{J})$ satisfying the usual counit and coassociativity relations.*

Proof. The counit $\varepsilon: \mathbf{B}(\mathcal{C}, \mathcal{J}) \rightarrow \mathcal{C}$ is defined componentwise by

$$f_0 \parallel \cdots \parallel f_{r+1} \mapsto \begin{cases} f_0 \circ f_1 & \text{if } r = 0 \\ 0 & \text{otherwise} \end{cases}.$$

The comultiplication $\Delta: \mathbf{B}(\mathcal{C}, \mathcal{J}) \rightarrow \mathbf{B}(\mathcal{C}, \mathcal{J}) \otimes_{\mathcal{C}} \mathbf{B}(\mathcal{C}, \mathcal{J})$ is defined componentwise by

$$f_0 \parallel \cdots \parallel f_{r+1} \mapsto \sum_{i=1}^{r+1} (-1)^{(r-i+1)(|f_0| + \cdots + |f_{i-1}|)} (f_0 \parallel \cdots \parallel f_{i-1} \parallel \text{Id}) \otimes (\text{Id} \parallel f_i \parallel \cdots \parallel f_{r+1}).$$

The counit and coassociativity axioms are easily checked. Furthermore, it is clear that Δ and ε are closed and commute with the \mathcal{C}, \mathcal{C} -bimodule structure on $\mathbf{B}(\mathcal{C}, \mathcal{J})$. \square

Let us give an alternate description of the two-sided bar complex. The bimodule $\mathcal{C} \otimes_{\mathcal{J}} \mathcal{C}$ is a coalgebra object $C_{\mathcal{J}}$ in \mathcal{C} , \mathcal{C} -bimodules, and $\mathbf{B}(\mathcal{C}, \mathcal{J})$ is the counital idempotent $\mathbf{P}_{C_{\mathcal{J}}}$ associated to $C_{\mathcal{J}}$ [Hog17].

If $\mathcal{J} \subset \mathcal{J}' \subset \mathcal{C}$, then the counit of $C_{\mathcal{J}'}$ factors through the counit of $C_{\mathcal{J}}$, so $C_{\mathcal{J}} \leq C_{\mathcal{J}'}$ in the notation of [Hog17]. The following is an immediate consequence of this.

Proposition 5.4. *If $\mathcal{J} \subset \mathcal{J}' \subset \mathcal{C}$ are unital subcategories then*

$$\mathbf{B}(\mathcal{C}, \mathcal{J}) \otimes_e \mathbf{B}(\mathcal{C}, \mathcal{J}') \simeq \mathbf{B}(\mathcal{C}, \mathcal{J}') \otimes_e \mathbf{B}(\mathcal{C}, \mathcal{J}).$$

5.2. Shrinking the bar complex.

Definition 5.5. We say that a full subcategory $\mathcal{B} \subset \mathcal{C}$ *generates* \mathcal{C} if for all dg functors $F : \mathcal{C} \rightarrow \mathbb{k}\text{-dgm}$, $F(X) \simeq 0$ for all $X \in \mathcal{B}$ implies $F(X) \simeq 0$ for all $X \in \mathcal{C}$.

Proposition 5.6. *Suppose every object of \mathcal{C} is homotopy equivalent to a finite one-sided twisted complex constructed from objects of \mathcal{B} . Then \mathcal{B} generates \mathcal{C} in the sense of Definition 5.5.*

Proof. Let $F : \mathcal{C} \rightarrow \mathbb{k}\text{-dgm}$ be a dg functor such that $F(X) \simeq 0$ for all $X \in \mathcal{B}$, and let $Z = \text{tw}_{\delta}(\bigoplus_i \Sigma^{a_i} X_i)$ be a finite one-sided twisted complex. Then $F(Z)$ is isomorphic to a one-sided twisted complex $\text{tw}_{F(\delta)}(\bigoplus_i \Sigma^{a_i} F(X_i))$. It follows that $F(Z) \simeq 0$ since finite one-sided twisted complexes built from contractible complexes are contractible. \square

Remark 5.7. The statement of Proposition 5.6 remains true if we replace “finite” with “bounded above”.

Proposition 5.8. *If $\mathcal{B} \subset \mathcal{C}$ generates, then the natural inclusion $\mathbf{B}(\mathcal{C}, \mathcal{J}_{\mathcal{B}}) \rightarrow \mathbf{B}(\mathcal{C}, \mathcal{J}_{\mathcal{C}})$ is the section of a deformation retract.*

Proof. Let $\mathcal{J} = \mathcal{J}_{\mathcal{C}} \subset \mathcal{C}$ be the subcategory of identity morphisms in \mathcal{C} , and let $\mathcal{J}' = \mathcal{J}_{\mathcal{B}} \subset \mathcal{C}$ be the subcategory of identity morphisms in \mathcal{B} . Let $C = \mathcal{C} \otimes_{\mathcal{J}} \mathcal{C}$ and $D = \mathcal{C} \otimes_{\mathcal{J}'} \mathcal{C}$.

Note that $C = D \oplus E$, where $E = \bigoplus_{Y \notin \mathcal{B}} \mathcal{C}Y \otimes Y\mathcal{C}$.

Consider the \mathcal{C} , \mathcal{C} -bimodules \mathbf{P}_C , \mathbf{P}_D and $\mathbf{A}_D = \text{Cone}(\mathbf{P}_D \rightarrow \mathcal{C})$. Since $C = D \oplus E$, it follows that \mathbf{P}_C can be expressed as a one-sided twisted complex constructed from one copy of \mathbf{P}_D and terms of the form

$$\mathbf{A}_D \otimes_e E \otimes_e \mathbf{A}_D \otimes_e \cdots \otimes_e E \otimes_e \mathbf{A}_D.$$

But one of the features of \mathbf{A}_D is $Y\mathbf{A}_D \simeq 0 \simeq \mathbf{A}_DY$ for all $Y \in \mathcal{B}$. Since \mathcal{B} generates, it follows that $Y\mathbf{A}_D \simeq 0 \simeq \mathbf{A}_DY$ for all $Y \in \mathcal{C}$. In particular $E \otimes_e \mathbf{A}_D \simeq 0 \simeq \mathbf{A}_D \otimes_e E$. Contracting the contractible terms yields the desired deformation retract

$$\mathbf{P}_C = \mathbf{P}_{D \oplus E} \rightarrow \mathbf{P}_D. \quad \square$$

5.3. The bar complex of the pretriangulated hull. Let \mathcal{C} be a dg category and $\text{Pretr}(\mathcal{C})$ its pretriangulated hull. We have a homotopy equivalence of bimodules

$$\mathbf{B}(\text{Pretr}(\mathcal{C}), \mathcal{J}_{\text{Pretr}(\mathcal{C})}) \rightarrow \mathbf{B}(\text{Pretr}(\mathcal{C}), \mathcal{J}_{\mathcal{C}})$$

implicitly constructed in the previous section. It will be useful to understand this map explicitly.

We first consider the relation between the two-sided bar complexes of \mathcal{C} and $\Sigma\mathcal{C}$. There is a map of $\Sigma\mathcal{C}$, $\Sigma\mathcal{C}$ -bimodules

$$\Phi: \mathbf{B}(\Sigma\mathcal{C}, \mathcal{J}_{\Sigma\mathcal{C}}) \rightarrow \mathbf{B}(\Sigma\mathcal{C}, \mathcal{J}_{\mathcal{C}})$$

defined as follows: given objects $X_0, \dots, X_r \in \Sigma\mathcal{C}$ of the form

$$X_i = \bigoplus_j \Sigma^{a_{i,j}} X_{ij}$$

and a sequence of composable morphisms

$$g_1 \otimes \cdots \otimes g_r \in X_0(\Sigma\mathcal{C})X_1 \otimes \cdots \otimes X_{r-1}(\Sigma\mathcal{C})X_r,$$

we define $\Phi(\text{Id} \|g_1\| \cdots \|g_r\| \text{Id})$ to be the sum of terms of the form

$$\pm \sigma_{j_0} \|(g_1)_{j_0, j_1}\| \|(g_2)_{j_1, j_2}\| \cdots \|(g_r)_{j_{r-1}, j_r}\| \pi_{j_r},$$

where $(g_i)_{j_{i-1}, j_i}$ indicates the component of g_i living in $X_{i-1, j_{i-1}} \mathcal{C} X_{i, j_i}$ and σ_{j_0} denotes the inclusion of X_{0, j_0} into X_0 and π_{j_r} denotes the projection of X_r onto X_{r, j_r} . It is an exercise to find the correct signs such that Φ defines a deformation retract.

Next we consider the relation between the two-sided bar complex of \mathcal{C} and $\text{Tw}(\mathcal{C})$. We define a map of \mathcal{C} , \mathcal{C} -bimodules

$$\Psi: \mathbf{B}(\text{Tw}(\mathcal{C}), \mathcal{J}_{\text{Tw}(\mathcal{C})}) \rightarrow \mathbf{B}(\text{Tw}(\mathcal{C}), \mathcal{J}_{\mathcal{C}})$$

as follows. Given objects $\text{tw}_{\alpha_i}(X_i) \in \text{Tw}(\mathcal{C})$ ($0 \leq i \leq r$) and an element

$$\text{Id} \|f_1\| \cdots \|f_r\| \text{Id} \in \mathbf{B}(\text{Tw}(\mathcal{C}), \mathcal{J}_{\text{Tw}(\mathcal{C})}),$$

we define $\Psi(\text{Id} \|f_1\| \cdots \|f_r\| \text{Id})$ to be the sum of terms of the form

$$\pm \phi \underbrace{\|\alpha_0\| \cdots \|\alpha_0\|}_{i_0} \|f_1\| \underbrace{\|\alpha_1\| \cdots \|\alpha_1\|}_{i_1} \cdots \underbrace{\|\alpha_{r-1}\| \cdots \|\alpha_{r-1}\|}_{i_{r-1}} \|f_r\| \underbrace{\|\alpha_r\| \cdots \|\alpha_r\|}_{i_r} \phi^{-1}$$

where $i_0, \dots, i_r \geq 0$ and ϕ and ψ denote Id_{X_0} and Id_{X_r} , regarded as degree zero (not necessarily closed) maps $X_0 \rightarrow \text{tw}_{\alpha_0}(X_0)$ and $X_r \rightarrow \text{tw}_{\alpha_r}(X_r)$. It is an exercise to find the correct signs and check that this defines a deformation retract.

Combining these gives a bimodule map relating the two-sided bar complex of \mathcal{C} and its pretriangulated hull; this map is a deformation retract and, in particular, a homotopy equivalence.

5.4. The semi-orthogonal bar complex.

Definition 5.9. Let Γ be a poset, and let $\mathcal{B}_\gamma \subset \mathcal{C}$ be full dg subcategories of \mathcal{C} , indexed by $\gamma \in \Gamma$ such that

- (1) every object $Z \in \mathcal{C}$ is homotopy equivalent to a one-sided twisted complex $\text{tw}_\delta(\bigoplus_{\gamma \in \Gamma} Z_\gamma)$ where $Z_\gamma \in \mathcal{B}_\gamma$, only finitely many Z_γ are nonzero and the twist δ is strictly lower triangular with respect to the partial order on Γ .
- (2) $\text{Hom}_{\mathcal{C}}(Y_\gamma, Y_{\gamma'})$ is contractible unless $\gamma \leq \gamma'$.

Then we say that \mathcal{B}_γ defines a Γ -indexed semi-orthogonal decomposition of \mathcal{C} .

Note that if $X \in \mathcal{B}_\gamma \cap \mathcal{B}_{\gamma'}$ for $\gamma \neq \gamma'$ then X is contractible. Thus, we usually assume, without loss of generality, that the subcategories \mathcal{B}_γ have no objects in common.

Let $\{\mathcal{B}_\gamma\}_{\gamma \in \Gamma}$ be a semi-orthogonal decomposition of \mathcal{C} . Define the full dg subcategory $\mathcal{B} := \bigcup_{\gamma \in \Gamma} \mathcal{B}_\gamma \subset \mathcal{C}$, which generates \mathcal{C} by Proposition 5.6.

For $X, Y \in \mathcal{B}$ we write $X \leq Y$ if $X \in \mathcal{B}_\gamma$ and $Y \in \mathcal{B}_{\gamma'}$ with $\gamma \leq \gamma'$. This relation is transitive and reflexive, but not anti-symmetric: $X \leq Y$ and $Y \leq X$ holds if and only if $X, Y \in \mathcal{B}_\gamma$ for some γ .

The *semi-orthogonal two-sided bar complex* of \mathcal{C} is defined to be the subcomplex $\mathbf{B}^\Gamma(\mathcal{C}, \mathcal{J}_\mathcal{B}) \subset \mathbf{B}(\mathcal{C}, \mathcal{J}_\mathcal{B})$ spanned by elements of the form

$$f_0 \parallel \cdots \parallel f_{r+1} \text{ in } \mathcal{C}X_1 \otimes X_1\mathcal{C}X_2 \otimes \cdots \otimes X_{r+1}\mathcal{C}$$

with $X_1 \geq X_2 \geq \cdots \geq X_{r+1}$.

Proposition 5.10. *Retain notation as above. The inclusion $\mathbf{B}^\Gamma(\mathcal{C}, \mathcal{J}_\mathcal{B}) \hookrightarrow \mathbf{B}(\mathcal{C}, \mathcal{J}_\mathcal{B})$ is the section of a deformation retract; the data of this deformation retract are \mathcal{C} , \mathcal{C} -bilinear.*

Proof. The complex $\mathbf{B}(\mathcal{C}, \mathcal{J}_\mathcal{B})$ is a one-sided twist of

$$\bigoplus_{r \geq 0} \bigoplus_{X_1, \dots, X_{r+1} \in \mathcal{B}} \Sigma^r \left(\mathcal{C}X_1 \otimes_{\mathbb{k}} X_1\mathcal{C}X_2 \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} X_{r+1}\mathcal{C} \right)$$

But $X_i\mathcal{C}X_{i+1} = \text{Hom}_{\mathcal{C}}(X_{i+1}, X_i)$ is contractible unless $X_i \geq X_{i+1}$. Contracting all such terms gives the desired deformation retract. Since the contractions are all of the form

$$\mathcal{C}X_1 \otimes_{\mathbb{k}} \underbrace{X_1\mathcal{B}X_2 \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} X_r\mathcal{B}X_{r+1}}_{\text{contract some factor}} \otimes X_{r+1}\mathcal{C},$$

(i.e. we only ever contract an ‘‘internal’’ tensor factor) the data of the deformation retract commute with the left and right \mathcal{C} -actions. \square

Since \mathcal{B} generates \mathcal{C} , Proposition 5.8 implies the following.

Theorem 5.11. *Retain notation as above. The full bar complex $\mathbf{B}(\mathcal{C})$ deformation retracts onto the semi-orthogonal bar complex $\mathbf{B}^\Gamma(\mathcal{C}, \mathcal{J}_\mathcal{B})$. The data of this deformation retract are \mathcal{C} , \mathcal{C} -bilinear.*

Corollary 5.12. *The semi-orthogonal bar complex $\mathbf{B}^\Gamma(\mathcal{C}, \mathcal{J}_\mathcal{B})$ has the structure of an A_∞ algebra object in \mathcal{C} , \mathcal{C} -bimodules, c.f. Remark 2.12. \square*

5.5. Hochschild homology (vertical trace). If \mathcal{A} is an additive \mathbb{k} -category then the (vertical) trace of \mathcal{A} is the \mathbb{k} -module

$$\text{HH}_0(\mathcal{A}) := \bigoplus_{X \in \mathcal{A}} \text{End}(X) / \text{span}_{\mathbb{k}} \{g_0 \circ g_1 - g_1 \circ g_0\}$$

where (g_0, g_1) runs over pairs of morphisms which are composable in either order.

Notation 5.13. *The class of $f \in \text{End}(X)$ in $\text{HH}_0(\mathcal{A})$ will be denoted $[f]$. We also write $[X] := [\text{Id}_X]$ for the associated class in $\text{HH}_0(\mathcal{A})$.*

Remark 5.14. If \mathcal{A} is monoidal, then $\mathrm{HH}_0(\mathcal{A})$ inherits the structure of an algebra via $[f] \cdot [g] := [f \otimes g]$.

It is natural to consider a derived version of the vertical trace, in which the relation $g_0 \circ g_1 = g_1 \circ g_0$ is not strictly enforced, but is achieved by the formal adjunction of a homotopy $h(g_0, g_1)$. In this setting it is also natural to allow our additive category to be an arbitrary dg category \mathcal{C} . After adjoining such homotopies, symbols of the form $h(g_0 g_1, g_2) - h(g_0, g_1 g_2) + h(g_2 g_0, g_1)$ are automatically closed. In order avoid creating such new closed elements, one is forced to adjoin higher homotopies $h(g_0, g_1, g_2)$. Continuing in this fashion, one obtains a complex

$$(11) \quad \mathbf{C}(\mathcal{C}) = \bigoplus_{r \geq 0} \bigoplus_{X_1, \dots, X_{r+1} \in \mathcal{C}} \Sigma^r \left(X_1 \mathcal{C} X_2 \otimes \cdots \otimes X_{r+1} \mathcal{C} X_1 \right)$$

where the direct sum is over objects $X_1, \dots, X_{r+1} \in \mathcal{C}$. We write $\|f_1\|f_2\| \cdots \|f_{r+1}$ with $f_i \in X_i \mathcal{C} X_{i+1}$ (with indices taken modulo $r+1$) for elementary tensors in the degree r part of (11).

The differential of such an element is

$$\begin{aligned} d_{\mathbf{C}(\mathcal{C})}(\|f_1\|f_2\| \cdots \|f_{r+1}) &:= (-1)^r \sum_{i=1}^{r+1} (-1)^{|f_1| + \cdots + |f_{i-1}|} \|f_1\| \cdots \|d_{\mathcal{C}}(f_i)\| \cdots \|f_{r+1} \\ &+ \sum_{i=1}^r (-1)^i \|f_1\| \cdots \|f_i f_{i+1}\| \cdots \|f_{r+1} + (-1)^{|f_1|(r-1+|f_2|+\cdots+|f_{r+1}|)} \|f_2\| \cdots \|f_{r+1} f_1. \end{aligned}$$

Note that the differential $d_{\mathbf{C}}$ splits as $d_{\oplus} + \delta$ where d_{\oplus} is the direct sum of differentials on the terms in (11), shown in the first line, while δ is an additional contribution shown in the second line.

The complex just constructed is called the cyclic bar complex of \mathcal{C} .

Definition 5.15. The *cyclic bar complex* of \mathcal{C} (relative to a unital subcategory $\mathcal{J} \subset \mathcal{C}$) is the dg \mathbb{k} -module $\mathbf{C}(\mathcal{C}, \mathcal{J})$ obtained as the quotient of the two-sided bar complex $\mathbf{B}(\mathcal{C}, \mathcal{J})$ modulo the \mathbb{k} -span $[\mathcal{C}, \mathbf{B}(\mathcal{C})]$ of elements of the form $f \cdot m - (-1)^{|m||f|} m \cdot f$ for all $m \in \mathbf{B}(\mathcal{C}, \mathcal{J})$ and all morphisms f in \mathcal{C} .

The *Hochschild homology* $\mathrm{HH}(\mathcal{C}, \mathcal{J})$ is defined to be the homology of $\mathbf{C}(\mathcal{C}, \mathcal{J})$. Hochschild homology is written with homological convention for gradings, and so we write $\mathrm{HH}_k(\mathcal{C}, \mathcal{J}) := H^{-k}(\mathbf{C}(\mathcal{C}, \mathcal{J}))$ for $k \geq 0$.

Remark 5.16. Instead of (11), the definition describes the cyclic bar complex as:

$$(12) \quad \mathbf{C}(\mathcal{C}) = \bigoplus_{r \geq 0} \bigoplus_{X_0, X_1, \dots, X_{r+1} \in \mathcal{C}} \Sigma^r \left(X_0 \mathcal{C} X_1 \otimes X_1 \mathcal{C} X_2 \otimes \cdots \otimes X_{r+1} \mathcal{C} X_0 \right) / \sim$$

where the linear relation \sim is generated by

$$(-1)^{|f_0|(r+|f_1|+\cdots+|f_{r+1}|)} f_0 \|f_1\|f_2\| \cdots \|f_{r+1} \sim \mathrm{Id} \|f_1\|f_2\| \cdots \|f_{r+1} f_0$$

The previously introduced notation for elements of the cyclic bar complex simply drops the leading identity term after such a rewrite—in the example above: $\|f_1\|f_2\|\cdots\|f_{r+1}f_0$.

Remark 5.17. When $\mathcal{J} = \mathcal{J}_e$ is the subcategory of identity morphisms, then we have $\mathbf{C}(\mathcal{C}, \mathcal{J}) = \mathbf{C}(\mathcal{C})$ as in (11). If $\mathcal{B} \subset \mathcal{C}$ is a full subcategory and $\mathcal{J} = \mathcal{J}_{\mathcal{B}}$ is the subcategory of identity morphisms in \mathcal{B} , then $\mathbf{C}(\mathcal{C}, \mathcal{J}_{\mathcal{B}}) = \mathbf{C}(\mathcal{B})$.

Remark 5.18. If \mathcal{C} has trivial grading and trivial differential then $\mathrm{HH}_0(\mathcal{C})$ coincides with the vertical trace defined earlier.

An inclusion of subcategories $\mathcal{J} \subset \mathcal{J} \subset \mathcal{C}$ gives a chain map $\mathbf{C}(\mathcal{C}, \mathcal{J}) \rightarrow \mathbf{C}(\mathcal{C}, \mathcal{J})$. In particular, if $\mathcal{B} \subset \mathcal{C}$ is a full subcategory then we have a canonical inclusion $\mathbf{C}(\mathcal{B}) \hookrightarrow \mathbf{C}(\mathcal{C})$.

The following is our main tool for computing $\mathrm{HH}(\mathcal{C})$.

Theorem 5.19. *If $\mathcal{B} \subset \mathcal{C}$ generates \mathcal{C} (see Definition 5.5) then the natural inclusion $\mathbf{C}(\mathcal{B}) \hookrightarrow \mathbf{C}(\mathcal{C})$ is the section of a deformation retract. More generally, if Γ is a finite poset and $\{\mathcal{B}_\gamma\}_{\gamma \in \Gamma}$ gives a semi-orthogonal decomposition of \mathcal{C} then $\mathbf{C}(\mathcal{C})$ deformation retracts onto $\bigoplus_{\gamma \in \Gamma} \mathbf{C}(\mathcal{B}_\gamma)$.*

Proof. For the first assertion, recall from Proposition 5.8 that $\mathbf{B}(\mathcal{C})$ deformation retracts onto $\mathbf{B}(\mathcal{C}, \mathcal{J}_{\mathcal{B}})$, which induces a deformation retract

$$\mathbf{C}(\mathcal{C}) = \mathbf{B}(\mathcal{C})/[\mathcal{C}, \mathbf{B}(\mathcal{C})] \xrightarrow{\cong} \mathbf{B}(\mathcal{C}, \mathcal{J}_{\mathcal{B}})/[\mathcal{C}, \mathbf{B}(\mathcal{C}, \mathcal{J}_{\mathcal{B}})] = \mathbf{C}(\mathcal{B})$$

by Remark 5.17 since \mathcal{B} is a full subcategory of \mathcal{C} .

To prove the second assertion, we set $\mathcal{B} = \bigcup_{\gamma} \mathcal{B}_\gamma$ and recall from Theorem 5.11 that $\mathbf{B}(\mathcal{C})$ deformation retracts onto the semi-orthogonal bar complex $\mathbf{B}^\Gamma(\mathcal{C}, \mathcal{J}_{\mathcal{B}})$ in the category of \mathcal{C} , \mathcal{C} -bimodules. There is an induced deformation retract

$$\mathbf{C}(\mathcal{C}) = \mathbf{B}(\mathcal{C})/[\mathcal{C}, \mathbf{B}(\mathcal{C})] \xrightarrow{\cong} \mathbf{B}^\Gamma(\mathcal{C}, \mathcal{J}_{\mathcal{B}})/[\mathcal{C}, \mathbf{B}^\Gamma(\mathcal{C}, \mathcal{J}_{\mathcal{B}})].$$

The complex $\mathbf{B}^\Gamma(\mathcal{C}, \mathcal{J}_{\mathcal{B}})$ is a one-sided twist of

$$\bigoplus_{r \geq 0} \bigoplus_{X_1, \dots, X_{r+1} \in \mathcal{B}} \Sigma^r \left(\mathcal{C}X_1 \otimes X_1\mathcal{C}X_2 \otimes \cdots \otimes X_{r+1}\mathcal{C} \right)$$

with $X_1 \geq X_2 \geq \cdots \geq X_{r+1} =: X_0$. By semi-orthogonality, the summands with $X_1 > X_0$ become contractible in the quotient $\mathbf{B}^\Gamma(\mathcal{C}, \mathcal{J}_{\mathcal{B}})/[\mathcal{C}, \mathbf{B}^\Gamma(\mathcal{C}, \mathcal{J}_{\mathcal{B}})]$. Contracting these induces a deformation retract onto the subcomplex where $X_1, \dots, X_r, X_0 \in \mathcal{B}_\gamma$ for some γ . For each γ the contribution of all such terms is the two-sided bar complex $\mathbf{B}(\mathcal{B}_\gamma)$. By inspection the differential preserves summands, and we obtain the direct sum decomposition in the statement. \square

The following corollary was proved in [Kuz09].

Corollary 5.20. *Suppose $\{\mathcal{B}_\gamma \subset \mathcal{C}\}$ defines a Γ -indexed semi-orthogonal decomposition of \mathcal{C} . Then we have*

$$\mathrm{HH}(\mathcal{C}) \cong \bigoplus_{\gamma \in \Gamma} \mathrm{HH}(\mathcal{B}_\gamma).$$

Corollary 5.21. *If \mathcal{C} is a dg category, then the cyclic bar complex of $\text{Pretr}(\mathcal{C})$ deformation retracts onto the cyclic bar complex of \mathcal{C} , and we have a natural isomorphism $\text{HH}(\mathcal{C}) \cong \text{HH}(\text{Pretr}(\mathcal{C}))$ induced by the inclusion $\mathcal{C} \hookrightarrow \text{Pretr}(\mathcal{C})$. \square*

Example 5.22. *If \mathcal{A} is an additive category (regarded as a dg category trivially) then we have the natural isomorphism $\text{HH}(\mathcal{A}) \cong \text{HH}(\text{Ch}^b(\mathcal{A}))$ induced by the inclusion $\mathcal{A} \hookrightarrow \text{Ch}^b(\mathcal{A})$.*

If $X \in \mathcal{C}$ is an object, then we write $[X] := [\text{Id}_X]$ for its class in $\text{HH}_0(\mathcal{C}) \subset \text{HH}(\mathcal{C})$. If X is a one-sided twisted complex constructed from objects X^i , it is natural to ask for the relation between the classes $[X]$ and $[X^i]$.

Lemma 5.23. *For objects $X, Y \in \mathcal{C}$ and $f: X \rightarrow Y$, we have:*

$$(13a) \quad [X \oplus Y] = [X] + [Y]$$

$$(13b) \quad [\Sigma^1 X] = -[X]$$

$$(13c) \quad [\text{Cone}(f: X \rightarrow Y)] = [Y] - [X]$$

Proof. It is sufficient to prove (13c). Let $Z = \text{Cone}(X \xrightarrow{f} Y)$, let i_X, i_Y denote the inclusions of X and Y into Z and let π_X, π_Y denote the projections of Z onto X and Y . Note that i_X and π_X have degrees ± 1 , and

$$\begin{aligned} d_{\mathcal{C}}(i_X) &= i_Y f, & d_{\mathcal{C}}(i_Y) &= 0, & d_{\mathcal{C}}(\pi_X) &= 0, & d_{\mathcal{C}}(\pi_Y) &= -f \pi_X, \\ \pi_X i_X &= \text{Id}_X, & \pi_Y i_Y &= \text{Id}_Y, & i_X \pi_X + i_Y \pi_Y &= \text{Id}_Z. \end{aligned}$$

So we compute

$$\begin{aligned} d_{\mathcal{C}}(\|i_X\|\pi_X) &= -\|\text{Id}_X - \|i_X\|\pi_X - \|i_Y f\|\pi_X, \\ d_{\mathcal{C}}(\|i_Y\|\pi_Y) &= \|\text{Id}_Y - \|i_Y\|\pi_Y + \|i_Y\|f\pi_X, \\ d_{\mathcal{C}}(\|i_Y f\|\pi_X) &= 0 - \|i_Y f\|\pi_X + \|i_Y\|f\pi_X, \end{aligned}$$

since $\pi_X i_Y = 0$ (each of the above has cohomological degree -1 before applying $d_{\mathcal{C}}$). Therefore

$$d_{\mathcal{C}}(\|i_Y f\|\pi_X - \|i_X\|\pi_X - \|i_Y\|\pi_Y) = \|\text{Id}_Z - (\|\text{Id}_Y - \|\text{Id}_X\|)\|,$$

which shows that $[\text{Id}_Z] \simeq [\text{Id}_Y] - [\text{Id}_X]$ in $\text{HH}_0(\mathcal{C})$. \square

Corollary 5.24. *Let \mathcal{C} be a dg category and consider an object $X = \text{tw}_{\alpha}(\bigoplus_i X^i)$ in $\text{Pretr}(\mathcal{C})$, then under the identification $\text{HH}(\text{Pretr}(\mathcal{C})) \cong \text{HH}(\mathcal{C})$ the class of X is given by the Euler characteristic*

$$[X] = \sum_i (-1)^i [X^i].$$

5.6. Connes differential and HKR isomorphism. The cyclic bar complex of a dg category has a canonical differential \mathbb{B} of cohomological degree -1 [LQ84] that we now describe. Given $\|f_1\| \cdots \|f_{r+1}\| \in \mathbf{C}^{-r}(\mathcal{C})$ with $f_1 \in X_1 \mathcal{C} X_2$, we define

$$\begin{aligned} t(\|f_1\| \cdots \|f_{r+1}\|) &= (-1)^{|f_{r+1}|(|f_1|+\cdots+|f_r|)} \|f_{r+1}\| \|f_1\| \cdots \|f_r\| \\ s(\|f_1\| \cdots \|f_{r+1}\|) &= \|\mathrm{Id}_{X_1}\| \|f_1\| \cdots \|f_{r+1}\| \end{aligned}$$

Now we define $N = 1 + \dots + t^r$ and $\mathbb{B} = (1 - t)sN$. The operator \mathbb{B} is sometimes called Connes differential.

Example 5.25. For $\|f_1\| \in X_1 \mathcal{C} X_1 \subset \mathbf{C}^0(\mathcal{C})$ we have

$$\mathbb{B}(\|f_1\|) = \|\mathrm{Id}_{X_1}\| \|f_1\| - \|f_1\| \|\mathrm{Id}_{X_1}\| \in \mathbf{C}^{-1}(\mathcal{C}).$$

For $\|f_1\| \|f_2\| \in \Sigma^1(X_1 \mathcal{C} X_2 \otimes X_2 \mathcal{C} X_1) \subset \mathbf{C}^{-1}(\mathcal{C})$ we have

$$\begin{aligned} \mathbb{B}(\|f_1\| \|f_2\|) &= (1 - t) (\|\mathrm{Id}_{X_1}\| \|f_1\| \|f_2\| + (-1)^{|f_2||f_2|} \|\mathrm{Id}_{X_2}\| \|f_2\| \|f_1\|) \\ &= \|\mathrm{Id}_{X_1}\| \|f_1\| \|f_2\| + (-1)^{|f_2||f_1|} \|\mathrm{Id}_{X_2}\| \|f_2\| \|f_1\| - (-1)^{|f_2||f_1|} \|f_2\| \|\mathrm{Id}_{X_1}\| \|f_1\| - \|f_1\| \|\mathrm{Id}_{X_2}\| \|f_2\|. \end{aligned}$$

The following is well known.

Lemma 5.26 ([LQ84]). *We have $d_{\mathbf{C}}\mathbb{B} + \mathbb{B}d_{\mathbf{C}} = 0$ and $\mathbb{B}^2 = 0$.*

Theorem 5.27 ([HKR62, LQ84]). *Let A be the ring of functions on a smooth affine scheme X . Then there is an algebra isomorphism*

$$\Omega^*(X) \simeq \mathrm{HH}_*(A),$$

which identifies the de Rham differential D on the algebra of differential forms $\Omega^(X)$ on the left with the (induced) Connes differential \mathbb{B} on the right hand side.*

Example 5.28. We have $\mathrm{HH}_1(A) = A \otimes A / (ab \otimes c - a \otimes bc + ac \otimes b)$. We can identify this with $\Omega^1(X)$ by sending $a \otimes b \rightarrow aD(b)$. Indeed, $aD(bc) = abD(c) + acD(b)$. Now $\mathbb{B}(a) = 1 \otimes a - a \otimes 1 \in A \otimes A$ is identified with $D(a) - aD(1) = D(a) \in \Omega^1(X)$.

Example 5.29. Let $R = \mathbb{C}[x_1, \dots, x_n]$ be the algebra of functions on \mathbb{C}^n . Then

$$\mathrm{HH}_k(R) \simeq \Omega^k(\mathbb{C}^n) \simeq \mathbb{C}[x_1, \dots, x_n] \otimes \wedge^k[\theta_1, \dots, \theta_n],$$

where $\theta_i = D(x_i)$. The de Rham differential can be written as

$$(14) \quad D = \sum_i \theta_i \frac{\partial}{\partial x_i}.$$

6. THE DG MONOIDAL CENTER AND TRACE

6.1. Monoidal dg categories and the shuffle product. Note that any (dg) algebra A can be regarded as a dg category with one object. By the same token, a commutative (dg) algebra A can be viewed as a *monoidal* dg category with one object.

A monoidal dg category is a dg category \mathcal{C} equipped with an object $\mathbb{1} \in \mathcal{C}$, a functor $\star : \mathcal{C} \otimes_{\mathbb{k}} \mathcal{C} \rightarrow \mathcal{C}$, and closed degree zero natural isomorphisms (*associator* and *unitors*)

$$(X \star Y) \star Z \cong X \star (Y \star Z), \quad \mathbb{1} \star X \cong X \cong X \star \mathbb{1}$$

satisfying the usual coherence relations for monoidal categories (on the nose; not up to homotopy). The associator and unitor isomorphisms will usually be suppressed from the notation, and we will refer to $(\mathcal{C}, \star, \mathbb{1})$ as a dg monoidal category. If the associators and unitors are in fact identity morphisms, then $(\mathcal{C}, \star, \mathbb{1})$ is said to be *strictly monoidal*. The usual Eckmann-Hilton argument shows that $\text{End}_{\mathcal{C}}(\mathbb{1})$ is always a commutative dg algebra.

We say that a dg monoidal category *has duals* if every object X has a left dual X^* and a right dual *X with evaluation and coevaluation maps

$$\text{ev}_X: X^* \star X \rightarrow \mathbb{1}, \quad \text{coev}_X: \mathbb{1} \rightarrow X \star X^*, \quad \text{ev}'_X: X \star {}^*X \rightarrow \mathbb{1}, \quad \text{coev}'_X: \mathbb{1} \rightarrow {}^*X \star X$$

satisfying the usual *string-straightening* axioms. Note that the existence of duals is a property of \mathcal{C} .

For a dg monoidal category \mathcal{C} with duals, the operations $(-)^*$ and ${}^*(-)$ extend to (contravariant) monoidal dg functors. A *pivotal structure* on \mathcal{C} is a monoidal natural isomorphism between $(-)^*$ and ${}^*(-)$ (equivalently, a monoidal natural isomorphism between $\text{Id}_{\mathcal{C}}$ and $(-)^{**}$).

We say a dg monoidal category \mathcal{C} is *strictly pivotal* if it is strictly monoidal and we have $X^* = {}^*X$ for every object X , with the identity natural transformation as pivotal structure.

A monoidal structure on \mathcal{C} endows the two-sided bar complex $\mathbf{B}(\mathcal{C}, \mathcal{J})$ with the additional structure of an algebra, via the so-called *shuffle product* (e.g. [LQ84, EZ53]) which we recall below.

Suppose \mathcal{C} is a dg monoidal category. Let

$$\underline{f} = f_0 \parallel \cdots \parallel f_{r+1} \in \Sigma^r (X_0 \mathcal{C} X_1 \otimes \cdots \otimes X_{r+1} \mathcal{C} X_{r+2}) \subset \mathbf{B}_r(\mathcal{C}, \mathcal{J})$$

and

$$\underline{g} = g_0 \parallel \cdots \parallel g_{s+1} \in \Sigma^s (Y_0 \mathcal{C} Y_1 \otimes \cdots \otimes Y_{s+1} \mathcal{C} Y_{s+2}) \subset \mathbf{B}_s(\mathcal{C}, \mathcal{J})$$

be two elements in $\mathbf{B}(\mathcal{C}, \mathcal{J})$. To define the product $\underline{f} * \underline{g}$ we set

$$e_i = \begin{cases} f_i \star \text{Id} & 1 \leq i \leq r \\ \text{Id} \star g_{i-r} & r < i \leq r+s \end{cases}$$

where we keep an open mind about what objects the identity morphisms are associated with. Now we set

$$\underline{f} * \underline{g} := (-1)^{|\underline{f}|s} \sum_{\pi \in S_{(r,s)}} (-1)^{w\sigma(\pi, \underline{f}, \underline{g})} (f_0 \star g_0) \parallel e_{\pi(1)} \parallel \cdots \parallel e_{\pi(r+s)} \parallel (f_{r+1} \star g_{s+1})$$

where $S_{(r,s)} \subset S_{r+s}$ denotes the set of shuffle permutations, and $w\sigma(\pi, \underline{f}, \underline{g})$ denotes the *weighted sign* of the permutation π , to which a transposition of $f_i \star \text{Id}$ and $\text{Id} \star g_j$ contributes $(-1)^{|f_i||g_j|}$. For the $e_{\pi(i)}$ in the summands of this formula, we implicitly choose the identity morphism factor which makes the sequence of morphisms in the summand composable. Note that different summands require different choices, although this is suppressed in the notation.

The shuffle product together with the coproduct Δ give the two-sided bar complex $B(\mathcal{C}, \mathcal{J})$ the structure of a bialgebra.

6.2. The quadmodule associated to a dg monoidal category. Let \mathcal{C} now be a dg monoidal category. Fix objects $X_1, X_2, Y_1, Y_2 \in \mathcal{C}$, and let $f \in \text{Hom}_{\mathcal{C}}(X_1 \star X_2, Y_1 \star Y_2)$ be given. We also consider objects $X'_i, Y'_i \in \mathcal{C}$ ($i = 1, 2$) and morphisms

$$a_i \in \text{Hom}_{\mathcal{C}}(Y_i, Y'_i), \quad b_i \in \text{Hom}_{\mathcal{C}}(X'_i, X_i).$$

Then we define the following operations:

$$\begin{aligned} a_1 \circ_1 f &:= (a_1 \star \text{Id}_{Y_2}) \circ f, & a_2 \circ_2 f &:= (\text{Id}_{Y_1} \star a_2) \circ f, \\ f \circ_1 b_1 &:= f \circ (b_1 \star \text{Id}_{X_2}), & f \circ_2 b_2 &:= f \circ (\text{Id}_{X_1} \star b_2). \end{aligned}$$

These operations give a $(\mathcal{C} \otimes_{\mathbb{k}} \mathcal{C}), (\mathcal{C} \otimes_{\mathbb{k}} \mathcal{C})$ -bimodule structure on

$$\mathcal{X} := \bigoplus_{X_1, X_2, Y_1, Y_2} (Y_1 \star Y_2) \mathcal{C}(X_1 \star X_2).$$

We may regard \mathcal{X} just defined as a *quadmodule* over \mathcal{C} . By combining the monoidal structure and composition of morphisms in \mathcal{C} , we have morphisms

$$\mu_{\swarrow}, \mu_{\nearrow} : \mathcal{X} \otimes_{\mathbb{k}} \mathcal{X} \rightarrow \mathcal{X}$$

defined by

$$\mu_{\swarrow}(f, g) := (f \star \text{Id}) \circ (\text{Id} \star g)$$

and

$$\mu_{\nearrow}(f, g) := (\text{Id} \star f) \circ (g \star \text{Id})$$

whenever these compositions make sense. These operations interact with the quadmodule structure according to

$$\begin{aligned} \mu_{\swarrow}(f \circ_2 a, g) &= \mu_{\swarrow}(f, a \circ_1 g) \\ \mu_{\swarrow}(a \circ_1 f, g \circ_2 b) &= a \circ_1 \mu_{\swarrow}(f, g) \circ_2 b \\ \mu_{\swarrow}(a \circ_2 f, a' \circ_2 g) &= (a \star a') \circ_2 \mu_{\swarrow}(f, g) \\ \mu_{\swarrow}(f \circ_1 b, g \circ_1 b') &= \mu_{\swarrow}(f, g) \circ_1 (b \star b'), \end{aligned}$$

with similar identities involving μ_{\nearrow} (swapping the roles of \circ_1 and \circ_2).

From \mathcal{X} we obtain a \mathcal{C}, \mathcal{C} -bimodule by forgetting the “northeast” and “southwest” actions of \mathcal{C} . Precisely, \mathcal{X}_{12} equals \mathcal{X} , but with \mathcal{C}, \mathcal{C} -bimodule structure

$$a \otimes f \otimes b \mapsto a \circ_1 f \circ_2 b, \quad a, b \in \mathcal{C}, \quad f \in \mathcal{X}_{12}.$$

One may define a bimodule \mathcal{X}_{21} in a similar fashion, but we will not need it.

Remark 6.1. The map μ_{\swarrow} makes \mathcal{X}_{12} into an algebra object in \mathcal{C}, \mathcal{C} -bimodules, with unit $\mathcal{C} \rightarrow \mathcal{X}_{12}$ given by the bimodule map sending $\text{Id}_X \in \mathcal{C}$ to the canonical isomorphism $\mathbb{1} \star X \rightarrow X \star \mathbb{1}$, regarded as an element of $X \mathcal{X}_{12} X$.

Remark 6.2. The map μ_{\succ} also defines an associative multiplication on \mathcal{X}_{12} which interacts with the bimodule structure in a nonstandard way:

$$\begin{aligned}\mu_{\succ}(f \circ_1 a, g) &= \mu_{\succ}(f, a \circ_2 g) \\ \mu_{\succ}(a \circ_2 f, g \circ_1 b) &= a \circ_2 \mu_{\succ}(f, g) \circ_1 b \\ \mu_{\succ}(a \circ_1 f, a' \circ_1 g) &= (-1)^{|a'|(|a|+|f|)}(a' \star a) \circ_1 \mu_{\succ}(f, g) \\ \mu_{\succ}(f \circ_2 b, g \circ_2 b') &= (-1)^{|b|(|b'|+|g|)}\mu_{\succ}(f, g) \circ_2 (b' \star b),\end{aligned}$$

By fixing $Y_2 = Z'$ and $X_1 = Z$, we obtain sub-bimodules of \mathcal{X}_{12} of the form

$$\mathcal{X}_{12}(Z', Z) := \bigoplus_{X, Y \in \mathcal{C}} (Y \star Z') \mathcal{C}(Z \star X) \subset \mathcal{X}_{12}$$

Note that μ_{\nwarrow} restricts to morphisms of \mathcal{C} , \mathcal{C} -bimodules

$$\mu_{\nwarrow} : \mathcal{X}_{12}(Z', Z) \otimes_{\mathcal{C}} \mathcal{X}_{12}(U', U) \rightarrow \mathcal{X}_{12}(Z' \star U', Z \star U),$$

while μ_{\succ} restricts to morphisms

$$\mu_{\succ} : \mathcal{X}_{12}(Z'', Z') \otimes_{\mathbb{k}} \mathcal{X}_{12}(Z', Z) \rightarrow \mathcal{X}_{12}(Z'', Z)$$

In particular the bimodule $\mathcal{X}_{12}(Z) := \mathcal{X}_{12}(Z, Z)$ inherits an associative multiplication (which respects the dg \mathbb{k} -module structure and is compatible with the \mathcal{C} , \mathcal{C} -bimodule structure).

6.3. The dg monoidal centralizer. Let \mathcal{C} be a dg monoidal category, and fix an object $Z \in \mathcal{C}$. We would like to discuss what it means for Z to be central in \mathcal{C} . Just as in the usual Drinfeld center, this is not a *property* enjoyed by Z , but rather additional *structure* which must be provided.

Actually, we will consider the slightly broader problem of defining what it means for Z to centralize a full dg monoidal subcategory $\mathcal{M} \subset \mathcal{C}$. For Z to centralize \mathcal{M} (up to homotopy) requires the following data:

- (1) for each object $X \in \mathcal{M}$, a degree zero closed morphism, called *half-braiding*, $\tau_X \in (X \star Z) \mathcal{C}(Z \star X)$.
- (2) for each closed morphism $f \in Y \mathcal{M} X$ a homotopy $h_f \in (X \star Z) \mathcal{C}(Z \star X)$ with $d_{\mathcal{C}}(h_f) = f \circ_1 \circ \tau_X - \tau_Y \circ_2 f$
- (3) certain higher homotopies.

To get a feeling for the sort of higher homotopies required, observe that for each pair of closed morphisms $f_0 \in X_0 \mathcal{M} X_1$ and $f_1 \in X_1 \mathcal{M} X_2$ we have *two ways* of commuting $f_0 \circ f_1$ past Z . First, we have the homotopy $h_{f_0 \circ f_1}$. But we also have $h_{f_0} \circ_2 f_1 + (-1)^{|f_0|} f_0 \circ_1 \circ h_{f_1}$. We should require the difference of these two homotopies (which is a closed morphism of degree -1) to be null-homotopic. The various higher homotopies required are, in fact, already organized for us in the form of the two-sided bar complex.

Definition 6.3. Let \mathcal{C} be a dg monoidal category and $\mathcal{M} \subset \mathcal{C}$ a subcategory. The *dg monoidal centralizer of \mathcal{M} in \mathcal{C}* is the dg category $\mathcal{Z}_{\mathcal{C}}^{\text{dg}}(\mathcal{M})$ whose objects are pairs (Z, τ) where $Z \in \mathcal{C}$ and $\tau : \mathbf{B}(\mathcal{M}) \rightarrow \mathcal{X}_{12}(Z)$ is a map of \mathcal{C} , \mathcal{C} -bimodules as well as a map

of dg algebras. The complex of morphisms in $\mathcal{Z}_c^{\text{dg}}(\mathcal{M})$ from (Z, τ) to (Z', τ') is the subcomplex of $\text{Hom}_c(Z, Z')$ consisting of those morphisms which commute strictly with the *structure maps* in the images of $\mathbf{B}(\mathcal{M})$. I.e. those $z \in \text{Hom}_c(Z, Z')$ such that for $f \in Y\mathbf{B}(\mathcal{M})X$ we have

$$z \circ_2 \tau(f) = \tau'(f) \circ_1 z$$

The dg Drinfeld center is defined to be $\mathcal{Z}^{\text{dg}}(\mathcal{C}) := \mathcal{Z}_c^{\text{dg}}(\mathcal{C})$.

Remark 6.4. It is perhaps better to weaken the condition on morphisms in $\mathcal{Z}^{\text{dg}}(\mathcal{C})$ to strictly commute with the structure maps from $\mathbf{B}(\mathcal{C})$. This would bring us into the world of A_∞ categories, which we choose to avoid for the moment. In the rest of this paper, we only consider objects of the dg Drinfeld center and not morphisms.

The dg Drinfeld center $\mathcal{Z}^{\text{dg}}(\mathcal{C})$ has a tensor product defined by

$$(Z, \tau) \star (Z', \tau') := (Z \star Z', \tau''),$$

where τ'' is the composition of maps

$$\mathbf{B}(\mathcal{C}) \rightarrow \mathbf{B}(\mathcal{C}) \otimes_c \mathbf{B}(\mathcal{C}) \rightarrow \mathcal{X}_{12}(Z) \otimes_c \mathcal{X}_{12}(Z') \rightarrow \mathcal{X}_{12}(Z \star Z').$$

Here, the first map is the coproduct on the two-sided bar complex, and the second map sends

$$f \otimes f' \mapsto (f \star \text{Id}_{Z'}) \star (\text{Id}_Z \star f') \in X''\mathcal{X}_{12}(Z \star Z')X$$

for all $f' \in X'\mathcal{X}_{12}(Z')X$ and all $f \in X''\mathcal{X}_{12}(Z)X'$.

The following is immediate.

Proposition 6.5. *There is a natural forgetful functor $\mathcal{Z}^{\text{dg}}(\mathcal{C}) \rightarrow \mathcal{C}$. It is monoidal.*

Remark 6.6. It is well known that the Drinfeld center of a monoidal category is braided. It is natural to expect that dg Drinfeld center is braided in a dg sense, that is, the braiding is natural up to homotopy. To define such a structure, it seems likely that passage to the world of A_∞ (braided monoidal) categories and functors is unavoidable. We save such explorations for future work.

Remark 6.7. If \mathcal{C} is a monoidal category, considered as a dg category with trivial differential, then the Drinfeld center $\mathcal{Z}^0(\mathcal{C})$ is a monoidal subcategory of $\mathcal{Z}^{\text{dg}}(\mathcal{C})$.

In the following remarks we spell out the meaning of some of the structure maps that are part of the data of a central object (Z, τ) .

Remark 6.8. For each $X \in \mathcal{C}$ the two-sided bar complex has a distinguished degree zero closed element $\text{Id}_X \parallel \text{Id}_X$. The image of this element in $\mathcal{X}_{12}(Z)$ under τ will be denoted τ_X and is called the half-braiding of X with Z . Then each degree zero element $a \parallel b$ gets sent to $a \circ_1 \tau_X \circ_2 b$ where X is the codomain of b (same as the domain of a), since τ commutes with the bimodule structures on $\mathbf{B}(\mathcal{C})$ and $\mathcal{X}_{12}(Z)$.

Remark 6.9. In $\mathbf{B}(\mathcal{C})$ we have $(\mathrm{Id}_X \parallel \mathrm{Id}_X) * (\mathrm{Id}_Y \parallel \mathrm{Id}_Y) = \mathrm{Id}_{X*Y} \parallel \mathrm{Id}_{X*Y}$. By definition of the center, τ is a map of dg algebras, which implies

$$\tau_{X*Y} = (\mathrm{Id}_X * \tau_Y) \circ (\tau_X * \mathrm{Id}_Y)$$

and one recovers the familiar compatibility between half-braiding morphisms and the monoidal structure in \mathcal{C} .

Remark 6.10. For each morphism $f \in Y\mathcal{C}X$ of degree l the two-sided bar complex has a degree $l - 1$ element of the form $\mathrm{Id}_Y \parallel f \parallel \mathrm{Id}_X$, whose image under τ we denote by h_f . If f is closed then we have

$$d(h_f) = d(\tau(\mathrm{Id}_Y \parallel f \parallel \mathrm{Id}_X)) = \tau(f \parallel \mathrm{Id}_X - \mathrm{Id}_Y \parallel f) = f \circ_1 \tau_X - \tau_Y \circ_2 f,$$

since τ commutes with the differentials in $\mathbf{B}(\mathcal{C})$ and $\mathcal{X}_{12}(Z)$. This means h_f is a homotopy for commuting f through the half-braiding with Z .

Similarly for each sequence of composable (closed) morphisms f, g we have an element $h_{f,g} = \tau(\parallel f \parallel g \parallel)$ (dropping explicit occurrences of identity maps) satisfying

$$\begin{aligned} d(h_{f,g}) &= d(\tau(\parallel f \parallel g \parallel)) \\ &= \tau(d(\parallel f \parallel g \parallel)) \\ &= \tau(f \parallel g \parallel - \parallel f \circ g \parallel + \parallel f \parallel g) \\ &= (-1)^{|f|} f \circ_1 h_g - h_{f \circ g} + h_f \circ_2 g \end{aligned}$$

and so on. In summary, we see that τ gives the data of a half-braiding with Z that is natural up to coherent homotopy.

Remark 6.11. If \mathcal{C} is an arbitrary dg category, there is a functor from $Z^{dg}(\mathcal{C})$ to the usual Drinfeld center of the homotopy category $Z^0(H^0(\mathcal{C}))$. It sends a central object (Z, τ) to (Z, τ_X) and forgets all higher homotopies.

Lemma 6.12. *If (Z, τ) is an object of $Z^{dg}(\mathcal{C})$ and X an object of \mathcal{C} that has a right dual, then $\tau_X : X * Z \rightarrow X * Z$ is a homotopy equivalence.*

Proof. If $*X$ is a right dual of X with

$$\mathrm{ev}' : X * *X \rightarrow \mathbb{1}, \quad \mathrm{coev}' : \mathbb{1} \rightarrow *X * X$$

then we set

$$\tau_X^{-1} := (\mathrm{ev}' * \mathrm{Id}_{Z*X}) \circ (\mathrm{Id}_X * \tau_{*X} * \mathrm{Id}_{*X}) \circ (\mathrm{Id}_{X*Z} * \mathrm{coev}')$$

and one can check that $\tau_X \tau_X^{-1} \simeq \mathrm{Id}_{X*Z}$ and $\tau_X^{-1} \tau_X \simeq \mathrm{Id}_{Z*X}$. \square

Remark 6.13. The dg monoidal center $Z^{dg}(\mathcal{C})$ embeds as a monoidal dg subcategory of $Z^{dg}(\mathrm{Pretr}(\mathcal{C}))$. To see this, let (Z, τ) be an object of $Z^{dg}(\mathcal{C})$. The half-braiding morphism τ extends to $\Sigma\mathcal{C}$ in a trivial fashion, so (Z, τ) can be thought of as an

object of $\mathcal{Z}^{\text{dg}}(\Sigma\mathcal{C})$. Next, given a morphism $f: X \rightarrow Y$, the half-braiding of the cone $\text{Cone}(f) := [X \xrightarrow{f} Y]$ past Z is defined by the following morphism:

$$\tau_{\text{Cone}(f)} : \begin{array}{ccc} [Z \star X & \xrightarrow{\text{Id}_Z \star f} & Z \star Y \\ \downarrow \tau_X & \searrow h_f & \downarrow \tau_Y \\ [X \star Z & \xrightarrow{f \star \text{Id}_Z} & Y \star Z \end{array}$$

Furthermore, the (higher) homotopies for half-braidings for cones are analogously determined by τ . Iterating this construction, we see that (Z, τ) is derived central for one-sided twisted complexes, i.e. it represents an object in $\mathcal{Z}^{\text{dg}}(\text{Pretr}(\mathcal{C}))$.

Remark 6.14. Given an object (Z, τ) in $\mathcal{Z}^{\text{dg}}(\mathcal{C})$, it is instructive to compute the half-braiding of twisted complexes past Z explicitly. Thus, suppose we have a twisted complex $\text{tw}_\alpha(X)$ with $X \in \mathcal{C}$ (or $X \in \Sigma\mathcal{C}$).

Note that $Z \star \text{tw}_\alpha(X) = \text{tw}_{\text{Id}_Z \star \alpha}(Z \star X)$ and $\text{tw}_\alpha(X) \star Z = \text{tw}_{\alpha \star \text{Id}_Z}(X \star Z)$. The half-braiding morphism $Z \star \text{tw}_\alpha(X) \rightarrow \text{tw}_\alpha(X) \star Z$ can then be constructed using standard homological perturbation theory as:

$$(15) \quad \begin{aligned} \tau_{\text{tw}_\alpha(X)} &: Z \otimes \text{tw}_\alpha(X) \rightarrow \text{tw}_\alpha(X) \otimes Z, \\ \tau_{\text{tw}_\alpha(X)} &:= \sum_{r \geq 0} (-1)^{\binom{r+1}{2}} \tau(\text{Id}_X \parallel \underbrace{\alpha \parallel \dots \parallel \alpha}_r \parallel \text{Id}_X). \end{aligned}$$

(This is well-defined if X has the structure of a one-sided twisted complex.) To see that this is a closed degree zero morphism in $\text{Tw}(\Sigma\mathcal{C})$, we must show that

$$d(\tau_{(X, \alpha)}) + (\alpha \otimes \text{Id}_Z) \circ \tau_{(X, \alpha)} - \tau_{\text{tw}_\alpha(X)} \circ (\text{Id}_Z \otimes \alpha) = 0.$$

We compute:

$$\begin{aligned} d(\tau_{\text{tw}_\alpha(X)}) &= \sum_{r \geq 0} (-1)^{\binom{r+1}{2}} d(\tau(\text{Id}_X \parallel \underbrace{\alpha \parallel \dots \parallel \alpha}_r \parallel \text{Id}_X)) \\ &= \sum_{r \geq i \geq 1} (-1)^{\binom{r+1}{2} + r + i - 1} \text{Id}_X \parallel \underbrace{\alpha \parallel \dots \parallel \alpha}_{i-1} \parallel d(\alpha) \parallel \underbrace{\alpha \parallel \dots \parallel \alpha}_{r-i} \parallel \text{Id}_X \\ &\quad + \sum_{r \geq 1} (-1)^{\binom{r+1}{2} + r - 1} \alpha \circ \tau(\text{Id}_X \parallel \underbrace{\alpha \parallel \dots \parallel \alpha}_{r-1} \parallel \text{Id}_X) \\ &\quad + \sum_{r > i \geq 1} (-1)^{\binom{r+1}{2} + i} \text{Id}_X \parallel \underbrace{\alpha \parallel \dots \parallel \alpha}_{i-1} \parallel \alpha^2 \parallel \underbrace{\alpha \parallel \dots \parallel \alpha}_{r-i-1} \parallel \text{Id}_X \\ &\quad + \sum_{r \geq 1} (-1)^{\binom{r+1}{2} + r} \tau(\text{Id}_X \parallel \underbrace{\alpha \parallel \dots \parallel \alpha}_{r-1} \parallel \text{Id}_X) \circ \alpha \\ &= -\alpha \circ \tau_{\text{tw}_\alpha(X)} + (-1)^{|\tau_{\text{tw}_\alpha(X)}|} \tau_{\text{tw}_\alpha(X)} \circ \alpha. \end{aligned}$$

Since $\binom{r+1}{2} + r - 1 \equiv \binom{r+2}{2} \pmod{2}$ and $d(\alpha) + \alpha^2 = 0$ the terms in the first and third line cancel and we get $d(\tau_{(X,\alpha)}) = -(\alpha \otimes \text{Id}_Z) \circ \tau_{(X,\alpha)} + \tau_{(X,\alpha)} \circ (\alpha \otimes \text{Id}_Z)$ as desired.

Remark 6.15. It an exercise to check that the braiding morphisms which commute Z past one-sided twisted complexes satisfy the required compatibility with the monoidal structure.

6.4. The dg monoidal trace. Let \mathcal{C} be a dg monoidal category (which we will soon assume to be strictly monoidal). First, recall the \mathcal{C}, \mathcal{C} -bimodule \mathcal{X}_{12} with

$$\mathcal{X}_{12} := \bigoplus_{X,Y,Z,Z' \in \mathcal{C}} \text{Hom}(X \star Z, Z' \star Y) \quad \left(= (Z' \star Y)\mathcal{C}(X \star Z) \right),$$

with bimodule structure defined by

$$a \cdot f \cdot b := a \circ_1 f \circ_2 b \quad \forall a, b \in \mathcal{C}, \quad f \in \mathcal{X}_{12}.$$

Definition 6.16. We define a dg category $\text{Tr}(\mathcal{C})$, the (*dg horizontal*) *trace* of \mathcal{C} , as follows. First, define the \mathcal{C}, \mathcal{C} bimodule

$$\mathbf{B}(\mathcal{C}) \otimes_{\mathcal{C}} \mathcal{X}_{12}.$$

Elements of this bimodule are linear combinations of symbols of the form $c_0 \| c_1 \| \cdots \| c_r \| (c_{r+1} \circ_1 f)$ where $c_0, \dots, c_{r+1} \in \mathcal{C}$ is a sequence of composable morphisms and $f \in \mathcal{X}_{12}$. Then we identify the left and right \mathcal{C} -actions by forming the quotient

$$\mathbf{C}(\mathcal{C}, \mathcal{X}_{12}) := (\mathbf{B}(\mathcal{C}) \otimes_{\mathcal{C}} \mathcal{X}_{12}) / \sim$$

with respect to the relations of the form

$$(-1)^s c_0 \| c_1 \| \cdots \| c_r \| f \simeq \text{Id} \| c_1 \| \cdots \| c_r \| (f \circ_2 c_0),$$

where the sign is determined by $s = |c_0|(|c_1| + \cdots + |c_r| + |f|) + |c_0|r$. As usual we will typically drop leftmost identity map from the notation, writing

$$\| c_1 \| \cdots \| c_r \| f = \text{Id} \| c_1 \| \cdots \| c_r \| f$$

Note that $\mathbf{C}(\mathcal{C}, \mathcal{X}_{12})$ is the Hochschild chain complex of \mathcal{C} with coefficients in the bimodule \mathcal{X}_{12} . Recall that each pair of objects $X, X' \in \mathcal{C}$ determines a subbimodule

$$\mathcal{X}_{12}(X', X) := \bigoplus_{Y, Y'} \text{Hom}_{\mathcal{C}}(X \star Y, Y' \star X') \subset \mathcal{X}_{12}$$

and so we have subcomplexes

$$\mathbf{C}(\mathcal{C}, \mathcal{X}_{12}(X', X)) \subset \mathbf{C}(\mathcal{C}, \mathcal{X}_{12}).$$

Now, define a dg category $\text{Tr}(\mathcal{C})$ as follows. Objects of $\text{Tr}(\mathcal{C})$ are the same as objects of \mathcal{C} , though to avoid confusion we will write $\text{Tr}(X)$ for $X \in \mathcal{C}$ regarded as an object of $\text{Tr}(\mathcal{C})$. The complex of morphisms is given by

$$\text{Hom}_{\text{Tr}(\mathcal{C})}(\text{Tr}(X), \text{Tr}(X')) := \mathbf{C}(\mathcal{C}, \mathcal{X}_{12}(X', X)),$$


with composition induced induced by the shuffle product on the two-sided bar complex and composition in \mathcal{C} .

Explicitly, this means that $\mathrm{Hom}_{\mathrm{Tr}(\mathcal{C})}(\mathrm{Tr}(X), \mathrm{Tr}(X'))$ has basis given by formal symbols of the form

$$(\underline{c}, f) := \|c_1\| \cdots \|c_r\| f,$$

where $r \geq 0$, $c_i \in Y_i \mathcal{C} Y_{i+1}$ and $f \in (Y_{r+1} \star X') \mathcal{C} (X \star Y_1)$ for objects $Y_1, \dots, Y_{r+1} \in \mathcal{C}$, and we abbreviate by writing $\underline{c} = (c_1, \dots, c_r)$. When $r = 0$, the sequence \underline{c} is empty, and we will use the notation (\emptyset, f) and $\|f\|$ interchangeably.

We also allow formal symbols of the form $c_0 \|c_1\| \cdots \|c_r\| f$ but modulo the relations imposed on Hochschild chains, such an expression equals $\pm \|c_1\| \cdots \|c_r\| (f \circ_2 c_0)$. We picture these symbols as follows.



The cohomological grading is

$$\mathrm{deg}(\underline{c}, f) = -r + |f|$$

We say that (\underline{c}, f) has *bar degree* r .

The differential is the usual bar differential (an alternating sum of ways of deleting bars) plus the terms involving the differentials of the individual components:


$$\begin{aligned} d_{\mathrm{Tr}(\mathcal{C})}(\|c_1\| \cdots \|c_r\| f) &= c_1 \|c_2\| \cdots \|c_r\| f \\ &+ \sum_{i=1}^{r-1} (-1)^i \|c_1\| \cdots \|c_i \circ c_{i+1}\| \cdots \|c_r\| f \\ &+ (-1)^r \|c_1\| \cdots \|c_{r-1}\| (c_r \circ_1 f) \\ &+ \sum_{i=1}^r (-1)^{s(i)} \|c_1\| \cdots \|d(c_i)\| \cdots \|c_r\| f \\ &+ (-1)^{s(r+1)} \|c_1\| \cdots \|c_r\| d(f). \end{aligned}$$

where $s(i) = |c_0| + \cdots + |c_{i-1}|$. When $r = 0$ the above generates to $d(\|f\|) = \|d(f)\|$.

The composition of morphisms is defined by

$$(\underline{c}, f) \circ (\underline{d}, g) := (\underline{d} * \underline{c}, \mu_{\nearrow}(f, g))$$

where $*$ denotes the shuffle product from Section 6.1, and $\mu_{\succ}(f, g)$ is as defined in Section 6.2. The composition can be pictured as follows.



Remark 6.17. The identity endomorphism of $\mathrm{Tr}(X)$ in $\mathrm{Tr}(\mathcal{C})$ is given by (\emptyset, ϕ) where ϕ is the canonical isomorphism $X \star \mathbb{1} \rightarrow \mathbb{1} \star X$. Then $(\emptyset, \phi) \circ (\emptyset, \phi)$ in $\mathrm{Tr}(\mathcal{C})$ is by definition (\emptyset, ψ) where ψ is the canonical isomorphism $X \star (\mathbb{1} \star \mathbb{1}) \rightarrow (\mathbb{1} \star \mathbb{1}) \star X$. This ψ is homotopic, but not *equal* to ϕ . Thus, strictly speaking (\emptyset, ϕ) only acts as the identity of $\mathrm{Tr}(X)$ up to homotopy! There is a similar problem concerning the associativity of composition in $\mathrm{Tr}(\mathcal{C})$. The essential issue is that the two-sided bar complex of \mathcal{C} is too large. These annoyances do not arise when \mathcal{C} is strict monoidal, and so this will be assumed *en force* in the sequel. Without this assumption, $\mathrm{Tr}(\mathcal{C})$ should be regarded as an A_∞ category, not a dg category.

We make some observations about $\mathrm{Tr}(\mathcal{C})$ below. First, note that there is a dg functor $\mathrm{Tr}: \mathcal{C} \rightarrow \mathrm{Tr}(\mathcal{C})$ sending $X \mapsto \mathrm{Tr}(X)$ and $f \mapsto \parallel f$.

Remark 6.18. If \mathcal{A} is an ordinary monoidal category (regarded as a dg category with trivial grading and differential) then $H^0(\mathrm{Tr}(\mathcal{A}))$ is isomorphic to the usual *horizontal trace* $\mathrm{Tr}_0(\mathcal{A})$, as defined in [BHLv17, Section 2.4]. In this sense, $\mathrm{Tr}(\mathcal{A})$ is a derived version of $\mathrm{Tr}_0(\mathcal{A})$, much as Hochschild homology of an algebra A is a derived version of its trace $A/[A, A]$.

Lemma 6.19. *If \mathcal{A} is a \mathbb{k} -linear monoidal category, considered as a dg category with trivial grading and differential, then there is a functor $\mathrm{Tr}(\mathcal{A}) \rightarrow \mathrm{Tr}_0(\mathcal{A})$ defined on objects by $\mathrm{Tr}(X) \mapsto \mathrm{Tr}_0(X)$ and on morphisms by $\parallel f \mapsto f$ and $(\underline{c}, f) \mapsto 0$ for all sequences of \underline{c} of length $r \geq 1$.*

The well-known relationship between vertical and horizontal traces (turn head by 90 degrees) transfers to the derived setting as follows. The following is clear from the definitions.

Proposition 6.20. *If \mathcal{C} is a dg monoidal category then*

$$\mathbf{C}_*(\mathcal{C}) = \mathrm{End}_{\mathrm{Tr}(\mathcal{C})}(\mathrm{Tr}(\mathbb{1})).$$

as dg algebras.

Strictly speaking, $\text{End}_{\text{Tr}(\mathcal{C})}(\text{Tr}(\mathbb{1}))$ is given by Hochschild chains of \mathcal{C} with coefficients in the bimodule

$$\bigoplus_{X,Y \in \mathcal{C}} \text{Hom}(\mathbb{1} \star X, Y \star \mathbb{1}),$$

while $\mathbf{C}_*(\mathcal{C})$ is given by Hochschild cochains of \mathcal{C} with coefficients in \mathcal{C} . If \mathcal{C} is strict monoidal, then these two bimodules are equal; otherwise they are isomorphic via the unitor maps.

The following says that the natural functor $\mathcal{C} \rightarrow \text{Tr}(\mathcal{C})$ satisfies a categorical “trace-like” property, provided that \mathcal{C} has duals.

Lemma 6.21. *Given two objects X, Y in \mathcal{C} , we define the traciator*

$$w_{X,Y} : \text{Tr}(X \star Y) \rightarrow \text{Tr}(Y \star X)$$

to be the degree zero closed morphism associated to the identity map (or associator in the non-strict monoidal case) $(X \star Y) \star X \rightarrow X \star (Y \star X)$. If X has a right dual in \mathcal{C} , then this map is a homotopy equivalence.

Proof. Suppose *X is the right dual to X , with structure maps $\text{coev}_X : \mathbb{1} \rightarrow {}^*X \star X$ and $\text{ev}_X : X \star {}^*X \rightarrow \mathbb{1}$. Define

$$w_{X,Y}^{-1} : \text{Tr}(Y \star X) \rightarrow \text{Tr}(X \star Y)$$

to be the composite of

$$\|(\text{coev}_X \star \text{Id}_Y \star \text{Id}_X) : \text{Tr}(Y \star X) \rightarrow \text{Tr}({}^*X \star X \star Y \star X),$$

followed by


$$w_{{}^*X, X \star Y \star X} : \text{Tr}({}^*X \star X \star Y \star X) \rightarrow \text{Tr}(X \star Y \star X \star {}^*X)$$

and finally

$$\|(\text{Id}_X \star \text{Id}_Y \star \text{ev}_X) : \text{Tr}(X \star Y \star X \star {}^*X) \rightarrow \text{Tr}(X \star Y).$$

One can now check that $\text{Id}_{X \star Y} - w_{X,Y}^{-1} w_{X,Y}$ and $\text{Id}_{Y \star X} - w_{X,Y} w_{X,Y}^{-1}$ are exact. \square

These morphisms can be pictures as follows.



Remark 6.22. It is easy to see that the traciator $w_{X,Y}$ is natural in Y , but it is natural in X only up to coherent homotopy. For any closed map $f : X \rightarrow X'$ the composition $w_{X',Y} \circ (f \star \text{Id}_Y) : \text{Tr}(X \star Y) \rightarrow \text{Tr}(Y \star X')$ is represented by the map $X \star Y \star X' \xrightarrow{f \star \text{Id} \star \text{Id}} X' \star Y \star X'$, while the composition $(\text{Id}_Y \star f) \circ w_{X,Y}$ is represented by the map $X \star Y \star$

$X \xrightarrow{f \star \text{Id} \star \text{Id}} X \star Y \star X'$. The difference of these two maps is given by the boundary of the morphism

$$(16) \quad w(f; Y) := \|f\| \text{Id}_{X \star Y \star X'}$$

More generally, for any sequence of composable morphisms $f_i \in X_i \mathcal{C} X_{i+1}$ for $1 \leq i \leq r$ the derived trace contains *higher traciators*

$$w(f_1 \| \dots \| f_r; Y) := \|f_1\| \cdots \|f_r\| \text{Id}_{X_{r+1} \star Y \star X_1} \in \text{Hom}_{\text{Tr}}(\text{Tr}(X_{r+1} \star Y), \text{Tr}(Y \star X_1))$$

which provide the (higher) naturality data for the traciator.

Remark 6.23. For an object X in \mathcal{C} we define the *rotator* $w_X: \text{Tr}(X) \rightarrow \text{Tr}(X)$ to be the endomorphism given by $\|\text{Id}_{X \star X}\|$. Note that this agrees with the traciator $w_{X,1}$ as defined in the proof of Lemma 6.21 if \mathcal{C} is strict monoidal. If X has a right dual then w_X is invertible up to homotopy.

In this way, the assignment $X \mapsto w_X$ is a degree zero endomorphism of the canonical dg functor $\mathcal{C} \rightarrow \text{Tr}(\mathcal{C})$, natural up to coherent homotopy in the sense of Remark 6.22.

Remark 6.24. Similar to Remark 6.13, the higher traciators (from (16)) for $\text{Tr}(\mathcal{C})$ already carry enough information to determine traciators in $\text{Tr}(\text{Pretr}(\mathcal{C}))$. For example, given a morphism $f: X \rightarrow X'$ and its cone $\text{Cone}(f) := [X \rightarrow X']$, the traciator $w_{\text{Cone}(f), Y}: \text{Tr}(\text{Cone}(f) \star Y) \rightarrow \text{Tr}(Y \star \text{Cone}(f))$ is represented by the morphism

$$(17) \quad w_{\text{Cone}(f), Y} = \begin{array}{ccc} [\text{Tr}(X \star Y) & \xrightarrow{\text{Tr}(f \star \text{Id}_Y)} & \text{Tr}(X' \star Y)] \\ \downarrow w_{X, Y} & \searrow w(f; Y) & \downarrow w_{X', Y} \\ [\text{Tr}(Y \star X) & \xrightarrow{\text{Tr}(\text{Id}_Y \star f)} & \text{Tr}(Y \star X')] \end{array}$$

It is very important to note that the dg category $\text{Tr}(\mathcal{C})$ is additive but not triangulated. Indeed, there are lots of morphisms in $\text{Tr}(\mathcal{C})$ which do not exist in \mathcal{C} , so their cones do not exist as objects in \mathcal{C} or in $\text{Tr}(\mathcal{C})$. However, we can consider its pretriangulated hull as in section 3, we will denote it by $\text{Pretr}(\text{Tr}(\mathcal{C}))$.

Lemma 6.25. *If \mathcal{C} is a dg monoidal category, then $\text{Pretr}(\text{Tr}(\mathcal{C})) \simeq \text{Pretr}(\text{Tr}(\text{Pretr}(\mathcal{C})))$.*

Proof. As it is a dg functor Tr sends twisted complexes to twisted complexes. Thus we have a dg functor

$$\text{Pretr}(\text{Tr}(\text{Pretr}(\mathcal{C}))) \hookrightarrow \text{Pretr}(\text{Pretr}(\text{Tr}(\mathcal{C}))) \simeq \text{Pretr}(\text{Tr}(\mathcal{C})).$$

Conversely we have a dg functor $\text{Pretr}(\text{Tr}(\mathcal{C})) \hookrightarrow \text{Pretr}(\text{Tr}(\text{Pretr}(\mathcal{C})))$ induced by $\mathcal{C} \hookrightarrow \text{Pretr}(\mathcal{C})$ which is a quasi-inverse. \square

6.5. Homotopy trace-like functors. In the introduction we highlighted that the (underived) trace functor $\mathrm{Tr}_0: \mathcal{C} \rightarrow \mathrm{Tr}_0(\mathcal{C})$ for a monoidal category (or bicategory) \mathcal{C} with left duals is initial among all trace-like functors from \mathcal{C} to another category \mathcal{D} . Here we consider a dg analog of this situation.

Let \mathcal{C} be a dg monoidal category and \mathcal{D} a dg category. We are interested in dg functors $\phi: \mathcal{C} \rightarrow \mathcal{D}$ that are *homotopy trace-like* in the sense that for every pair of objects X, Y in \mathcal{C} we get natural (up to coherent homotopy) maps $\phi(X \star Y) \rightarrow \phi(Y \star X)$, possibly even isomorphisms. We define a *dg monoidal trace* on \mathcal{C} with values in \mathcal{D} to be a dg functor $\phi: \mathcal{C} \rightarrow \mathcal{D}$ that factors through $\mathrm{Tr}: \mathcal{C} \rightarrow \mathrm{Tr}(\mathcal{C})$. I.e. there exists a dg functor $\phi': \mathrm{Tr}(\mathcal{C}) \rightarrow \mathcal{D}$ such that $\phi = \phi' \circ \mathrm{Tr}$.

Note that a dg monoidal trace ϕ not only contains the data of specific morphisms $\phi'(w_{X,Y}): \phi(X \star Y) \rightarrow \phi(Y \star X)$, but also homotopies enforcing natural compatibility relations between these morphisms. In particular, by Remark 6.22 $\phi'(w_{X,Y})$ is natural in Y but it is natural in X only up to coherent higher homotopies $\phi'(w(f_1 \parallel \dots \parallel f_r; Y))$.

Remark 6.26. For an underived trace-like functor $\phi = \phi' \circ \mathrm{Tr}_0$ one requires (see e.g. [Zhu18]) that the traciators are compatible with associators in the sense that diagrams of the following type commute:

$$(18) \quad \begin{array}{ccccc} \phi((X \star Y) \star Z) \xrightarrow{\phi'(w_{X \star Y, Z})} \phi(Z \star (X \star Y)) & \longrightarrow & \phi((Z \star X) \star Y) \\ \downarrow & & \uparrow \\ \phi(X \star (Y \star Z)) & \longrightarrow & \phi((Y \star Z) \star X) & \longrightarrow & \phi(Y \star (Z \star X)). \end{array}$$

In the derived case, similar conditions are required for the images of higher traciators $\phi'(w(f_r \parallel \dots \parallel f_1; Y))$.

Remark 6.27. It is likely that if \mathcal{C} has left duals then knowing $\phi'(w_{X,Y})$ and the naturality data $\phi'(w(f_1 \parallel \dots \parallel f_r; Y))$ satisfying (18) and its analogues is enough to prove that reconstruct the functor ϕ' and thus verify that ϕ is trace-like, but we do not prove it here.

Examples of dg monoidal traces are the following.

Example 6.28. Any endofunctor of $\mathrm{Tr}(\mathcal{C})$ gives rise to a dg monoidal trace on \mathcal{C} by pre-composition with the universal trace $\mathcal{C} \rightarrow \mathrm{Tr}(\mathcal{C})$. A natural source of endofunctors of $\mathrm{Tr}(\mathcal{C})$ is the dg monoidal center, see Section 6.6.

Example 6.29. Let X be an object of $\mathrm{Tr}(\mathcal{C})$, then another dg monoidal trace is given by the universal trace $\mathcal{C} \rightarrow \mathrm{Tr}(\mathcal{C})$ composed with the representable functor $\mathrm{Hom}_{\mathrm{Tr}(\mathcal{C})}(X, -)$. The target is $\mathrm{dgmod}\text{-}\mathrm{End}_{\mathrm{Tr}(\mathcal{C})}(X)$. A particularly interesting case is $X = \mathrm{Tr}(\mathbb{1})$, for which Proposition 6.20 identifies the target with $\mathrm{dgmod}\text{-}\mathbf{C}_*(\mathcal{C})$.

Example 6.30. If \mathcal{C} is a pivotal dg monoidal category, then $\mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, -)$ is a trace-like functor with target $\mathrm{dgmod}\text{-}\mathrm{End}_{\mathcal{C}}(\mathbb{1})$. To see that $\mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, -)$ is trace-like, we compute:

$$\mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, X \star Y) \cong \mathrm{Hom}_{\mathcal{C}}(X^*, Y) \cong \mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, Y \star X^{**}) \cong \mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, Y \star X)$$

where the last isomorphism uses pivotality. Since $\mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, -)$ is trace-like (strictly, not just up to homotopy), it factors through the dg monoidal trace, providing a dg functor

$$\mathrm{Tr}(\mathcal{C}) \rightarrow \mathrm{dgmod}\text{-}\mathrm{End}_{\mathcal{C}}(\mathbb{1})$$

Example 6.31. A similar computation for pivotal \mathcal{C} shows that if (Z, τ) is central, then the dg functor $\mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, Z \star -)$ is trace-like. More generally, if (Z', τ') is also central, then $\mathrm{Hom}_{\mathcal{C}}(Z', Z \star -)$ is again trace-like. An alternative way to construct these traces is to first factor $\mathrm{Hom}_{\mathcal{C}}(\mathbb{1}, -)$ through the trace $\mathcal{C} \rightarrow \mathrm{Tr}(\mathcal{C})$ and then act by the dg monoidal center.

Example 6.32. The full twist FT and its powers FT^k (with suitable half-braiding data) are objects in dg Drinfeld center of the category of complexes of Soergel bimodules [EH]. By applying Example 6.31 we see that

$$\mathrm{Hom}_{\mathrm{SBim}}(\mathbb{1}, \mathrm{FT}^k \star -)$$

is a trace-like functor for all k . Such functors play a prominent role in the work of the first and second author, Neguț and Rasmussen [GNR16, GH17].

Example 6.33. If \mathcal{C} is a monoidal category, then any (ordinary) trace-like functor $\phi: \mathcal{C} \rightarrow \mathcal{D}$ lifts to a dg monoidal trace. To see this, we factor $\phi = \phi' \circ \mathrm{Tr}_0$ for $\mathrm{Tr}_0: \mathcal{C} \rightarrow \mathrm{Tr}_0(\mathcal{C})$ and some functor $\phi': \mathrm{Tr}_0(\mathcal{C}) \rightarrow \mathcal{D}$. The induced dg monoidal trace is obtained by precomposing ϕ' with the functor $\mathrm{Tr}(\mathcal{C}) \rightarrow \mathrm{Tr}_0(\mathcal{C})$ from Lemma 6.19 and the universal trace $\mathrm{Tr}: \mathcal{C} \rightarrow \mathrm{Tr}(\mathcal{C})$.

6.6. Action of the dg monoidal center on the trace. It is a basic observation that the center $Z(A)$ of an associative algebra A acts on the trace $A/[A, A]$. Similarly, if we think of the trace $\mathrm{Tr}_0(\mathcal{C})$ of a monoidal category as \mathcal{C} integrated over an annulus, the Drinfeld center $\mathcal{Z}^0(\mathcal{C})$ acts by “cutting open” the annulus and “gluing in” central objects or morphisms, before “resealing the annulus”. In this section we describe the analogous action of the dg monoidal center $\mathcal{Z}^{\mathrm{dg}}(\mathcal{C})$ on the dg monoidal trace $\mathrm{Tr}(\mathcal{C})$.

Let (Z, τ) be an object of $\mathcal{Z}^{\mathrm{dg}}(\mathcal{C})$. The structure map τ is a map of \mathcal{C} , \mathcal{C} -bimodules $\mathbf{B}(\mathcal{C}) \rightarrow \mathcal{X}_{12}(Z, Z)$. Using this map we have a map of \mathcal{C} , \mathcal{C} -bimodules given by composing:


$$\begin{aligned} \mathbf{B}(\mathcal{C}) \otimes_{\mathcal{C}} \mathcal{X}_{12}(X', X) &\rightarrow \mathbf{B}(\mathcal{C}) \otimes_{\mathcal{C}} \mathbf{B}(\mathcal{C}) \otimes_{\mathcal{C}} \mathcal{X}_{12}(X', X) \\ &\rightarrow \mathbf{B}(\mathcal{C}) \otimes_{\mathcal{C}} \mathcal{X}_{12}(Z, Z) \otimes_{\mathcal{C}} \mathcal{X}_{12}(X', X) \\ &\rightarrow \mathbf{B}(\mathcal{C}) \otimes_{\mathcal{C}} \mathcal{X}_{12}(Z \star X', Z \star X). \end{aligned}$$

The first of these maps is the comultiplication on the two-sided bar complex of \mathcal{C} , the second is an application of τ , and the last is the algebra structure on the bimodule \mathcal{X}_{12} (see Remark 6.1). Applying the functor which identifies the left and right actions of \mathcal{C} , we obtain a map of complexes

$$\mathrm{Hom}_{\mathrm{Tr}(\mathcal{C})}(\mathrm{Tr}(X), \mathrm{Tr}(X')) \rightarrow \mathrm{Hom}_{\mathrm{Tr}(\mathcal{C})}(\mathrm{Tr}(Z \star X), \mathrm{Tr}(Z \star X')).$$

In this way, (Z, τ) determines an endofunctor $\Xi_{(Z, \tau)}: \text{Tr}(\mathcal{C}) \rightarrow \text{Tr}(\mathcal{C})$ defined on objects by $\text{Tr}(X) \mapsto \text{Tr}(Z \star X)$ and on morphisms by the above chain map. That this map respects composition of morphisms follows from the assumption that τ is a morphism of dg algebras and the compatibility of the shuffle product and coproduct on the two-sided bar complex.

The action on morphism complexes can be pictured as follows.



Remark 6.34. Morphisms $f: (Z, \tau) \rightarrow (Z', \tau')$ in $\mathcal{Z}^{\text{dg}}(\mathcal{C})$ should give natural transformations $\Xi_f: \Xi_{Z, \tau} \rightarrow \Xi_{Z', \tau'}$ that assemble into a monoidal functor

$$\Xi: \mathcal{Z}^{\text{dg}}(\mathcal{C}) \rightarrow \text{End}(\text{Tr}(\mathcal{C}))$$

from $\mathcal{Z}^{\text{dg}}(\mathcal{C})$ to the endofunctors of $\text{Tr}(\mathcal{C})$. We will not pursue this in detail, see also Remark 6.4.

7. TRACES OF THE SOERGEL CATEGORY

7.1. Soergel bimodules. Let W be a Coxeter group with simple reflections $S \subset W$, length function ℓ , and Bruhat order \leq . We consider realizations $(V, \{\alpha_s^V\}, \{\alpha_s\})$ of W [EW16] over \mathbb{C} , which we assume to be reflection faithful and balanced unless stated otherwise. As usual, we consider the polynomial ring $R = \mathbb{C}[V] := \text{Sym}^\bullet(V^*)$, which is graded by declaring elements in V^* to be of degree 2. In particular, we have $\alpha_s \in R$, and these elements generate R if they span V^* .

Remark 7.1. In type A we have $W = S_n$ with simple reflections (transpositions) indexed by $i \in \{1, \dots, n-1\}$. We will consider the realization with $V = \mathbb{C}^n$, on which S_n acts by permuting standard basis vectors, and we identify $R = \mathbb{C}[x_1, \dots, x_n]$ and $\alpha_i = x_i - x_{i+1}$ for $1 \leq i \leq n-1$.

We now describe the category of Soergel bimodules $\text{SBim}(W)$ associated to W and its chosen realization. For each simple reflection $s \in S$ we let

$$B_s := R \otimes_{R^s} R(1),$$

where $R^s = \{f \in R \mid s(f) = f\}$ and (1) is the “downward” grading shift. Then the category of Soergel bimodules $\text{SBim}(W)$ is the smallest full monoidal subcategory of graded R, R -bimodules containing the B_s , which is closed under grading shift, isomorphism, direct sums, and direct summands. The monoidal structure on

$\text{SBim}(W)$ is denoted by \star and tensor products of bimodules of the form B_s are called *Bott–Samelson bimodules*.

In this paper we will occasionally use the diagrammatic Hecke category \mathcal{D}_W of Elias–Williamson [EW16], whose Karoubi completion is equivalent to the category of Soergel bimodules under the assumptions taken. The diagrammatic category \mathcal{D}_W has the advantages of being strictly monoidal and manifestly strictly pivotal (by the balanced assumption) besides being well-behaved for a larger class of realizations (which will not be relevant here).

Let $\mathcal{C}(W) := \text{Ch}^b(\text{SBim}(W))$ be the dg monoidal category of bounded chain complexes of Soergel bimodules for W . If $\underline{\beta}$ is a word in the alphabet σ_s, σ_s^{-1} with $i \in S$, then we have a finite complex $F(\underline{\beta}) \in \mathcal{C}(W)$ defined by

$$(19) \quad \begin{aligned} F(\sigma_s) &:= (\underline{B}_s \rightarrow R(1)), & F(\sigma_s^{-1}) &:= (R(-1) \rightarrow \underline{B}_s), \\ F(\underline{\beta} \cdot \underline{\beta}') &:= F(\underline{\beta}) \star F(\underline{\beta}'), \end{aligned}$$

where the maps $B_s \rightarrow R(1)$ and $R(-1) \rightarrow B_s$ are the canonical bimodule maps, defined by

$$1 \otimes 1 \mapsto 1, \quad 1 \mapsto \frac{1}{2}(\alpha_s \otimes 1 - 1 \otimes \alpha_s),$$

respectively. It is well known [Rou04, Rou06] that the complexes $F(\underline{\beta})$ satisfy the braid relations up to homotopy equivalence. We often abusively write $F(\beta)$ where $\beta \in \text{Br}(W)$ is an element of the braid group associated to W (when in reality the complex $F(\beta)$ depends on a choice of braid word $\underline{\beta}$ representing β).

Definition 7.2. For each $w \in W$, let Δ_w and ∇_w denote the Rouquier complex of the positive and negative braid lift of a chosen reduced expression of w .

In particular, we have $\Delta_w \star \nabla_{w^{-1}} \simeq \mathbb{1} \simeq \nabla_{w^{-1}} \star \Delta_w$ for any $w \in W$. We will sometimes write $\Delta_w^{-1} := \nabla_{w^{-1}}$.

7.2. Hochschild homology of the Soergel category — linear structure. The following is well known to experts.

Proposition 7.3. *The complexes $\Delta_w, w \in W$, generate $\mathcal{C}(W)$ with respect to cones, shifts, sums, and homotopy equivalences (and similarly for ∇_w). These complexes satisfy*

$$\text{Hom}_{\mathcal{C}(W)}(\Delta_v, \Delta_w) \simeq 0 \simeq \text{Hom}_{\mathcal{C}(W)}(\nabla_w, \nabla_v) \quad \text{unless} \quad v \leq w.$$

In other words, $\{\Delta_v\}_{v \in W}$ and $\{\nabla_v\}_{v \in W}$ each generate a semi-orthogonal decomposition of $\mathcal{C}(W)$ (the latter one with the opposite poset structure).

Proof. By the main result of [LW14], one has

$$\text{Hom}_{\mathcal{C}(W)}(\Delta_v, \nabla_w) \simeq \begin{cases} R & \text{if } v = w, \\ 0 & \text{otherwise.} \end{cases}$$

On the other hand, it is easy to see that Δ_w is filtered by ∇_u for $u \leq w$. Therefore if $\text{Hom}_{\mathcal{C}(W)}(\Delta_v, \Delta_w) \not\simeq 0$ then $\text{Hom}_{\mathcal{C}(W)}(\Delta_v, \nabla_u) \not\simeq 0$ for some $u \leq w$, hence $v = u \leq w$. See also [GHMN19, Appendix] for more details. \square

Thus the following is an immediate consequence of Theorem 5.11.

Lemma 7.4. *There is a deformation retract from the cyclic bar complex $\mathbf{C}(\mathrm{SBim}(W))$ to $\bigoplus_{w \in W} \mathbf{C}(\mathrm{End}(\Delta_w))$. \square*

Recall that we have chosen a realization V of W . Consider the following complex of \mathbb{C} -vector spaces

$$Z := \left(V^*(-2) \rightarrow \underline{V^*(-2) \oplus V^*(-2)} \right), \quad \phi \mapsto (\phi, -\phi).$$

Complexes of \mathbb{C} -vector spaces form a symmetric monoidal (dg) category, and so we have Schur functors. The symmetric algebra of Z is just

$$\mathrm{Sym}^\bullet(V^*(-2) \oplus V^*(-2) \oplus \Sigma^1 V^*(-2)) \cong R \otimes R \otimes \Lambda,$$

with its differential inherited from Z . After choosing a basis of V and letting $x_1, \dots, x_n \in V^*$ denote the dual basis, we can identify

$$\mathrm{Sym}^\bullet(Z) \cong \mathbb{C}[x_1, \dots, x_n, x'_1, \dots, x'_n, \theta_1, \dots, \theta_n], \quad d(\theta_i) = x_i - x'_i.$$

That is to say, $\mathrm{Sym}^\bullet(Z)$ is the Koszul resolution of R as a bimodule over itself. We can use this resolution to compute $\mathrm{HH}_\bullet(R)$:

$$\mathrm{HH}_\bullet(R) \cong R \otimes \Lambda.$$

We have a dg algebra map $R = \mathrm{End}_{\mathcal{C}(W)}(\mathbb{1}) \rightarrow \mathrm{End}_{\mathcal{C}(W)}(\Delta_w)$ sending $f \mapsto \mathrm{Id}_{\Delta_w} \star f$. The homotopy equivalence

$$\mathrm{Hom}_{\mathcal{C}(W)}(\Delta_w, \Delta_w) \simeq \mathrm{Hom}_{\mathcal{C}(W)}(\Delta_w^{-1} \star \Delta_w, \mathbb{1}) \simeq \mathrm{End}_{\mathcal{C}(W)}(\mathbb{1}) = R$$

is clearly R -linear ($f \in R$ acts by $-\star f$ on each hom space). The homotopy equivalence $\Delta_w^{-1} \star \Delta_w \rightarrow \mathbb{1}$ is in fact a deformation retract in $\mathcal{C}(W)$. Thus, we have proved the following.

Lemma 7.5. *Let $\mathrm{Id} \star R$ denote the subalgebra of $\mathrm{End}_{\mathcal{C}(W)}(\Delta_w)$ spanned by elements of the form $\mathrm{Id} \star f$ with $f \in R$. Then there is a deformation retract $\mathrm{End}_{\mathcal{C}(W)}(\Delta_w) \rightarrow \mathrm{Id} \star R$, the data of which are R -linear.*

Thus, combining everything up to this point gives:

Corollary 7.6. *There is an R -linear deformation retract from the cyclic bar complex $\mathbf{C}_*(\mathcal{C}(W))$ to its homology $R \otimes \Lambda \rtimes \mathbb{C}[W]$. Here the generators x_i of R have degree $(2, 0)$, the generators θ_i of Λ have degree $(2, -1)$ and W has degree $(0, 0)$.*

7.3. Hochschild homology of the Soergel category — algebra structure. In this section we prove Theorem 1.3. Recall that by Theorem 2.10 any deformation retract $\mathbf{C}_* \twoheadrightarrow H$ induces an A_∞ structure on H .

Recall that the dg algebra \mathbf{C}_* is called formal if all higher Massey products vanish (i.e. the higher A_∞ maps on H are trivial).

Theorem 7.7. *We have an isomorphism of algebras*

$$\mathrm{HH}_*(\mathrm{SBim}(W)) \cong R \otimes \Lambda \rtimes W$$

with multiplication μ_2 .

Proof. By Theorem 5.19 we have a canonical isomorphism

$$\mathrm{HH}_*(\mathrm{SBim}(W)) \cong \mathrm{HH}_*(\mathcal{C}(W)),$$

and we will compute the latter. All homs in this proof are taken in $\mathcal{C}(W)$ and so the subscripts in $\mathrm{Hom}_{\mathcal{C}}(-, -)$ and $\mathrm{End}_{\mathcal{C}(W)}(-)$ will be omitted. First we observe that $\mathrm{HH}_*(\mathrm{End}(\mathbb{1})) \cong \mathrm{HH}_*(R) \cong R \otimes \Lambda$ is a subalgebra of $\mathrm{HH}_*(\mathcal{C}(W))$ by definition. Second, let σ_w denote the class of the identity morphism of $\mathrm{End}(\Delta_w)$ in $\mathrm{HH}_*(\mathcal{C}(W))$. Then $\mathrm{End}(\Delta_w) \simeq R$ is a bimodule over $\mathrm{End}(\mathbb{1}) \cong R$ where the right action of R is standard and the left action of R is twisted by w , and the same holds for the actions of $\mathrm{HH}_*(R)$ on $\mathrm{HH}_*(\mathrm{End}(\Delta_w)) \simeq \mathrm{HH}_*(R)$. In other words,

$$f(x, \theta) \cdot \sigma_w = \sigma_w \cdot f(w^{-1}(x), w^{-1}(\theta)).$$

It remains to prove that $w \mapsto \sigma_w$ is a group homomorphism. For this, we construct an action of σ_w on $\mathrm{End}(\mathbb{1})$ by the following composition of homotopy equivalences (where all homs are taken in $\mathcal{C}(W)$):

$$\sigma_w : \mathrm{End}(\mathbb{1}) \simeq \mathrm{Hom}(\mathbb{1}, \Delta_w \Delta_w^{-1}) \simeq \mathrm{Hom}(\Delta_w, \Delta_w) \simeq \mathrm{Hom}(\mathbb{1}, \Delta_w^{-1} \Delta_w) \cong \mathrm{End}(\mathbb{1}).$$

Here the middle isomorphism follows from the fact that Δ_w and Δ_w^{-1} are biadjoint in $\mathcal{C}(W)$. One can check that this action is compatible with the shuffle multiplication in $\mathrm{HH}_0(\mathcal{C}(W))$ and σ_w acts as a permutation $(-1)^{\ell(w)} w$ on $\mathrm{End}(\mathbb{1})$ (the signs that appear here depend on the homological shift conventions for Rouquier complexes, see [GW19, Section 4.3 and Remark 4.26] for a discussion). Since the representation of W on $\mathrm{End}(\mathbb{1})$ is faithful, we get that the σ_w generate a copy of W inside $\mathrm{HH}_0(\mathcal{C}(W))$. \square

Remark 7.8. The main result of [EL16] states that $\mathrm{HH}_0(\mathrm{SBim}(W)) \cong R \rtimes \mathbb{C}[W]$, so our theorem is a natural generalization. The methods of proof in [EL16], however, were completely different and used cellularity of $\mathrm{SBim}(W)$.

Remark 7.9. One can check that the classes $\sigma_w = [\Delta_w]$ generate a copy of W in a more direct way. Indeed, for simple reflections s the objects Δ_s satisfy braid relations in $\mathcal{C}(W)$, and hence $[\Delta_s]$ satisfy them too. Furthermore,

$$[\Delta_s]^2 = [(\Delta_s)^2] \simeq [B_s(-1) \rightarrow B_s(1) \rightarrow R(2)] = [R] - [B_s] + [B_s] = [R] = 1.$$

The middle equation follows from (13c).

Theorem 7.10. *All higher Massey products on $\mathrm{HH}_*(\mathrm{SBim}(W))$ vanish.*

Proof. By Corollary 7.6 all retractions and homotopies are R -linear. Therefore, all Massey products μ_k are R -linear as well, and it is sufficient to compute the products $\mu_k(\alpha_1, \dots, \alpha_k)$ for α_s in $\Lambda \rtimes \mathbb{C}[W]$, $1 \leq s \leq k$.

Recall that R is generated by elements x_i of bidegree $(2, 0)$, Λ is generated by θ_i of bidegree $(2, -1)$ and W is supported in bidegree $(0, 0)$. Therefore any homogeneous element in $\mathrm{HH}_*(\mathrm{SBim}(W))$ has bidegree (i, j) such that $i + 2j \geq 0$ while all homogeneous elements of $\Lambda \rtimes \mathbb{C}[W]$ satisfy $i + 2j = 0$.

Suppose that α_s for $1 \leq s \leq k$ are homogeneous elements of $\Lambda \rtimes \mathbb{C}[W]$ of bidegree (i_s, j_s) with $i_s + 2j_s = 0$. Then $\mu_k(\alpha_1, \dots, \alpha_k)$ has bidegree $(i_1 + \dots + i_k, j_1 + \dots + j_k + 2 - k)$. However, the collapsed degree

$$(i_1 + \dots + i_k) + 2(j_1 + \dots + j_k + 2 - k) = \sum_s (i_s + 2j_s) + 2(2 - k) = 2(2 - k)$$

of $\mu_k(\alpha_1, \dots, \alpha_k)$ is negative for $k > 2$, which implies $\mu_k(\alpha_1, \dots, \alpha_k) = 0$. \square

We can also describe the action of the Connes differential on Hochschild homology.

Proposition 7.11. *The action of the Connes differential \mathbb{B} on $\mathrm{HH}_*(\mathrm{SBim}(W))$, pulled back to $R \otimes \Lambda \rtimes \mathbb{C}[W]$ via the isomorphism from Theorem 7.7, is given by*

$$(20) \quad \mathbb{B} = \sum_i \theta_i \frac{\partial}{\partial x_i},$$

where x_i, θ_i are dual bases in V, V^* .

Proof. Recall the isomorphisms

$$\mathrm{HH}_*(\mathrm{SBim}(W)) \cong \mathrm{HH}_*(\mathcal{C}(W)) \cong \bigoplus_{w \in W} \mathrm{HH}_*(\mathrm{End}(T_w)) \cong \bigoplus_{w \in W} \mathrm{HH}_*(R).$$

Observe that the second isomorphism is obtained by certain retractions which commute with the action of the Connes differential \mathbb{B} , so it is sufficient to know the action of \mathbb{B} on $\mathrm{HH}_*(\mathrm{End}(T_w)) = \mathrm{HH}_*(R)$. The differential \mathbb{B} on $\mathrm{HH}_*(R)$ is given by (14), and it is obviously W -invariant. \square

7.4. The dg monoidal trace of the Soergel category. In this section, we let W be an arbitrary finite Coxeter group and again write $\mathcal{C}(W) = \mathrm{Ch}^b(\mathrm{SBim}(W))$ for the dg monoidal category of bounded chain complexes of Soergel bimodules for W with a given realisation.

Any object in $\mathrm{Tr}(\mathcal{C}(W))$ is homotopy equivalent to a twisted complex built out of finite direct sums of $\mathrm{Tr}(B_w)$. The same applies to any object in $\mathrm{Pretr}(\mathrm{Tr}(\mathcal{C}(W)))$. Indeed, any object X in $\mathcal{C}(W)$ is a complex built out of finite direct sums of B_w , which we can write as an iterated cone. Since Tr is a dg functor, we can write $\mathrm{Tr}(X)$ as an iterated cone built out of $\mathrm{Tr}(B_w)$ which is a twisted complex. Note that Tr indeed sends a complex to a twisted complex in general, see Example 8.16.

Proposition 7.12. *Let X and Y be two objects in $\mathcal{C}(W)$. Then $\mathrm{Tr}(Y) \mathrm{Tr}(\mathcal{C}(W)) \mathrm{Tr}(X)$ is homotopy equivalent to a bounded complex of free finitely generated R -modules.*

In particular, the homs in $\mathrm{Tr}(\mathcal{C}(W))$ are finite-dimensional in each bidegree.

Proof. Here we use the semiorthogonal decomposition from Proposition 7.3 again, and we write $\mathcal{C} = \mathcal{C}(W)$. As in Section 5.4, the hom complexes in $\mathrm{Tr}(\mathcal{C})$ retract onto semi-orthogonal hom complexes, i.e. for objects X and Y we have a retraction

$$\mathrm{Tr}(Y) \mathrm{Tr}(\mathcal{C} \mathrm{Tr}(X)) = \mathbf{B}(\mathcal{C} \otimes_{\mathcal{C} \boxtimes \mathcal{C}^{\mathrm{op}}} \mathcal{X}_{12}(Y, X)) \rightarrow \mathbf{B}^{\Gamma}(\mathcal{C}, \mathcal{J}_{\mathcal{B}}) \otimes_{\mathcal{C} \boxtimes \mathcal{C}^{\mathrm{op}}} \mathcal{X}_{12}(Y, X),$$

and further onto a complex which has chain groups $\bigoplus_{w_0} (\Delta_{w_0} Y) \mathcal{C}(X \Delta_{w_0})$ in degree $r = 0$ and for $r > 0$:

$$\bigoplus_r \bigoplus_{w_0, \dots, w_r} \Sigma^r \left(\Delta_{w_0} \mathcal{C} \Delta_{w_1} \otimes \cdots \otimes \Delta_{w_{r-1}} \mathcal{C} \Delta_{w_r} \otimes (\Delta_{w_r} Y) \mathcal{C}(X \Delta_{w_0}) \right)$$

with $w_0 \leq w_1 \leq \dots \leq w_r$ in Bruhat order. This still leaves us with an infinite complex since w_i could repeat arbitrarily. However, such infinite repeats form a copy of the two-sided bar complex of $\mathrm{End}(\Delta_w) = R$, which can be retracted to a finite complex. Since W was assumed to be finite, which implies that chains in Bruhat order are finite, performing all such retractions we arrive at a homotopy equivalent bounded complex. Freeness now follows since morphism spaces in \mathcal{C} are free over R , see [Soe07] or [EW14, Theorem 3.6]. \square

8. THE SOERGEL CATEGORY IN TYPE A

In this section we describe the derived horizontal trace of the Soergel category in type A, using the computation of its derived vertical trace in Section 7.

In Proposition 6.20 we have seen that the (derived) vertical trace $\mathbf{C}_*(\mathcal{C})$ of a monoidal (dg) category \mathcal{C} can be identified with the endomorphisms of $\mathrm{Tr}(\mathbb{1})$ in the (derived) horizontal trace $\mathrm{Tr}(\mathcal{C})$. That this in fact determines the entire horizontal trace for Soergel bimodules of type A is the upshot of the following theorem.

Theorem 8.1. *The Karoubi completion of the triangulated hull of $\mathrm{Tr}(\mathrm{SBim}_n)$ is split-generated by $\mathrm{Tr}(\mathbb{1})$. We have*

$$\mathrm{Kar}^{dg}(\mathrm{Pretr}(\mathrm{Tr}(\mathrm{SBim}_n))) \simeq \mathrm{Perf}(\mathbb{C}[x_1, \dots, x_n, \theta_1, \dots, \theta_n] \rtimes \mathbb{C}[S_n]\text{-mod}).$$

Here in the right hand side we have the category of perfect dg modules over the algebra

$$R \otimes \Lambda \rtimes \mathbb{C}[S_n] = \mathbb{C}[x_1, \dots, x_n, \theta_1, \dots, \theta_n] \rtimes \mathbb{C}[S_n],$$

where R and S_n are supported in cohomological degree zero and the variables θ_i have cohomological degree -1 .

The proof of this theorem will occupy the rest of this section. Here we outline the strategy of the proof. First, we use the technology of Frobenius extensions to prove that $\mathrm{Tr}(B_I)$ is a direct summand in a direct sum of several copies of $\mathrm{Tr}(\mathbb{1})$, where B_I is the Soergel bimodule corresponding to the longest element in a parabolic subgroup $W_I \subset W$, for any subset $I \subset S$ of simple transpositions. Next, we use an explicit “annular simplification” algorithm to present $\mathrm{Tr}(B)$ as a direct summand in a direct sum of $\mathrm{Tr}(B_I)$ for any Soergel bimodule B . Finally, we show that any complex of

Soergel bimodules is mapped by the trace functor to a summand in a finite twisted complex built out of $\mathrm{Tr}(\mathbb{1})$, thus completing the proof.

8.1. Frobenius extensions and horizontal trace. In this self-contained section we show that Soergel bimodules B_I associated to longest elements of finite parabolic subgroups W_I of a Coxeter group W always have traces $\mathrm{Tr}_0(W_I)$ isomorphic to summands of $\mathrm{Tr}_0(\mathbb{1})$, after Karoubi completion. We prove this in a slightly more general setting, using the language of Frobenius extensions.

Definition 8.2. A *Frobenius extension* is an extension of commutative rings $\iota: A \hookrightarrow B$, such that B is free and finitely generated as an A -module, equipped with a non-degenerate A -linear map $\partial: B \rightarrow A$, called the *trace*. Here, *non-degeneracy* asserts the existence of A -linear *dual bases* $\{x_\alpha\}$ and $\{y_\alpha\}$ for B such that $\partial(x_\alpha y_\beta) = \delta_{\alpha,\beta}$ (the Kronecker delta).

Example 8.3. Let (W, S) be a Coxeter system of finite rank as in §7 and let R again denote the base ring associated with a reflection faithful balanced realization over \mathbb{C} . For every finite parabolic subgroup W_I , let R^I denote the subring of W_I -invariants in R . Let $w_I \in W_I$ be the longest element. Then $\iota: R^I \rightarrow R$ is a graded Frobenius extension of rank $|W_I|$ with trace $\partial = \partial_I = \partial_{s_1} \cdots \partial_{s_r}$ where $w_I = s_1 \cdots s_r$ is a reduced expression and

$$\partial_s(f) = \frac{f - s(f)}{\alpha_s}.$$

See e.g. [Wil08, Section 3].

Remark 8.4. The subsets I corresponding to finite parabolic subgroups W_I are called *finitary*. Note that we do not need W to be finite in Example 8.3.

Example 8.5. In type A , we have $R = \mathbb{C}[X_1, \dots, X_n]$ and $W = S_n$ and $R^{S_n} \hookrightarrow R$ is a graded Frobenius extension of rank $n!$. An R^{S_n} -linear basis of R is given by the monomials $X_1^{a_1} X_2^{a_2} \cdots X_{n-1}^{a_{n-1}}$ where $0 \leq a_i \leq n - i$. Then we have

$$\partial(X_1^{a_1} X_2^{a_2} \cdots X_{n-1}^{a_{n-1}}) = \begin{cases} 1 & \text{if } a_i = n - i \text{ for } 1 \leq i \leq n, \\ 0 & \text{otherwise} \end{cases}.$$

The basis dual to the monomial basis has elements $\prod_{k=1}^{n-1} (-1)^{b_k} e_{b_k}(X_{n+1-k}, \dots, X_n)$ where $b_k = k - a_{n-k}$ for $1 \leq k \leq n - 1$.

Given a Frobenius extension $\iota: A \rightarrow B$ with trace ∂ , we have the following maps of B, B -bimodules, which are best encoded diagrammatically:

- the “multiplication” $\blacktriangleright: B \otimes_A B \otimes_A B \rightarrow B \otimes_A A \otimes_A B \cong B \otimes_A B$,
- the “inclusion” $\blacktriangleleft: B \rightarrow B \otimes_A B$ defined by $1 \mapsto \sum_\alpha x_\alpha \otimes y_\alpha$, and
- the “trace” $\blacktriangleright: B \otimes_A B \rightarrow B$ given by $x \otimes y \mapsto xy$,

which exhibit $B \otimes_A B$ as a Frobenius extension of B . This is an instance of Jones’ basic construction [Jon83]. Further we have:

- $\blacktriangleright: B \otimes_A B \rightarrow B \otimes_A B \otimes_A B$ given by $1 \otimes 1 \mapsto 1 \otimes 1 \otimes 1$.

These satisfy the relations:

$$\begin{array}{c} \text{X} \\ \text{H} \\ \text{N} \end{array} = \begin{array}{c} \text{H} \\ \text{N} \end{array}, \quad \begin{array}{c} \text{I} \\ \text{J} \\ \text{K} \\ \text{L} \\ \text{M} \end{array} = \begin{array}{c} \text{I} \\ \text{J} \\ \text{K} \\ \text{L} \\ \text{M} \end{array}, \quad \begin{array}{c} \text{P} \\ \text{Q} \end{array} = \sum_{\alpha} x_{\alpha} | y_{\alpha}, \quad \begin{array}{c} \text{R} \\ \text{S} \end{array} = \partial(p) |$$

We also define the following shorthand notation:

$$\begin{array}{c} \text{A} \\ \text{B} \\ \text{C} \end{array} := \begin{array}{c} \text{A} \\ \text{B} \\ \text{C} \end{array}, \quad \begin{array}{c} \text{D} \\ \text{E} \end{array} := \begin{array}{c} \text{D} \\ \text{E} \end{array}, \quad \begin{array}{c} \text{F} \\ \text{G} \end{array} := \begin{array}{c} \text{F} \\ \text{G} \end{array}$$

These morphisms satisfy the expected string-straightening and vertex rotation relations.

Definition 8.6. In the following, we will write $B_I := R \otimes_{R^I} \otimes R$ and call this a *generalised Bott–Samelson bimodule*. See also [Eli16].

Lemma 8.7. *With the same assumptions as in Example 8.3, we have an isomorphism*

$$\mathrm{Tr}_0(B_I) \cong \oplus_{|W_I|} \left(\mathrm{Tr}_0(\mathbb{1}), \frac{[\mathrm{Id}_{B_I}]}{|W_I|} \right)$$

in $\mathrm{Kar} \mathrm{Tr}_0(\mathrm{SBim}(W))$, where we have used the identification

$$\mathrm{End}_{\mathrm{Tr}_0(\mathrm{SBim}(W))}(\mathrm{Tr}_0(\mathbb{1})) \cong \mathrm{HH}_0(\mathrm{SBim}(W))$$

to describe the idempotents appearing on the right-hand side.

Proof. The Frobenius extension provides an isomorphism $\phi: B_I \star B_I \cong \oplus_{|W_I|} B_I$, which admits a convenient diagrammatic description, c.f. [Eli16, Section 4]. To make it explicit, let $\{x_{\alpha}\}$ and $\{y_{\alpha}\}$ denote dual bases of R as a free R^I -module, where $\alpha \in W_I$. Then $\phi = \oplus_{\alpha \in W_I} \phi_{\alpha}$ with

$$\phi_{\alpha}: B_I \star B_I \rightarrow B_I, \quad \phi_{\alpha}((r_1 \star r_2) \star (r_3 \star r_4)) := \partial(x_{\alpha} r_2 r_3)(r_1 \star r_4), \quad \phi_{\alpha} = \begin{array}{c} \text{A} \\ \text{B} \end{array}$$

and $\phi^{-1} = \sum_{\beta \in W_I} \phi_{\beta}^{-1}$ with

$$\phi_{\beta}^{-1}: B_I \rightarrow B_I \star B_I, \quad \phi_{\beta}^{-1}(r_1 \star r_2) := \sum_{\alpha \in W_I} (r_1 \star 1) \star (y_{\beta} \star r_2), \quad \phi_{\beta}^{-1} = \begin{array}{c} \text{C} \\ \text{D} \end{array}$$

Now in $\mathrm{Kar} \mathrm{Tr}_0(\mathrm{SBim}(W))$ we have:

$$\begin{aligned} \mathrm{Tr}_0(B_I) &= \begin{array}{c} \text{E} \\ \text{F} \end{array} = \sum_{\alpha} \begin{array}{c} \text{G} \\ \text{H} \end{array} = \sum_{\alpha, \beta} \frac{1}{|W_I|} \begin{array}{c} \text{I} \\ \text{J} \end{array} \\ &= \sum_{\alpha, \beta} \frac{1}{|W_I|} \begin{array}{c} \text{K} \\ \text{L} \end{array} = \sum_{\alpha} \frac{1}{|W_I|} \begin{array}{c} \text{M} \\ \text{N} \end{array} \end{aligned}$$

Conversely we have:

$$\frac{1}{|W_I|} \begin{array}{c} \text{O} \\ \text{P} \end{array} = \frac{1}{|W_I|} \begin{array}{c} \text{Q} \\ \text{R} \end{array} = \frac{\delta_{\alpha, \beta}}{|W_I|} \begin{array}{c} \text{S} \\ \text{T} \end{array}$$

This implies that $\frac{1}{|W_I|} \overline{\mathbb{1}}^{x_\alpha}$ and $\overline{\mathbb{1}}^{y_\alpha}$ are the components of inverse isomorphisms

$$\mathrm{Tr}_0(B_I) \leftrightarrow \oplus_{|W_I|} \left(\mathrm{Tr}_0(\mathbb{1}), \frac{[\mathrm{Id}_{B_I}]}{|W_I|} \right).$$

□

Remark 8.8. An analogous argument in the derived setting shows that

$$\mathrm{Tr}(B_I) \cong \oplus_{|W_I|} \left(\mathrm{Tr}(\mathbb{1}), \frac{[\mathrm{Id}_{B_I}]}{|W_I|} \right)$$

in $\mathrm{Kar} \mathrm{Tr}(\mathrm{SBim}(W))$. The main difference is that isotopies of diagrammatic morphisms through the seam of the annulus are now only possible only up to homotopy. One thus arrives at homotopy idempotents, which split in the Karoubi envelope, as described in Section 4.

We can describe the idempotent $\frac{[\mathrm{Id}_{B_I}]}{|W_I|}$ more explicitly. Indeed, it is well known that B_I can be presented as a twisted complex consisting of $\Delta_w[\ell(w)]$ for $w \in W_I$, so by (13c) we get

$$[\mathrm{Id}_{B_I}] = \sum_{w \in W_I} (-1)^{\ell(w)} [\mathrm{Id}_{\Delta_w}] = \sum_{w \in W_I} w$$

where we identified $[\mathrm{Id}_{\Delta_w}]$ with $(-1)^{\ell(w)} w$ in $\mathrm{HH}_0(\mathrm{SBim}_W) = \mathrm{HH}_0(\mathcal{K}^b(\mathrm{SBim}_W))$ using Theorem 7.7. We get the following

Corollary 8.9. *We have*

$$\mathrm{Tr}(B_I) \cong \oplus_{|W_I|} \left(\mathrm{Tr}(\mathbb{1}), \frac{1}{|W_I|} \sum_{w \in W_I} w \right).$$

Example 8.10. Let $W = S_n$ and $W_I = S_{k_1} \times \cdots \times S_{k_r}$ for $k_1 + \cdots + k_r = n$. Then

$$\mathrm{Tr}(B_I) \simeq \bigoplus_{\lambda} (\mathrm{Tr}(\mathbb{1}), e_\lambda)^{\oplus x_{\lambda, k_1, \dots, k_r}}$$

where the sum is over all partitions λ of n , e_λ is an idempotent in $\mathbb{C}[S_n]$ corresponding to the irreducible representation V_λ and the graded multiplicities are given by

$$x_{\lambda, k_1, \dots, k_r} := [k_1]! \cdots [k_r]! c_{k_1, \dots, k_r}^\lambda$$

where $c_{k_1, \dots, k_r}^\lambda$ denotes the multiplicity of the Schur function s_λ in the product $h_{k_1} \cdots h_{k_r}$.

8.2. Explicit annular simplification in type A. Throughout this section, we work with type A Soergel bimodules corresponding to the action of $W = S_n$ on $V = \mathbb{C}^n$. We abbreviate the notation $B_i := B_{s_i}$ for the Bott-Samelson bimodules associated to the simple reflections $s_i \in S_n$. Moreover, if $|i - j| = 1$, we also consider the Soergel bimodule $B_{iji} = B_{jij} := R \otimes_{R^{(s_i, s_j)}} R$. Further, to declutter many expressions in this subsection, we will omit the \star indicating the composition of bimodules.

Lemma 8.11. *Any Bott-Samelson bimodule for S_n is isomorphic to a direct summand of a direct sum of Bott-Samelson bimodules, in each of which B_{n-1} appears at most once.*

Proof. This is a standard argument which uses $B_i B_i \cong B_i(1) \oplus B_i(-1)$, $B_i B_j \cong B_j B_i$ if $|i - j| > 1$, and $B_i B_j B_i \cong B_{iji} \oplus B_i$ if $|i - j| = 1$. \square

For each subset $I = \{i_1 < \dots < i_k\} \subset S = \{1, \dots, n-1\}$, we define the Coxeter-Bott-Samelson bimodule $\underline{B}_I := B_{i_1} B_{i_2} \dots B_{i_k}$.

Lemma 8.12. *For any Soergel bimodule B the trace $\text{Tr}(B)$ is isomorphic to a direct summand in the direct sum of traces of Coxeter-Bott-Samelson bimodules as defined above.*

Proof. The proof is by induction on $n \geq 1$. The base case is trivial. By Lemma 8.11 we can present B as a direct summand in the direct sum of objects of the form X or $X B_{n-1} Y$, where $X, Y \in \text{SBim}_{n-1}$. The first case is taken care of by induction. In the second case we can use Lemma 6.21 to replace $\text{Tr}(X B_{n-1} Y)$ by $\text{Tr}(Y X B_{n-1})$. Now by Lemma 8.11 we can either write $Y X \subset_{\oplus} X'$ or $Y X \subset_{\oplus} X' B_{n-2} Y'$ for some $X', Y' \in \text{SBim}_{n-2}$. Thus we either have $Y X B_{n-1} \subset_{\oplus} X' B_{n-1}$ or

$$\text{Tr}(Y X B_{n-1}) \subset_{\oplus} \text{Tr}(X' B_{n-2} Y' B_{n-1}) \cong \text{Tr}(Y' X' B_{n-2} B_{n-1}),$$

which is taken care of by induction. \square

For the following let L_n denote the indecomposable Soergel bimodule corresponding to the longest element in S_n .

Lemma 8.13. *In SBim_{n+1} we have:*

$$L_n B_n L_n \cong [n-1]! L_{n+1} \oplus [n]! L_n$$

where we use quantum numbers to indicate multiple direct summands with grading shifts.

Example 8.14. For $n = 2$ we have $X_2 = B_1$, so $B_1 B_2 B_1 \cong B_{121} \oplus B_1$.

Proof. Let $R = \mathbb{C}[X_1, \dots, X_{n+1}]$

$$\begin{aligned} L_n B_n L_n &= R \otimes_{R^{S_n \times S_1}} R \otimes_{R^{S_1 \times \dots \times S_1 \times S_2}} R \otimes_{R^{S_n \times S_1}} R \\ &\cong [n-1]! R \otimes_{R^{S_n \times S_1}} R^{S_{n-1} \times S_1 \times S_1} \otimes_{R^{S_{n-1} \times S_2}} R^{S_{n-1} \times S_1 \times S_1} \otimes_{R^{S_n \times S_1}} R \\ &\cong [n-1]! R \otimes_{R^{S_{n+1}}} R \oplus [n-1]! [n] R \otimes_{R^{S_n \times S_1}} R \\ &= [n-1]! L_{n+1} \oplus [n]! L_n \end{aligned}$$

Here we have used the well-known “square-switch” isomorphism of singular Bott-Samelson bimodules to proceed to the third line. This can, for example, be deduced from [Wu14, Lemma 11.2]— the corresponding statement for matrix factorizations— by taking homology with respect to the positive differential and forgetting the negative differential, in the terminology of *loc. cit.* \square

In the previous proof we have used the fact that $\mathbb{C}[X_1, \dots, X_{n-1}]$ is a free module of (graded) rank $[n-1]!$ of $\mathbb{C}[X_1, \dots, X_{n-1}]^{S_{n-1}}$ (in fact, a Frobenius extension).

Lemma 8.15. *In $\text{Kar Tr}(\text{SBim}_n)$ we have that $\text{Tr}(B_1 \dots B_{n-1})$ is isomorphic to a summand in a direct sum of traces of Soergel bimodules corresponding to the longest elements in parabolic subgroups in S_n .*

Where we abbreviate $x := (x_1 - x_2)/2$ and $\theta := (\theta_2 - \theta_1)/2$. Since we work over \mathbb{C} , we may also rescale to get $x = x_1 - x_2$ and $\theta = \theta_1 - \theta_2$. (Note that the second line indeed shows a twisted complex, and the dashed arrows encode a degree zero closed invertible morphism between twisted complexes.) In the underived horizontal trace the long arrow would be zero, and the complex would split into three direct summands.

8.3. Proof of Theorem 8.1. In this section we prove Theorem 8.1. First we specialize some general facts about the dg category $\text{Perf}(A)$ proved in section 4.6 to the case of $A := R \otimes \Lambda \rtimes \mathbb{C}[S_n]$.

Recall that the generators θ_i of Λ have cohomological degree -1 , so A is indeed supported in nonpositive cohomological degrees. Since A has zero differential, we have $\text{Kar}^{dg}\langle A \rangle = \text{Kar}\langle A \rangle$ where $\langle A \rangle$ is the category of free A modules of finite rank. Because we work over a field of characteristic zero, $\text{Kar}\langle A \rangle$ is a semisimple category with finitely many indecomposable objects labeled by irreducible representations of S_n . The category $\text{Perf}(A) = \text{Pretr}(\text{Kar}\langle A \rangle)$ consists of twisted complexes built out of these objects. By Theorem 4.26 the category $\text{Perf}(A)$ is homotopy idempotent complete.

Proof of Theorem 8.1. The quasi-equivalence between $\text{Tr}(\text{SBim}_n)$ and $\text{Perf}(A)$ is constructed in two steps.

First, recall that the endomorphism algebra of $\text{Tr}(\mathbb{1})$ can be identified with the cyclic bar complex $\mathbf{C}(\text{SBim}_n)$. We have therefore a functor

$$(21) \quad \text{Hom}_{\text{Tr}}(\text{Tr}(\mathbb{1}), -) : \text{Tr}(\text{SBim}_n) \rightarrow \text{mod-}\mathbf{C}(\text{SBim}_n)$$

By Lemma 8.12 we can resolve any object in $\text{Tr}(\text{SBim}_n)$ by the traces of Coxeter-Bott-Samelson bimodules. By Lemma 8.15 the trace of every such bimodule is equivalent to a summand in the sum of traces of indecomposable Soergel bimodules corresponding to the longest elements of parabolic subgroups. Finally, by Lemma 8.7 the trace of any such bimodule is equivalent to a summand in the direct sum of several copies of the trace of the identity bimodule.

This means that any object in the essential image of (21) is homotopy equivalent to a direct summand in a free $\mathbf{C}(\text{SBim}_n)$ -module, and (21) defines a quasi-fully faithful quasi-functor³

$$\text{Hom}_{\text{Tr}}(\text{Tr}(\mathbb{1}), -) : \text{Tr}(\text{SBim}_n) \rightarrow \text{Kar}^{dg}\langle \mathbf{C}(\text{SBim}_n) \rangle$$

The corresponding functor

$$(22) \quad \text{Kar}^{dg}(\text{Pretr}(\text{Tr}(\text{SBim}_n))) \rightarrow \text{Kar}^{dg}(\text{Pretr}(\text{Kar}^{dg}\langle \mathbf{C}(\text{SBim}_n) \rangle))$$

$$(23) \quad = \text{Kar}^{dg}(\text{Perf}(\mathbf{C}(\text{SBim}_n)))$$

obtained by successively applying Pretr and Kar^{dg} on both sides is then a quasi-equivalence.

³Note that the category of $\mathbf{C}(\text{SBim}_n)$ -modules homotopy equivalent to direct summands of free modules is quasi-equivalent to $\text{Kar}^{dg}\langle \mathbf{C}(\text{SBim}_n) \rangle$

For the second step, observe that by Theorem 7.7 the dg algebra $\mathbf{C}(\text{SBim}_n)$ deformation retracts onto A and by Theorem 7.10 all higher A_∞ operations on A induced by this deformation retract vanish. As in Lemma 4.28, we get quasi-equivalences $\langle \mathbf{C}(\text{SBim}_n) \rangle \simeq \langle A \rangle$ and

$$\text{Kar}^{dg}(\text{Perf}(\mathbf{C}(\text{SBim}_n))) \simeq \text{Kar}^{dg}(\text{Perf}(A)).$$

Since $\text{Perf}(A)$ is Karoubian, $\text{Kar}^{dg}(\text{Perf}(A)) \simeq \text{Perf}(A)$.

By combining (22) with all these quasi-equivalences, we conclude that

$$\text{Kar}^{dg}(\text{Pretr}(\text{Tr}(\text{SBim}_n))) \simeq \text{Perf}(A). \quad \square$$

Corollary 8.17. *We have the following equivalence of categories:*

$$\text{Kar}(\text{Pretr}(\text{Tr}_0(\text{SBim}_n))) \simeq \text{Ch}^b(\text{Kar}(\langle A \rangle)) \simeq \text{Perf}(R \rtimes \mathbb{C}[S_n]).$$

In particular, after taking the pretriangulated hull and Karoubi completion the functor $\text{Tr}(\text{SBim}_n) \rightarrow \text{Tr}_0(\text{SBim}_n)$ can be identified with the forgetful functor $\epsilon: \text{Perf}(A) \rightarrow \text{Ch}^b(A)$ defined in section 4.6.

Proof. The algebra A is supported in nonpositive cohomological degrees, with $R \rtimes \mathbb{C}[S_n]$ in cohomological degree zero. Therefore

$$\text{Ch}^b(\text{Kar}(\langle A \rangle)) \simeq \text{Ch}^b(\text{Kar}(R \rtimes \mathbb{C}[S_n])) = \text{Perf}(R \rtimes \mathbb{C}[S_n]). \quad \square$$

8.4. A derived annular Khovanov-Rozansky invariant. In [QR18, GW19] the (un-derived) traces of web categories were related to annular Khovanov-Rozansky invariants, and to the Khovanov-Rozansky homology of links in \mathbb{R}^3 . In this subsection we review this construction using the category SBim_n and its trace.

Given a braid word $\underline{\beta}$ on n strands, let $F(\underline{\beta})$ again denote the Rouquier complex for $\underline{\beta}$ as defined in Section 7.1, which we now consider as an object in $K^b(\text{SBim}_n)$, the bounded homotopy category of SBim_n . We have already mentioned that these complexes satisfy braid relations up to (canonical) homotopy equivalence. In fact, braid cobordisms induce natural chain maps (up to homotopy) between Rouquier complexes [EK10].

Consider the underived trace functor $\text{Tr}_0: \text{SBim}_n \rightarrow \text{Tr}_0(\text{SBim}_n)$, with the target considered as embedded in the Karoubi completion $\text{Kar}(\text{Tr}_0(\text{SBim}_n))$. Recall that the latter is equivalent to the category of graded projective $R \rtimes \mathbb{C}[S_n]$ -modules. In the following definition we use the functor Tr_0 , extended to the bounded homotopy categories of the source and target.

Definition 8.18. The *annular Khovanov–Rozansky invariant* of a braid word $\underline{\beta}$ on n strands is defined as

$$\text{AKhR}(\underline{\beta}) := \text{Hom}_{\text{Tr}_0}(\text{Tr}_0(\mathbb{1}), \text{Tr}_0(F(\underline{\beta}))) \in \text{Perf}(R \rtimes \mathbb{C}[S_n])$$

By virtue of factoring through the underived trace, AKhR is a categorical invariant of braid conjugacy classes (a.k.a. annular links with a coherent orientation) which is natural under annular link cobordisms (preserving the coherent orientation). More

precisely, AKhR is defined on n -strand braid words for each $n \geq 0$ separately, but these invariants fit together to give a *monoidal* annular link invariant, see [GW19] for details. Relatives of this notion of annular Khovanov–Rozansky invariant have previously been studied in [QR18, QRS18]. Another ahistorical aspect of our presentation here is that the interest in *annular* Khovanov–Rozansky invariants rose well after the construction of the triply-graded Khovanov–Rozansky homology of links in \mathbb{R}^3 [KR08, Kho07], which categorifies the HOMFLY-PT polynomial. We will comment on their relationship in the next section.

We are now ready to define a derived annular Khovanov–Rozansky invariant.

Definition 8.19. The *derived annular Khovanov–Rozansky invariant* is defined on closures of n -strand braid words $\underline{\beta}$ as

$$\text{AKhR}_{\text{dg}}(\hat{\underline{\beta}}) := \text{Hom}_{\text{Tr}}(\text{Tr}(\mathbb{1}), \text{Tr}(F(\underline{\beta}))) \in \text{Perf}(R \otimes \Lambda \rtimes \mathbb{C}[S_n])$$

where $F(\underline{\beta})$ is the Rouquier complex of $\underline{\beta}$, Tr is the universal dg monoidal trace, and we use that $\text{Hom}_{\text{Tr}}(\text{Tr}(\mathbb{1}), -)$ realises the equivalence from Theorem 8.1. As for the underived annular Khovanov–Rozansky invariant, this construction is functorial under braid-like annular link cobordisms between braid closures up to homotopy.

One important feature of $\text{AKhR}_{\text{dg}}(\hat{\underline{\beta}})$ is that the 2π rotation of the annular link $\hat{\underline{\beta}}$ typically induces an interesting endomorphism of $\text{AKhR}_{\text{dg}}(\hat{\underline{\beta}})$, while it always induces the identity map on $\text{KhR}(\hat{\underline{\beta}})$ and its annular version.

Recall that by (20) the Connes differential $\mathbb{B} = \sum \theta_i \frac{\partial}{\partial x_i}$ defines a derivation on the algebra $\text{HH}_*(\text{SBim}_n) \cong R \otimes \Lambda \rtimes \mathbb{C}[S_n]$. This allows us to define an interesting endofunctor on the category of twisted $R \otimes \Lambda \rtimes S_n$ -modules. Given a twisted complex (A, d) we have:

$$0 = \mathbb{B}(d^2) = \mathbb{B}(d)d + d\mathbb{B}(d),$$

so that $\mathbb{B}(d)$ is always a closed (degree 1) endomorphism of (A, d) .

Conjecture 8.20. *The action of the rotator w_X (defined in Remark 6.23) on twisted complexes X built out of summands of $\text{Tr}(\mathbb{1})$ is homotopic to $\text{Id} + \mathbb{B}$, where \mathbb{B} is the action of Connes differential defined above.*

Example 8.21. Let f be an arbitrary polynomial in $R = \text{End}_{\text{SBim}_n}(\mathbb{1})$, let $X = \text{Cone}(f)$. Since the rotator $w_{\mathbb{1}} : \text{Tr}(\mathbb{1}) \rightarrow \text{Tr}(\mathbb{1})$ is trivial, by (17) the rotator w_X is given by the morphism

$$\begin{array}{ccc} [\text{Tr}(\mathbb{1}) & \xrightarrow{f} & \text{Tr}(\mathbb{1}) \\ \downarrow \text{Id} & \searrow w(f; \mathbb{1}) & \downarrow \text{Id} \\ [\text{Tr}(\mathbb{1}) & \xrightarrow{f} & \text{Tr}(\mathbb{1}) \end{array}$$

It is easy to see that $w(f; \mathbb{1}) = \|f\| \text{Id}$ is homotopic to $\mathbb{B}(f)$.

Another important feature is that the derived annular Khovanov–Rozansky invariant of an annular link has an action of the derived center $\mathcal{Z}^{\text{dg}}(\text{SBim}_n)$. The second author has shown with Ben Elias that the Rouquier complex of the full twist braid FT_n (together with suitable half-braiding data) is an object of the derived Drinfeld center [EH]. The corresponding endofunctor of the derived trace sends $\text{AKhR}_{\text{dg}}(\hat{\beta})$ to $\text{AKhR}_{\text{dg}}(\widehat{\text{FT}_n\beta})$, i.e. it cuts the thickened annulus containing the annular link $\hat{\beta}$ and re-glues it after a 2π twist to create $\widehat{\text{FT}_n\beta}$.

This operation suggests that AKhR_{dg} should be considered as an invariant of (coherently oriented) links in $S^1 \times D^2$, which can be computed by choosing an I -bundle structure on $S^1 \times D^2$, but which comes with the data necessary to change I -bundle structure.

8.5. Triply graded homology and the Hochschild cohomology of Soergel bimodules. In this section we focus on the Hochschild cohomology of individual Soergel bimodules M in $\text{SBim}(W)$ (we will soon specialise to SBim_n), not on the Hochschild (co)homology of the category SBim_n .

We define $\text{HH}^i(M) := \text{Ext}^i(R, M)$ where the Ext groups are computed in the category of R, R -bimodules. In particular, $\text{HH}^0(M) = \text{Hom}(R, M)$.

Given a complex $C = [\dots \rightarrow M_j \rightarrow M_{j+1} \rightarrow \dots]$ of Soergel bimodules, we define a complex of graded vector spaces

$$\text{HH}^i(C) := [\dots \rightarrow \text{HH}^i(M_j) \rightarrow \text{HH}^i(M_{j+1}) \rightarrow \dots].$$

Definition 8.22. Given a braid word $\underline{\beta}$ on n strands, the triply-graded *Khovanov–Rozansky homology* of the braid closure $\hat{\beta}$ is defined as

$$\text{KhR}(\hat{\beta}) := H^*(\oplus_i \text{HH}^i(F(\underline{\beta}))),$$

where $F(\underline{\beta})$ is the associated Rouquier complex of Soergel bimodules in SBim_n defined in Section 7.1. The vector space $\text{KhR}(\hat{\beta})$ is triply-graded by q -degree, cohomological degree in the complex $F(\underline{\beta})$, and Hochschild degree i .

As defined, KhR is a braid conjugacy invariant, i.e. an invariant of links in a thickened annulus that are obtained from braid closures. However, after an overall grading shift, KhR becomes invariant under the second Markov move, and thus an invariant of links in \mathbb{R}^3 . Since braid cobordisms induce natural chain maps (up to homotopy) between Rouquier complexes, one also has induced morphisms between the Khovanov–Rozansky homologies. If we also take into account the morphisms induced by braid conjugation on the level of Hochschild homology, this can be summarised by saying that KhR is functorial under braidlike annular link cobordisms between braid closures.

Next we explain how the triply-graded Khovanov–Rozansky homology can be recovered from the annular Khovanov–Rozansky invariant (derived or underived). For an alternative but related approach see [QRS18].

It is well-known that for arbitrary Soergel bimodules M and N there is a natural isomorphism

$$(24) \quad \mathrm{HH}^i(M \otimes N) \cong \mathrm{HH}^i(N \otimes M)$$

of q -graded vector spaces. In other words, HH^i is a trace-like linear functor and hence factors through Tr_0 . Moreover, in type A Rasmussen proved [Ras15, Proposition 4.6] that $\mathrm{HH}^i(M)$ is free over R . (More generally this is known for all Weyl groups, see Webster–Williamson [WW11]).

Lemma 8.23. *For a Soergel bimodule M in SBim_n we have*

$$\mathrm{HH}^i(M) \cong \mathrm{Hom}_{\mathrm{Tr}_0}(\mathbf{e}_{i,1^{n-i}} \mathrm{Tr}_0(\mathbb{1}), \mathrm{Tr}_0(M)) \oplus \mathrm{Hom}_{\mathrm{Tr}_0}(\mathbf{e}_{i-1,1^{n-i+1}} \mathrm{Tr}_0(\mathbb{1}), \mathrm{Tr}_0(M)).$$

where $\mathbf{e}_{i,1^{n-i}}$ is the idempotent in $\mathbb{C}[S_n]$ corresponding to the hook partition $(i, 1^{n-i})$. In particular,

$$\mathrm{Hom}(\mathbb{1}, M) \cong \mathrm{HH}^0(M) \cong \mathrm{Hom}_{\mathrm{Tr}_0}(\mathbf{e}_- \mathrm{Tr}_0(\mathbb{1}), \mathrm{Tr}_0(M))$$

where $\mathbf{e}_- = \mathbf{e}_{1^n}$ is the antisymmetrizer in $\mathbb{C}[S_n]$. The isomorphisms above are natural in M .

Proof. By (24) the functor HH^i is a trace-like functor from SBim_n to graded vector spaces, so as discussed in Section 6.5 it factors through $\mathrm{Tr}(\mathrm{SBim}_n)$ and defines a functor $\mathrm{HH}^i: \mathrm{Tr}(\mathrm{SBim}_n) \rightarrow \mathrm{Vect}$. By Theorem 8.1 we have that $\mathrm{Tr}(M)$ is isomorphic to the direct sum of direct summands of $\mathrm{Tr}(\mathbb{1})$, so it is sufficient to check it for $M = \mathrm{Tr}(\mathbb{1})$. Now

$$\mathrm{HH}^i(R) \cong R \otimes \wedge^i(\mathbb{C}^n) \cong R \otimes (V_{i,1^{n-i}} \oplus V_{i-1,1^{n-i+1}}).$$

while

$$\mathrm{Hom}_{\mathrm{Tr}_0}(\mathrm{Tr}_0(\mathbb{1}), \mathrm{Tr}_0(\mathbb{1}))\mathbf{e}_\lambda \cong (R \rtimes \mathbb{C}[S_n])\mathbf{e}_\lambda \cong R \otimes V_\lambda$$

for any irreducible representation V_λ of S_n . \square

Theorem 8.24. *For any complex C of Soergel bimodules one has the following isomorphism of complexes of graded vector spaces:*

$$\mathrm{HH}^i(C) \cong \mathrm{Hom}_{\mathrm{Tr}_0}(\mathbf{e}_{i,1^{n-i}} \mathrm{Tr}_0(\mathbb{1}), \mathrm{Tr}_0(C)) \oplus \mathrm{Hom}_{\mathrm{Tr}_0}(\mathbf{e}_{i-1,1^{n-i+1}} \mathrm{Tr}_0(\mathbb{1}), \mathrm{Tr}_0(C)).$$

Proof. The functor Tr_0 sends C to

$$\mathrm{Tr}_0(C) = [\dots \rightarrow \mathrm{Tr}_0(M_j) \rightarrow \mathrm{Tr}_0(M_{j+1}) \rightarrow \dots]$$

with no higher differentials. Now the statement follows from Lemma 8.23. \square

Theorem 8.24 implies that Khovanov–Rozansky homology can be computed by first applying the functor Tr_0 to $F(\underline{\beta})$, then running the annular simplification from Theorem 8.1.

Proposition 8.25. *For a braid word $\underline{\beta}$ on n strands we have:*

$$\mathrm{KhR}(\hat{\beta}) \cong \bigoplus_i \mathrm{Hom}_{\mathrm{Tr}_0}((\mathbf{e}_{i,1^{n-i}} + \mathbf{e}_{i-1,1^{n-i+1}}) \mathrm{Tr}_0(\mathbb{1}), \mathrm{Tr}_0(F(\underline{\beta})))$$

Remark 8.26. In [GW19] the first and third authors defined the *evaluation* functor $\mathrm{Tr}_0(\mathrm{SBim}_n) \rightarrow \mathrm{Vect}$ which sends $\mathrm{Tr}_0(\mathbb{1})$ to R as a module over $R \rtimes \mathbb{C}[S_n]$. This is equivalent to the above since

$$\mathrm{Hom}_{\mathrm{Tr}_0}(\mathbf{e}_- \mathrm{Tr}_0(\mathbb{1}), \mathrm{Tr}_0(\mathbb{1})) = (R \rtimes \mathbb{C}[S_n])\mathbf{e}_- = R.$$

Similarly, higher Khovanov-Rozansky homology HH^i can be computed by evaluating $\mathrm{Tr}(\mathbb{1})$ to $\mathrm{HH}^i(R) = R \otimes \wedge^i(\mathbb{C}^n)$ as a module over $R \rtimes \mathbb{C}[S_n]$.

REFERENCES

- [AF19] David Ayala and John Francis. A factorization homology primer. In *Handbook of Homotopy Theory*. Chapman and Hall/CRC, 2019. [arXiv:2019arXiv190310961A](#).
- [BFO09] Roman Bezrukavnikov, Michael Finkelberg, and Victor Ostrik. On tensor categories attached to cells in affine Weyl groups. III. *Israel J. Math.*, 170:207–234, 2009. [MR2506324](#) [DOI:10.1007/s11856-009-0026-9](#) [arXiv:math/0605628](#).
- [BGHT99] François Bergeron, Adriano M Garsia, Mark Haiman, and Glenn Tesler. Identities and positivity conjectures for some remarkable operators in the theory of symmetric functions. *Methods and Applications of Analysis*, 6(3):363–420, 1999.
- [BHLv17] Anna Beliakova, Kazuo Habiro, Aaron D. Lauda, and Marko Živković. Trace decategorification of categorified quantum \mathfrak{sl}_2 . *Math. Ann.*, 367(1-2):397–440, 2017. [MR3606445](#) [DOI:10.1007/s00208-016-1389-y](#) [arXiv:1404.1806](#).
- [BKR01] Tom Bridgeland, Alastair King, and Miles Reid. The McKay correspondence as an equivalence of derived categories. *J. Amer. Math. Soc.*, 14(3):535–554, 2001. [MR1824990](#) [DOI:10.1090/S0894-0347-01-00368-X](#) [arXiv:math/9908027](#).
- [BN93] Marcel Bökstedt and Amnon Neeman. Homotopy limits in triangulated categories. *Compositio Math.*, 86(2):209–234, 1993. [MR1214458](#).
- [BN09] David Ben-Zvi and David Nadler. *The Character Theory of a Complex Group*, 2009. [arXiv:0904.1247](#).
- [BPW19] Anna Beliakova, Krzysztof K. Putyra, and Stephan M. Wehrli. Quantum link homology via trace functor I. *Invent. Math.*, 215(2):383–492, 2019. [MR3910068](#) [DOI:10.1007/s00222-018-0830-0](#) [arXiv:1605.03523](#).
- [BV08] Alexander Beilinson and Vadim Vologodsky. A DG guide to Voevodsky’s motives. *Geom. Funct. Anal.*, 17(6):1709–1787, 2008. [MR2399083](#) [DOI:10.1007/s00039-007-0644-5](#) [arXiv:math/0604004](#).
- [BZFN10] David Ben-Zvi, John Francis, and David Nadler. Integral transforms and Drinfeld centers in derived algebraic geometry. *J. Amer. Math. Soc.*, 23(4):909–966, 2010. [MR2669705](#) [DOI:10.1090/S0894-0347-10-00669-7](#) [arXiv:0805.0157](#).
- [EH] Benjamin Elias and Matthew Hogancamp. In preparation.
- [EK95] David E. Evans and Yasuyuki Kawahigashi. On Ocneanu’s theory of asymptotic inclusions for subfactors, topological quantum field theories and quantum doubles. *Internat. J. Math.*, 6(2):205–228, 1995. [MR1316301](#) [DOI:10.1142/S0129167X95000468](#).
- [EK10] Ben Elias and Dan Krasner. Rouquier complexes are functorial over braid cobordisms. *Homology Homotopy Appl.*, 12(2):109–146, 2010. [MR2721032](#) [euclid.hha/1296223879](#) [arXiv:0906.4761](#).
- [EL16] Ben Elias and Aaron D. Lauda. Trace decategorification of the Hecke category. *J. Algebra*, 449:615–634, 2016. [MR3448186](#) [DOI:10.1016/j.jalgebra.2015.11.028](#) [arXiv:1504.05267](#).
- [Eli16] Ben Elias. Thicker Soergel calculus in type A . *Proc. Lond. Math. Soc. (3)*, 112(5):924–978, 2016. [MR3502025](#) [DOI:10.1112/plms/pdw012](#) [arXiv:1009.2120](#).

- [EW14] Ben Elias and Geordie Williamson. The Hodge theory of Soergel bimodules. *Ann. of Math. (2)*, 180(3):1089–1136, 2014. [MR3245013](#) [DOI:10.4007/annals.2014.180.3.6](#) [arXiv:1212.0791](#).
- [EW16] Ben Elias and Geordie Williamson. Soergel calculus. *Represent. Theory*, 20:295–374, 2016. [MR3555156](#) [DOI:10.1090/ert/481](#) [arXiv:1309.0865](#).
- [EZ53] Samuel Eilenberg and J. A. Zilber. On products of complexes. *Amer. J. Math.*, 75:200–204, 1953. [MR52767](#) [DOI:10.2307/2372629](#).
- [GH17] Eugene Gorsky and Matthew Hogancamp. Hilbert schemes and y -ification of khovanov-rozansky homology, 2017. [arXiv:1712.03938](#).
- [GHMN19] Eugene Gorsky, Matthew Hogancamp, Anton Mellit, and Keita Nakagane. Serre duality for Khovanov–Rozansky homology. 2019. [arXiv:1902.08281](#).
- [GNR16] Eugene Gorsky, Andrei Neguț, and Jacob Rasmussen. Flag Hilbert schemes, colored projectors and Khovanov–Rozansky homology, 2016. [arXiv:1608.07308](#).
- [GW19] Eugene Gorsky and Paul Wedrich. Evaluations of annular Khovanov–Rozansky homology, 2019. [arXiv:1904.04481](#).
- [Hai01] Mark Haiman. Hilbert schemes, polygraphs and the Macdonald positivity conjecture. *J. Amer. Math. Soc.*, 14(4):941–1006, 2001. [MR1839919](#) [DOI:10.1090/S0894-0347-01-00373-3](#) [arXiv:math/0010246](#).
- [HKR62] G. Hochschild, Bertram Kostant, and Alex Rosenberg. Differential forms on regular affine algebras. *Trans. Amer. Math. Soc.*, 102:383–408, 1962. [MR142598](#) [DOI:10.2307/1993614](#).
- [Hog17] Matthew Hogancamp. Idempotents in triangulated monoidal categories, 2017. [arXiv:1703.01001](#).
- [HPT16] André Henriques, David Penneys, and James Tener. Categorified trace for module tensor categories over braided tensor categories. *Doc. Math.*, 21:1089–1149, 2016. [MR3578212](#) [arXiv:1509.02937](#).
- [HSS17] Marc Hoyois, Sarah Scherotzke, and Nicolò Sibilla. Higher traces, noncommutative motives, and the categorified Chern character. *Adv. Math.*, 309:97–154, 2017. [MR3607274](#) [DOI:10.1016/j.aim.2017.01.008](#) [arXiv:1511.03589](#).
- [Jon83] Vaughan F. R. Jones. Index for subfactors. *Invent. Math.*, 72(1):1–25, 1983. [MR696688](#) [DOI:10.1007/BF01389127](#).
- [Kad80] Tornike V Kadeishvili. On the homology theory of fibre spaces. *Russian Mathematical Surveys*, 35(3):231, 1980. [MR580645](#).
- [Kel01] Bernhard Keller. Introduction to A -infinity algebras and modules. *Homology Homotopy Appl.*, 3(1):1–35, 2001. [MR1854636](#) [DOI:10.4310/hha.2001.v3.n1.a1](#) [arXiv:math/9910179](#).
- [Kel06] Bernhard Keller. On differential graded categories. In *International Congress of Mathematicians. Vol. II*, pages 151–190. Eur. Math. Soc., Zürich, 2006. [MR2275593](#).
- [Kho07] Mikhail Khovanov. Triply-graded link homology and Hochschild homology of Soergel bimodules. *Internat. J. Math.*, 18(8):869–885, 2007. [MR2339573](#) [DOI:10.1142/S0129167X07004400](#) [arXiv:math/0510265](#).
- [KR08] Mikhail Khovanov and Lev Rozansky. Matrix factorizations and link homology. II. *Geom. Topol.*, 12(3):1387–1425, 2008. [MR2421131](#) [arXiv:math.QA/0505056](#) [DOI:10.2140/gt.2008.12.1387](#).
- [Kuz09] Alexander Kuznetsov. Hochschild homology and semiorthogonal decompositions, 2009. [arXiv:0904.4330](#).
- [LH03] Kenji Lefèvre-Hasegawa. *Sur les A-infini catégories*. PhD thesis, U. Paris 7, 2003. [arXiv:math/0310337](#).

- [LQ84] Jean-Louis Loday and Daniel Quillen. Cyclic homology and the Lie algebra homology of matrices. *Comment. Math. Helv.*, 59(4):569–591, 1984. [MR780077](#) [DOI:10.1007/BF02566367](#).
- [Lus84] G. Lusztig. Intersection cohomology complexes on a reductive group. *Invent. Math.*, 75(2):205–272, 1984. [MR732546](#) [DOI:10.1007/BF01388564](#).
- [LW14] Nicolas Libedinsky and Geordie Williamson. Standard objects in 2-braid groups. *Proc. Lond. Math. Soc.* (3), 109(5):1264–1280, 2014. [MR3283617](#) [DOI:10.1112/plms/pdu022](#) [arXiv:1205.4206](#).
- [Mer99] S. A. Merkulov. Strong homotopy algebras of a Kähler manifold. *Internat. Math. Res. Notices*, (3):153–164, 1999. [MR1672242](#) [DOI:10.1155/S1073792899000070](#) [arXiv:math/9809172](#).
- [MV10] Marco Mackaay and Pedro Vaz. The diagrammatic Soergel category and $\mathfrak{sl}(N)$ -foams, for $N \geq 4$. *Int. J. Math. Math. Sci.*, pages Art. ID 468968, 20, 2010. [MR2671770](#) [arXiv:0911.2485](#).
- [MWW19] Scott Morrison, Kevin Walker, and Paul Wedrich. Invariants of 4-manifolds from Khovanov-Rozansky link homology, 2019. [arXiv:1907.12194](#).
- [Ocn94] Adrian Ocneanu. Chirality for operator algebras. In *Subfactors (Kyuzeso, 1993)*, pages 39–63. World Sci. Publ., River Edge, NJ, 1994. [MR1317353](#).
- [PS13] Kate Ponto and Michael Shulman. Shadows and traces in bicategories. *J. Homotopy Relat. Struct.*, 8(2):151–200, 2013. [MR3095324](#) [DOI:10.1007/s40062-012-0017-0](#) [arXiv:0910.1306](#).
- [PVdB19] Alexander Polishchuk and Michel Van den Bergh. Semiorthogonal decompositions of the categories of equivariant coherent sheaves for some reflection groups. *J. Eur. Math. Soc. (JEMS)*, 21(9):2653–2749, 2019. [MR3985610](#) [DOI:10.4171/JEMS/890](#) [arXiv:1503.04160](#).
- [QR16] Hoel Queffelec and David E. V. Rose. The \mathfrak{sl}_n foam 2-category: a combinatorial formulation of Khovanov-Rozansky homology via categorical skew Howe duality. *Adv. Math.*, 302:1251–1339, 2016. [MR3545951](#) [DOI:10.1016/j.aim.2016.07.027](#) [arXiv:1405.5920](#).
- [QR18] Hoel Queffelec and David E. V. Rose. Sutured annular Khovanov-Rozansky homology. *Trans. Amer. Math. Soc.*, 370(2):1285–1319, 2018. [MR3729501](#) [DOI:10.1090/tran/7117](#) [arXiv:1506.08188](#).
- [QRS18] Hoel Queffelec, David E. V. Rose, and Antonio Sartori. Annular evaluation and link homology, 2018. [arXiv:1802.04131](#).
- [QW18] Hoel Queffelec and Paul Wedrich. Khovanov homology and categorification of skein modules, 2018. [arXiv:1806.03416](#).
- [Ras15] Jacob Rasmussen. Some differentials on Khovanov–Rozansky homology. *Geom. Topol.*, 19(6):3031–3104, 2015. [MR3447099](#) [DOI:10.2140/gt.2015.19.3031](#) [arXiv:math/0607544](#).
- [Rid13] Laura Rider. Formality for the nilpotent cone and a derived Springer correspondence. *Adv. Math.*, 235:208–236, 2013. [MR3010057](#) [DOI:10.1016/j.aim.2012.12.001](#) [arXiv:1206.4343](#).
- [Rou04] Raphaël Rouquier. Categorification of the braid groups, 2004. [arXiv:0409593](#).
- [Rou06] Raphaël Rouquier. Categorification of \mathfrak{sl}_2 and braid groups. In *Trends in representation theory of algebras and related topics*, volume 406 of *Contemp. Math.*, pages 137–167. Amer. Math. Soc., Providence, RI, 2006. [MR2258045](#) [DOI:10.1090/conm/406/07657](#).
- [RR16] Laura Rider and Amber Russell. Perverse sheaves on the nilpotent cone and Lusztig’s generalized Springer correspondence. In *Lie algebras, Lie superalgebras, vertex algebras and related topics*, volume 92 of *Proc. Sympos. Pure Math.*, pages 273–292. Amer. Math. Soc., Providence, RI, 2016. [MR3644235](#) [arXiv:1409.7132](#).

- [RW16] David E.V. Rose and Paul Wedrich. Deformations of colored $\mathfrak{sl}(N)$ link homologies via foams. *Geom. Topol.*, 20(6):3431–3517, 2016. [MR3590355](#) [DOI:10.2140/gt.2016.20.3431](#) [arXiv:1501.02567](#).
- [Sei08] Paul Seidel. *Fukaya categories and Picard-Lefschetz theory*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008. [MR2441780](#) [DOI:10.4171/063](#).
- [Soe90] Wolfgang Soergel. Kategorie \mathcal{O} , perverse Garben und Moduln über den Koinvarianten zur Weylgruppe. *J. Amer. Math. Soc.*, 3(2):421–445, 1990. [MR1029692](#) [DOI:10.2307/1990960](#).
- [Soe92] Wolfgang Soergel. The combinatorics of Harish-Chandra bimodules. *J. Reine Angew. Math.*, 429:49–74, 1992. [MR1173115](#) [DOI:10.1515/crll.1992.429.49](#).
- [Soe07] Wolfgang Soergel. Kazhdan-Lusztig-Polynome und unzerlegbare Bimoduln über Polynomringen. *J. Inst. Math. Jussieu*, 6(3):501–525, 2007. [MR2329762](#) [DOI:10.1017/S1474748007000023](#).
- [Tab05] Goncalo Tabuada. Une structure de catégorie de modèles de Quillen sur la catégorie des dg-catégories. *C. R. Math. Acad. Sci. Paris*, 340(1):15–19, 2005. [MR2112034](#) [DOI:10.1016/j.crma.2004.11.007](#) [arXiv:math/0407338](#).
- [Toë07] Bertrand Toën. The homotopy theory of dg-categories and derived Morita theory. *Invent. Math.*, 167(3):615–667, 2007. [MR2276263](#) [DOI:10.1007/s00222-006-0025-y](#) [arXiv:math/0408337](#).
- [Wal] Kevin Walker. Topological quantum field theories. Available at <http://canyon23.net/math/>.
- [Wed19] Paul Wedrich. Exponential growth of colored HOMFLY-PT homology. *Adv. Math.*, 353:471–525, 2019. [MR3982970](#) [DOI:10.1016/j.aim.2019.06.023](#) [arXiv:1602.02769](#).
- [Wil08] Geordie Williamson. Singular Soergel bimodules, 2008. Dissertation, Freiburg, <http://people.mpim-bonn.mpg.de/geordie/GW-thesis.pdf>.
- [Wu14] Hao Wu. A colored $\mathfrak{sl}(N)$ homology for links in S^3 . *Dissertationes Math.*, 499:217, 2014. [MR3234803](#) [DOI:10.4064/dm499-0-1](#) [arXiv:0907.0695](#).
- [WW11] Ben Webster and Geordie Williamson. The geometry of Markov traces. *Duke Math. J.*, 160(2):401–419, 2011. [MR2852120](#) [DOI:10.1215/00127094-1444268](#) [arXiv:0911.4494](#).
- [Zhu18] Xinwen Zhu. Geometric Satake, categorical traces, and arithmetic of Shimura varieties. In *Current developments in mathematics 2016*, pages 145–206. Int. Press, Somerville, MA, 2018. [MR3837875](#) [arXiv:1810.07375](#).

E.G.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT DAVIS, ONE SHIELDS AVENUE, DAVIS CA 95616

E-mail address: egorskiy@math.ucdavis.edu

M.H.: DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, 360 HUNTINGTON AVE, BOSTON, MA 02115

E-mail address: m.hogancamp@northeastern.edu

P.W.: MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY

P.W.: MATHEMATICAL INSTITUTE, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY

P.W.: MATHEMATICAL SCIENCES RESEARCH INSTITUTE, 17 GAUSS WAY, BERKELEY, CA 94720. PAUL.WEDRICH.AT

E-mail address: p.wedrich@gmail.com