

UC Riverside

UC Riverside Previously Published Works

Title

High yield C-derivatization of weakly coordinating carborane anions.

Permalink

<https://escholarship.org/uc/item/47c5n0dd>

Journal

Inorganic chemistry, 49(11)

ISSN

0020-1669

Authors

Nava, Matthew J
Reed, Christopher A

Publication Date

2010-06-01

DOI

10.1021/ic100053u

Peer reviewed

High Yield C-Derivatization of Weakly Coordinating Carborane Anions

Matthew J. Nava and Christopher A. Reed*

Center for S and P Block Chemistry, Department of Chemistry, University of California, Riverside, California 92651

Received January 9, 2010

Unlike the “parent” carborane anion $\text{CHB}_{11}\text{H}_{11}^-$, halogenated carborane anions such as $\text{CHB}_{11}\text{H}_5\text{Br}_6^-$ can be readily C-functionalized in high yield and purity, enhancing their utility as weakly coordinating anions.

B-halogenated icosahedral carborane anions such as $\text{CHB}_{11}\text{H}_5\text{X}_6^-$ and $\text{CHB}_{11}\text{X}_{11}^-$ (X = halogen; Figure 1) are particularly useful members of a class of exceptionally inert, weakly coordinating anions^{1–5} whose versatility might be further tailored by suitable C-derivatization chemistry. Long-chain hydrocarbon “tails” formed by C-alkylation should lower the lattice energies of salts, increase their solubilities in low dielectric solvents, and allow better exploitation of reactive cations in catalysis. Applications in surfactant chemistry can also be envisioned. Similarly, C-fluorocarbon tails should improve the solubility of carborane ion pairs in fluorocarbon solvents, where catalytic applications have been reported.⁶ Attachment of a carborane anion to a polymer has allowed exploitation of immobilization chemistry in cation-selective sensor technology.⁷ C-arylation takes advantage of the unique scaffold of a carborane anion in rigid rod supramolecular chemistry.^{8,9}

Despite these promising applications, the C-derivatization chemistry of carborane anions has progressed rather slowly. Most work has been performed on the “parent” carborane anion, $\text{CHB}_{11}\text{H}_{11}^-$, but is frequently hampered by modest yields and difficult separations from starting material. Being ionic rather than neutral, the chromatographic separation of different carborane anions is not trivial on a synthetic scale. The activation of $\text{CHB}_{11}\text{H}_{11}^-$ via C-lithiation with butyl




Figure 1. Carborane anions $\text{CHB}_{11}\text{H}_5\text{X}_6^-$ and $\text{CHB}_{11}\text{X}_{11}^-$ (gray = C, white = H, red = H or halogen, green = halogen).

lithium appears to be essentially quantitative,¹⁰ but the partial regeneration of starting material during subsequent reactions with electrophiles is common, despite careful control of the conditions. Alkylation of $1\text{-Li-CB}_{11}\text{H}_{11}^-$ with alkyl halides gives mixed results. While the yields for methylations are frequently quite high,^{2,11–14} they drop to 63% for ethylation of $\text{CHB}_{11}\text{H}_{11}^-$ and are even lower for most other alkylations, silylations, phosphinations, and metalations.^{2,10} The yields of C-monohalogenated products, $1\text{-X-CB}_{11}\text{H}_{11}^-$, have been raised to 81–96% by careful attention to conditions, but chromatography purification is still recommended for most derivatives.¹¹

We now report that, when these C-functionalization reactions are performed on an already halogenated carborane anion such as $\text{CHB}_{11}\text{H}_5\text{Br}_6^-$, rather than on the unfunctionalized parent $\text{CHB}_{11}\text{H}_{11}^-$, isolated yields are generally excellent and compound purity is sufficiently high that chromatographic purification is unnecessary. High yield C-cyanation of undecahalogenated carboranes has very recently been reported.¹⁵ These findings makes sense within

*To whom correspondence should be addressed. E-mail: chris.reed@ucr.edu.

(1) Reed, C. A. *Acc. Chem. Res.* 2010, 43, 121–128.
(2) Koerber, S.; Schreiber, P. J.; Michl, J. *Chem. Rev.* 2006, 106, 5208–5249.
(3) Krossing, I.; Raabe, I. *Angew. Chem., Int. Ed.* 2004, 43, 2066–2090.
(4) Reed, C. A. *Acc. Chem. Res.* 1998, 31, 133–139.
(5) Strauss, S. H. *Chem. Rev.* 1993, 93, 927–942.
(6) Douvris, C.; Ozerov, O. V. *Science* 2008, 321, 1188–1190.
(7) Qin, Y.; Bakker, E. *Anal. Chem.* 2003, 75, 6002–6010.
(8) Schwab, P. F. H.; Levin, M. D.; Michl, J. *Chem. Rev.* 1999, 99, 1863–1933.
(9) Franken, A.; Kilner, C. A.; Kennedy, J. D. S. *Inorg. Chem. Commun.* 2003, 6, 1104–1108.

(10) Jelinek, T.; Baldwin, P.; Scheidt, W. R.; Reed, C. A. *Inorg. Chem.* 1993, 32, 1982–1990.
(11) King, B. T.; Koerber, S.; Schreiber, P. J.; Clayton, J.; Nemcova, A.; Havlas, Z.; Vyakaranam, K.; Fete, M. G.; Zharov, I.; Ceremuga, J.; Michl, J. *J. Am. Chem. Soc.* 2007, 129, 12960–12980.
(12) Ivanov, S. V.; Rockwell, J. J.; Polyakov, O. G.; Gaudinski, C. M.; Anderson, O. P.; Solntsev, K. A.; Strauss, S. H. *J. Am. Chem. Soc.* 1998, 120, 4224–4225.
(13) Xie, Z.; Tsang, C.-W.; Sze, E. T.-P.; Yang, Q.; Chan, D. T. W.; Mak, T. C. W. *Inorg. Chem.* 1998, 37, 6444–6451.
(14) Kueppers, T.; Bernhardt, E.; Eujen, R.; Willner, H.; Lehmann, C. *Angew. Chem.* 2007, 46, 6346–6349.
(15) Finze, M.; Sprenger, J.; Schaack, B. *Dalton Trans.* 2010, 39, 2708–2716.

which yielded a white precipitate, which was isolated on a fine frit, washed with water, and vacuum-dried (0.152 g, 92%). ^1H NMR (d_6 -acetone): δ 3.46 [s, 12H, Me₄N], 2.57 [d, 2H, methylene, J_{H} 7.2 Hz], 4.99 [m, 2H, vinyl], 5.63 [m, 1H, vinyl]. ^{11}B NMR: -2.29 [s, 1B, B(12)], -8.85 [s, 5B, B(7–11)], -16.97 [d, 5B, B(2–6), J_{BH} 522 Hz]. m/z calcd for $\text{C}_3\text{H}_5\text{-CB}_{11}\text{H}_5\text{Br}_6^-$: 656.6917. Found: 656.6913.

[Me₄N][*closo*-1-CH₂(C₆H₅)-CB₁₁H₅Br₆] was prepared from Li[1-Li-CB₁₁H₅Br₆] generated from [Me₃NH][CHB₁₁H₅Br₆] (0.1401 g, 0.207 mmol) and benzyl bromide (~0.25 mL) in a similar manner to [Me₄N][*closo*-1-C₃H₅-CB₁₁H₅Br₆] above (0.243 g, 90%). ^1H NMR (d_6 -acetone): δ 3.46 [s, 12H, Me₄N], 3.15 [s, 2H, methylene], 7.12 [m, 2H, phenyl], 7.27 [m, 3H, phenyl]. ^{11}B NMR: -2.29 [s, 1B, B(12)], -8.79 [s, 5B, B(7–11)], -16.84 [d, 5B, B(2–6), J_{BH} 501.8 Hz]. m/z calcd for $\text{C}_7\text{H}_7\text{-CB}_{11}\text{H}_5\text{Br}_6^-$: 706.7074. Found: 706.7061.

Li[*closo*-1-(*t*-butyl)₂P-CB₁₁H₅Br₆] was prepared from Li[1-Li-CB₁₁H₅Br₆] generated from [Me₃NH][CHB₁₁H₅Br₆] (0.2350 g, 0.347 mmol) and di-*t*-butylchlorophosphine (0.065 mL). The reaction was allowed to stir for 1 h at room temperature, followed by the removal of ca. 85% of the solvent. Only NMR data were collected due to the extreme sensitivity of the product to oxygen. ^1H NMR (d_3 -acetonitrile): δ 1.18 [d, 18 H, *t*-butyl, J_{PH} 32.2 Hz]. ^{31}P NMR: -151.40 [s, 1P]. ^{11}B NMR: -1.13 [s, 1B, B(12)], -8.69 [s, 5B, B(7–11)], -15.20 [d, 5B, B(2–6), J_{BH} 551.9 Hz].

[Me₄N][*closo*-1-C₆F₅-CB₁₁H₅Br₆] was prepared from Li[1-Li-CB₁₁H₅Br₆] generated from [Me₃NH][CHB₁₁H₅Br₆] (0.2708 g, 0.400 mmol) and C₆F₆ (~1 mL) in a similar manner to [Me₄N][*closo*-1-Me₃Si-CB₁₁H₅Br₆] above, except that the crude yellow product was extracted into diethylether (45 mL) for the initial filtration before dissolving in water (45 mL). (If a large excess of C₆F₆ is not used, multiple substitution occurs.) Yield of off-white solid: 0.323 g, 94%. ^{11}B NMR: $+0.049$ [s, 1B, B(12)], -8.55 [s, 5B, B(7–11)], -15.85 [d, 5B, B(2–6), J_{BH} 495.9 Hz]. ^{19}F NMR: -133.53 [broad, *o*-F], -154.70 [tt, *p*-F, J_{FF} 22.9, 5.1 Hz], -162.9531 [m, *m*-F]. m/z calcd for $\text{C}_6\text{F}_5\text{-CB}_{11}\text{H}_5\text{Br}_6^-$: 782.6446. Found: 782.6459.

[Me₄N][*closo*-1-C₆F₁₁-CB₁₁H₅Br₆] was prepared from Li[1-Li-CB₁₁H₅Br₆] generated from [Me₃NH][CHB₁₁H₅Br₆] (0.3051 g, 0.451 mmol) and perfluoro-1-hexene (0.150 g, 0.5 mmol) in a similar manner to [Me₄N][*closo*-1-Me₃Si-CB₁₁H₅Br₆] above except that the crude brown product was extracted into diethylether (45 mL) for the initial filtration before evaporation and dissolution in water (45 mL). An additional extraction of the final product into dichloromethane (10 mL) followed filtration through a fine frit, and evaporation of the solvent gave an oil that solidified to a brown solid (0.3682 g, 91.1%). ^{11}B NMR: $+0.049$ [s, 1B, B(12)], -8.55 [s, 5B, B(7–11)], -15.85 [d, 5B, B(2–6), J_{BH} 487.5 Hz]. ^{19}F NMR: -157.73 [d, alkene, J_{FF} 144.5 Hz], -146.70 [d, alkene, J_{FF} 137.37 Hz], -127.70 [m, CF₂], -118.75 [m, CF₂], -84.67 [dd,

CF₂, J_{FF} 31.7, 9.93 Hz], -80.73 [t, CF₃, J_{FF} 8.6 Hz]. m/z calcd for $\text{C}_6\text{F}_{11}\text{CB}_{11}\text{H}_5\text{Br}_6^-$: 896.6351. Found: 893.6390.

[Me₄N][*closo*-1-C₃F₅-CB₁₁H₅Br₆]. **Caution!** *Extreme care should be exercised when confining hexafluoropropene (B.Pt. -28°C) to glass Schlenkware. All reactions must be maintained at dry ice temperatures to avoid explosion.* In heavy walled Schlenkware, hexafluoropropene (~1 mL) was precondensed at -78°C and then transferred to a heavy walled reaction vessel at dry ice temperature containing THF (5 mL) and Li[*closo*-1-Li-CB₁₁H₅Br₆] generated from [Me₃NH][CHB₁₁H₅Br₆] (0.4510 g, 0.666 mmol). The resulting yellow solution was allowed to stir for 3 h before gradually removing the excess hexafluoropropene under a vacuum. The reaction was then allowed to warm to room temperature and the solvent removed under reduced pressure to give a colorless residue. This was extracted into diethylether (50 mL) and filtered through a fine frit and the filtrate evaporated to dryness. The crude product was dissolved in water (35 mL), and Me₄NCl (200 mg, 1.8 mmol) was added. The resulting white precipitate was isolated by filtration onto a fine frit, washed with water and vacuum-dried (0.4608 g, 92.7%). ^{11}B NMR: $+1.82$ [s, 1B, B(12)], -8.60 [s, 5B, B(7–11)], -17.52 [d, 5B, B(2–6), J_{BH} 504.2 Hz]. ^{19}F NMR: -67.56 [dd, CF₂(sp³), J_{FF} 23.5, 10.7 Hz], -122.60 [dm, CF, J_{FF} 146.1, 23.7 Hz], -160.60 [d, CF₂(sp²), J_{FF} 145.3 Hz]. m/z calcd for $\text{C}_3\text{F}_5\text{CB}_{11}\text{H}_5\text{Br}_6^-$: 746.6446. Found: 746.6458.

Cs[*closo*-1-Merrifield Peptide Resin-CB₁₁H₅Br₆]. To a THF solution [Li][1-Li-CB₁₁H₅Br₆] generated from [Me₃NH][CHB₁₁H₅Br₆] (0.2330 g, 0.344 mmol) was added Merrifield's Peptide Resin (0.185 g, 1.95 mmol Cl/g, 1.05 equiv). The reaction was allowed to stir for 1 week, yielding a yellow powder which was filtered onto a medium frit and was washed thoroughly with three aliquots of THF (50 mL). All filtrates were collected and the solvent evaporated. To the residue was added deionized water (20 mL). The ^{11}B NMR of this solution showed no detectable signals. To the filtrate were added two drops of HNO₃, followed by AgNO₃ (1 mL, 0.5 M). The resulting white precipitate of AgCl was filtered onto a fine frit and dried (0.048 g, 0.99 equiv). The resin was suspended in water and CsCl (1 g, 6 mmol) added. The suspension was stirred for 1 day before collecting the resin by filtration onto a medium frit and washing with water. The faintly pale product was oven-dried for 2 h at 90°C before obtaining an IR spectrum.

Acknowledgment. We thank the National Science Foundation (CHE-0841428 and 0742001) for support. M.J.N. was a MARCU* trainee supported by award T34GM062756 from the National Institute of General Medical Sciences.

Supporting Information Available: General experimental conditions, NMR and mass spectra for all new compounds (18 pages). This material is available free of charge via the Internet at <http://pubs.acs.org>.