

UC Riverside

UC Riverside Previously Published Works

Title

Presurgical Biomarker Performance in the Detection of Gleason Upgrading in Prostate Cancer

Permalink

<https://escholarship.org/uc/item/48k7g5gk>

Journal

Cancer Epidemiology Biomarkers & Prevention, 25(12)

ISSN

1055-9965

Authors

Wittig, Kristina
Yamzon, Johnathan L
Smith, David D
[et al.](#)

Publication Date

2016-12-01

DOI

10.1158/1055-9965.epi-16-0488

Peer reviewed



HHS Public Access

Author manuscript

Cancer Epidemiol Biomarkers Prev. Author manuscript; available in PMC 2017 December 01.

Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2016 December ; 25(12): 1643–1645. doi:
10.1158/1055-9965.EPI-16-0488.

Pre-Surgical Biomarker Performance in the Detection of Gleason Upgrading in Prostate Cancer

Kristina Wittig[§], Johnathan L. Yamzon[§], David D. Smith[#], Daniel R. Jeske^{#,†}, and Steven S. Smith^{*,§,¶}

[§]Division of Urology, Department of Surgery, City of Hope, Duarte, CA

[#]Division of Biostatistics, Department of Information Sciences, City of Hope, Duarte CA

[¶]Beckman Research Institute of the City of Hope, Duarte CA

[†]Department of Statistics, University of California – Riverside, Riverside CA

Abstract

Background—Gleason Score (GS) upgrading is generally considered a trigger for exit to definitive treatment during active surveillance (AS). Predicting the potential for GS upgrading would be of value in assessing AS eligibility.

Methods—We assessed the performance of biomarkers in pre-surgical specimens of expressed prostatic secretion (EPS) in this setting.

Results—Although EPS volume, total recovered RNA, and RNA expression biomarkers (TMPRSS2: ERG, PCA3, PSA) have been successful in both biopsy outcome prediction, and in the prediction of upstaging in active surveillance eligible patients, they were unable to predict upgrading in patients eligible for active surveillance under NCCN guidelines.

Conclusions—These biomarkers do not improve the prediction of upgrading over indications from standard clinical parameters.

Impact—Additional biomarkers will be needed in this area.

Keywords

Prostate Cancer; Gleason Score; Upgrading; Active Surveillance; Biomarkers

Introduction

Current practice during AS is to monitor low risk patients by repeating biopsies periodically. Patients with an initial biopsy GS of 6 are offered definitive treatment if a repeat biopsy during surveillance yields a GS ≥ 7 . This suggests that the cancer may have progressed

Correspondence to: Kristina Wittig.

***Corresponding Author:** Steven S. Smith, Ph.D., Division of Urology, Department of Surgery, Beckman Research Institute of the City of Hope, Fox North, Rm 1100A, City of Hope, 1450 E. Duarte Rd., Duarte, CA 91010, Office: 626-301-8316, Fax: 626-256-8774, ssmith@coh.org.

Conflict of Interest Disclosure: The authors declare that they have no potential conflicts of interest.

during the interval from the previous biopsy. Even so, a sizable proportion of GS 7 patients may have been misclassified GS 6 due to sampling error on low volume tumors (1, 2). For this reason, a non-invasive test that could detect this relatively subtle change in Gleason sum would be of value.

Results with post-DRE urine (3, 4), have consistently mirrored those with (EPS) (5, 6), suggesting that either specimen can be used in the detection and classification of prostate cancer. In previous work, improvements in AUC values of 0.1 (3-5) have been obtained for the prediction that cancer will be found at biopsy using these specimens. In this study, we sought to assess the performance of biomarkers in predicting Gleason upgrading in cohorts of men who underwent radical prostatectomy (RP). Our sample sizes were chosen for a statistical power of greater than 90% for an AUC improvement over baseline of 0.1.

Materials and Methods

EPS specimens from were obtained by milking the urethra after an attentive DRE before the patients underwent a Robot-Assisted Radical Prostatectomy (RARP). Out of 646 men who consented under an Institutional Review Board-approved protocol for the collection and evaluation of biomarkers in EPS, 642 had complete clinical data, 224 met the 2012 NCCN criteria and 206 met the 2014 NCCN criteria for active surveillance based on their preoperative characteristics.

EPS specimen processing was as described previously (7). Standard clinico-pathologic data (age, Gleason sum, ethnicity, pre-surgical PSA, T stage, DRE and race.), direct measurements of EPS volume, total recovered RNA, and quantitative reverse transcription polymerase chain reaction values for TMPRSS2:ERG, PCA3, and PSA RNAs were obtained.

Receiver Operator Characteristic (ROC) analysis of biomarker performance was carried out in an attempt to predict upgrading. Statistical power for AUC differences attributable to biomarkers was estimated with PASS software. Minimal models for each cohort emerged from stepwise logistic regression analysis on six clinical and ten laboratory variables (Table 1).

Results

For the 2012 NCCN AS cohort EPS volume, Total RNA and RNA expression biomarkers did not significantly increase the AUC value beyond the 0.6969 AUC achieved by clinical variables for ethnicity and serum PSA in the prediction of upgrading post surgery (Figure 1A). For the 2014 NCCN AS cohort, these same biomarkers failed to significantly enhance the AUC value beyond the 0.7222 value achieved with age, ethnicity and serum PSA (Figure 1B). For the 2012 cohort with complete clinical and laboratory data and sample sizes of (125 not upgraded and 99 upgraded), the power to detect an increase in the AUC of .05 and .10 was approximately 40% and 90%, respectively. For the 2014 cohort with 206 patients with complete clinical and laboratory data, the sample sizes (112 not upgraded and 94 upgraded), the power to detect an increase in the AUC of .05 and .10 is also approximately 40% and 90%, respectively.

Discussion

For NCCN AS cohorts, the AUC values achieved with standard clinical markers remained unchanged by the inclusion of biomarkers that have been successful in the prediction of biopsy outcome and upstaging. Although an increase in AUC value of 0.05 due to these biomarkers cannot be excluded, it is unlikely that such an increase would warrant the clinical use of these biomarkers in counseling patients in this setting in spite of their success in other areas of treatment. Moreover, the general concordance between results obtained with EPS specimens and those obtained with urine samples obtained after an attentive DRE argues that the results obtained with EPS will mirror those obtained with post DRE urine. We conclude that in this cohort of men, models for predicting upgrading are dominated by the standard clinicopathologic parameters with no significant contribution from current biomarkers of RNA expression or prostate function tested in EPS.

Acknowledgments

Financial Support: NIH NCI Grants CA 102521 and CA 136055 to Steven Smith.

References

1. Chism DB, Hanlon AL, Troncoso P, Al-Saleem T, Horwitz EM, Pollack A. The Gleason score shift: score four and seven years ago. *International journal of radiation oncology, biology, physics.* 2003; 56:1241–7.
2. Corcoran NM, Hovens CM, Hong MK, Pedersen J, Casey RG, Connolly S, et al. Underestimation of Gleason score at prostate biopsy reflects sampling error in lower volume tumours. *BJU Int.* 2012; 109:660–4. [PubMed: 21895937]
3. Laxman B, Morris DS, Yu J, Siddiqui J, Cao J, Mehra R, et al. A first-generation multiplex biomarker analysis of urine for the early detection of prostate cancer. *Cancer Res.* 2008; 68:645–9. [PubMed: 18245462]
4. Tomlins SA, Aubin SM, Siddiqui J, Lonigro RJ, Sefton-Miller L, Miick S, et al. Urine TMPRSS2:ERG fusion transcript stratifies prostate cancer risk in men with elevated serum PSA. *Sci Transl Med.* 2011; 3:94ra72.
5. Clark JP, Munson KW, Gu JW, Lamparska-Kupsik K, Chan KG, Yoshida JS, et al. Performance of a Single Assay for both Type III-Type VI TMPRSS2:ERG fusions in Non-Invasive Prediction of Prostate Biopsy Outcome. *Clinical Chemistry.* 2008; 54:2007–17. [PubMed: 18948370]
6. Whelan C, Kawachi M, Smith DD, Linehan J, Babilonia G, Mejia R, et al. Expressed Prostatic Secretion Biomarkers Improve Stratification of National Comprehensive Cancer Network Active Surveillance Candidates: Performance of Secretion Capacity and TMPRSS2:ERG Models. *The Journal of urology.* 2013
7. Clark JP, Munson KW, Gu JW, Lamparska-Kupsik K, Chan KG, Yoshida JS, et al. Performance of a single assay for both type III and type VI TMPRSS2:ERG fusions in noninvasive prediction of prostate biopsy outcome. *Clin Chem.* 2008; 54:2007–17. [PubMed: 18948370]




Figure 1.

ROC for Multivariate Analysis of Biomarkers in the prediction of upgrading **A.** NCCN 2012 Cohort of 225 Patients. 2012 NCCN guidelines permit low and very low risk patients with a Gleason sum (GS) ≤ 6 , a clinical diagnosis of T1-T2a, and a serum PSA value $<10\text{ng/ml}$ to enter active surveillance. Patients who have less than 10 years life expectancy are permitted to enter active surveillance if they exhibit only one of the following risk factors: GS 7, or T2b-T2c or PSA 10-20. Baseline GS, serum PSA and Ethnicity gave an AUC value of 0.6969 with or without EPS volume, total recovered RNA, and quantitative reverse transcription polymerase chain reaction values for RNA expression levels from TMPRSS2:ERG, PCA3 and PSA. Patient demographics: Ethnicity: Hispanic, 6.3%; Non-Hispanic, 92%; Unknown, 1.8%. Race: African American, 3.6%; Asian American, 6.3%; Caucasian American, 87.5%; American Indian, 0.4%; Other, 2.2%.

B. NCCN 2014 Cohort of 206 Patients. NCCN guidelines issued in 2014 permit only low and very low risk patients with a GS ≤ 6 , a clinical diagnosis of T1-T2a, and a serum PSA value $<10\text{ng/ml}$ to enter active surveillance. Baseline GS, serum PSA, age and Ethnicity gave an AUC value of 0.7222 with or without EPS volume, total recovered RNA, and quantitative reverse transcription polymerase chain reaction values for RNA expression levels from TMPRSS2:ERG, PCA3 and PSA. Patient demographics: Ethnicity: Hispanic, 6.3%; Non-Hispanic, 91.7%; Unknown, 1.9%. Race: African American, 3.9%; Asian American, 5.8%; Caucasian American, 87.8%; American Indian, 0.5%; Other, 1.9%.

Table 1

Logistic regression model p-values for prognostic factors for Gleason upgrading. Univariate (UV) p-values are from single variable models. Multivariate (MV) p-values are from the final model selected from a stepwise selection procedure.

Variable	NCCN Cohorts*			
	2012 (N=224, Yes=99, No=125)		2014 (N=206, Yes=94, No=112)	
	UV	MV	UV	MV
Pretreatment Gleason Score				
Ethnicity	.0598	.0460*	.0702	.0395*
Race	.2404		.4513	
Age	.1590		.0080*	.0066*
DRE	.1026		.1462	
T Stage	.0860		.0934	
Prebiopsy Serum PSA	.0001**	<.0001**	.0002**	<.0001**
PCA3 RNA	.6811		.7230	
PCA3 RNA/ Input RNA	.6914		.7215	
PCA3 RNA/PSA mRNA	.1946		.1906	
TMPRSS2:ERG	.5614		.5773	
TMPRSS2:ERG/Input RNA	.5888		.5956	
TMPRSS2:ERG/PSA mRNA	.5188		.5385	
PSA_mRNA	.2326		.2563	
PSA mRNA/Input RNA	.2356		.2320	
Expressed Volume Prostatic Fluid (μL)	.7702		.5221	
Total RNA in specimen (ng)	.5982		.5917	