

UC Irvine

UC Irvine Previously Published Works

Title

Hippocampal gene expression patterns linked to late-life physical activity oppose age and AD-related transcriptional decline

Permalink

<https://escholarship.org/uc/item/492258c5>

Authors

Berchtold, Nicole C

Prieto, G Aleph

Phelan, Michael

et al.

Publication Date

2019-06-01

DOI

10.1016/j.neurobiolaging.2019.02.012

Peer reviewed



Published in final edited form as:

Neurobiol Aging. 2019 June ; 78: 142–154. doi:10.1016/j.neurobiolaging.2019.02.012.

Hippocampal gene expression patterns linked to late-life physical activity oppose age and AD-related transcriptional decline.

Nicole C. Berchtold, Ph.D^{1,*}, G. Aleph Prieto, Ph.D¹, Michael Phelan, Ph.D², Daniel L. Gillen, Ph.D³, Pierre Baldi⁴, David A. Bennett, M.D.⁵, Aron S. Buchman, M.D.⁵, Carl W. Cotman, Ph.D^{1,6}

¹Institute for Memory Impairments and Neurological Disorders, University of California-Irvine, Irvine, CA, 92697-4540

²Dept of Statistics, University of California, Irvine, CA, 92697-1250

³Statistics Dept., University of California, Irvine, CA 92697-1250

⁴Institute for Genomics and Bioinformatics, School of Information and Computer Sciences, University of California, Irvine, CA 92697-3435

⁵Department of Neurological Sciences, Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL 60612

⁶Department of Neurology, University of California-Irvine, Irvine, CA

Abstract

Exercise has emerged as a powerful variable that can improve cognitive function and delay age-associated cognitive decline and Alzheimer's disease (AD), however, the underlying mechanisms are poorly understood. To determine if protective mechanisms may occur at the transcriptional level, we used microarrays to investigate the relationship between physical activity levels and gene expression patterns in the cognitively-intact aged human hippocampus. In parallel, hippocampal gene expression patterns associated with Aging and AD were assessed using publicly available microarray data profiling hippocampus from young (20–59 yrs), cognitively-intact aging (73–95 yrs) and age-matched AD cases. To identify “anti-Aging/AD” transcription patterns associated with physical activity, probesets significantly associated with both physical activity and Aging/AD were identified and their directions of expression change in each condition were compared. Remarkably, of the 2210 probesets significant in both datasets, nearly 95% showed opposite transcription patterns with physical activity compared to Aging/AD. The majority (>70%) of these anti-Aging/AD genes showed increased expression with physical activity and decreased expression in Aging/AD. Enrichment analysis of the anti-Aging/AD genes showing increased

*CORRESPONDING AUTHOR: Nicole C. Berchtold 1206 Gillespie Neuroscience Facility Institute for Memory Impairments and Neurological Disorders, University of California-Irvine Irvine, CA, 92697-4540 Tel: 949 824-6071 Fax: 949 824-2071 nbercto@uci.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

expression in association with physical activity revealed strong overrepresentation of mitochondrial energy production and synaptic function, along with axonal function and myelin integrity. Synaptic genes were notably enriched for synaptic vesicle priming, release and recycling, glutamate and GABA signaling and spine plasticity. Anti-Aging/AD genes showing decreased expression in association with physical activity were enriched for transcription-related function. These data reveal that physical activity is associated with a more youthful profile in the hippocampus across multiple biological processes, providing a potential molecular foundation for how physical activity can delay age- and AD-related decline of hippocampal function.

Keywords

exercise; microarray; plasticity; synaptic vesicle trafficking; mitochondria; white matter; myelin; axon

1. INTRODUCTION

Lifestyle factors have a striking ability to empower the adult brain - strengthening plasticity, building cognitive reserve, and preventing functional decline. Physical activity in particular has emerged as a powerful life-style factor that can improve cognitive function (Buchman et al., 2012; Buchman et al., 2008; Colcombe and Kramer, 2003; Colcombe et al., 2003; Erickson et al., 2010; Erickson et al., 2011; Kramer and Erickson, 2007a, b; Smith et al., 2010; Stubbs et al., 2017; Suwabe et al., 2018), reduce the risk of age-associated cognitive decline and AD (Buchman et al., 2012; Erickson et al., 2012; Hillman et al., 2008; Kirk-Sanchez and McGough, 2014; Korol et al., 2013; Prakash et al., 2015; Weuve et al., 2004; Yaffe et al., 2001), counteract age- and AD-related losses of grey and white matter (Baker et al., 2010; Best et al., 2017; Bherer et al., 2013; Colcombe et al., 2003; Erickson et al., 2014; Lautenschlager et al., 2008; Okonkwo et al., 2014; Suzuki et al., 2012; Voss et al., 2013; Zlatar et al., 2015) and counteract age- and AD-related declines in cognitive neural network function (Colcombe et al., 2004; Huang et al., 2016; Voss et al., 2010). While the benefits of physical activity are evident, the underlying mechanisms by which physical activity supports brain function and health and slows Aging- and AD-related declines in the human brain are poorly understood.

Gene expression studies to address this question have been precluded by a paucity of human autopsy cases that are well-defined for late-life lifestyle factors, and in particular, where physical activity has been objectively documented in a manner that avoids recall bias. Participants in the Rush Memory and Aging Project (MAP) meet these criteria. MAP participants have been followed longitudinally for up to 18 years, until death, with detailed annual assessments of cognitive function and multiple lifestyle variables including quantitative assessment of physical activity (Bennett et al., 2012). Importantly, physical activity was continuously monitored over multiple days with actigraphy, capturing both exercise and non-exercise activity (e.g., light physical activity) to provide a highly accurate account of the total daily activity levels (Buchman et al., 2012).

In this study, we use post-mortem hippocampal tissue from cognitively-healthy, aged (76–100 years) MAP cases that captured a broad spectrum of physical activity level in

combination with a microarray-based genome-wide approach to investigate mechanisms by which physical activity may promote cognitive health and hippocampal function, and counteract Aging- and AD-related transcription patterns in the human brain. The data suggest that gene expression patterns associated with physical activity have the potential to extensively counteract transcriptional changes that occur in the human hippocampus with Aging or AD. In particular, physical activity may slow Aging/AD-related declines in hippocampal health and function by enhancing mitochondrial energy production and synaptic function, improving axonal function and myelin integrity, and maintaining appropriate transcriptional control mechanisms.

2. MATERIALS AND METHODS

Samples

To investigate the relationship between physical activity and gene expression patterns in the cognitively intact aged human hippocampus, frozen post-mortem hippocampal brain tissue was obtained from 47 clinically well-characterized cognitively intact cases (aged 76–100 years) from MAP, an ongoing longitudinal, study of older individuals without known dementia recruited from northeastern Illinois. All tissue was obtained from the mid-hippocampus region containing all hippocampal subfields. Participants were diagnosed as cognitively-intact based on annual assessment of cognitive status using a battery of 19 cognitive tests, at which time they were additionally assessed for other lifestyle factors including cognitive frequency, social frequency and depression, among other variables (Bennett et al., 2012). Participants underwent annual assessment of physical activity using actigraphs worn continuously over multiple days. Post-mortem tissue was assessed for pathology as previously reported (Bennett et al., 2012), and cases with Lewy Bodies, infarcts, or hippocampal sclerosis were excluded from the study. Hippocampal RNA from 35 participants (76–100 yrs, average 87.62 ± 6.36 yrs) was of high quality sufficient to process on microarrays and an additional 12 samples were included in the Nanostring analysis.

Quantification of Physical activity

Physical activity was measured with actigraphy (Actical: Mini Mitter, Bend OR). Total daily exercise and non exercise physical activity was measured 24 hours per day for up to 10 days, with actigraphs worn on the non-dominant wrist. Average daily physical activity was calculated as the average across days of the total daily activity counts recorded. Average physical activity ranged from 0.13 to 5.6×10^5 activity counts/day (Supplemental Figure 1). Data were converted to approximate metabolic equivalents (METs) using the distance conversion metrics reported by Hall et al. (Hall et al., 2013) and standard MET equivalent values (Ainsworth et al., 1993). Using these metrics, 1.0×10^5 counts/day corresponded to 20 min/day moderate-high activity (walking, 3.5 mph) (equivalent to 1.5 MET-hrs of activity/day), and 5.0×10^5 counts/day was the equivalent of 100 min/day of moderate-high activity (7.5 MET-hrs/day). Previously, Weuve et al. (Weuve et al., 2004), described a dose-dependent effect of physical activity on cognitive performance, with highest cognitive performance in individuals with activity estimates above 26 MET-hrs/week.

RNA extraction and Microarray quality control

RNA from approximately 25 mg tissue (mid-level hippocampus containing all hippocampal subregions) was extracted using Trizol, treated with DNase, and purified using Qiagen spin columns as described previously (Berchtold et al., 2008). RNA quality was assessed using the Agilent Bioanalyzer and RNA integrity number (RIN), and samples with RIN ≥ 6.5 were further processed on microarrays. CEL file quality control was assessed using the affyQC automated R workflow available through ArrayAnalysis (<http://www.arrayanalysis.org>). Quality control parameters were: beta-actin 3'/5' ratio < 3 , GAPDH 3'/5' ratio < 1.25 , internal hybridization control signal (BioB=Present), Percent Present $\geq 49\%$, Percent Present spread $\leq 10\%$, Background spread $\leq 20\%$ and log scale factor spread ≤ 3 .

Actual samples:

Of the 47 hippocampal tissues, high quality RNA was obtained from 35 samples (RIN: 7.36 ± 0.58 , range 6.5–8.7) which were processed individually on Affymetrix HgU133plus 2.0 microarrays by the Genomic high-Throughput Facility at UC Irvine, following manufacturer's recommendations. CEL file quality control was assessed using the affyQC workflow in ArrayAnalysis (<http://www.arrayanalysis.org>). One microarray did not meet all quality control parameters in ArrayAnalysis and was excluded. Sample characteristics and quality control measures are provided in Supplemental Tables 1A and 1D. The remaining 34 CEL files were submitted to GeneSpring software GS version 14.5 (GS14.5) (Agilent Technologies) for preprocessing by MAS and GC-RMA. MAS and GC-RMA values were quantile normalized and underwent baseline transformation to the median of all samples across all genes as implemented in GS14.5. The CEL files and microarray data are available through the public gene expression omnibus (GEO) (<http://www.ncbi.nlm.nih.gov/geo>), dataset accession number GSE110298.

Young, Aged, and AD samples:

To characterize gene expression changes associated with Aging and AD, 58 CEL files profiling gene expression patterns in hippocampus from young (20–59 yrs), cognitively intact aging cases (69–99 yrs), and AD cases (73–99 yrs) were downloaded from the public gene expression omnibus (<http://www.ncbi.nlm.nih.gov/geo>) dataset, accession numbers GSE11882 and GSE48350. CEL file quality control was assessed using the affyQC workflow in ArrayAnalysis (<http://www.arrayanalysis.org>) and identified 48 microarrays that met all quality control and study parameters (young (n=16), cognitively intact aging cases (n=18), and AD cases (n=14)). Sample characteristics and quality control measures are provided in Supplemental Tables 1B and 1E.

Microarray data analysis – identifying genes associated with physical activity

Poorly performing probesets were eliminated by filtering the array data based on flags (Present, Marginal, Absent) using MAS preprocessing, retaining only probesets flagged Present on all samples of at least one physical activity group (low, moderate, high activity: Supplemental Figure 1) (23,712 probesets) and that overlapped with the probeset list used for the analysis of Aging and AD (21,002 probesets). Using the programming environment R (<http://www.R-project.org/>), multiple linear regression was applied to GC-RMA

normalized values using $\text{lm}()$ to adjust for technical factors that were potential confounding variables (batch, RIN, and PMI) and to evaluate the relationship between gene expression and late-life physical activity levels across the 34 samples (Miller et al., 2017). For each probeset, the estimated regression coefficient for physical activity, its standard error, and the t-ratio and P-value for a two-sided test were extracted using $\text{infer}()$ with robust standard errors. Significance was set at $p = 0.05$ and q values were calculated with Graph Pad Prism 7 statistical software using the two-stage step-up method of Benjamini, Krieger and Yekutiessli. To estimate fold changes across low, moderate and high physical activity tiers, multiple linear regression was applied to each 2 group comparison, and p and q values were calculated. Log values were transformed to linear scale using Excel software. All calculated values are provided in Supplemental Table 2.

Microarray analysis – Identifying genes associated with Aging and AD

The 54,675 probesets on the array were filtered using MAS preprocessing derived flags to retain only probesets flagged Present on all samples of at least one treatment group (21,002 probesets) and that overlapped with the probeset list used for the analysis of gene expression associated with physical activity (21,002). Using R, multiple linear regression was applied to GC-RMA normalized values to adjust for batch, RIN, and PMI and to evaluate gene expression changes in Aging (Aged vs. young) and AD (AD vs. Aged). For each probeset, the estimated regression coefficient for physical activity, its standard error, and the t-ratio and P-value for a two-sided test were extracted using $\text{infer}()$ with robust standard errors. Significance was set at $p = 0.05$ and q values were calculated using Graph Pad Prism 7 statistical software. Log values were transformed to linear scale using Excel software. All calculated values are provided in Supplemental Table 2

Anti-Aging/AD patterns associated with exercise

To investigate mechanisms by which physical activity may protect the brain from Aging and AD-related change, probesets significantly associated with both physical activity and Aging or AD were selected, patterns of expression change were compared for each probeset, and probesets showing opposite patterns of change with physical activity versus Aging or AD were identified (henceforth referred to as “anti-Aging/AD probesets”) (See Methods in Figure 1). Significance of the overlap patterns between physical activity and Aging-AD (eg, opposite direction of gene expression change, or similar direction) was assessed using the Fisher exact test. 2 approaches were used to estimate the number of probesets expected to occur by chance in the comparisons (a) physical activity and aging or (b) physical activity and AD. The first approach assumes normality of the p value distribution and independence of the comparisons, and was calculated as $(2 \times (0.05^2) - (0.05)^3) \times 21,002$ probesets tested. This approach estimated the number of putative false positives to be 102 probesets (e.g., $p = 0.004875 \times 21,002$), corresponding to a false positive rate (FDR) estimate of 4.6% (e.g., 102 false discoveries/2210 true discoveries). Permutation analysis was used as a second approach where normality of the p value distribution is not assumed. Using the R function “ $\text{transform}()$ ” activity labels were randomized across samples in the physical activity dataset, and across the Young, Aged, and AD samples for the Aging-AD analysis, and the number of overlapping significant probes was calculated for the permuted data. 100 permutation analyses were performed, resulting in estimations ranging from 10 – 1,057 false

positives, with an average \pm standard deviation estimate of 176 ± 192 and a median estimate of 109 false positives. The computed average false positives (176) corresponds to a false positive rate of 7.9% and the median false positive estimate (109) corresponds to a false positive rate of 4.9%, similar to the FDR estimate (4.6%) calculated under the assumption of normal distribution of p values.

DAVID Functional enrichment analysis

Functional enrichment analysis of the Anti-Aging/AD genes was used to identify biological modules of Aging/AD that show the opposite pattern of change with physical activity. Significant gene lists were submitted to the high-throughput data mining bioinformatics resources DAVID (the Database for Annotation, Visualization and Integrated Discovery, <https://david.ncifcrf.gov>) (Dennis et al., 2003; Huang da et al., 2009a, b), to identify over-represented biological functions and pathways. Significant probesets were translated to DAVID ids followed by mapping of unambiguous DAVID ids to terms in the Gene Ontology (GO) databases (Cell Component (GOTERM-CC), Biological Process (GOTERM-BP), Molecular Function (GOTERM-MF), KEGG and BioCarta pathways, or the UniProt keywords database. Enrichment was assessed against a background gene set of the 21,002 probesets with reliable expression on both datasets, EASE score threshold $p = 0.01$, and Benjamini multiple testing correction $p = 0.10$. Benjamini p-values are indicated in the results.

Nanostring nCounter™ gene expression system

Microarray expression changes associated with physical activity were validated using the NanoString nCounter™ gene expression system to assess gene expression on an expanded set of human hippocampal tissues, for experimental and independent replication of the microarray data. Experimental replication included 8 additional low activity cases and 4 additional high physical activity cases, in addition to low (n=10) and high activity (n=12) cases included in the microarray analysis (Supplemental Tables 1C, 1D). One low activity sample (sample 1_1-E) included in the microarray analysis was not included in the nanostring run due to sample degradation. NanoString provides similar sensitivity to TaqMan-based RQ-PCR and SYBR Green I fluorescent dye-based RQ-PCR but does not involve reverse transcription of mRNA and subsequent cDNA amplification, thus eliminating amplification bias (Geiss et al., 2008; Veldman-Jones et al., 2015). Gene expression was assessed in low activity (n= 18) and high activity (n=16) cases for 13 synaptic genes that showed decreased expression with Aging/AD and increased expression with physical activity (GABRG2, GLRB, GRIN2a, SLC1a6, SYN2, VAMP2, SNAP25, USP14, NRXN1, NRXN3, SYT4, SSTR1, Munc13–3), and for 10 genes where microarray analysis demonstrated decreased expression with Aging/AD and trends for increased expression with physical activity (GABBR1, GABRA2, GABRA4, GABRB3, GABRG1, GRIA2, DOC2a, SCN2b, DLG3/SAP102, AKAP5). 7 putative internal control genes (ATF6, DDX52, METTL20, GAPDH, SLC38a7, ZNF780b, ZNF791) were first identified based on microarray data demonstrating expression homogeneity across physical activity tier and low variance, and were evaluated using Nanostring for suitability. Putative control genes were excluded if Nanostring gene expression showed a trend ($p = 0.1$) for a significant correlation with physical activity across cases or a trend for a significant t-test comparing low vs. high

physical activity groups. ZNF780b and DDX52 were identified as suitable control genes. Nanostring expression values underwent sample normalization to the geometric mean of the control genes and Graph Pad Prism 7 statistical software was used to calculate means, standard error of the mean (sem) and 2 group comparisons of the low activity (n=18) and high activity (n=16) groups across the genes.

For independent validation of the microarray data, only the new Nanostring cases were included in the analysis (n=8 low activity, n=4 high activity). The Mann-Whitney test was used to determine whether the nanostring-measured genes, as a group, tended to move in the same direction as those genes did on the array. For each gene, the average nanostring value for the low and high activity groups were calculated, and the Mann-Whitney test was applied using Graph pad Prism 7 statistical software.

3. RESULTS

3.1 Physical activity is associated with anti-Aging/AD gene expression patterns in the hippocampus

Hippocampal tissue from 34 cognitively-intact aged cases well-characterized for physical activity levels were run on microarrays to investigate gene expression patterns significantly associated with physical activity levels. In parallel, microarray data profiling hippocampal gene signatures from young (20–59 yrs), cognitively intact aging (69–99 yrs) and AD (73–99) cases were analyzed to identify gene expression patterns associated with Aging and AD (gene expression omnibus (GEO) dataset accession number GSE11882, <http://www.ncbi.nlm.nih.gov/geo>). To investigate mechanisms by which physical activity may protect the brain from Aging and AD-related change, probesets significantly associated with both physical activity and Aging or AD were selected patterns of expression change were compared for each probeset, and probesets showing opposite patterns of change with physical activity versus Aging or AD were identified (henceforth referred to as “anti-Aging/AD probesets”) (Figure 1).

Of the 21,002 probesets with reliable expression on both datasets, the analysis identified 2210 probesets that were significantly associated with both physical activity and Aging/AD, a number far greater than expected to occur by chance (e.g. ~102 expected false positives/ 2210 significant probesets, FDR = 4.6%), with the significance of the overlap calculated to be $p < 0.0001$ (Fisher exact test). Remarkably, of these 2210 probesets, 94.7% (2,093 probesets) showed the opposite patterns of change in association with physical activity compared with Aging/AD (henceforth called “anti-Aging/AD probesets”). The majority (>70%) of these anti-Aging/AD probesets showed increased expression with physical activity (1,472 probesets), while ~30% (621 probesets) showed decreased expression in association with physical activity. These data suggest that gene expression patterns associated with physical activity extensively counteract gene expression changes that occur in the human hippocampus with Aging or AD, particularly with respect to those genes undergoing transcriptional decline.

3.2 Anti-Aging/AD genes are enriched for several core biological functions

To investigate potential repercussions of physical activity for hippocampal health and function, we investigated if the anti-Aging/AD probesets significantly associated with physical activity were enriched for biological functions using the integrated data-mining bioinformatics resource DAVID version 6.8 (Dennis et al., 2003; Huang da et al., 2009a, b). The anti-Aging/AD probeset lists (1472 with increased expression and 621 with decreased expression in association with physical activity) were uploaded and translated to DAVID ids followed by mapping of unambiguous DAVID ids to terms in the GO databases, KEGG pathways or UniProt keywords database. Benjamini p-values are indicated below.

For anti-Aging/AD probesets showing increased expression in association with physical activity, the probeset list translated into 1176 unique and unambiguous DAVID ids of which 1117 genes could be mapped to database terms. DAVID analysis revealed robust enrichment for several functional categories key to cellular health and synaptic function (Table 1). Highly significant enrichment was found for mitochondrial-related genes (e.g., UniProt keyword “Mitochondrion”, $p = 2 \times 10^{-5}$) especially for electron transport chain function, as well as for genes associated with the synapse (e.g., UniProt keyword “Synapse”, $p = 5 \times 10^{-7}$), myelin sheath (GOTERM “myelin sheath”, $p = 1 \times 10^{-4}$), and axon (GOTERM “axon”, $p = 3 \times 10^{-3}$). These data suggest that Aging/AD is accompanied by diminishing gene expression for these core biological functions, and that physical activity may slow Aging/AD-related declines in hippocampal health and function by enhancing mitochondrial energy production, synaptic health and axon function.

A different enrichment pattern was observed for anti-Aging/AD probesets showing decreased expression in association with physical activity. The 621 anti-Aging/AD translated into 408 unique and unambiguous DAVID ids of which 378 genes could be mapped to an enrichment term. Anti-Aging/AD genes showing decreased expression with physical activity were significantly enriched primarily for transcription-related function (e.g., UniProt keyword “Transcription”, $p = 7 \times 10^{-3}$), notably for negative regulation of transcription (e.g., GOTERM “negative regulation of transcription from RNA polymerase II promoter”, $p = 8.8 \times 10^{-2}$) (Table 2). These data suggest that Aging/AD is accompanied by increased expression of genes that suppress transcription, and that physical activity may restrain this transcriptional dysregulation to maintain appropriate transcriptional control of gene expression in the Aging and AD hippocampus.

3.2.1 Physical activity-associated gene expression patterns oppose Aging/AD-related declines in mitochondrial function and energy production—

DAVID analysis revealed that amongst anti-Aging/AD genes increased with physical activity, mitochondria-related genes were the most highly enriched class and represented 12–13% of the gene list, a 1.5-fold enrichment (GOTERM-CC, $p = 7.0 \times 10^{-6}$). These included pronounced enrichment for oxidative phosphorylation (KEGG pathway, 3.17-fold enrichment, $p = 5.5 \times 10^{-7}$) along with components of the mitochondrial inner membrane (GOTERM-CC: 1.93-fold enrichment, $p = 8.2 \times 10^{-6}$) and mitochondrial outer membrane (GOTERM-CC: 2.08-fold enrichment, $p = 9.6 \times 10^{-2}$). These mitochondrial-associated genes were next investigated in greater detail to determine which aspects of mitochondrial

function that decline with Aging/AD may be particularly amenable to improvement with physical activity. Genes in these enriched mitochondria-related categories have been compiled in Supplemental Table 3.

Approximately 30% of the mitochondrial anti-Aging/AD genes were associated with energy production via oxidative phosphorylation and the tricarboxylic acid cycle (TCA). Anti-Aging/AD genes were 3.4-fold enriched for cellular components of the mitochondrial respiratory chain complex I (GOTERM CC, $p = 4.8 \times 10^{-3}$) and 4.0-fold enriched for mitochondrial respiratory chain complex V (ATP synthase complex) (GOTERM CC, $p = 9.67 \times 10^{-2}$). Overlay of all genes significantly associated with physical activity onto the KEGG pathway “oxidative phosphorylation” (3.17-fold enriched, $p = 5.5 \times 10^{-7}$) reveals the extensive engagement of genes critical for energy production. Increased expression was found for approximately 30% of the genes encoding mitochondrial complex I - the entry point to the electron transport pathway - along with 35% of the genes encoding Complex IV (cytochrome C oxidase) and Complex V (ATP synthase), as well as 2 components (SLC25A27, SLC25A14) of the uncoupling protein (Figure 2). The broadly increased expression of oxidative phosphorylation genes was accompanied by increased expression of key genes driving pyruvate metabolism and the TCA cycle, which generates energy from pyruvate metabolism and provides NADH and FADH₂ reducing equivalents for the electron transport chain. These included dihydrolipoamide dehydrogenase (DLD), malate dehydrogenase 1 (MDH1), malate dehydrogenase 2 (MDH2), malic enzyme 1 (ME1), malic enzyme 3 (ME3), pyruvate dehydrogenase (PDHA1) and succinate CoA ligase (beta subunit, SUCLA2). In addition to the numerous genes involved in energy production, other notably mitochondrial anti-Aging/AD genes included increased expression in association with physical activity of 16 mitochondrial ribosomal structural genes and 8 genes that negatively regulate neuronal apoptosis (B-Raf proto-oncogene serine/threonine kinase (BRAF), DnaJ heat shock protein family Hsp40 member C5 (DNAJC5), PTEN induced putative kinase 1 (PINK1), oxidation resistance 1 (OXR1), peptidylprolyl cis/trans isomeras (PIN1), syntaxin binding protein 1 (STXB1), synucleins alpha and beta (SNCA, SNCB), among other genes (Supplemental Table 3).

Taken together, these data suggest that multiple components necessary for mitochondrial health and energy production undergo declining gene expression with Aging/AD, and that physical activity may slow Aging/AD-related declines in hippocampal function by counteracting these gene expression patterns to enhance mitochondrial health and capacity for energy production.

3.2.2 Physical activity-associated gene expression patterns oppose Aging/AD-related declines in synaptic gene expression—Genes related to synaptic and neuronal function constituted the second most significantly enriched class of genes amongst anti-Aging/AD genes increased with physical activity. For example, genes in the Uniprot Keyword category “Synapse” represented ~5% of the gene list (60 genes), a 2.28-fold enrichment ($p = 5.0 \times 10^{-7}$) (Table 1). Significant genes-related to synaptic function could be grouped into the following main functional categories: (1) synaptic vesicle priming, release and recycling; (2) Glutamate and GABA signaling and receptor trafficking; and (3) synapse formation, plasticity, and axon health. Significant genes in enriched terms related to

synapse (Uniprot Keyword, $p = 5.0 \times 10^{-7}$), synaptic vesicle membrane (GOTERM-CC, 3.9-fold enriched, $p = 4.9 \times 10^{-4}$), chemical synaptic transmission (GOTERM-BP 2.2-fold enriched, $p = 9.4 \times 10^{-2}$) and post-synaptic membrane (GOTERM-CC, 2.1-fold enriched, $p = 1.4 \times 10^{-2}$) are listed in Supplemental Table 4.

3.2.2.1 Synaptic vesicle priming, release, and recycling: Our data revealed that numerous anti-Aging/AD genes showing increased activity with physical activity were associated with synaptic vesicle priming, regulating the readily releasable and reserve pools of synaptic vesicles, and synaptic vesicle recycling and recovering (Table 1). Key enriched terms associated with these functions included synaptic vesicle membrane (GOTERM-CC, 3.9-fold enriched, $p = 4.9 \times 10^{-4}$) and chemical synaptic transmission (GOTERM-BP, $p = 9.4 \times 10^{-2}$).

Several key genes that regulate synaptic vesicle priming and docking at the presynaptic active zone were present in the anti-Aging/AD genes showing increased expression with physical activity. Synaptic vesicle priming and docking require the assembly of 3 essential SNARE proteins into a complex, followed by binding of the SNARE complex to Sec1/Munc18-like (SM) proteins and other chaperones to catalyze vesicle fusion to the presynaptic membrane. Our data revealed that anti-Aging/AD genes included 2 of the 3 core SNARE proteins (synaptosome associated protein 25 (SNAP25), synaptobrevin/VAMP2) that are essential for active zone priming and docking, as well as several SNARE chaperones that included syntaxin-binding proteins (STXBP1/Munc18-1, STXBP5), amyloid precursor protein (APP), the RIM-binding protein ERC2, alpha and beta synuclein (SNCA, SNCB) and rab-3 interacting molecules (RIMS2, RIMS3).

In parallel, many of the anti-Aging/AD genes increased with physical activity are key regulators of presynaptic vesicle availability. For example, among the anti-Aging/AD genes were several active zone scaffolding proteins that regulate the balance between the reserve and the readily releasable pools of neurotransmitter, including synapsin II (SYN2), cyclin-dependent kinase 5 (CDK5) and STXBP5. In addition, there were numerous genes that regulate endocytosis and recovery of synaptic vesicles following neurotransmitter release, highly important steps for maintaining presynaptic synaptic vesicle availability, particularly in the hippocampus. These genes included increased expression of dynamins (DNM1L, DNM3), which are fundamental to synaptic vesicle endocytosis and recycling, along with synaptotagmin1 (SYNJ1) and endophilin1 (SH3GL2), which work in concert and are required for ultrafast endocytosis and clathrin uncoating of endocytosed synaptic vesicles. Several additional anti-Aging/AD genes associated with synaptic vesicle endocytosis and recovery were the clathrin adaptor protein complex 2 (AP2M1), the Bloc-1 subunit pallidin (BLOC1S6), Syndapin/PACSIN1, amphiphysin (AMPH), DNAJC5, secretory carrier membrane protein 5 (SCAMP5), syntaxin 12, TorsinA (TOR1A) and multiple V-ATPase subunits for APT6V1 (subunits A, C1, D, E1, G2) (Figure 3).

Taken together, these data suggest that Aging/AD are accompanied by declining expression of multiple components of the synaptic vesicle release, trafficking, and recycling machinery, and that physical activity may counteract these gene expression patterns and maintain a more effective presynaptic machinery.

3.2.2.2 Glutamate and GABA signaling: neurotransmitter levels, receptors, and receptor trafficking: Further analysis of the anti-Aging/AD genes showing increased expression with physical activity revealed numerous genes that regulate postsynaptic signaling efficiency of glutamatergic and GABAergic neurotransmission. These included several genes that regulate glutamate availability (glutaminase (GLS), high affinity glutamate transporter SLC1A6), glutamate receptor levels (NMDA receptor subunit 2a (GRIN2a), kainate receptor 2 (GRIK2), AMPA receptor subunit 4 (GRIA4)) and auxiliary components that modulate glutamate receptor signaling kinetics (ferric chelate reductase 1 like (FRRS1L), neuropilin and tolloid like 1 and 2 (NETO1, NETO2), homer scaffolding protein 1 (HOMER1)). In addition, there were several genes that regulate glutamate receptor availability in the post-synaptic density (PSD), primarily genes that promote removal of AMPA GluR1 from the PSD (glutamate receptor interacting protein 1 (GRIP1), ATPase family AAA domain containing 1 (ATAD1), SYNJ1, VAMP2), but also genes that increase AMPA GluR1 anchoring (sortilin related VPS10 domain containing receptor 3 (SORCS3), leucine rich repeat transmembrane neuronal 4 (LRRTM4)), GRIN2A clustering in the PSD (rabphilin 3A: RPH3A) and GRIN2B localization in the PSD (lin-7 homologs A and B: LIN7A, LIN7B). At the same time, anti-Aging/AD genes included genes regulating GABAergic neurotransmission, including increased expression of GABA_A receptors (GABRA1, GABRG2) and gephyrin (GPHN) - the neuronal assembly protein that anchors GABA receptors to the inhibitory PSD – along with several neuromodulators that augment GABA tone (glycine receptor beta (GLRB), somatostatin receptor 1 (SSTR1) and cholecystokinin (CCK)).

These data indicate that physical activity increases expression of many genes that regulate postsynaptic signaling efficiency of glutamatergic and GABAergic neurotransmission. The constellation of gene expression changes with increased physical activity may function to enhance excitatory signaling and at the same time to dampen neuronal responsiveness within neural circuits.

3.2.2.3 synapse formation, plasticity, and axon health: In parallel, genes showing increased expression in association with physical activity and decreased expression with Age/AD included central organizing molecules which promote new synapse formation and maturation, molecular pathways key for activity-dependent plasticity, and genes supporting axonal function and white matter integrity.

Notable categories associated with synapse formation, maturation and plasticity showing enrichment included a 2.24-fold enrichment for genes associated with the ‘post-synaptic cell membrane’ ($p = 7.5 \times 10^{-3}$) (Supplemental Table 5) and 3.6 fold enrichment for the BioCarta pathway “transcription factor CREB and its extracellular signals” ($p = 6.89 \times 10^{-2}$). Genes regulating spine genesis and synapse maturation included presynaptic neurexins (NRXN1, NRX3) and calstentenin 3 (CLSTN3) a synaptogenic adhesion molecule which works in concert with neurexins, along with leucine-rich repeat molecules (LRRTM4, LRRC4C, LRRC7), cell adhesion molecule 2 and 3 (CADM2, CADM3) and other genes: Rho guanine nucleotide exchange factor 7 (ARHGEF7), myocyte enhancer factor 2C (MEF2C), inositol-trisphosphate 3-kinase A (ITPKA), caytaxin (ATCAY), ATPase plasma membrane Ca²⁺ transporting 2 (ATPB2), phosphatase and actin regulator 1 (PHACTR1),

teneurin transmembrane protein 2 (TENM2), diacylglycerol kinase (DGK), glycoprotein M6A (GPM6A), cell division cycle 42 (CDC42), neuralized E3 ubiquitin protein ligase 1 (NEURL1). In addition, several of the anti-Aging/AD genes increased in association with physical activity are key molecules that enable dynamic plasticity critical for memory formation. These included kinases that drive the signaling in the nucleus required for sustained long-term potentiation (LTP) such as the catalytic and regulatory subunits of protein kinase A (PRKACB, PRKAR1A, PRKAR1B, PRKAR2B) and protein kinase C (beta subunit; PRKCB), genes regulating signaling through the CREB transcription factor (e.g., Adenylate cyclase 1 (ADCY1), calcium/calmodulin-dependent protein kinase delta (CAMK2D), GNAS complex (GNAS), growth factor receptor bound (GRB2)), and genes that facilitate trafficking and signaling of brain-derived neurotrophic factor (BDNF), a key plasticity molecule that facilitates LTP induction and stability and that is essential for hippocampus-dependent learning (e.g., endophilin (SH3GL2), synaptotagmin 4 (SYT4), ubiquitin specific peptidase 14 (USP14)).

Interestingly, genes showing increased expression in association with physical activity and decreased expression with Age/AD were also enriched for genes supporting axonal connectivity and function and white matter integrity. There was strong enrichment for the term 'growth cone' (2.38-fold enriched, $p=1.19 \times 10^{-2}$) (Supplemental Table 6), an actin-supported extension of a developing or regenerating neurite seeking its synaptic target, and a 1.79 fold enrichment for "Neuron projection" ($p = 9.56 \times 10^{-2}$). It is possible that this gene expression pattern reflects axonal outgrowth from newly generated neurons in the neurogenic subgranular zone of the hippocampus, as increased expression was seen for several genes that regulate axonal outgrowth and targeting (e.g., RAC1-activated kinases (PAK1, PAK3, PAK5, PAK6), ephrins (EFNB3, EFNA5 and its receptor EPHA3), slit guidance ligand 2 (SLIT2), roundabout guidance receptor 2 (ROBO2), SLIT-ROBO Rho GTPase activating protein (SRGAP3)), along with a number of important genes that enhance proliferation, survival and migration of immature neurons (Supplemental Table 7) (e.g., doublecortin (DCX), cyclin-dependent kinase 5 (CDK5), glycogen synthase kinase 3 beta (GSK3b), platelet activating factor acetylhydrolase (PAFAH1B1), TMF1-regulated nuclear protein 1 (TRNP1), ephexin 1 (NGEF), and ROBO2). However, while stimulation of neurogenesis with physical activity is one of the most consistently documented effects of exercise in the animal hippocampus, there is controversy concerning the capacity for neurogenesis in the adult human hippocampus. It is possible that these expression patterns are not indicative of active neurogenesis but rather reflect promotion of axonal health and function. Indeed, anti-Aging/AD genes undergoing decreased expression in the Aging/AD hippocampus but increased expression in association with physical activity showed strong enrichment for genes associated with the axon (GOTERM-CC, 2.03-fold enrichment, $p=3.53 \times 10^{-3}$) (Supplemental Table 8) as well as the myelin sheath (GOTERM-CC, 2.45-fold enriched, $p = 1.0 \times 10^{-4}$) (Supplemental Table 9).

Taken together, these data suggest that late-life physical activity may counteract Age/AD-associated declines in synaptic plasticity by enhancing the molecular machinery for spine formation and activity-dependent plasticity, and promoting white matter integrity along with axonal connectivity and function.

3.2.3 Anti-Aging/AD genes decreased in association with physical activity

are enriched for transcriptional control—DAVID analysis of the 621 anti-Aging/AD probesets showing decreased expression with physical activity identified 408 unique and unambiguous DAVID ids of which 378 genes could be mapped to an enrichment term. Anti-Aging/AD genes showing decreased expression with physical activity were significantly enriched primarily for transcription-related function (Table 2). Significantly enriched terms notably included “Transcription regulation” (UniProt keyword, 1.5-fold enriched, $p = 8.45 \times 10^{-3}$), “Repressor” (UniProt keyword, 2-fold enriched, $p = 1.9 \times 10^{-2}$), “Activator” (UniProt keyword, 1.9-fold enriched, $p = 4.37 \times 10^{-2}$) and “negative regulation of transcription from RNA polymerase II promoter (GOTERM, 2.11-fold enriched, $p=8.8 \times 10^{-2}$) (Supplemental Table 10). Transcriptional control genes that were increased with Aging/AD but decreased in association with physical activity included, among others, several epigenetic regulators of histones (histone deacetylase 1 (HDAC1), lysine-specific histone demethylase 1B (KDM1B), lysine 9 histone 3 demethylase (KDM3A), histone lysine methyltransferase 2C (KMT2C)), components of the HDAC complex (Sin3A associated proteins 18 and 30 (SAP18, SAP30), MDS1 And EVI1 Complex Locus Protein MDS1 (MECOM), bHLH Transcription Factor 1 (TAL1), Transformer-2 Alpha (TRA2A), numerous zinc finger proteins (ZFX3, ZNF24, ZNF37A, ZNF160, ZNF274, ZNF397, ZNF532, ZNF652, ZNF721, ZBTB20, ZBTB47), along with the transcription factor SP3, which recruits HDAC2 to suppress transcription of synaptic-plasticity associated genes. Importantly, these genes are ones that show increased expression with Aging and/or AD but decreased in association with physical activity, suggesting that physical activity may offset transcriptional dysregulation and help maintain appropriate transcriptional control of gene expression in the Aging and AD hippocampus.

3.3 Replication: Nanostring technology

NanoString nCounter™ gene expression system was used to assess expression levels of an expanded set of human hippocampal tissues that included 12 additional cases to those used in the microarray analysis. Gene expression was compared in low ($n=18$) vs high activity ($n=16$) cases for 13 synaptic genes that showed decreased expression with Aging/AD and significantly increased expression with physical activity (GABRG2, GLRB, GRIN2a, SLC1a6, SYN2, VAMP2, SNAP25, USP14, NRXN1, NRXN3, SYT4, SSTR1, Munc13–3), and for 10 genes where microarray analysis demonstrated decreased expression with Aging/AD and trends that did not reach statistical significance for increased expression with physical activity (GABBR1, GABRA2, GABRA4, GABRB3, GABRG1, GRIA2, DOC2a, SCN2b, DLG3/SAP102, AKAP5).

Nanostring analysis using an expanded set of samples demonstrated increased gene expression with high physical activity for GABRG2 ($p=0.038$), GRIN2a ($p=0.03$), GLRB ($p<0.03$), SLC1a6 ($p=0.013$), SYN2 ($p=0.01$), USP14 ($p=0.008$), NRXN1 ($p=0.007$), NRXN3 ($p=0.05$) and SYT4 ($p=0.022$) SSTR1 ($p=0.026$), and UNC13c/Munc13–3 ($p=0.039$), confirming the microarray data, while expression for VAMP2 ($p=0.08$), and SNAP25 ($p>0.10$) failed to reach statistical significance. The nanostring analysis additionally demonstrated increased gene expression with high physical activity for GABBR1 ($p=0.002$), GABRA2 ($p=0.001$), GABRA4 ($p=0.013$), GABRB3 ($p=0.009$),

GABRG1 ($p=0.001$), GRIA2 ($p=0.04$), DOC2a ($p=0.03$), SCN2b ($p=0.05$), DLG3/SAP102 ($p=0.03$), AKAP5 ($p=0.019$) (Figure 4).

Finally, we analyzed the Nanostring data using only the subset of Nanostring cases that represented the 12 new cases not used in the microarray ($n=8$ low activity, $n=4$ high activity) as an independent confirmation of the microarray data. 100% of the genes showed directional agreement between the Nanostring and microarray data. Mann-Whitney comparison of average values across the genes revealed significantly higher expression in the high activity group vs. low activity group (medians: 0.894 (low), 1.12 (high); $p<0.0001$). The data thus confirm in an independent set of experimental samples that expression levels in the nanostring-measured genes, as a group, moved in the same direction as in the microarray analysis.

4.0 DISCUSSION

Multiple studies have established that physical activity is associated with delayed age-related decline and onset of AD in humans (Best et al., 2017; Buchman et al., 2012; Erickson et al., 2012; Hillman et al., 2008; Kirk-Sanchez and McGough, 2014; Korol et al., 2013; Prakash et al., 2015; Suwabe et al., 2018; Weuve et al., 2004; Yaffe et al., 2001). This study sought to identify underlying mechanisms by which physical activity supports brain health and slows Aging/AD-related declines in hippocampal function. Using microarrays, hippocampal transcription patterns associated with late-life physical activity levels were compared to hippocampal gene expression changes associated with Aging and AD. The data demonstrate that late-life physical activity is associated with major reprogramming of hippocampal gene expression, with a striking number of these genes showing significant but opposite patterns of expression change in Aging/AD. The majority (>70%) of these anti-Aging/AD genes show increased expression in association with physical activity but decreased expression in Aging/AD, with an extensive representation of genes regulating mitochondrial energy production, synaptic plasticity at the structural and signaling levels, and axon function.

With age a decline in energy production has been reported across organs and multiple species (Lane et al., 2015; Lin and Beal, 2006). Our data show that physical activity is linked to increased gene expression that included multiple components of the electron transport chain along with key genes driving the TCA cycle, which operates in the mitochondrial matrix and provides key intermediaries (NADH and FADH₂) to the electron transport chain and enhances energy production. Importantly, the genes showing increased expression with physical activity are the same genes that undergo reduced expression in Aging and AD. Enhancement of mitochondrial function may serve as a core mechanism by which physical activity delays Age/AD-related cognitive decline, considering that a ready supply of energy is essential for the maintenance of synaptic fidelity (Harris et al., 2012; Hebert-Chatelain et al., 2016; Smith et al., 2016).

One of the major causes of cognitive decline in Aging and AD is synaptic dysfunction and ultimately failure that progressively disrupts connectivity patterns critical to ongoing functions of the brain (Koffie et al., 2011; Prieto et al., 2017; Selkoe, 2002). Our data suggest that gene expression patterns associated with physical activity may promote synaptic

fidelity in the aged hippocampus not only by targeting energy production but also by counteracting Age/AD-related declines in synaptic components and processes central to neurotransmission. Notably, anti-Aging/AD genes showing increased expression in association with physical activity included many presynaptic components that regulate synaptic vesicle priming, release and recycling, post-synaptic GABA and glutamate neurotransmitter receptors that control the excitatory/inhibitory balance, and PSD scaffolding molecules that tune NMDA and AMPA glutamate signaling, all of which are major mechanisms to regulate synaptic efficacy. Further, many of the anti-Aging/AD genes showing increased expression with physical activity target myelin and axon health, suggesting that physical activity may promote white matter integrity and axonal connectivity and function. The strong relationship between physical activity and expression of genes targeting axon functioning are important, given that white matter degeneration and reduced axonal efficacy are notable components of cognitive decline in both normal aging and in AD pathogenesis (Bartzokis et al., 2003; Fletcher et al., 2018; Kruggel et al., 2017).

Approximately 30% of the anti-Aging/AD genes showed decreased expression with physical activity, but increased expression in Aging/AD. Interestingly, these genes were primarily enriched for transcriptional regulation, including many genes driving negative regulation of transcription. Our data suggest that Aging/AD is accompanied by augmented transcriptional suppression, and that physical activity may offset transcriptional dysregulation and help maintain appropriate transcriptional control of gene expression in the Aging and AD hippocampus. The transcriptional control genes that were increased with Aging/AD but decreased in association with physical activity included several epigenetic regulators of histones including lysine-specific histone demethylases (KDM1B, KDM3A), histone lysine methyltransferase (KMT2C), histone deacetylases (HDAC1) and components of the HDAC complex (SAP18, SAP30, MECOM, TAL1), along with multiple transcriptional repressors and activators. The possibility that physical activity may reduce Aging/AD-associated transcriptional dysregulation would represent a powerful mechanism that could have far-reaching consequences for hippocampal function. For example, one gene of interest is the transcription factor SP3, which forms a chromatin modifying complex with HDAC2 that suppresses transcription of synaptic-plasticity associated genes, by removing acetylation from the promoter region (Yamakawa et al., 2017). Previous studies demonstrate that SP3 is elevated in brains of AD patients and in mouse models of AD, and that reducing SP3, or the SP3-HDAC2 interaction, improves synaptic plasticity and cognitive function without affecting HDAC2 function in other processes (Yamakawa et al., 2017). Our data reveal that SP3 is significantly elevated in the hippocampus already with Aging, shows a further increase with AD, and is reduced in association with physical activity, which may be one mechanism by which physical activity can facilitate synaptic gene expression in the hippocampus. Consistent with this idea, our data reveal that several synaptic target genes that are negatively co-regulated by HDAC2 and SP3 (Yamakawa et al., 2017) show increased expression in association with physical activity, including GRIK2, LIN7A, SYNGR3, and DLGAP1. Taken together, these data suggest that physical activity may offset Age/AD-associated transcriptional dysregulation and help maintain appropriate gene expression in the hippocampus.

While many aspects of human brain aging appear to be counteracted by transcriptional patterns associated with physical activity, an unexpected finding was that physical activity was not associated with extensive involvement of genes regulating immune function and inflammation. We had hypothesized that physical activity would be associated with decreased expression of this gene class based on the literature that the hippocampus undergoes extensive immune gene activation in aging and AD (Barrientos et al., 2015; Berchtold et al., 2008; Cribbs et al., 2012) and the literature that physical activity can constrain activation of immune/inflammatory processes in the hippocampus of animal models (Barrientos et al., 2011; Kohman et al., 2013; Littlefield et al., 2015). While analysis of the anti-Aging/AD genes showing decreased expression in association with physical activity revealed some over-representation of immune/inflammation related genes, the enrichment did not meet statistical significance. Nonetheless, we observed decreased expression in association with physical activity of a few notable genes that drive immune/inflammatory responses, including several activators of the NF- κ B pathway, such as adipocyte enhancer-binding protein 1 (AEBP1), CD40 and filamin A (FLNA), all of which have been implicated in the progression of AD pathology (Giunta et al., 2010; Shijo et al., 2018; Wang et al., 2012). Overall however, our data did not reveal a strong relationship between physical activity levels and expression levels of immune/inflammation-related genes, particularly for immune/inflammation-related genes that show increased expression with Aging and AD.

The transcriptional signatures identified in our study are supported by the human and animal literature reporting widespread benefits of lifestyle activity to brain structure and function. Imaging studies in humans have demonstrated that exercise increases hippocampal volume and density along with cortical gray and white matter volumes and integrity, notably in brain regions that undergo aging-associated atrophy and functional decline (Colcombe et al., 2006; Erickson et al., 2011; Kleemeyer et al., 2016; Ruscheweyh et al., 2011; Voss et al., 2013). Such changes are consistent with our transcriptional data suggesting that physical activity promotes synaptic and cellular health from augmented energy availability, synaptic function, and axon integrity. The findings in human studies are paralleled by the animal literature demonstrating structural and functional changes in the hippocampus and cortex with physical activity including synaptogenesis and increased dendritic complexity (Eadie et al., 2005; Stranahan et al., 2007), hippocampal neurogenesis (Erickson et al., 2013; Fabel et al., 2009; Gould et al., 1999; van Praag et al., 1999b) enhanced synaptic plasticity (Farmer et al., 2004; van Praag et al., 1999a), improved myelin integrity (Zhang et al., 2017; Zhou et al., 2018) and improved spatial learning and memory (Berchtold et al., 2010; Creer et al., 2010; Fordyce and Wehner, 1993; Intlekofer et al., 2013; van Praag et al., 2005). In addition, animal studies have demonstrated that exercise targets multiple aspects mitochondrial function (Cechella et al., 2017; Kim et al., 2010; Lee et al., 2014; Marques-Aleixo et al., 2012; Rampon et al., 2000; Stranahan et al., 2008; Tong et al., 2001). Finally, a handful of animal studies have investigated anti-Aging/AD benefits of physical activity, demonstrated that exercise reverses age or AD-related changes in hippocampal gene expression, most notably for genes associated with mitochondrial function, synaptic plasticity, and axon/myelin integrity (Boveris and Navarro, 2008; Choi et al., 2018; Kohman et al., 2011). We note that some of the findings of animal studies were not recapitulated in the human brain

(e.g., immune gene responses). The discrepancy between our data and the animal literature may be a function of differences in the extent or intensity of exercise participation between humans in this study vs. animals, with the exercise undertaken in animal studies generally being of higher frequency (daily), intensity, and distance (>2 km), but usually shorter duration (<4 weeks).

Taken together, our results suggest that physical activity may broadly shift multiple biological processes towards improved functioning; stimulating core functions that undergo Age/AD-related decline but that are critical for maintaining brain health and synaptic function. We note that most of these genes followed a pattern of only modest expression change in the course of cognitively normal aging followed by progressively greater expression change in AD, and conversely, small expression change with moderate physical activity followed by a relatively large expression change in the highest tier of physical activity. While a relatively modest expression change of any individual gene on its own may not significantly impact hippocampal plasticity and cognitive function, the cumulative effect of reduced expression of multiple genes in a particular functional category, and the gestalt effect of compromised function across multiple categories (e.g. energy production, synaptic efficacy, axon function, transcriptional control), may well be a root cause of age-related decline in hippocampal plasticity and cognitive function. Similarly, the well-established benefit of physical activity to delay age-related cognitive decline and onset of AD may well arise from the cumulative expression change of multiple genes across multiple categories of function. By preserving a more youthful profile across multiple biological processes, these transcriptional effects could empower the brain to strengthen plasticity and build cognitive reserve, providing several potential mechanisms for the cognitive-preserving effects of lifestyle. While the data presented here do not provide direct evidence of a causal relationship, these gene expression patterns may guide the combinatorial use of pharmaceutical agents that target biological domains untouched or minimally associated with physical activity in the human hippocampus transcription pattern.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

5.0 ACKNOWLEDGEMENTS

This work was made possible by National Institute on Aging (R01AG34667, R01AG051807, PPG AG000538, R01AG17917, R01AG15819, R01AG057558), NIH-NINDS (R01NS78009), the Illinois Department of Public Health, the Robert C. Borwell Endowment Fund, and through access to the Genomic High Throughput Facility Shared Resource of the Cancer Center Support Grant (CA-62203) at the University of California, Irvine. The authors declare no biomedical financial interests or potential conflicts of interest.

6. LITERATURE CITED

- Ainsworth BE, Haskell WL, Leon AS, Jacobs DR Jr., Montoye HJ, Sallis JF, Paffenbarger RS Jr., 1993 Compendium of physical activities: classification of energy costs of human physical activities. *Medicine and science in sports and exercise* 25(1), 71–80. [PubMed: 8292105]
- Baker LD, Frank LL, Foster-Schubert K, Green PS, Wilkinson CW, McTiernan A, Plymate SR, Fishel MA, Watson GS, Cholerton BA, Duncan GE, Mehta PD, Craft S, 2010 Effects of aerobic exercise

- on mild cognitive impairment: a controlled trial. *Archives of neurology* 67(1), 71–79. [PubMed: 20065132]
- Barrientos RM, Frank MG, Crysdale NY, Chapman TR, Ahrendsen JT, Day HE, Campeau S, Watkins LR, Patterson SL, Maier SF, 2011 Little exercise, big effects: reversing aging and infection-induced memory deficits, and underlying processes. *J Neurosci* 31(32), 11578–11586.
- Barrientos RM, Kitt MM, Watkins LR, Maier SF, 2015 Neuroinflammation in the normal aging hippocampus. *Neuroscience* 309, 84–99. [PubMed: 25772789]
- Bartzokis G, Cummings JL, Sultzer D, Henderson VW, Nuechterlein KH, Mintz J, 2003 White matter structural integrity in healthy aging adults and patients with Alzheimer disease: a magnetic resonance imaging study. *Archives of neurology* 60(3), 393–398. [PubMed: 12633151]
- Bennett DA, Schneider JA, Buchman AS, Barnes LL, Boyle PA, Wilson RS, 2012 Overview and findings from the rush Memory and Aging Project. *Curr Alzheimer Res* 9(6), 646–663. [PubMed: 22471867]
- Berchtold NC, Castello N, Cotman CW, 2010 Exercise and time-dependent benefits to learning and memory. *Neuroscience* 167(3), 588–597. [PubMed: 20219647]
- Berchtold NC, Cribbs DH, Coleman PD, Rogers J, Head E, Kim R, Beach T, Miller C, Troncoso J, Trojanowski JQ, Zielke HR, Cotman CW, 2008 Gene expression changes in the course of normal brain aging are sexually dimorphic. *Proceedings of the National Academy of Sciences of the United States of America* 105(40), 15605–15610. [PubMed: 18832152]
- Best JR, Rosano C, Aizenstein HJ, Tian Q, Boudreau RM, Ayonayon HN, Satterfield S, Simonsick EM, Studenski S, Yaffe K, Liu-Ambrose T, 2017 Long-term changes in time spent walking and subsequent cognitive and structural brain changes in older adults. *Neurobiology of aging* 57, 153–161. [PubMed: 28648916]
- Bherer L, Erickson KI, Liu-Ambrose T, 2013 A review of the effects of physical activity and exercise on cognitive and brain functions in older adults. *J Aging Res* 2013, 657508.
- Boveris A, Navarro A, 2008 Brain mitochondrial dysfunction in aging. *IUBMB life* 60(5), 308–314. [PubMed: 18421773]
- Buchman AS, Boyle PA, Yu L, Shah RC, Wilson RS, Bennett DA, 2012 Total daily physical activity and the risk of AD and cognitive decline in older adults. *Neurology* 78(17), 1323–1329. [PubMed: 22517108]
- Buchman AS, Wilson RS, Bennett DA, 2008 Total daily activity is associated with cognition in older persons. *Am J Geriatr Psychiatry* 16(8), 697–701. [PubMed: 18669949]
- Cechella JL, Leite MR, Pinton S, Zeni G, Nogueira CW, 2017 Neuroprotective Benefits of Aerobic Exercise and Organoselenium Dietary Supplementation in Hippocampus of Old Rats. *Mol Neurobiol*.
- Choi SH, Bylykbashi E, Chatila ZK, Lee SW, Pulli B, Clemenson GD, Kim E, Rompala A, Oram MK, Asselin C, Aronson J, Zhang C, Miller SJ, Lesinski A, Chen JW, Kim DY, van Praag H, Spiegelman BM, Gage FH, Tanzi RE, 2018 Combined adult neurogenesis and BDNF mimic exercise effects on cognition in an Alzheimer’s mouse model. *Science (New York, N.Y)* 361(6406).
- Colcombe S, Kramer AF, 2003 Fitness effects on the cognitive function of older adults: a meta-analytic study. *Psychol Sci* 14(2), 125–130. [PubMed: 12661673]
- Colcombe SJ, Erickson KI, Raz N, Webb AG, Cohen NJ, McAuley E, Kramer AF, 2003 Aerobic fitness reduces brain tissue loss in aging humans. *J Gerontol A Biol Sci Med Sci* 58(2), 176–180. [PubMed: 12586857]
- Colcombe SJ, Erickson KI, Scalf PE, Kim JS, Prakash R, McAuley E, Elavsky S, Marquez DX, Hu L, Kramer AF, 2006 Aerobic exercise training increases brain volume in aging humans. *J Gerontol A Biol Sci Med Sci* 61(11), 1166–1170. [PubMed: 17167157]
- Colcombe SJ, Kramer AF, McAuley E, Erickson KI, Scalf P, 2004 Neurocognitive aging and cardiovascular fitness: recent findings and future directions. *J Mol Neurosci*. 24(1), 9–14. [PubMed: 15314244]
- Creer DJ, Romberg C, Saksida LM, van Praag H, Bussey TJ, 2010 Running enhances spatial pattern separation in mice. *Proceedings of the National Academy of Sciences of the United States of America* 107(5), 2367–2372. [PubMed: 20133882]

- Cribbs DH, Berchtold NC, Perreau V, Coleman PD, Rogers J, Tenner AJ, Cotman CW, 2012 Extensive innate immune gene activation accompanies brain aging, increasing vulnerability to cognitive decline and neurodegeneration: a microarray study. *Journal of neuroinflammation* 9, 179. [PubMed: 22824372]
- Dennis G Jr., Sherman BT, Hosack DA, Yang J, Gao W, Lane HC, Lempicki RA, 2003 DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biol* 4(5), P3. [PubMed: 12734009]
- Eadie BD, Redila VA, Christie BR, 2005 Voluntary exercise alters the cytoarchitecture of the adult dentate gyrus by increasing cellular proliferation, dendritic complexity, and spine density. *The Journal of comparative neurology* 486(1), 39–47. [PubMed: 15834963]
- Erickson KI, Gildengers AG, Butters MA, 2013 Physical activity and brain plasticity in late adulthood. *Dialogues Clin Neurosci* 15(1), 99–108. [PubMed: 23576893]
- Erickson KI, Leckie RL, Weinstein AM, 2014 Physical activity, fitness, and gray matter volume. *Neurobiology of aging* 35 Suppl 2, S20–28. [PubMed: 24952993]
- Erickson KI, Raji CA, Lopez OL, Becker JT, Rosano C, Newman AB, Gach HM, Thompson PM, Ho AJ, Kuller LH, 2010 Physical activity predicts gray matter volume in late adulthood: the Cardiovascular Health Study. *Neurology* 75(16), 1415–1422. [PubMed: 20944075]
- Erickson KI, Voss MW, Prakash RS, Basak C, Szabo A, Chaddock L, Kim JS, Heo S, Alves H, White SM, Wojcicki TR, Mailey E, Vieira VJ, Martin SA, Pence BD, Woods JA, McAuley E, Kramer AF, 2011 Exercise training increases size of hippocampus and improves memory. *Proceedings of the National Academy of Sciences of the United States of America* 108(7), 3017–3022. [PubMed: 21282661]
- Erickson KI, Weinstein AM, Lopez OL, 2012 Physical activity, brain plasticity, and Alzheimer's disease. *Arch Med Res* 43(8), 615–621. [PubMed: 23085449]
- Fabel K, Wolf SA, Ehninger D, Babu H, Leal-Galicia P, Kempermann G, 2009 Additive effects of physical exercise and environmental enrichment on adult hippocampal neurogenesis in mice. *Front Neurosci* 3, 50. [PubMed: 20582277]
- Farmer J, Zhao X, van Praag H, Wodtke K, Gage FH, Christie BR, 2004 Effects of voluntary exercise on synaptic plasticity and gene expression in the dentate gyrus of adult male Sprague-Dawley rats in vivo. *Neuroscience* 124(1), 71–79. [PubMed: 14960340]
- Fletcher E, Gavett B, Harvey D, Farias ST, Olichney J, Beckett L, DeCarli C, Mungas D, 2018 Brain volume change and cognitive trajectories in aging. *Neuropsychology* 32(4), 436–449. [PubMed: 29494196]
- Fordyce DE, Wehner JM, 1993 Physical activity enhances spatial learning performance with an associated alteration in hippocampal protein kinase C activity in C57BL/6 and DBA/2 mice. *Brain research* 619(1–2), 111–119. [PubMed: 8374769]
- Geiss GK, Bumgarner RE, Birditt B, Dahl T, Dowidar N, Dunaway DL, Fell HP, Ferree S, George RD, Grogan T, James JJ, Maysuria M, Mitton JD, Oliveri P, Osborn JL, Peng T, Ratcliffe AL, Webster PJ, Davidson EH, Hood L, Dimitrov K, 2008 Direct multiplexed measurement of gene expression with color-coded probe pairs. *Nat Biotechnol* 26(3), 317–325. [PubMed: 18278033]
- Giunta B, Rezaei-Zadeh K, Tan J, 2010 Impact of the CD40-CD40L dyad in Alzheimer's disease. *CNS Neurol Disord Drug Targets* 9(2), 149–155. [PubMed: 20205645]
- Gould E, Beylin A, Tanapat P, Reeves A, Shors TJ, 1999 Learning enhances adult neurogenesis in the hippocampal formation [see comments]. *Nature neuroscience* 2(3), 260–265. [PubMed: 10195219]
- Hall KS, Howe CA, Rana SR, Martin CL, Morey MC, 2013 METs and accelerometry of walking in older adults: standard versus measured energy cost. *Medicine and science in sports and exercise* 45(3), 574–582. [PubMed: 23059862]
- Harris JJ, Jolivet R, Attwell D, 2012 Synaptic energy use and supply. *Neuron* 75(5), 762–777. [PubMed: 22958818]
- Hebert-Chatelain E, Desprez T, Serrat R, Bellocchio L, Soria-Gomez E, Busquets-Garcia A, Pagano Zottola AC, Delamarre A, Cannich A, Vincent P, Varilh M, Robin LM, Terral G, Garcia-Fernandez MD, Colavita M, Mazier W, Drago F, Puente N, Reguero L, Elezgarai I, Dupuy JW, Cota D, Lopez-Rodriguez ML, Barreda-Gomez G, Massa F, Grandes P, Benard G, Marsicano G, 2016 A

- cannabinoid link between mitochondria and memory. *Nature* 539(7630), 555–559. [PubMed: 27828947]
- Hillman CH, Erickson KI, Kramer AF, 2008 Be smart, exercise your heart: exercise effects on brain and cognition. *Nature reviews* 9(1), 58–65.
- Huang da W, Sherman BT, Lempicki RA, 2009a Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 37(1), 1–13. [PubMed: 19033363]
- Huang da W, Sherman BT, Lempicki RA, 2009b Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature protocols* 4(1), 44–57. [PubMed: 19131956]
- Huang P, Fang R, Li BY, Chen SD, 2016 Exercise-Related Changes of Networks in Aging and Mild Cognitive Impairment Brain. *Front Aging Neurosci* 8, 47. [PubMed: 27014055]
- Intlekofer KA, Berchtold NC, Malvaez M, Carlos AJ, McQuown SC, Cunningham MJ, Wood MA, Cotman CW, 2013 Exercise and Sodium Butyrate Transform a Subthreshold Learning Event into Long-Term Memory via a Brain-Derived Neurotrophic factor-Dependent Mechanism. *Neuropsychopharmacology* 38(10), 2027–2034. [PubMed: 23615664]
- Kim SE, Ko IG, Kim BK, Shin MS, Cho S, Kim CJ, Kim SH, Baek SS, Lee EK, Jee YS, 2010 Treadmill exercise prevents aging-induced failure of memory through an increase in neurogenesis and suppression of apoptosis in rat hippocampus. *Experimental gerontology* 45(5), 357–365. [PubMed: 20156544]
- Kirk-Sanchez NJ, McGough EL, 2014 Physical exercise and cognitive performance in the elderly: current perspectives. *Clin Interv Aging* 9, 51–62. [PubMed: 24379659]
- Kleemeyer MM, Kuhn S, Prindle J, Bodammer NC, Brechtel L, Garthe A, Kempermann G, Schaefer S, Lindenberger U, 2016 Changes in fitness are associated with changes in hippocampal microstructure and hippocampal volume among older adults. *NeuroImage* 131, 155–161. [PubMed: 26584869]
- Koffie RM, Hyman BT, Spiers-Jones TL, 2011 Alzheimer's disease: synapses gone cold. *Mol Neurodegener* 6(1), 63. [PubMed: 21871088]
- Kohman RA, Bhattacharya TK, Wojcik E, Rhodes JS, 2013 Exercise reduces activation of microglia isolated from hippocampus and brain of aged mice. *Journal of neuroinflammation* 10, 114. [PubMed: 24044641]
- Kohman RA, Rodriguez-Zas SL, Southey BR, Kelley KW, Dantzer R, Rhodes JS, 2011 Voluntary wheel running reverses age-induced changes in hippocampal gene expression. *PLoS ONE* 6(8), e22654.
- Korol DL, Gold PE, Scavuzzo CJ, 2013 Use it and boost it with physical and mental activity. *Hippocampus* 23(11), 1125–1135. [PubMed: 23996382]
- Kramer AF, Erickson KI, 2007a Capitalizing on cortical plasticity: influence of physical activity on cognition and brain function. *Trends Cogn Sci* 11(8), 342–348. [PubMed: 17629545]
- Kramer AF, Erickson KI, 2007b Effects of physical activity on cognition, well-being, and brain: human interventions. *Alzheimers Dement* 3(2 Suppl), S45–51. [PubMed: 19595974]
- Kruggel F, Masaki F, Solodkin A, 2017 Analysis of longitudinal diffusion-weighted images in healthy and pathological aging: An ADNI study. *J Neurosci Methods* 278, 101–115. [PubMed: 28057473]
- Lane RK, Hilsabeck T, Rea SL, 2015 The role of mitochondrial dysfunction in age-related diseases. *Biochim Biophys Acta* 1847(11), 1387–1400. [PubMed: 26050974]
- Lautenschlager NT, Cox KL, Flicker L, Foster JK, van Bockxmeer FM, Xiao J, Greenop KR, Almeida OP, 2008 Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: a randomized trial. *Jama* 300(9), 1027–1037. [PubMed: 18768414]
- Lee MC, Rakwal R, Shibato J, Inoue K, Chang H, Soya H, 2014 DNA microarray-based analysis of voluntary resistance wheel running reveals novel transcriptome leading robust hippocampal plasticity. *Physiol Rep* 2(11).
- Lin MT, Beal MF, 2006 Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature* 443(7113), 787–795. [PubMed: 17051205]
- Littlefield AM, Setti SE, Priester C, Kohman RA, 2015 Voluntary exercise attenuates LPS-induced reductions in neurogenesis and increases microglia expression of a proneurogenic phenotype in aged mice. *Journal of neuroinflammation* 12, 138. [PubMed: 26224094]

- Marques-Aleixo I, Oliveira PJ, Moreira PI, Magalhaes J, Ascensao A, 2012 Physical exercise as a possible strategy for brain protection: evidence from mitochondrial-mediated mechanisms. *Prog Neurobiol* 99(2), 149–162. [PubMed: 22940590]
- Miller JA, Guillozet-Bongaarts A, Gibbons LE, Postupna N, Renz A, Beller AE, Sunkin SM, Ng L, Rose SE, Smith KA, Szafer A, Barber C, Bertagnolli D, Bickley K, Brouner K, Caldejon S, Chapin M, Chua ML, Coleman NM, Cudaback E, Cuhaciyan C, Dalley RA, Dee N, Desta T, Dolbear TA, Dotson NI, Fisher M, Gaudreault N, Gee G, Gilbert TL, Goldy J, Griffin F, Habel C, Haradon Z, Hejazinia N, Hellstern LL, Horvath S, Howard K, Howard R, Johal J, Jorstad NL, Josephsen SR, Kuan CL, Lai F, Lee E, Lee F, Lemon T, Li X, Marshall DA, Melchor J, Mukherjee S, Nyhus J, Pendergraft J, Potekhina L, Rha EY, Rice S, Rosen D, Sapru A, Schantz A, Shen E, Sheffield E, Shi S, Sodt AJ, Thatra N, Tieu M, Wilson AM, Montine TJ, Larson EB, Bernard A, Crane PK, Ellenbogen RG, Keene CD, Lein E, 2017 Neuropathological and transcriptomic characteristics of the aged brain. *Elife* 6.
- Okonkwo OC, Schultz SA, Oh JM, Larson J, Edwards D, Cook D, Kosciak R, Gallagher CL, Dowling NM, Carlsson CM, Bendlin BB, LaRue A, Rowley HA, Christian BT, Asthana S, Hermann BP, Johnson SC, Sager MA, 2014 Physical activity attenuates age-related biomarker alterations in preclinical AD. *Neurology* 83(19), 1753–1760. [PubMed: 25298312]
- Prakash RS, Voss MW, Erickson KI, Kramer AF, 2015 Physical activity and cognitive vitality. *Annu Rev Psychol* 66, 769–797. [PubMed: 25251492]
- Prieto GA, Trieu BH, Dang CT, Bilousova T, Gyls KH, Berchtold NC, Lynch G, Cotman CW, 2017 Pharmacological Rescue of Long-Term Potentiation in Alzheimer Diseased Synapses. *J Neurosci* 37(5), 1197–1212. [PubMed: 27986924]
- Rampon C, Jiang CH, Dong H, Tang YP, Lockhart DJ, Schultz PG, Tsien JZ, Hu Y, 2000 Effects of environmental enrichment on gene expression in the brain. *Proceedings of the National Academy of Sciences of the United States of America* 97(23), 12880–12884.
- Ruscheweyh R, Willemer C, Kruger K, Duning T, Warnecke T, Sommer J, Volker K, Ho HV, Mooren F, Knecht S, Floel A, 2011 Physical activity and memory functions: an interventional study. *Neurobiology of aging* 32(7), 1304–1319. [PubMed: 19716631]
- Selkoe DJ, 2002 Alzheimer's disease is a synaptic failure. *Science (New York, N.Y)* 298(5594), 789–791. [PubMed: 12399581]
- Shijo M, Honda H, Suzuki SO, Hamasaki H, Hokama M, Abolhassani N, Nakabeppu Y, Ninomiya T, Kitazono T, Iwaki T, 2018 Association of adipocyte enhancer-binding protein 1 with Alzheimer's disease pathology in human hippocampi. *Brain Pathol* 28(1), 58–71. [PubMed: 27997051]
- Smith HL, Bourne JN, Cao G, Chirillo MA, Ostroff LE, Watson DJ, Harris KM, 2016 Mitochondrial support of persistent presynaptic vesicle mobilization with age-dependent synaptic growth after LTP. *Elife* 5.
- Smith PJ, Blumenthal JA, Hoffman BM, Cooper H, Strauman TA, Welsh-Bohmer K, Browndyke JN, Sherwood A, 2010 Aerobic exercise and neurocognitive performance: a meta-analytic review of randomized controlled trials. *Psychosomatic medicine* 72(3), 239–252. [PubMed: 20223924]
- Stranahan AM, Khalil D, Gould E, 2007 Running induces widespread structural alterations in the hippocampus and entorhinal cortex. *Hippocampus* 17(11), 1017–1022. [PubMed: 17636549]
- Stranahan AM, Lee K, Becker KG, Zhang Y, Maudsley S, Martin B, Cutler RG, Mattson MP, 2008 Hippocampal gene expression patterns underlying the enhancement of memory by running in aged mice. *Neurobiology of aging*.
- Stubbs B, Chen LJ, Chang CY, Sun WJ, Ku PW, 2017 Accelerometer-assessed light physical activity is protective of future cognitive ability: A longitudinal study among community dwelling older adults. *Experimental gerontology* 91, 104–109. [PubMed: 28263868]
- Suwabe K, Byun K, Hyodo K, Reagh ZM, Roberts JM, Matsushita A, Saotome K, Ochi G, Fukuie T, Suzuki K, Sankai Y, Yassa MA, Soya H, 2018 Rapid stimulation of human dentate gyrus function with acute mild exercise. *Proceedings of the National Academy of Sciences of the United States of America* 115(41), 10487–10492.
- Suzuki T, Shimada H, Makizako H, Doi T, Yoshida D, Tsutsumimoto K, Anan Y, Uemura K, Lee S, Park H, 2012 Effects of multicomponent exercise on cognitive function in older adults with amnesic mild cognitive impairment: a randomized controlled trial. *BMC Neurol* 12, 128. [PubMed: 23113898]

- Tong L, Shen H, Perreau V, Balazs R, Cotman CW, 2001 Effects of exercise on gene-expression profile in the rat hippocampus. *Neurobiology of disease* 8(6), 1046–1056. [PubMed: 11741400]
- van Praag H, Christie BR, Sejnowski TJ, Gage FH, 1999a Running enhances neurogenesis, learning, and long-term potentiation in mice. *PNAS* 96(23), 13427–13431.
- van Praag H, Kempermann G, Gage FH, 1999b Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus. *Nature Neurosci* 2(3), 266–270. [PubMed: 10195220]
- van Praag H, Shubert T, Zhao C, Gage FH, 2005 Exercise enhances learning and hippocampal neurogenesis in aged mice. *J Neurosci*. 25(38), 8680–8685. [PubMed: 16177036]
- Veldman-Jones MH, Brant R, Rooney C, Geh C, Emery H, Harbron CG, Wappett M, Sharpe A, Dymond M, Barrett JC, Harrington EA, Marshall G, 2015 Evaluating Robustness and Sensitivity of the NanoString Technologies nCounter Platform to Enable Multiplexed Gene Expression Analysis of Clinical Samples. *Cancer Res* 75(13), 2587–2593. [PubMed: 26069246]
- Voss MW, Heo S, Prakash RS, Erickson KI, Alves H, Chaddock L, Szabo AN, Mailey EL, Wojcicki TR, White SM, Gothe N, McAuley E, Sutton BP, Kramer AF, 2013 The influence of aerobic fitness on cerebral white matter integrity and cognitive function in older adults: results of a one-year exercise intervention. *Hum Brain Mapp* 34(11), 2972–2985. [PubMed: 22674729]
- Voss MW, Prakash RS, Erickson KI, Basak C, Chaddock L, Kim JS, Alves H, Heo S, Szabo AN, White SM, Wojcicki TR, Mailey EL, Gothe N, Olson EA, McAuley E, Kramer AF, 2010 Plasticity of brain networks in a randomized intervention trial of exercise training in older adults. *Front Aging Neurosci* 2.
- Wang HY, Bakshi K, Frankfurt M, Stucky A, Goberdhan M, Shah SM, Burns LH, 2012 Reducing amyloid-related Alzheimer's disease pathogenesis by a small molecule targeting filamin A. *J Neurosci* 32(29), 9773–9784. [PubMed: 22815492]
- Weuve J, Kang JH, Manson JE, Breteler MM, Ware JH, Grodstein F, 2004 Physical activity, including walking, and cognitive function in older women. *Jama*. 292(12), 1454–1461. [PubMed: 15383516]
- Yaffe K, Barnes D, Nevitt M, Lui LY, Covinsky K, 2001 A prospective study of physical activity and cognitive decline in elderly women: women who walk. *Archives of internal medicine* 161(14), 1703–1708. [PubMed: 11485502]
- Yamakawa H, Cheng J, Penney J, Gao F, Rueda R, Wang J, Yamakawa S, Kritskiy O, Gjoneska E, Tsai LH, 2017 The Transcription Factor Sp3 Cooperates with HDAC2 to Regulate Synaptic Function and Plasticity in Neurons. *Cell Rep* 20(6), 1319–1334. [PubMed: 28793257]
- Zhang L, Chao FL, Luo YM, Xiao Q, Jiang L, Zhou CN, Zhang Y, Lv FL, He Q, Ma J, Tang Y, 2017 Exercise Prevents Cognitive Function Decline and Demyelination in the White Matter of APP/PS1 Transgenic AD Mice. *Curr Alzheimer Res* 14(6), 645–655. [PubMed: 27978791]
- Zhou CN, Chao FL, Zhang Y, Jiang L, Zhang L, Luo YM, Xiao Q, Chen LM, Tang Y, 2018 Sex Differences in the White Matter and Myelinated Fibers of APP/PS1 Mice and the Effects of Running Exercise on the Sex Differences of AD Mice. *Front Aging Neurosci* 10, 243. [PubMed: 30174598]
- Zlatar ZZ, McGregor KM, Towler S, Nocera JR, Dzierzewski JM, Crosson B, 2015 Self-reported physical activity and objective aerobic fitness: differential associations with gray matter density in healthy aging. *Front Aging Neurosci* 7, 5. [PubMed: 25691866]

HIGHLIGHTS

- Physical activity is extensively associated with gene expression in the aged human hippocampus.
- Gene expression patterns associated with physical activity restore a more youthful state in the hippocampus, and opposed transcription patterns associated with Aging and Alzheimer's disease.
- Gene expression is primarily increased, with genes targeting multiple core cellular functions.
- Genes showed robust enrichment for mitochondrial energy production, synaptic signaling mechanisms, axon and white matter health, and control of transcription.




Figure 1.

Flowchart of the Methods to determine if physical activity-related gene expression patterns counter aging and AD-related expression change. 34 microarrays profiling gene expression in the aged human hippocampus were used to determine the relationship between gene expression and physical activity. Poorly performing probesets were eliminated by filtering the array data based on flags (Present, Marginal, Absent) using MAS preprocessing, retaining only probesets flagged Present on all samples of at least one group (low, moderate, high activity) (23,034 probesets). Using the programming environment R (<http://www.R-project.org/>), multiple linear regression was applied to GC-RMA normalized values using `lm()` to adjust for technical factors that were potential confounding variables (batch, RIN, and PMI) and to evaluate the relationship between gene expression and late-life physical activity levels across the 34 samples. In parallel, Aging and AD-related gene expression change was assessed using 51 CEL files from the public gene expression omnibus (<http://www.ncbi.nlm.nih.gov/geo>, dataset accession number GSE:11882) profiling expression patterns of 54,675 probesets in hippocampal samples from young cases (20–59 yrs, n=17), cognitively intact aged cases (73–99 yrs, n=20), and AD cases (73–99 yrs, n=14). The ~55,000 probesets on the array were filtered using MAS preprocessing derived flags to

retain only probesets flagged Present on all samples of at least one treatment group (21,002 probesets). Using R, multiple linear regression was applied to GC-RMA normalized values to adjust for batch, RIN, and PMI and to evaluate gene expression changes in Aging (aging vs young) and AD (AD vs aged). Probesets significantly associated with both physical activity ($p < 0.05$ and Aging or AD ($p < 0.05$) were selected, patterns of expression change were compared for each probeset, and probesets showing opposite patterns of change with physical activity versus aging or AD were identified. The number of probesets expected to occur by chance in the overlap of the 2 studies is 52 probesets, calculated by multiplying the product of the p-values (0.0025) by the number of probesets that were tested in both datasets (21,002). The analysis identified 2138 probesets significantly associated with both physical activity and Aging or AD, 94.9% (2029 probesets) of which showed the opposite pattern of change with physical activity compared to Aging/AD (“Anti-Aging/AD probesets”). These probesets were analyzed for functional enrichment using DAVID.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 2. The KEGG pathway “oxidative phosphorylation” (3.17-fold enriched, $p = 5.5 \times 10^{-7}$) overlaid with microarray data - blue highlight identifies genes showing significantly increased expression in association with physical activity and decreased expression in Aging or AD. Significant genes included approximately 30% of the genes encoding mitochondrial complex I (NADH:ubiquinone oxidoreductase) (NDUFA1, NDUFA7, NDUFA8, NDUFA9, NDUFAF5, NDUFB1, NDUFB8, NDUFB9, NDUFS1, NDUFS3, NDUFS5, NDUFV2 and NDUFV3) approximately 35% of the genes encoding Complex IV (cytochrome C oxidase) (subunits COX4A1, COX6A1, COX6C, COX7B, COX7A2, COX8A and COX11) and Complex V (ATP synthase) (ATP5A1, ATP5C1, ATP5O, ATP5G1, ATP5L, ATP51), as well as 2 components (SLC25A27, SLC25A14) of the uncoupling protein. No genes in the pathway showed decreased expression in association with physical activity.




Figure 3.

The synaptic vesicle KEGG pathway overlaid with microarray analysis with blue highlight identifying genes showing significantly increased expression in association with physical activity and decreased expression in Aging or AD. Significant genes are involved in synaptic vesicle trafficking, priming and docking at the active zone: SNAP25, synaptobrevin/VAMP2, synapsin 2 (SYN2), syntaxin binding protein 1 (STXBP1) as well as synaptic vesicle endocytosis and recycling (dynamin-like 1 (DNML1), DNML3, adaptor protein complex 2 (AP2M1). No genes in the pathway showed decreased expression in association with physical activity.




Figure 4.

The NanoString nCounter™ gene expression system was used to assess gene expression changes with physical activity, using an expanded set of human hippocampal samples (low activity, n=18; high activity, n=16) (A) 8 genes associated with GABA signaling: the G-protein coupled metabotropic GABA_B receptor 1 (GABBR1), ionotropic GABA_A receptor alpha subunits (GABRA2, GABRA4), GABA_A receptor beta subunit 3 (GABRB3), GABA_A receptor gamma subunits (GABRG1, GABRG2), glycine receptor beta (GLRB) and somatostatin receptor 1 (SSTR1). (B) 3 genes regulating glutamate signaling: Glutamate ionotropic Receptor AMPA Type Subunit 2 (GRIA2), Glutamate ionotropic Receptor NMDA Type Subunit 2A (GRIN2a), and the neuron specific Sodium-Dependent Glutamate/Aspartate Transporter (Solute Carrier Family 1 Member 6: SLC1a6). (C) 7 genes involved in synaptic vesicle trafficking and release: Synapsin II (SYN2), Unc-13 Homolog C (UNC13c/Munc13-3), Ubiquitin Specific Peptidase 14 (USP14), Double C2 Domain Alpha (DOC2a), voltage-gated sodium channel (type II, subunit B: SCN2B), Synaptosomal-associated protein 25 (SNAP25) and Vesicle Associated Membrane Protein 2 (VAMP2). (D) 5 genes involved in building new synapses and the maintenance and plasticity of the post-synaptic density: Neurexin 1 (NRXN1), Neurexin 3 (NRXN3), Discs Large Homolog 3 (DLG3, also known as synapse associated protein SAP102), A-Kinase Anchoring Protein 5 (AKAP5), Synaptotagmin 4 (SYT4), Mean ± SEM. * p 0.05, ** p 0.01, ***p 0.005.

Table 1

DAVID enrichment categories for unambiguous probesets with hippocampal expression significantly increased in association with physical activity and significantly decreased with Aging or AD. Note that significant genes may be contained with in multiple enriched terms within a category.

Mitochondria-related categories	Database	Term	Count	%	EASE PValue	List Total	Pop Hits	Pop Total	Fold Enrichment	Benjamini
Mitochondrion	UP_KEYWORDS		143	12.16	1.98E-07	1138	881	10612	1.51	2.04E-05
Mitochondrion	GOTERM_CC_DIRECT	GO:0005739	162	13.78	3.57E-08	1047	1026	9961	1.50	7.04E-06
Mitochondrion inner membrane	UP_KEYWORDS		55	4.68	2.12E-09	1138	218	10612	2.35	4.39E-07
Mitochondrion inner membrane	GOTERM_CC_DIRECT	GO:0005743	74	6.29	2.77E-08	1047	364	9961	1.93	8.20E-06
Mitochondrion outer membrane	UP_KEYWORDS		19	1.62	3.41E-03	1138	85	10612	2.08	9.60E-02
Respiratory chain	UP_KEYWORDS		17	1.45	1.41E-04	1138	55	10612	2.88	8.26E-03
oxidative phosphorylation	KEGG_PATHWAY	hsa00190	33	2.81	2.19E-09	380	107	3909	3.17	5.54E-07
mitochondrial respiratory chain complex I	GOTERM_CC_DIRECT	GO:0005747	15	1.28	5.70E-05	1047	42	9961	3.40	4.81E-03
ubiquinone	UP_KEYWORDS		10	0.85	1.88E-03	1138	28	10612	3.33	6.83E-02
mitochondrial electron transport, NADH to ubiquinone	GOTERM_BP_DIRECT	GO:0006120	15	1.28	6.51E-05	972	43	9380	3.37	9.24E-02
mitochondrial proton-transporting ATP synthase complex	GOTERM_CC_DIRECT	GO:0005753	8	0.68	2.23E-03	1047	19	9961	4.01	9.67E-02
Metabolic pathways	KEGG_PATHWAY	hsa01100	103	8.76	6.18E-04	380	792	3909	1.34	2.21E-02

Synapse, plasticity -related categories	Database	Term	Count	%	EASE PValue	List Total	Pop Hits	Pop Total	Fold Enrichment	Benjamini
Synapse	UP_KEYWORDS		60	5.10	1.23E-09	1138	245	10612	2.28	5.06E-07
synaptic vesicle membrane	GOTERM_CC_DIRECT	GO:0030672	16	1.36	4.16E-06	1047	39	9961	3.90	4.92E-04
chemical synaptic transmission	GOTERM_BP_DIRECT	GO:0007268	31	2.64	3.31E-05	972	134	9380	2.23	9.38E-02
Postsynaptic cell membrane	UP_KEYWORDS		26	2.21	1.46E-04	1138	108	10612	2.24	7.50E-03
Postsynaptic membrane	GOTERM_CC_DIRECT	GO:0045211	29	2.47	2.08E-04	1047	132	9961	2.09	1.36E-02
crebPathway:Transcription factor CREB and its extracellular signals	BIOCARTA		9	0.77	1.89E-03	102	26	1060	3.60	6.89E-02
axon	GOTERM_CC_DIRECT	GO:0030424	38	3.23	3.40E-05	1047	178	9961	2.03	3.35E-03
myelin sheath	GOTERM_CC_DIRECT	GO:0043209	36	3.06	7.04E-07	1047	140	9961	2.45	1.04E-04
growth cone	GOTERM_CC_DIRECT	GO:0030426	23	1.96	1.62E-04	1047	92	9961	2.38	1.19E-02
neuron projection	GOTERM_CC_DIRECT	GO:0043005	31	2.64	1.87E-03	1047	165	9961	1.79	9.56E-02

Synapse, plasticity -related categories	Database	Term	Count	%	EASE PValue	List Total	Pop Hits	Pop Total	Fold Enrichment	Benjamini
Cell_junction	UP_KEYWORDS		78	6.63	4.37E-06	1138	433	10612	1.68	3.61E-04
Cell_junction	GOTERM_CC_DIRECT		66	5.61	2.16E-08	1047	308	9961	2.04	1.28E-05
Other categories	Database	Term	Count	%	EASE PValue	List Total	Pop Hits	Pop Total	Fold Enrichment	Benjamini
gap_junction	KEGG_PATHWAY	hsa04540	16	1.36	1.25E-03	380	66	3909	2.49	3.12E-02
Circadian entrainment	KEGG_PATHWAY	hsa04713	17	1.45	6.86E-04	380	69	3909	2.53	2.15E-02
Transport	UP_KEYWORDS		172	14.63	4.71E-04	1138	1265	10612	1.27	2.14E-02
Transit peptide	UP_KEYWORDS		77	6.55	1.07E-05	1138	436	10612	1.65	7.33E-04
chrebpPathway:CHREBP regulation by carbohydrates and cAMP	BIOCARTA		9	0.77	3.51E-05	102	16	1060	5.85	6.62E-03
agprPathway:Attenuation of GPCR Signaling	BIOCARTA		7	0.60	2.02E-04	102	11	1060	6.61	1.89E-02
cskPathway:Activation of Csk by cAMP-dependent Protein Kinase Inhibits Signaling through the T Cell Receptor	BIOCARTA		7	0.60	1.03E-03	102	14	1060	5.20	6.30E-02
plcePathway:Phospholipase C-epsilon pathway	BIOCARTA		6	0.51	1.22E-03	102	10	1060	6.24	5.60E-02
raccPathway:Ion Channels and Their Functional Role in Vascular Endothelium	BIOCARTA		6	0.51	2.07E-03	102	11	1060	5.67	6.31E-02
nos1Pathway:Nitric Oxide Signaling Pathway	BIOCARTA		7	0.60	2.35E-03	102	16	1060	4.55	6.15E-02
gpcrPathway:Signaling Pathway from G-Protein Families	BIOCARTA		9	0.77	3.19E-03	102	28	1060	3.34	7.28E-02

Table 2

DAVID enrichment results for unambiguous probesets with hippocampal expression significantly decreased in association with physical activity and significantly increased with Aging or AD. Note that significant genes may be contained with in multiple enriched terms within a category.

Category	Database	Term	Count	%	EASE PValue	List Total	Pop Hits	Pop Total	Fold Enrichment	Benjamini
Nucleus	UP_KEYWORDS		155	37.99	3.08E-04	383	3398	10612	1.26	1.93E-02
Transcription	UP_KEYWORDS		81	19.85	9.04E-05	383	1480	10612	1.52	7.14E-03
Transcription regulation	UP_KEYWORDS		79	19.36	8.03E-05	383	1428	10612	1.53	8.45E-03
negative regulation of transcription from RNA polymerase II promoter	GOTERM_BP	GO:0000122	35	8.58	4.93E-05	357	436	9380	2.11	8.80E-02
Repressor	UP_KEYWORDS		31	7.60	3.64E-04	383	428	10612	2.01	1.91E-02
Activator	UP_KEYWORDS		30	7.35	1.13E-03	383	438	10612	1.90	4.37E-02
Polymorphism	UP_KEYWORDS		274	67.16	4.03E-08	383	6188	10612	1.23	1.28E-05
Alternative splicing	UP_KEYWORDS		269	65.93	9.26E-04	383	6639	10612	1.12	4.11E-02
Isopeptide bond	UP_KEYWORDS		48	11.76	1.94E-03	383	849	10612	1.57	6.60E-02

Category	Term	Count	%	PValue	List Total	Pop Hits	Pop Total	Fold Enrichment	Benjamini
UP_KEYWC	Polymorphism	274	67.15686	4.03E-08	383	6188	10612	1.226871	1.28E-05
UP_KEYWC	Phosphoprotein	259	63.48039	3.12E-07	383	5849	10612	1.226921	4.94E-05
UP_KEYWC	Transcription	81	19.85294	9.04E-05	383	1480	10612	1.516428	0.007139
UP_KEYWC	Transcription	79	19.36275	8.03E-05	383	1428	10612	1.532842	0.008448
UP_KEYWC	Repressor	31	7.598039	3.64E-04	383	428	10612	2.006857	0.019076
UP_KEYWC	Nucleus	155	37.9902	3.08E-04	383	3398	10612	1.263883	0.019342
UP_KEYWC	Alternative	269	65.93137	9.26E-04	383	6639	10612	1.12266	0.041106
UP_KEYWC	Activator	30	7.352941	0.001128	383	438	10612	1.897779	0.043743
UP_KEYWC	Isopeptide	48	11.76471	0.001936	383	849	10612	1.566506	0.065984
GOTERM_E	GO:000012	35	8.578431	4.93E-05	357	436	9380	2.109192	0.088027
UP_KEYWC	DNA-bindin	56	13.72549	0.003426	383	1061	10612	1.462417	0.103095