Lawrence Berkeley National Laboratory

Recent Work

Title

CARBON-13 NUCLEAR MAGNETIC RESONANCE IN SOLID AMMONIUM TARTRATE

Permalink

<https://escholarship.org/uc/item/4940w0jq>

Authors

Pines, A.
Chang, J.J.
Griffin, R.G.

Publication Date

1974-03-01

Submitted to Journal of
Chemical Physics

RECEIVED
LAWRENCE
RADIATION LABORATORY

LBL-2700
Preprint *e.2*

APR 29 1974

LIBRARY AND
DOCUMENTS SECTION

CARBON-13 NUCLEAR MAGNETIC RESONANCE IN
SOLID AMMONIUM TARTRATE

A. Pines, J. J. Chang and R. G. Griffin

March 1974

Prepared for the U. S. Atomic Energy Commission
under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

LBL-2700
e.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Carbon-13 Nuclear Magnetic Resonance
in Solid Ammonium Tartrate^{*}

A. Pines, J. J. Chang and R. G. Griffin[†]

Department of Chemistry, University of California

and

Inorganic Materials Research Division

Lawrence Berkeley Laboratory

Berkeley, California 94720

* Supported in part by a Cottrell Research Grant from Research Corporation and in part by the National Science Foundation.

† Visitor of the Inorganic Materials Research Division.

Permanent address: Francis Bitter National Magnet Laboratory,
Cambridge, Massachusetts 02139.

ABSTRACT

Chemical shielding tensors have been determined for ^{13}C in single crystals of ammonium D-tartrate using proton-enhanced nmr. The unit cell contains two molecules, and in some orientations eight lines are resolved from the eight magnetically inequivalent carbon nuclei. Details are provided of the experimental approach, equipment and analysis, and results are presented for three determinations on two single crystals. These indicate a precision of ± 2 ppm in the determination of the principal values and $\pm 3^\circ$ in the orientation of the principal axes for the carboxyl groups, with lower precision for the hydroxyl groups. The orientation of the principal axes of the tensors are related to the crystallographic (a, b, c^*) axes by employing orientation data from X-ray precession work in this study. The most plausible assignment of the tensors to particular ^{13}C nuclei in the unit cell indicates that the heaviest shielding for ^{13}C in the carboxyl groups occurs when H_0 is perpendicular to the carboxyl plane and the lightest shielding when H_0 is in the carboxyl plane bisecting the C-O bonds. For the hydroxyl ^{13}C , the heaviest shielding occurs with H_0 roughly along the C-O bond. The average principal shielding values are, for the carboxyl ^{13}C : $\sigma_{11} = -110.5$, $\sigma_{22} = -61.7$, $\sigma_{33} = 21.6$, and for the hydroxyl ^{13}C : $\sigma_{11} = 41.9$, $\sigma_{22} = 48.0$, $\sigma_{33} = 74.0$, in ppm (± 2) relative to an external reference of liquid benzene. The effects of hydrogen bonding on the tensors are discussed briefly.

I. INTRODUCTION

Nuclear magnetic resonance of ^{13}C is now accepted as a powerful tool for the study of structure and dynamics in liquids. (1) The advantages of observing ^{13}C rather than other nuclei such as ^1H , are well established, in particular the absence of magnetic couplings between the ^{13}C nuclei and the larger range of chemical shifts. This yields high resolution, easily interpretable spectra even for large molecules with many carbon nuclei. In contrast, the usefulness of ^{13}C nmr in solids to date has not been outstanding. In particular, this is due to extremely poor resolution and sensitivity. Despite the early realization that a great deal of information is potentially hidden in solid state spectra, it is only recently that such spectra have begun to emerge.

The quantity of prime interest at this stage is the chemical shielding tensor, of which only the trace is determined in liquid studies. (1,2) An expectation of both theoreticians and experimentalists approaching these problems is that studies of these tensors will aid in understanding magnetic shielding of nuclei and in obtaining orientational and structural information which is available only with difficulty or unavailable otherwise.

Several approaches have been taken to extracting high resolution ^{13}C nmr spectra from solids. All of them encounter the same problem, namely that of dipolar broadening by ^1H

spins normally present, except in special isolated cases.^(3,4)

In one approach, the dipolar broadening is reduced by substituting ^2H for ^1H .⁽⁵⁾ The latter has a much smaller magnetogyric ratio, but still leaves a problematic amount of broadening and will not yield very well resolved ^{13}C spectra for complex systems. This can be overcome by selective enrichment of ^{13}C , but introduces formidable and expensive preparative difficulties for general single crystal applications. Another approach employed by Schaefer and co-workers⁽⁶⁾ involves rapid rotation of samples at the 'magic angle', reducing the dipolar broadening.⁽⁷⁾ However, this also removes the chemical shielding anisotropy of ^{13}C , leaving only the trace of the tensors observable.

The most promising approach appears to be that of $^1\text{H} - ^{13}\text{C}$ nuclear magnetic double resonance, introduced recently in several versions.⁽⁸⁻¹⁰⁾ We have found the approach of direct detection with sensitivity enhancement offered by proton-enhanced nuclear induction spectroscopy⁽⁸⁾ to be most convenient and widely applicable. Using this approach, we have determined several ^{13}C tensors in molecules with carbon-oxygen bonds and from these data a picture of trends for particular functional groups is beginning to emerge. In this paper, we report details of the work in single crystals of ammonium D-tartrate. This material was selected because of several appealing features:

- (a) large single crystals can be grown with facility,
- (b) the crystals structure is known,

- (c) it contains two functional groups of interest, C-OH and COO⁻, for which full tensors have not been determined previously,
- (d) the spectra and analysis are complex enough to indicate the types of problems encountered in those experiments,
- (e) the analysis indicates a favorable case where the shielding tensors are related to the local symmetry at the ¹³C sites.

In section II a brief reminder of the basis for the double resonance approach is presented and in section III details of the experimental apparatus and analysis are described. Results are presented for work on single crystals in section IV, where the shielding values and orientational data are discussed and an indication of experimental precision is given.

II. CARBON-13-PROTON DOUBLE RESONANCE

In this section we describe briefly the approach employed and some sensitivity considerations. Detailed discussions of sensitivity enhancement in double resonance experiments have been provided by several authors.⁽⁸⁻¹³⁾ There are basically two difficulties with the extraction of high resolution ¹³C nmr spectra from solid samples, sensitivity and resolution. The problem of sensitivity is a perennial one and is encountered also in normal ¹³C nmr in liquids. In the case of solids, the problem is even more acute since resolution is lower, more lines are found and spin-lattice relaxation times are normally long.

The loss in resolution comes about primarily due to magnetic dipolar coupling of the ^{13}C nuclei to abundant ^1H nuclei usually present in the sample. For these reasons, until recently, cases of resolved solid state ^{13}C spectra were reported employing conventional techniques only on samples devoid of protons.

The double resonance techniques are based on concepts originally introduced by Bloch⁽¹⁴⁾ and Hahn and co-workers,⁽¹¹⁾ and have alleviated these problems to a large extent. Our approach is depicted schematically in figure 1 where the I spins are abundant (here ^1H) and the S spins are of rare nuclei (here ^{13}C) under observation. Sensitivity enhancement is effected in the 'cross-polarization' step by transferring nuclear magnetic polarization from the abundant to the rare spins. Subsequently, the resolution is enhanced by intense irradiation of the I nuclei at resonance, inducing spin decoupling during the observation of the S spin free precession. This process can be repeated until the I polarization is depleted, resulting in a large sensitivity enhancement over conventional Fourier transform spectroscopy.

To review briefly the considerations of sensitivity enhancement we consider two limiting versions of the experiment depicted in figure 2. The first is the approach we have employed most commonly, referred to as multiple cross-polarization. In this approach, the I spins, after reaching equilibrium in the magnetic field with a magnetization $M_I^{(0)}$ are spin-locked at resonance with an intense field of intensity H_{1I} . The cross-polarization

is effected by irradiating the S spins with a field H_{1s} such that the Hartmann-Hahn condition:

$$\gamma_I H_{1I} = \gamma_S H_{1s} \quad (1)$$

is satisfied; cross-relaxation is most efficient under these conditions. The S spin magnetization after this cross-polarization step is given by:

$$M_s^{(1)} = \left(\frac{\gamma_I}{\gamma_S} \right) M_s^{(0)} \quad (2)$$

where $M_s^{(0)}$ is the magnetization with the S spins in equilibrium with the lattice. The S spin free induction decay is now observed while the irradiation of the I spins continues and the whole process can be continued with the S signal accumulated and ultimately Fourier transformed. The optimal number N of cross-polarization steps depends on $T_{1\rho}$, the spin-lattice relaxation time of the I spins in the rotating frame. For the simple case $T_{1\rho} \rightarrow \infty$, the optimal enhancement in signal/noise voltage is realized for $N \sqrt{\frac{N_S}{N_I}} \approx 1$ for which

$$\left(\frac{S}{N} \right)_{MCP} \approx 0.6 \frac{\gamma_I}{\gamma_S} \sqrt{\frac{N_I}{N_S}} \left(\frac{S}{N} \right)^{(0)} \quad (3)$$

where N_I , N_S are the numbers of I and S spins, and $\left(\frac{S}{N} \right)^{(0)}$ is the voltage signal/noise for a normal ^{13}C free induction decay.

The advantages of this approach are its simplicity and the fact that the number of contacts can be adjusted optimally for $T_{1\rho}$. The main disadvantage in the case of long $T_{1\rho}$ is that in order to realize the full sensitivity enhancement,

prolonged irradiation of the I spins is necessary (for N cross-polarization and accumulation steps) and the problem of power dissipation becomes acute. For many cases, however, a single cross-polarization with the enhancement of equation (2) suffices and does not introduce severe problems of power dissipation.

The second limiting case, depicted schematically in figure 2(b) is the one we have referred to as 'total cross-polarization'.^(8,15) Here, by applying large H_{1S} fields, most or all of the 1H polarization can be transferred to the S spins in one step. However, since the Hartmann-Hahn condition is not satisfied in this case, the cross-relaxation is slow so that this is not always a profitable venture. The simplest version is that shown in the figure. Here, the 1H system is first adiabatically demagnetized in the rotating frame and the ^{13}C nuclei are then polarized by turning on a strong radiofrequency field near resonance. The ^{13}C polarization achievable is

$$M_S^{(1)} = \alpha \frac{\gamma_I}{\gamma_S} \sqrt{\frac{N_I}{N_S}} M_S^{(0)} \quad (4)$$

where $\alpha = 1$ if the ^{13}C field is turned on adiabatically and $\alpha = 1/2$ if it is turned on suddenly.

For cases where spin-lattice relaxation times are long and it is desirable to realize the full sensitivity enhancement, then this approach is useful as long as T_{IS} does not become excessively long, since the full experimental time involves

one cross-polarization and one observation. Pines and Shattuck⁽¹⁵⁾ have shown that $T_{IS} \sim 1$ sec for H_{1s} fields yielding large polarizations ($\times 15$ in adamantane even with $T_{1d} \sim 100$ msec) and thus this approach can be profitable.

For the present experiments, several approaches were investigated. Since $T_{1\rho}$ and T_{1d} for the 1H spins were found to be short the multiple cross-polarization was used with a small number of cross-polarization steps, normally one.

III. EXPERIMENTAL

Since this paper describes in detail the first in a series of experiments using our spectrometer on carbon-13 nmr in single crystals, we present more details on the experimental procedure and on the equipment required for the experiments. In addition, this section presents some details of the analysis employed to extract the chemical shielding tensors and assign them to molecules and bond directions. In general, our experimental procedure consists of the following steps:

- a) mount the crystal in a well machined reorientable cube with assigned axes (X_c, Y_c, Z_c).
- b) orient the crystal employing X-ray techniques. This gives the crystallographic axes (a, b, c) relative to the cube axes (X_c, Y_c, Z_c).
- c) obtain nmr spectra for orientations of the crystal about (X_c, Y_c, Z_c) perpendicular to H_0 .

- d) analyze the rotation plots to obtain the shielding tensors σ_c in the cube axes.
- e) diagonalize σ_c , yielding σ in the principal axes system (1, 2, 3). This gives the eigenvalues σ_{ii} and the orientation of (1, 2, 3) relative to (X_c , Y_c , Z_c) and thus relative to (a, b, c).
- f) assign the tensors σ to particular ^{13}C nuclei in the unit cell. This final step is not rigorous and involves some assumptions. However, this is the most interesting and valuable part of the experiment, particularly at this stage where so little is known about ^{13}C shielding tensors.

We describe in turn some details of the apparatus and analysis which permit the above procedure to be followed efficiently. This includes a description of the spectrometer, the probe and goniometer, sample orientation and data analysis.

Spectrometer

The spectrometer is a wideband double resonance device⁽¹⁷⁾ which operates in heterodyne mode with an intermediate frequency of 30 MHz. A schematic diagram is shown in figure 3. In principle, it is a double resonance version of the multiple-pulse spectrometer described elsewhere.⁽¹⁸⁾ The magnet is a wide bore superconducting solenoid. The radiofrequency sections are mostly homebuilt from commercial hybrids and the digital section is constructed around an on-line PDP 8/e minicomputer.

The low level radiofrequency circuitry is separated from the power amplifiers to facilitate shielding of the detectors from the intense radiation. The transmitter section contains four phase channels for each frequency with phase and amplitude adjustments. Special features of the detector section are good shielding and isolation, high sensitivity, wideband operation and flexibility. Frequency for the low level sections is selected by simply changing the source frequencies derived from synthesizers since all low level operation is wideband.

Power amplifiers are tuned and consist mainly of modified radio amateur transmitters. These are equipped with homebuilt power supplies designed for stable operation during the prolonged rf irradiation. The ¹H final stage, for example, is a modified Millen amplifier which operates in class C with a double glass tetrode. Anode voltage is 2000 V derived from a regulated power supply stable to 0.1%.

The digital section contains the minicomputer, pulse programmer, and a data acquisition system described briefly elsewhere. ⁽¹⁹⁾ The latter is exceedingly simple and employs the computer directly as a fast real time signal averager through the 'direct add to memory' option. The pulse programmer is a homebuilt device controlled by the computer. It operates asynchronously with an independent memory and with facilities to control sixteen pulse channels and two channels for adiabatic demagnetization and remagnetization. The computer handles pulse programming, data acquisition, data storage and filing, display, data analysis, etc.

An example of the sensitivity enhancement provided by the spectrometer is given in figure 4. This provides a demonstration of the accumulative cross-polarization processes for ^1H - ^{13}C double resonance in a polycrystalline sample of adamantane. For the present experiments on ammonium D-tartrate, good signal/noise was obtained by averaging over several ^1H polarization cycles, typically in less than five minutes per spectrum.

Probe and Goniometer

Figure 5 shows a schematic diagram of the probe employed for the geometry of the superconducting solenoid, with details of the probe head containing the variable temperature sample chamber, radiofrequency coils and goniometer. The ^{13}C coil is a short horizontal solenoid of diameter 6 mm and the ^1H coils is wound orthogonal in a Helmholtz configuration. The double coil is mounted on leads supported in a Kel-F base, and connected to rigid coaxial cable via shielded tuning and matching circuits.

The goniometer consists of precision Kel-F gears with a shaft rotatable remotely. The lower end of the shaft is connected to a gear and calibrated dial permitting the angle of the upper gear to be read directly. The crystal holder is shown in the detail of figure 5(b) and is keyed onto the vertical gear so that its position and orientation are precisely reproducible. The cube containing the crystal is cemented onto the end of the crystal holder and is positioned in the center

of the coils when the holder is keyed in position and fixed with the set screws.

The tuning and matching electronics consist primarily of homebuilt capacitors to withstand the high radiofrequency voltages across the coils. These are housed beneath the Kel-F base in shielded tubes and consist of polished smooth copper tubes with Teflon dielectric. The capacitance is adjustable remotely as shown in the figure to permit tuning with the probe in position inside the magnet bore.

Sample Orientation

Crystals of ammonium D-tartrate were grown from aqueous solution by slow evaporation. The crystals were mounted in Kel-F cubes machined to match a crystal holder as shown in figure 5(b). The cubes holding the crystals are assigned axes (X_c , Y_c , Z_c). The crystal holders were keyed onto the end of an X-ray goniometer, maintaining the orientation of the crystal between the nmr goniometer and the X-ray equipment.

The crystal structure of ammonium D-tartrate has been determined previously:⁽²⁰⁾ space group $P2_1$, with two molecules in the unit cell. The crystallographic a , b , c axis were located in an X-ray precession experiment, and the orientation of (X_c , Y_c , Z_c) relative to the orthogonal system (a , b , c^*) were then determined directly.

Rotation Plots and Data Analysis

The chemical shift σ is related to the zz component of the shielding tensor in the laboratory frame, where z is the direction

of the magnetic field. For a rotation about the axis k of the crystal cube, it is easily shown that: (21)

$$\sigma = \frac{1}{2} (\sigma_{ii} + \sigma_{jj}) - \frac{1}{2} (\sigma_{ij} + \sigma_{ji}) \sin 2\theta + (\sigma_{ii} - \sigma_{jj}) \cos 2\theta$$

where $\{i, j, k\} \equiv \{X_c, Y_c, Z_c\}$ with cyclic permutations. (5)

The coefficients containing the elements of $\underline{\sigma}$ are determined for each rotation by fitting the rotation data to equation (5) and from the three rotations, nine equations are obtained for the six parameters of the shielding tensor; the latter are extracted by least squares fitting. At this stage the shielding tensor elements (the symmetric part) in the cube axes are fully determined. This is now diagonalised by standard procedures to yield the principal values σ_{ii} and the orientation of the principal axes of $\underline{\sigma}$ in the cube axis system.

IV. RESULTS

As indicated previously, three independent sets of measurements were performed on two different crystals. Results are presented here for the three determinations to give an idea of the experimental precision available at this preliminary stage of the development of our apparatus and procedure. Several sources of error contribute towards the small fluctuations in results, including X-ray orientation, goniometer misalignment, errors in cube reorientation and spectral resolution. We describe first the limited results

of a powder spectrum and then provide details on spectra from crystals, orientation plots and determination of the tensors.

Powder Spectrum

Powder spectra are simple to obtain by our method and in favorable cases can yield directly the principal elements of the chemical shielding tensors.⁽²²⁾ Clearly, the orientational information is normally lost. In cases where there is severe overlap between lines, analysing the spectrum is not a trivial problem; it can be approached by several methods aside from full single crystal studies, including partial cross-relaxation, selective ^{13}C enrichment, and detailed lineshape analysis.

Figure 6 shows the ^{13}C proton-enhanced free induction decay and Fourier transform nmr spectrum obtained from a polycrystalline sample of ammonium D-tartrate. In this case, the assignment is simple, the two peaks arising from the carboxyl and hydroxyl carbon nuclei. The principal values of the shielding tensors extracted directly from this spectrum are given in table 1 and are compared later with values obtained from the single crystal work. Note that within experimental error the shielding tensor for the $^{13}\text{C-OH}$ groups is axially symmetric.

Single Crystal Spectra

The unit cell of ammonium D-tartrate contains two molecules, as shown in figure 7. In isotropic liquid solution only two

lines are observed from the magnetically inequivalent carboxyl and hydroxyl nuclei. In the crystal, all eight carbon nuclei can be magnetically inequivalent and we expect to see several lines. Figure 8 shows proton-enhanced ^{13}C spectra from a single crystal of ammonium D-tartrate in different orientations, displaying the resolution of lines from all the nuclei in the unit cell. The resolution is limited primarily by magnetic field inhomogeneity and by dipolar coupling to ^{14}N spins.

Orientation Dependence

As explained in section III a full analysis of the data is performed by taking spectra such as those in figure 8 for rotations about the cube axes (X_c , Y_c , Z_c). The orientation dependence of the line position for such a rotation on one of the crystals is given in figure 9. Similar data are obtained from the other determinations. The relevant information is provided in the caption. The data are fitted to equation (5) and the tensors in the cube axes are then determined by combining sets of lines from the three rotations. The sets of lines are easily assigned in this case by the requirement that they match up at equivalent orientations of the cube, and by the requirement that they provide 'reasonable' tensors (see following discussions). Due to problems of resolution and line assignment for the hydroxyl lines, the errors are much larger as manifested later in the orientation determination.

Shielding Tensors

Since there are eight inequivalent carbon nuclei in the unit cell (see figure 7) we expect to obtain eight different shielding tensors from the analysis. Employing the data from X-ray crystal orientation work, the principal axes are then known with respect to the crystallographic axes. This part of the analysis is rigorous and involves no assumptions. In the next section, assumptions are made to relate σ to the molecules, since clearly the tensors can be assigned in several ways to specific ^{13}C nuclei, subject only to the constraint that they must conform to the transformations of the crystallographic symmetry.

The shielding values are shown in table 2 for the four $^{13}\text{COO}^-$ and four ^{13}COH tensors determined by diagonalization of the tensors in the cube axis system. As mentioned previously, three independent experiments were performed, one experiment on one crystal and two on a second crystal. The three entries are for these three determinations and the average indicates an average over all twelve values for the four tensors and three determinations. It is hard to attach any significance to the difference between the tensors from similar carbon nuclei, since they appear to be in the range of experimental error. The precision is good (± 2 ppm) and the values are in agreement with the determination from the powder spectrum (figure 6, table 1) within experimental error.

Of particular interest is the orientation of the principal axes of σ . This is summarized in table 3 for the three determinations,

as direction cosines of (1, 2, 3) relative (a, b, c^{*}), for each of the eight tensors.

Assignment to Specific ¹³C Nuclei

The only rigorous orientational output of our experiment is the orientation of σ in the crystallographic axes. However, one of the reasons for undertaking this whole project is to see if σ is related to local structure and symmetry. For an isolated COO⁻ group with equal C-O bonds we expect one axis of σ to be perpendicular to the plane, one to bisect the CO bonds and the third perpendicular to these two. For an isolated CH₃O group analogous to the hydroxyl carbon, we expect a unique axis along the CO bond and two others degenerate and perpendicular. Indeed, one assignment of tensors to carbon nuclei conforms quite closely to these requirements. Our assumption that this is the correct assignment is strengthened by other experiments in which one assignment yields the heaviest shielding for ¹³COO⁻ when H₀ is perpendicular to the plane. (23,24) The orientations are summarized in table 4 as angles between the principal axes of the tensors and bonds to the ¹³C to which they are assigned.

We see that with this assignment σ conforms reasonably well to the local symmetry with σ_{33} (COO⁻) being perpendicular to the COO⁻ plane and σ_{33} (COH) being along the CO bond. The orientation for the ¹³COO⁻ group is summarized in figure 10.

Comments

The shielding values and orientation of the tensors for the carboxyl groups are determined quite precisely in this study, with a reproducibility of ± 2 ppm and $\pm 3^\circ$ between the three determinations. The data for the hydroxyl groups are less satisfactory because of the limited resolution amongst the lines but are still instructive as far as the main features of the tensors are concerned. The determinations can be made considerably more precise by employing higher proton decoupling fields and a more uniform dc magnetic field.

Based on these results and others determined in our laboratory, several comments can be made in summary. For carboxyl groups the most shielded element (σ_{33}) of the ^{13}C tensor corresponds to an axis perpendicular to the $^{13}\text{COO}^-$ plane. For the carboxyl carbon in benzophenone, σ_{33} was also found to be perpendicular to the nodal plane of the CO π molecular orbital.⁽⁵⁾ The orientations of the other two axes of the tensor depend on structure. It is interesting to observe the trend in orientation of the tensor as we proceed from the case of a symmetrical ketone to a symmetrical carboxyl group. This is depicted in figure 11 from results on benzophenone,⁽⁵⁾ dimethyl oxalate⁽²³⁾ and other work on carboxyl groups in our laboratory. Since the difference between (c) and (d) derives from the effects of hydrogen bonding, we feel that ^{13}C shielding tensors will provide a sensitive tool for the study of such bonding.

The shielding values for the hydroxyl carbon are in agreement with those determined for methanol;⁽²⁵⁾ thus, they appear to be characteristic for this type of functional group. The tensor conforms to the local symmetry of an isolated R_3CO group with the heaviest shielding along the CO bond.

REFERENCES

1. J. B. Stothers, Carbon-13 NMR Spectroscopy, (Academic Press, New York, 1972); George C. Levy and Gordon L. Nelson, Carbon-13 Nuclear Magnetic Resonance for Organic Chemists (Wiley-Interscience, 1972).
2. F. Ramsey, Phys. Rev. 78, 699 (1950); J. W. Emsley, P. Feeney, and L. H. Sutcliffe, High Resolution NMR Spectroscopy, (Pergamon, London, 1967); R. Ditchfield and P. D. Ellis, Topics in Carbon-13 NMR, Vol. 1, in press.
3. P. C. Lauterbur, Phys. Rev. Lett. 1, 343 (1958).
4. A. Pines, W. K. Rhim and J. S. Waugh, J. Chem. Phys. 54, 5438 (1971).
5. J. Kempf, H. W. Spiess, U. Haeberlen and H. Zimmermann, Chem. Phys. Lett. 17, 39 (1972).
6. J. Schaefer, presented at the 14th Experimental NMR Conference, Boulder, Colorado, 1973.
7. E. R. Andrew in Progress in Nuclear Magnetic Resonance Spectroscopy, Vol. 8, ed. by J. W. Emsley, J. Feeney and L. H. Sutcliffe (Pergamon Press, 1971).
8. A. Pines, M. G. Gibby and J. S. Waugh, J. Chem. Phys. 56, 1776 (1972); A. Pines, M. G. Gibby and J. S. Waugh, J. Chem. Phys. 59, 569 (1973).
9. H. E. Bleich and A. G. Redfield, J. Chem. Phys. 55, 5405 (1971); C. S. Yannoni and H. E. Bleich, J. Chem. Phys. 55, 5401 (1971).

10. P. Mansfield and D. K. Grannell, J. Phys. C., L177 (1971).
11. S. R. Hartmann and E. L. Hahn, Phys. Rev. 128, 2042 (1962);
D. A. McArthur, E. L. Hahn and R. Walstedt, Phys. Rev. 188,
609 (1969).
12. E. P. Jones and S. R. Hartmann, Phys. Rev. B. 6, 757 (1972).
13. F. M. Lurie and C. P. Slichter, Phys. Rev. 133, A1108 (1964).
14. F. Bloch, Phys. Rev. 111, 841 (1958); L. R. Sarles and R. M.
Cotts, Phys. Rev. 111, 853 (1958).
15. A. Pines and T. W. Shattuck, Chem. Phys. Lett. 23, 614 (1973);
A. Pines, Proceedings of the First Specialized Colloque Ampère,
Krakow, Poland, August 1973, in press.
16. A. Pines and T. W. Shattuck, to be published; A. Pines and T. W.
Shattuck, presented at the Philadelphia APS meeting, March,
1974, Bull. Am. Phys. Soc. 19, 351 (1974).
17. A. Pines and D. D. Wilkinson, to be published.
18. J. D. Ellett, Jr., et al., Adv. Magn. Reson. 5, 117 (1971).
19. T. Merrick, S. Smiriga and A. Pines, J. Mag. Res., in press.
20. V. S. Yadova and V. M. Padnamabhan, Acta. Cryst. B29, 493
(1973).
21. M. Mehring, R. G. Griffin and J. S. Waugh, J. Chem. Phys.
55, 746 (1971).
22. N. Bloembergen and T. J. Rowland, Phys. Rev. 97, 1679 (1955);
N. Bloembergen and T. J. Rowland, Acta Metall. 1, 731 (1953).
23. A. Pines and E. Abramson, to be published.
24. A. Pines, paper presented at the Fifteenth Experimental NMR
Conference, Raleigh, N. C., April 1974.

25. A. Pines, M. G. Gibby and J. S. Waugh, Chem. Phys. Lett.
15, 373 (1972).

ACKNOWLEDGMENT

We are indebted to D. D. Wilkinson and D. N. Shirley for extensive assistance with the equipment and computer programs and to Dr. A. Zalkin and Dr. G. Chapuis for assistance with the X-ray precession work. A. P. is particularly grateful to Professor D. H. Templeton for some valuable conversations on shielding tensors and crystal structure. Financial support from the National Science Foundation, Research Corporation, and the U. S. Atomic Energy Commission through the Inorganic Materials Research Division of the Lawrence Berkeley Laboratory is gratefully acknowledged.

Table 1

Elements of σ in Ammonium D-tartrate from Powder Spectrum, in ppm
Relative to External Reference of Liquid Benzene^(a).

	σ_{11}	σ_{22}	σ_{33}
$^{13}\text{COO}^-$	-116.0	-70.0	15.5
^{13}COH	39.2	39.2	72.1

(a) Error Range is ± 6 ppm

Table 2

Elements of σ from Single Crystal Work, in ppm Relative to
External Reference of Liquid Benzene.

		<u>Tensor 1</u>	<u>Tensor 2</u>	<u>Tensor 3</u>	<u>Tensor 4</u>	<u>Average</u>
$^{13}\text{COO}^-$	σ_{11}	-110.1	-107.4	-110.8	-109.6	-110.5
		-111.2	-105.1	-109.4	-109.4	
		-115.8	-114.4	-112.0	-110.2	
	σ_{22}	-66.4	-57.8	-62.2	-56.5	-61.7
		-63.7	-58.7	-63.4	-56.3	
		-68.1	-61.6	-65.3	-59.8	
	σ_{33}	26.2	20.2	23.3	22.1	21.6
		22.7	19.9	21.8	21.7	
		22.5	18.9	18.2	21.2	

		<u>Tensor 5</u>	<u>Tensor 6</u>	<u>Tensor 7</u>	<u>Tensor 8</u>	
^{13}COH	σ_{11}	46.0	42.6	41.5	44.0	41.9
		41.2	43.8	42.7	42.4	
		39.9	40.9	39.0	39.0	
	σ_{22}	49.7	49.5	50.6	45.9	48.0
		49.7	46.2	49.9	47.9	
		46.8	47.2	46.8	45.4	
	σ_{33}	75.9	74.8	77.7	76.0	74.0
		76.0	74.3	74.7	73.7	
		70.9	70.1	72.8	70.7	

Table 3

Direction Cosines of Principal Axes of Tensors Relative to Crystallographic Axes

		Tensor 1			Tensor 2			Tensor 3			Tensor 4		
		a	b	c*	a	b	c*	a	b	c*	a	b	c*
¹³ COO-	σ ₁₁	-.8062	-.5305	-.2618	.7996	.5840	.1399	.8671	-.3776	.3249	-.8833	.4528	-.3185
		-.8099	-.4883	-.3250	.8164	.5312	.2266	.8399	-.4359	.3235	-.7863	.5383	-.3031
		-.8281	-.4658	-.3120	.8045	.5348	.2585	.8766	-.4206	.2349	-.8254	.5271	-.2025
	σ ₂₂	.5653	-.8213	-.0768	-.4005	.3448	.8500	-.4446	-.8806	.1632	.4385	.1882	-.8788
		.5488	-.8264	-.1258	-.4133	.2631	.8719	-.4871	-.8683	.0947	.4752	.2135	-.8536
		.5398	-.8129	-.2191	-.4277	.2198	.8768	-.4509	-.8875	.0950	.4014	.2955	-.8670
	σ ₃₃	.1743	.2099	-.9620	-.4477	.7349	-.5094	-.2246	.2860	.9317	.3380	.8716	.3552
		.2072	.2802	-.9373	-.4036	.8053	-.4342	-.2396	.2371	.9415	.3947	.8152	.4238
		.1516	.3498	-.9244	-.4121	.8159	.4055	-.1670	.1885	.9678	.3971	.7968	.4554
		Tensor 5			Tensor 6			Tensor 7			Tensor 8		
		a	b	c*	a	b	c*	a	b	c*	a	b	c*
¹³ COH	σ ₁₁	.4357	.8395	-.3247	-.8458	-.4646	.2625	-.5170	.8447	.1384	.9807	-.1372	-.1393
		.4689	.8794	-.0818	-.9928	-.0925	.0758	-.4237	.9014	.0891	.8663	-.4354	-.2448
		.4860	.8722	-.0547	-.9964	-.0050	.0854	-.4635	.8837	.0645	.8094	-.5081	-.2943
	σ ₂₂	-.1578	.4264	.8907	-.5336	.7387	-.4118	.0621	.1982	-.9782	.1768	.9267	.3318
		-.1358	.1633	.9772	-.1165	.8908	-.4394	.1123	.1500	-.9823	.4994	.7658	.4051
		-.0888	.1149	.9898	-.0355	.9320	-.3608	.1058	.1274	-.9861	.5871	.7107	.3877
	σ ₃₃	.8862	-.3368	.3182	-.0026	-.4884	-.8776	-.8539	-.4971	-.1550	.0836	-.3501	.9330
		.8728	-.4471	.1959	-.0269	-.4451	-.8951	-.8989	-.4062	-.1648	.0111	-.4732	.8808
		.8695	-.4762	.1316	-.0778	-.3625	-.9288	-.8797	-.4503	-.1526	.0122	-.4866	.8736

Table 4

Orientation of Principal Axes of ^{13}C Shielding Tensors Relative to Molecular Bond Directions*

		Tensor 1			Tensor 2			Tensor 3			Tensor 4		
Bond		$\text{C}_5-^{13}\text{C}_1$ (a)	$\parallel(^{13}\text{COO})_1$ (b)	$\perp(^{13}\text{COO})_1$ (c)	$\text{C}_6-^{13}\text{C}_2$	$\parallel(^{13}\text{COO})_2$	$\perp(^{13}\text{COO})_2$	$\text{C}_7-^{13}\text{C}_3$	$\parallel(^{13}\text{COO})_3$	$\perp(^{13}\text{COO})_3$	$\text{C}_8-^{13}\text{C}_4$	$\parallel(^{13}\text{COO})_4$	$\perp(^{13}\text{COO})_4$
$^{13}\text{COO}^-$	σ_{11}	3.2	86.9	87.9	7.2	86.1	39.8	7.9	95.7	82.9	5.7	86.9	100.5
		4.7	88.6	83.9	1.3	90.4	94.5	5.3	92.1	83.5	4.4	85.7	95.1
		3.9	90.4	84.5	1.9	88.4	94.7	4.3	94.5	88.5	2.3	91.9	94.5
	σ_{22}	93.1	3.6	92.0	87.0	8.8	81.7	84.4	5.7	88.8	93.0	3.4	88.3
		91.4	1.6	89.4	89.7	4.5	85.5	87.8	3.1	92.1	94.4	6.0	85.5
		90.1	5.8	84.2	91.6	3.2	87.1	85.5	5.3	92.6	88.3	6.2	84.3
	σ_{33}	90.6	88.1	2.9	96.5	98.3	8.3	95.6	90.5	7.1	85.2	91.2	10.7
		94.5	90.7	6.1	91.3	94.6	6.4	94.8	87.6	6.8	90.5	94.1	6.8
		93.9	95.8	8.1	91.0	92.8	5.6	89.6	87.3	2.9	91.6	95.9	7.3

		Tensor 5			Tensor 6			Tensor 7			Tensor 8		
Bond		$^{13}\text{C}_5-\text{O}_2$ (d)	$^{13}\text{C}_5-\text{C}_1$	$^{13}\text{C}_5-\text{C}_6$	$^{13}\text{C}_6-\text{O}_6$	$^{13}\text{C}_6-\text{C}_2$	$\text{C}_5-^{13}\text{C}_6$	$^{13}\text{C}_7-\text{O}_7$	$^{13}\text{C}_7-\text{C}_3$	$^{13}\text{C}_7-\text{C}_8$	$^{13}\text{C}_8-\text{O}_8$	$^{13}\text{C}_8-\text{C}_7$	$\text{C}_6-^{13}\text{C}_8$
^{13}COH	σ_{11}	103.0	133.9	26.6	96.0	150.7	132.7	98.6	144.2	38.3	99.5	147.6	113.3
		101.7	143.3	40.6	97.5	147.9	108.8	104.6	140.6	39.6	96.0	151.4	130.5
		100.6	145.0	42.3	100.4	143.4	105.7	102.1	143.4	41.5	96.2	149.0	136.3
	σ_{22}	108.3	145.3	107.2	92.5	99.1	46.9	95.0	106.7	123.8	99.3	75.5	33.7
		97.0	132.6	102.0	92.6	75.2	26.2	95.5	102.8	127.1	93.3	96.6	45.2
		93.8	103.0	129.0	87.1	68.7	27.8	94.4	102.6	128.3	93.3	102.2	50.5
	σ_{33}	22.7	131.0	105.6	6.5	117.6	74.4	10.0	120.7	105.8	13.4	118.2	67.0
		13.7	124.3	105.2	8.0	117.8	71.9	15.7	126.5	102.0	6.9	117.7	73.5
		11.0	121.9	103.8	12.7	118.3	67.7	12.8	123.7	103.5	7.0	118.0	74.4

* The numbering of the atoms corresponds to that in Figure 7. The three entries are from the three independent determinations.

(a) $\text{C}_5-^{13}\text{C}_1$ is along the $\text{C}_5-^{13}\text{C}_1$ bond bisecting the $^{13}\text{C}_5\text{O}_2$ bonds.(b) $\parallel(^{13}\text{COO})_1$ is in the $^{13}\text{C}_5\text{O}_2\text{O}_1$ plane perpendicular to the $\text{C}_5-^{13}\text{C}_1$ bond.(c) $\perp(^{13}\text{COO})_1$ is perpendicular to the $^{13}\text{C}_5\text{O}_2\text{O}_1$ plane.(d) $^{13}\text{C}_5-\text{O}_2$ is along the hydroxyl carbon $^{13}\text{C}_5-\text{O}_2$ bond.

FIGURE CAPTIONS

Figure 1 General scheme for high sensitivity and high resolution ^{13}C nmr in solids using proton-enhanced nuclear induction spectroscopy.

Figure 2 Schemes for multiple cross-polarization and single total cross-polarization versions of double resonance. (a) In this version the Hartmann-Hahn condition is matched and the ^1H polarization is transferred in steps to the ^{13}C nuclei, the ^{13}C free precession signal being accumulated during ^1H irradiation. (b) Here the ^{13}C nuclei are polarized with a large H_1 field from the ^1H system after adiabatic demagnetization in the rotating frame. This yields larger ^{13}C polarizations in one step but the cross-polarization times are longer.

Figure 3 Schematic diagram of double-resonance spectrometer.

Figure 4 Demonstration of sensitivity enhancement for carbon-13 spins. The oscillograph photographs on the left depict the output of the ^{13}C phase detector for various numbers (n) of $^1\text{H} \rightarrow ^{13}\text{C}$ cross-polarization steps in solid adamantane. The positive pulses are from rf receiver blocking during the cross-polarization, and they are followed by proton-decoupled ^{13}C free precession signals. It is these which are rapidly transferred and accumulated by the computer. The cross-polarization pluses are 5 msec long. The traces on the right show the accumulated

signals and acquisition times corresponding to the number of cross-polarization steps on the left. These were recorded further away from resonance.

Figure 5 Schematic diagram of double-resonance probe and goniometer.

Figure 6 Fourier transform proton-enhanced ^{13}C nmr spectrum in polycrystalline ammonium D-tartrate at 23° C. The horizontal scale is in ppm relative to an external reference of liquid benzene. The low field peak is from the carboxyl carbon nuclei and the high field from hydroxyl carbon nuclei; both display an anisotropy of the chemical shift. Values for the principal elements of the shielding tensors extracted from this spectrum appear in table 1.

Figure 7 Perspective view along the crystallographic b axis of two tartrate ions, showing the carbon and oxygen atoms in the unit cell of ammonium D-tartrate. The labelling of the atoms corresponds to that used in the discussion of tensor orientations relative to molecular bond directions.

Figure 8 Fourier transform proton-enhanced ^{13}C nmr spectra of single crystal of ammonium D-tartrate in three different orientations. The horizontal scale is relative to an external reference of liquid benzene.

Figure 9 Dependence of line positions in ^{13}C spectra of ammonium D-tartrate on crystal orientation about the crystallographic axis.

Figure 10 Orientation of principal axes of ^{13}C chemical shielding tensor in carboxyl groups of ammonium D-tartrate. The ^{13}C nucleus is most shielded when H_0 is perpendicular to the $^{13}\text{COO}^-$ plane and least shielded when H_0 is along the $\text{C}-^{13}\text{C}$ bond bisecting the ^{13}CO bonds.

Figure 11 Trends in the orientation of the principal axes of the ^{13}C chemical shielding tensor for various carbonyl groups. The most shielded axis, corresponding to σ_{33} , is perpendicular to the planes of the groups shown. As we proceed from the symmetrical ketone (a) through an ester group (b), an unsymmetrical (c) and a symmetrical (d) carboxyl group, axes 1 and 2 rotate systematically in the plane. The cases (a) - (d) are exemplified respectively by benzophenone, dimethyl oxalate, ammonium hydrogen malonate and the diammonium salts of oxalic and tartaric acids.

a

b

XBL 7312-7169

Fig. 1

XBL 743-5732

Fig. 2

XBL743-5734

Fig. 3

XBB 742-900

Fig. 4

a

b

XBL 743-5752

Fig. 5

XBL 743-5694

Fig. 6

XBL 743-5733

Fig. 7

XBL 743-5693

Fig. 8

XBL743-5729

Fig. 9

XBL 742-5591

Fig. 10

XBL 743-5784

Fig. 11

—LEGAL NOTICE—

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720