

Lawrence Berkeley National Laboratory

Recent Work

Title

DISILANYLPHOSPHINE AND BISILYLPHOSPHINE

Permalink

<https://escholarship.org/uc/item/498965mf>

Authors

Gokhale, Sudarshan D.
Jolly, William L.

Publication Date

1964-03-01

UCRL-11342

University of California
Ernest O. Lawrence
Radiation Laboratory

DISILANYLPHOSPHINE AND DISILYLPHOSPHINE

Sudarshan D. Gokhale and William L. Jolly

March 1964

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California
AEC Contract No. W-7405-eng-48

DISILANYLPHOSPHINE AND DISILYLPHOSPHINE

Sudarshan D. Gokhale and William L. Jolly

March 1964

Contribution from the Department of Chemistry, and
Inorganic Materials Research Division of the Lawrence
Radiation Laboratory, University of California
Berkeley, California

Disilanylphosphine and Disilylphosphine

By Sudarshan D. Gokhale¹ and William L. Jolly

Disilanylphosphine, $\text{Si}_2\text{H}_5\text{PH}_2$, and disilylphosphine, $(\text{SiH}_3)_2\text{PH}$, have been prepared and identified by nmr spectroscopy. Disilanylphosphine has been isolated and characterized by determination of its infrared spectrum, its vapor pressure, and the products of its hydrolysis and pyrolysis.

(1) Permanent Address: Chemistry Division, Atomic Energy Establishment,
Trombay, Bombay.


Introduction

Silylphosphine, SiH_3PH_2 , was first prepared by the pyrolysis of a mixture of silane and phosphine.² Trisilylphosphine, $(\text{SiH}_3)_3\text{P}$, was

(2) G. Fritz, Z. Naturforsch., 8b, 776 (1953); Z. anorg. allgem. chem., 28, 332 (1955).

prepared by the reaction between bromosilane and potassium dihydrogen phosphide:³

(3) E. Amberger and H. Boeters, Angew. Chem., 74, 32 (1962).



Both silylphosphine and a material of empirical composition Si_2PH_7 were detected in the products of the decomposition of a mixture of silane and phosphine in a silent electric discharge.⁴ The compound Si_2PH_7

(4) J. E. Drake and W. L. Jolly, Chem. and Ind., 1470 (1962).

intrigued us because of the possibility of the existence of two structural isomers. The purpose of the present investigation was to prepare macroscopic amounts of the material and, if possible, to identify, separate, and characterize the isomers.

Experimental

A mixture of silane and phosphine in a mole ratio of 2:1 was circulated in a closed system through an ozonizer-type electric discharge tube at -78° followed by a trap at -95° (toluene slush) until practically all the reaction gases were consumed. A similar method had been used earlier for the preparation of higher silanes,^{5,6} germanes,⁷

-
- (5) E. J. Spanier and A. G. MacDiarmid, Inorg. Chem., 1, 432 (1962).
(6) S. D. Gokhale and W. L. Jolly, UCRL Report 11078, November 1963.
(7) J. E. Drake and W. L. Jolly, Proc. Chem. Soc., 379 (1961); J. Chem. Soc., 2807 (1962).

and "mixed" (ternary) hydrides.^{4,8} The products of the reaction were

-
- (3) E. J. Spanier and A. G. MacDiarmid, Inorg. Chem., 2, 215 (1963).

first subjected to fractional condensation in a series of traps at -63° (chloroform slush), -95° (toluene slush), -130° (n-pentane slush) and -196° (liquid nitrogen). The unreacted silane and phosphine condensed in the -196° trap, and disilane and silylphosphine condensed in the -130° trap. Diphosphine, trisilane and part of the Si_2PH_7 condensed in the -95° trap. The trap at -63° contained the higher silanes, higher ternary hydrides, and the remainder of the Si_2PH_7 . The condensate in the -95° trap was allowed to stand at room temperature for a week, during which

time the diphosphine decomposed almost completely. After pumping out a little hydrogen which had formed, the mixture was fractionally condensed in a trap at -83° (ethyl acetate slush) followed by a trap at -196° . Phosphine, trisilane, and only a trace of Si_2PH_7 collected in the -196° trap. The nmr spectrum of the material collected in the -83° trap showed clearly the presence of both $(\text{SiH}_3)_2\text{PH}$ and $\text{Si}_2\text{H}_5\text{PH}_2$ in approximately equal amounts, as well as some trisilane. The condensate in the -63° trap was passed through a -45° (chlorobenzene slush) in order to remove most of the higher molecular weight compounds from the Si_2PH_7 . The Si_2PH_7 -rich condensates were injected into a gas chromatographic column containing silicone oil on fire-brick at 70° . We were never able to elute disilylphosphine from the column; we assume that it was thermally decomposed on the column. The observed presence of small amounts of phosphine and silylphosphine in all the elution fractions is consistent with this interpretation. The method of injecting the mixture in the chromatographic apparatus and of collecting the different fractions was similar to that used in the preparation of the tetrasilanes.⁶ It was found that ordinary injection needles became clogged in spite of all reasonable precautions. Gas-tight syringes with Teflon-tipped plungers and 22- or 23-gauge needles were more satisfactory. When syringes with removable needles were used, a little grease had to be used at the joints to make them gas tight. The chromatograms showed that the retention times of Si_3H_8 and $\text{Si}_2\text{H}_5\text{PH}_2$ were in the proportion of roughly 1:2. The peak for $\text{Si}_2\text{H}_5\text{PH}_2$ precede those of the tetrasilanes. It was found that the $\text{SiH}_3\text{SiH}_2\text{PH}_2$ was contaminated with a small amount of $(\text{Si}_2\text{H}_5)_2\text{O}$. By keeping

the contaminated product at -78° in a trap and by pumping out the vapors into a trap at -196° , it was possible to recover pure disilanylphosphine. Starting with 250 mmoles of the $\text{SiH}_4 - \text{PH}_3$ mixture, about 3 mmoles of $\text{Si}_2\text{H}_5\text{PH}_2$ was obtained.

The nmr spectra were obtained with a Varian Associates A-60 nmr spectrometer, and the infrared spectra with Perkin-Elmer Infracord spectrophotometers (NaCl Model 137B and KBr Model 137). The nmr spectrum for $\text{SiH}_3\text{SiH}_2\text{PH}_2$ is given in Figure 1. Although $(\text{SiH}_3)_2\text{PH}$ was not obtained in pure form, the relevant portions of the spectrum of a mixture containing $(\text{SiH}_3)_2\text{PH}$ are given in Figures 2 and 3. The infrared spectrum of disilanylphosphine shows absorptions, as seen in Figure 4, at the following frequencies (cm^{-1}): 2300(w), 2150(s), 1070(sh,w), 990(w), 942(m), 932(m), 833(s), 877(s), 790(sh,s), 784(s), 736(w), 728(w), 724(w), 637(w), 628(w), and 475 to 515(w, broad).

The vapor pressure of disilanylphosphine was found to be 31 ± 1 mm at 0° and $31 + 1$ mm at 20° . It was found during the fractional condensation that the isomers $(\text{SiH}_3)_2\text{PH}$ and $\text{Si}_2\text{H}_5\text{PH}_2$ could not be separated from each other even partially. Thus we conclude that the volatilities of the two isomers are very similar. Gas-density molecular weights of 92.1, 95.5 and 96.0 were obtained for three different samples. (Calculated for Si_2PH_7 : 94.2) The mass spectrum of disilanylphosphine showed bands corresponding to the fragments $\text{Si}_2\text{P}^+ - \text{Si}_2\text{PH}_7^+$, $\text{Si}_2^+ - \text{Si}_2\text{H}_6^+$, $\text{SiP}^+ - \text{SiPH}_5^+$ (uncertain because of ambiguity with Si_2H_x^+), $\text{Si}^+ - \text{SiH}_4^+$, and $\text{P}^+ - \text{PH}_3^+$.

In several experiments, $\text{Si}_2\text{H}_5\text{PH}_2$ vapor was allowed to remain in contact with an excess of water vapor for several hours. Analysis showed that the main hydrolysis products were phosphine and bisdisilanyl oxide, and that

very small amounts of disilane and silane were formed. The glass walls of the reaction tubes became coated with a very thin white deposit. In one experiment 0.204 mmole of $\text{Si}_2\text{H}_5\text{PH}_2$ and 0.147 mmole of H_2O were allowed to react for 4 hours at room temperature, yielding 0.208 mmole of PH_3 , 0.126 mmole of a mixture of H_2O and $(\text{Si}_2\text{H}_5)_2\text{O}$ (it was not possible to separate these two), and a trace of Si_2H_6 and SiH_4 . The products were identified by their infrared spectra.

Traces of phosphine and disilane were formed when disilanylphosphine was stored for one day at room temperature while exposed to ordinary room light. When kept as the vapor in a sealed tube at either 50° or 110° for one day, about 5% of the compound decomposed to give phosphine and disilane. No other products were found. When kept at 210° for one day, 32% of the compound decomposed to give disilane, silylphosphine and (by mass spectroscopy) Si_3PH_9 and possibly $\text{Si}_2\text{P}_2\text{H}_8$. No phosphine or hydrogen was found in the 210° pyrolysis. A faint solid film formed on the inner wall of the container.

Discussion

Nmr spectra.- The chemical shift data are presented in Table I. When the nmr spectrum of trisilane⁹ is compared with that of disilanylphosphine

(9) S. D. Gokhale and W. L. Jolly, to be published.

(Figure 1), it is noticed that the relative positions of the resonances due to the SiH_3 and SiH_2 groups are reversed. The protons in the SiH_2 group of $\text{SiH}_3\text{SiH}_2\text{PH}_2$ are less shielded than those in the SiH_2 group of $\text{SiH}_3\text{SiH}_2\text{SiH}_3$

because phosphorus is more electronegative than silicon. For the same reason, the protons of the SiH_3 group in $(\text{SiH}_3)_2\text{PH}$ are less shielded than those of the SiH_3 group in $\text{SiH}_3\text{SiH}_2\text{PH}_2$. Although we cannot explain the fact that the SiH_3 protons in trisilane are less shielded than those in disilanylphosphine, this datum is consistent with the observed variations of the chemical shift of the SiH_3 group in disilanyl halides¹⁰ and the

(10) M. Abedini, C. H. Van Dyke, and A. G. MacDiarmid, unpublished data.

higher silanes.⁹ The proton of the PH group in $(\text{SiH}_3)_2\text{PH}$ is more shielded than the protons of the PH_2 group in $\text{Si}_2\text{H}_5\text{PH}_2$ because silicon is less electronegative than hydrogen.

The coupling constant data are presented in Table II. The SiH_3 proton signal of $\text{SiH}_3\text{SiH}_2\text{PH}_2$ is split into a pair of triplets by the SiH_2 protons and the phosphorus atom. This pattern is not clearly observable in the main resonance, but the ^{29}Si satellite resonance on the high-field side of the main resonance very clearly shows this pattern when seen at high gain. The SiH_2 proton signal of $\text{SiH}_3\text{SiH}_2\text{PH}_2$ is split into a pair of triplets, each line of which is split into a quartet, by the phosphorus nucleus and the two other types of protons in the molecule. It happens that the SiH_3SiH_2 coupling constant is approximately half of the SiH_2PH_2 coupling constant; therefore there is a near-superposition of lines, as shown in Figure 1, causing the signal to look something like a pair of octets. At lower sweep widths, the individual lines are resolved. The PH_2 proton signal of $\text{SiH}_3\text{SiH}_2\text{PH}_2$ is split into a pair of triplets by the phosphorus nucleus and the SiH_2 protons.

The SiH_3 proton signal of $(\text{SiH}_3)_2\text{PH}$ is split into a pair of doublets by the phosphorus atom and the PH proton. This spectrum is shown in Figure 2, in which the weaker peaks are due to $\text{Si}_2\text{H}_5\text{PH}_2$ impurity. The PH proton signal of $(\text{SiH}_3)_2\text{PH}$ is split into a pair of septets by the phosphorus nucleus and the six SiH_3 protons. Because of this multiple splitting of a weak signal, the lines do not show up prominently. With very high gain, it is possible to see five of the peaks of each septet. The high-field multiplet is shown in Figure 3. We observe an intensity ratio of 3.0 : 7.4 : 10 : 7.6 : 2.9, which corresponds much more closely to that expected for the middle five lines of a septet (3 : 7.5 : 10 : 7.5 : 3) than to that expected for a simple quintet (1.7 : 6.7 : 10 : 6.7 : 1.7).

The ^{31}P nmr spectrum of $\text{SiH}_3\text{SiH}_2\text{PH}_2$ was observed to be a triplet of triplets, the splitting being due to the PH_2 and SiH_2 protons. The P-H coupling constants were, within experimental error, the same as those obtained from the proton nmr spectrum.

Infrared Spectra.— The infrared spectrum of disilanylphosphine is presented in Figure 4. The absorptions at 2300 and 2150 cm^{-1} can be easily assigned to the PH and SiH stretches, respectively. By comparing the spectrum with that of trisilane,¹¹ we tentatively assign the bands at 942 and 932 cm^{-1}

(11) E. J. Spanier and A. G. MacDiarmid Inorg. Chem., 1, 433 (1962).

and those at 888 and 878 cm^{-1} to the SiH_3 asymmetrical and symmetrical deformation modes, and the bands at 736, 728, and 724 cm^{-1} to SiH_3 wagging and SiH_2 bending modes. In their studies of the infrared spectra of

diphosphine and silylphosphine, Nixon¹² and Linton and Nixon¹³ assigned

(12) E. R. Nixon J. Phys. Chem., 60, 1054 (1956).

(13) H. R. Linton and E. R. Nixon Spectrochim. Acta 15, 146 (1959).

bands at 1081 and 1068.5 cm^{-1} to PH_2 deformation. Nixon assigned strong bands at 633 and 792 cm^{-1} in the spectrum of diphosphine, to PH_2 wagging modes, and much weaker bands at 743 and 827 cm^{-1} to PH_2 twisting modes. We therefore suggest that the weak band at 1073 cm^{-1} in the spectrum of disilanylphosphine is due to PH_2 deformation, while the bands at 784 and 790 cm^{-1} and 637 and 628 cm^{-1} are due to PH_2 wagging and twisting. Linton and Nixon assigned the strong band at 454 cm^{-1} in the spectrum of SiH_3PH_2 to the Si-P stretch in SiH_3PH_2 . In case of $\text{SiH}_3\text{SiH}_2\text{PH}_2$ a weak band is observed in the region of 515 to 475 which could be due to the Si-P stretch.

Reactions- Disilanylphosphine reacts with an excess of water as follows:



Thus the hydrolysis is strictly analogous to the hydrolysis of silylphosphine, in which disiloxane and phosphine are formed. Disilylphosphine is so unstable that it is completely decomposed on a chromatographic column at 70°, whereas disilanylphosphine is less than half decomposed after one day at 210°. Perhaps the instability of disilylphosphine is similar in cause to the presumed instability of disilylamine.¹⁴

(14) F. G. A. Stone, "Hydrogen Compounds of the Group IV Elements",
Prentice-Hall, Englewood Cliffs, N.J., 1962, pp. 27-33.

Acknowledgement: We wish to thank Dr. B. Holder and Dr. J. Happe for the ^{31}P spectrum, and Dr. A. Newton and Mr. A. Sciamanna for the mass spectra. This research was supported by the United States Atomic Energy Commission.

Table I

Chemical Shifts

To Lower Field of TMS in ppm \pm 0.1

	$(\text{SiH}_3)_2\text{PH}$	$\text{SiH}_3\text{SiH}_2\text{PH}_2$
SiH_3	3.70	3.19
SiH_2		3.57
PH_2		1.25
PH	-0.40	


Table II

Coupling Constants (cps)

J_{P-H}				
$\overline{\text{SiH}_3\text{PH}}$	$-\overline{\text{SiH}_2\text{PH}_2}$	$-\overline{\text{SiH}_2\text{PH}_2}$	$\overline{(\text{SiH}_3)_2\text{PH}}$	$\overline{\text{SiH}_3\text{SiH}_2\text{PH}_2}$
184.0 ± 0.5	182.5 ± 0.5	18.5 ± 0.2	17.5 ± 0.2	2.0 ± 0.1
J_{H-H}				
	$\overline{(\text{SiH}_3)_2\text{PH}}$	$-\overline{\text{SiH}_2\text{PH}_2}$	$\overline{\text{SiH}_3\text{SiH}_2-}$	
	5.1 ± 0.1	5.8 ± 0.1	3.0 ± 0.1	

Figure Captions


- Figure 1. Nmr spectrum of disilanylphosphine at 60 Mc.
Chemical shifts to low field of external TMS.
- Figure 2. Portion of nmr spectrum of disilylphosphine at 60 Mc.
Chemical shift to low field of external TMS.
- Figure 3. Portion of nmr spectrum of disilylphosphine at 60 Mc.
The septet of the PH proton on the high-field side
of external TMS.
- Figure 4. Infrared spectrum of disilanylphosphine. (10 mm
pressure in 5 cm cell)
Lower curve at $\sim 3000 \text{ cm}^{-1}$ is the polystyrene calibration.
Weak bands at 825 and 838 cm^{-1} are due to impurities.



MU-33914


FIG 1

2 517



MU-33915


FIG 2



MU-33916

FIG 3

Q 1



MU-33063

FIG 4

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

