

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

EFFECT OF SERUM ON THE GROWTH OF BALB 3T3 A31 MOUSE FIBROBLASTS and AN SV40-TRANSFORMED DERIVATIVE

### Permalink

<https://escholarship.org/uc/item/4b33b3cj>

### Author

Bartholomew, James C.

### Publication Date

1975-07-01

Submitted to Journal of Cell Biology

LBL-3893  
Preprint c.1

EFFECT OF SERUM ON THE GROWTH OF BALB 3T3 A31  
MOUSE FIBROBLASTS AND AN SV40-TRANSFORMED DERIVATIVE


James C. Bartholomew, Hisao Yokota, and Priscilla Ross

July 1975

Prepared for the U. S. Energy Research and  
Development Administration under Contract W-7405-ENG-48

**For Reference**

Not to be taken from this room



LBL-3893  
c.1

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

0 0 0 0 4 3 0 4 8 8 5

EFFECT OF SERUM ON THE GROWTH OF BALB 3T3 A31  
MOUSE FIBROBLASTS AND AN SV40-TRANSFORMED  
DERIVATIVE

James C. Bartholomew, Hisao Yokota, and Priscilla Ross  
Laboratory of Chemical Biodynamics  
University of California  
Berkeley, California 94720

Send Proofs to: Dr. James C. Bartholomew  
Laboratory of Chemical Biodynamics  
University of California  
Berkeley, California 94720  
(415) 843-2740, Ext. 6131

## ABSTRACT

The effect of serum on the growth properties of non-transformed Balb 3T3 A31 and SV40-transformed Balb 3T3 A31 was studied. The concentration of serum in the growth medium of non-transformed cells had little effect on the initial population doubling time, but did regulate the cell density at which the population became quiescent in  $G_1$ . The doubling time of transformed cells, however, was increased significantly as the concentration of serum was decreased below 4%. This effect on the growth of transformed cells was seen at serum concentrations so low that non-transformed cells did not complete one population doubling. Flow microfluorimetric analysis of these populations indicated that the primary effect of different serum concentrations on the non-transformed cells was to modulate the average residence time in  $G_1$ ; whereas, all the cell cycle phases of the transformed cells were affected by serum. At saturation densities, the non-transformed cells became quiescent in  $G_1$ , but the transformed cells still traversed the cell cycle and their saturation density appeared to be a balance between cell production and cell death occurring primarily in the  $G_1$  phase of the cell cycle.

## INTRODUCTION

The growth properties of mammalian cells in culture are thought to reflect the corresponding properties in vivo (1, 22). Cell lines that have "tight" growth regulatory properties in culture generally do not give tumors when injected back into animals (1). Balb 3T3 A31 mouse fibroblast cells exhibit "tight" growth control in culture, but when transformed by SV40 virus their growth potential in culture under a given set of conditions increases as does their ability to form tumors in animals (1). These observations suggest that an understanding of the mechanisms for the regulation of growth of mammalian cells in culture will be relevant to growth control in vivo.

One of the elements that has been shown to play a roll in the regulation of growth of mammalian cells in culture is serum. Holley and Kiernan (9) found that the concentration of serum in the growth medium determined the final cell density of both transformed and non-transformed mouse 3T3 cells in culture. This final cell density or saturation density was much higher for transformed cells than non-transformed cells at a given serum concentration. Nilausen and Green (14) demonstrated that when non-transformed mouse fibroblasts reached their saturation density cell cycle traverse was arrested in  $G_1$ . This block, however, is reversible and cell cycle traverse can be stimulated by the addition of fresh serum (25). Factors other than serum concentration can regulate cell growth by restricting passage through  $G_1$  (17,20,24) and it has been postulated that the growth of non-transformed cells is controlled by a restriction point in  $G_1$  that is sensitive to suboptimal growth conditions (15).

The effect of limitation of serum or other medium components on the growth of transformed cells has not been clearly defined. There is some evidence that as a population of transformed cells reaches its saturation

density, and presumably depletes some medium components, the mitotic index drops (6). Paul (11) has demonstrated that with SV3T3 cells low levels of leucine inhibit cell cycle traverse; but, in contrast to non-transformed cells, the transformed cells were not arrested in one phase of the cell cycle.

In this report we describe the effect of serum on the doubling time of Balb 3T3 A31 and a closely derived SV40 virus transformed line, FNE. Our results indicate that limitation of serum has less effect on the initial population doubling time of Balb 3T3 A31 than on FNE. Furthermore, serum regulates the growth of Balb 3T3 A31 by increasing the average residence time in  $G_1$ , but regulates FNE growth by lengthening the average residence time of all phases of the cell cycle.

## MATERIALS AND METHODS

### Cell Culture Techniques

The cells used in this study were obtained from Helene Smith of the Naval Biomedical Laboratory, Oakland, California. Balb 3T3 A31 was cloned prior to use in these experiments to give the derivative, Balb 3T3 A31 HYF. FNE is a derivative of Balb 3T3 A31 which has been transformed with SV40 virus. It is producing SV40 T-antigen and unlike Balb 3T3 A31 is highly tumorigenic when injected as a suspension in newborn Balb/c mice<sup>1</sup>. It was passaged nineteen times as described below before use in these experiments.

Both cell lines were carried in 100 mm plastic dishes (Falcon, Oxnard, Calif.) and incubated in a 10% CO<sub>2</sub> incubator at 37°. The cells were grown in Vogt and Dulbecco's modification of Eagle's medium (27), containing 10% newborn calf serum (GIBCO, Santa Clara, Calif.). The cells were transferred

---

<sup>1</sup> Smith, H., personal communication.

when reaching confluency by removal from the dishes with 0.05% trypsin (Difco, 1:250; Detroit, Mich.) in 25 mM Tris buffer, pH 7.4 containing 140 mM NaCl, 5 mM KCl, and 0.7 mM  $\text{Na}_2\text{HPO}_4$  (isotonic Tris buffer). The seeding density was 1/10 of the saturation density.

### Growth Curves

The growth of the cultures in media containing different concentrations of calf serum was monitored by determining the increase in the number of cells per culture dish as a function of time. On day 0 four 35 mm plastic dishes were seeded for each time point to be taken. The dishes were seeded with  $5.0 \times 10^4$  cells in 1.5 ml of test medium. At each time point the cells were removed from the dishes by treatment with 2.0 ml 0.01% trypsin in isotonic Tris buffer for 20 min at 37°. The number of cells per dish was determined by counting an aliquot of the trypsin solution in a Coulter counter model Fn (Coulter Electronics, Hialeah, Florida). The values were corrected for coincidence counting and averaged.

### Autoradiography

To determine the percentage of the population in S at each time point, the cultures were pulsed for 15 min with 1  $\mu\text{Ci/ml}$   $^3\text{H}$ -thymidine (20.1 Ci/mM; New England Nuclear, Boston, Mass.). At the end of the pulse the cells were washed on the dish 2 times in isotonic Tris buffer, swollen with 0.075 M KCl and fixed with methanol-acetic acid (3:1)(19). After air drying the dishes were coated with Kodak Nuclear Track Emulsion NTB-2 (Eastman Kodak, Rochester, N.Y.), exposed in the dark at room temperature for 14 days, developed, and fixed. The cells were stained with crystal violet and the percentage of labeled nuclei was determined by counting a total of 2000 cells for each time point.



### Flow Microfluorimetry

The amount of DNA per individual cell was quantified by staining the cells with acriflavine (26) and measuring the amount of fluorescence per cell by flow microfluorimetry (10). These measurements were done in collaboration with Drs. Joe Gray and Marvin Van Dilla of the Lawrence Livermore Laboratory. In this technique, the stained cells are passed individually through the beam of an argon-ion laser (Spectra-Physics, Mountain View, Calif.) tuned to 488 nm. The pulse of fluorescent light was filtered to reduced scattered light and detected by a photomultiplier tube positioned at right angles to the laser beam. The resulting signal is amplified electronically and recorded in the memory of a pulse height analyzer (Northern Scientific, Middletown, Wisconsin). As the data accumulates it generates a histogram as shown in Figure 1. The data was stored on magnetic tape and processed by a program written for a Sigma 2 computer (Xerox, Rochester, N.Y.).<sup>2</sup> As seen in Figure 1 the integration of these plots gives the percentage of cells in the population in  $G_1$ , S, and  $G_2$  plus  $M(G_2+M)$ . The procedure used to obtain the proportion of the population in each of the phases is described in a separate report (2).

### RESULTS

#### Effect of Serum on the Growth of Balb 3T3 A31 HYF and FNE

Figure 2 shows that the growth of Balb 3T3 A31 HYF and FNE is dependent on the concentration of serum in the medium. As the serum concentration increased, the doubling time of the population decreased (Table 1) and the saturation density increased. The doubling times in Table 1 were computed from the cell counts using a least squares curve fit program. Only values

---

<sup>2</sup> Wong, S., K. Wiley and A. Salmon, unpublished procedure.

that had correlation coefficients greater than 0.97 are reported. The initial doubling time of Balb 3T3 A31 HYF at serum concentrations below 8% was difficult to determine because of the small amount of growth seen with these serum concentrations, and above 8% the effect of serum concentration on the initial doubling time was minimal. The doubling time of FNE, on the other hand, was clearly dependent on the concentration of serum in the medium. Accurate population doubling times for FNE could be measured at serum concentrations as low as 1%. A comparison of the doubling times of Balb 3T3 A31 HYF and FNE at serum concentrations above 8% indicates that the two cell lines have the same minimum doubling time. Throughout the course of these experiments, the cultures were monitored microscopically for cells floating in the medium as an indication of cell death. The only evidence that cell death was occurring was with FNE after it had reached its saturation density.

Cell Cycle Distributions After Two Days Growth in Medium Containing Different Serum Concentrations

Flow microfluorimetry (FMF) can determine the distribution of a population in the cell cycle by measuring the DNA content of individual cells. Cells that are in  $G_1$  have half the amount of DNA of cells that are in  $G_2$  or mitosis. Cells that are in S have DNA contents between that of  $G_1$  cells and  $G_2$  cells depending on how far through S they have progressed. Assuming a random distribution of cells around the cell cycle, the proportion of the cells in a particular phase of the cell cycle is a function of the average residence time in that phase. Using the doubling time of the population and the distribution of cells in the various phases, the average

residence time of each phase can be calculated (12). The technique assumes a negligible death rate which appears to be true for Balb 3T3 A31 HYF and FNE growing logarithmically (see above). We have used this procedure to determine how different serum concentrations in the growth medium affect the average residence time of the two populations in the individual phases of the cell cycle.

The FMF distributions of Balb 3T3 A31 HYF after two days of growth in medium containing different serum concentrations are shown in Fig. 3. It was not possible to analyze the cells before day 2 because the cell cycle distributions were still affected by the state of the population prior to setting up the experiment (3). Table 2 gives the values for the integrated areas under the DNA histograms. These calculations gave values for the percentage of the population in S that agreed within 5% of those obtained by incorporation of a 15 min pulse of  $^3\text{H}$ -thymidine followed by autoradiography to detect DNA synthesizing cells. As the serum concentration decreased below 8% the percentage of the population in  $G_1$  increased while the other phases decreased. At low serum concentrations there were essentially no cells in S.  $^3\text{H}$ -thymidine incorporation followed by autoradiography of cells growing in medium containing less than 2% serum confirmed that less than 3% of the cells were synthesizing DNA on day 2.

There appeared to be some cells with a DNA content of  $G_2+M$  cells even at low serum concentrations. It should be pointed out, however, that binucleated or aggregated cells would be recorded as having a DNA content equivalent to  $G_2+M$  cells. However, if aggregation or multinucleation were affecting the results, peaks representing higher multiples of  $G_2+M$  should be present in the histograms shown in Fig. 3. The absence of these peaks suggests that the  $G_2+M$  peak is not contaminated with a significant amount of aggregates or multinucleates.

The growth of Balb 3T3 A31 HYF was logarithmic on day 2 only in medium containing 8% serum or greater. Using the doubling time of 16 hr for these logarithmic populations (Table 1), the average residence time of cells in each of the cell cycle phases was calculated for the different growth conditions. These values are given in Table 2 and indicate that although the doubling times were identical for a range of serum concentrations, the average residence times varied. As the serum concentration increased, the value for  $G_1$  decreased and the value for S increased. The average residence time for  $G_2+M$  remained approximately constant.

Figure 4 gives the FMF distributions of FNE growing in medium with different concentrations of serum. Integrations of the areas under the histograms gave the values in Table 3 for the distribution of cells in the cell cycle. Unlike Balb 3T3 A31 HYF, FNE even in medium with very low serum concentrations had a significant number of cells in all phases of the cell cycle. As an example, approximately 10% of the population was in S when the serum concentration was only 0.1%. This was confirmed by  $^3\text{H}$ -thymidine incorporation followed by autoradiography. As seen in Fig. 2, FNE grew logarithmically at serum concentrations as low as 1%. When the average residence time of the cell cycle phases was calculated for the logarithmic populations it was apparent that as the doubling time increased the length of all phases increased.

#### Cell Cycle Distributions at Saturation Density

Using FMF we analyzed the cell cycle distributions of cultures of Balb 3T3 A31 HYF and FNE at their saturation density in medium containing different concentrations of serum. Balb 3T3 A31 HYF was predominately (>93%) in  $G_1$  at saturation density regardless of the concentration of serum in the medium.  $^3\text{H}$ -thymidine incorporation followed by autoradiography

indicated that less than 0.5% of the population was making DNA at any of the serum concentrations. FNE gave cell cycle distributions at saturation densities in media with different levels of serum as seen in Fig. 5. Even though the cultures had stopped increasing in density there were still cells in all parts of the cell cycle. The cells were still incorporating  $^3\text{H}$ -thymidine into DNA at all serum concentrations.

As mentioned above when cultures of FNE reached saturation density some cells could be seen in the medium. These cells were not viable when replated into fresh medium and when analyzed using the FMF technique they had DNA contents equivalent to  $G_1$  cells.

#### DISCUSSION

The doubling time of a population of cells is dependent on three variables: (a) the length of the cell cycle; (b) the fraction of the population traversing the cycle; and (c) the rate of cell loss (4). A number of models have been proposed using these variables to explain the growth properties of mammalian cells. Prescott (18) has suggested that the growth rate of cells in culture is determined primarily by the length of  $G_1$ . Smith and Martin (21), on the other hand, have recently proposed that the doubling time of a population in a given environmental situation is dependent on the probability of cell cycle transit. Both of these models propose that the length of S,  $G_2$  and M are constant, despite the evidence that in vivo at least the length of S and  $G_2$  does vary (5, 11).

Understanding the mechanism of growth control becomes particularly important when considering malignant cells. Analysis of the growth properties of malignant cells in vivo suggests that they do not necessarily grow with a shorter doubling time than normal cells, but appear to be able to grow under conditions where normal cells will not (11). Under such

conditions the fraction of proliferating cells (13) is high in the case of malignant cells, but low with normal cells.

In the experiments reported here we have analyzed the growth properties of Balb 3T3 A31 HYF, a non-transformed mouse fibroblast with low tumorigenic potential, and a highly tumorigenic derivative, FNE, in media containing growth limiting amounts of serum. As seen by Holley and Kiernan (8) serum concentration primarily affects the cell density at which a population of non-transformed cells becomes quiescent. This stringent effect on the saturation density makes analysis of the effect of serum on the rate of population growth difficult. However, in other experiments (3) we have used a  $G_1$  synchronized population to show that the level of serum in the medium has little effect on the time of transit from  $G_1$  through S to  $G_2$  and M, but does have an effect on the fraction of the population capable of being stimulated. These experiments suggest that the primary effect of serum on the growth of Balb 3T3 A31 HYF is in determining the fraction of the population capable of cell cycle transit. A secondary effect of serum may be on other variables determining the growth properties of the population, but these effects must be small and are difficult to separate from the primary effect of serum. These experiments are consistent with the model of Smith and Martin (21) and suggest that serum acts by determining the probability of cell cycle transit.

The doubling time of FNE, on the other hand, is very much dependent upon the concentration of serum in the medium at concentrations of serum below 4% (Table 1). These effects do not seem to be due to cell loss since microscopic observation of the cultures did not reveal a significant increase in floating dead cells as the doubling time increased. At concentrations of serum where the doubling time is lengthened the average residence time in each of the cell cycle phases is also lengthened (Table 3). As the

concentration of serum in the medium dropped to 0.4% or lower, the population accumulated in  $G_2+M$ . It is important to note that these effects of serum on the length of individual phases of the cell cycle of FNE were seen at serum concentrations so low that Balb 3T3 A31 HYF did not grow significantly. Thus, it is possible that serum may affect the average residence time of Balb 3T3 A31 HYF in the cell cycle phases other than  $G_1$ , but at concentrations where this effect may occur the population has become quiescent in  $G_1$ .

The cell cycle distributions at the saturation density of Balb 3T3 A31 HYF indicated that no matter what the concentration of serum in the medium, the cells stopped growing by becoming quiescent in  $G_1$ . FNE, on the other hand, did not become quiescent at its saturation density, but continued to traverse the cell cycle. Microscopic observations indicated that at saturation density dead floating cells were accumulating. FMF analysis of the floating cells indicated that death had occurred predominately in  $G_1$ .

These studies indicate that at serum concentrations where Balb 3T3 A31 HYF can grow significantly there is little effect of serum on the doubling time of the population. The primary effect is on determining the saturation density at which the population becomes quiescent. When this cell line is transformed with SV40 virus, the transformant grows at lower serum concentrations, but with doubling times that are dependent on serum concentration. These longer doubling times are the result of lengthening the average residence time in each of the cell cycle phases.

## ACKNOWLEDGEMENTS

We would like to thank especially Drs. Joe Gray and Marvin Van Dilla for their assistance in performing the flow microfluorimetric analyses, and Dr. Helene Smith of the Naval Biomedical Laboratory for supplying the cells. In addition, we would like to thank Evie Litton and Margaret Emrich for helping with the preparation of this manuscript.

These studies were supported by NCI Grant number CA14828-02 and by the United States Energy Resources and Development Administration.



REFERENCES

1. Aaronson, S. A., and G. J. Todaro (1968) "Basis for the acquisition of malignant potential by mouse cells cultivated in vitro", *Science*, 162:1024-1026.
2. Bartholomew, J. C., N. T. Neff and P. A. Ross (1975) "Stimulation of WI-38 cell cycle transit--effect of serum concentration and cell density", manuscript submitted.
3. Bartholomew, J. C., H. Yokota, R. Kahn and P. Ross (1974) "Effect of serum on the cell cycle parameters of 3T3 and SV3T3", *J. Cell Biol.* 63:16a.
4. Baserga, R. (1971) in *The Cell Cycle and Cancer*, ed. R. Baserga (Marcel Dekker, Inc., New York), p. 55.
5. Baserga, R. and F. Wiebel (1969) "The cell cycle of mammalian cells", *Int. Rev. Exptl. Path.* 7:1-30.
6. Castor, L. N. (1970) "Flattening, movement and control of division of epithelial-like cells", *J. Cell Physiol.* 75:57-64.
7. Glinos, A. D. and R. J. Werrlein (1972) "Density dependent regulation of growth in suspension cultures of L-929 cells", *J. Cell Physiol.* 79:79-90.
8. Holley, R. W. and J. A. Kiernan (1968) "Contact inhibition of cell division in 3T3 cells", *Proc. Natl. Acad. Sci. USA* 60:300-304.
9. Holley, R. W. and J. A. Kiernan (1971), "Studies of serum factors required by 3T3 and SV3T3 cells" in *CIBA Foundation Symposium on Growth Control in Cell Cultures*, ed. G. E. W. Wolstenholme and J. Knight (J. A. Churchill, London), p. 3-15.
10. Holm, D. M. and L. S. Cram (1973) "An improved flow microfluorometer for rapid measurement of cell fluorescence", *Exp. Cell Res.* 80:105-110.

11. Lala, P. K. (1971) "Studies on tumor cell population kinetics" in Methods in Cancer Research, Vol. VI (Academic Press, N.Y.) ed., H. Bush, pp. 3-95.
12. Mak, S. (1965) "Mammalian cell cycle analysis using microspectrophotometric combined with autoradiography", Exp. Cell Res. 39:286-308.
13. Mendelsohn, M. L. (1962) "Autoradiographic analysis of cell proliferation in spontaneous breast cancer of C3H mouse. III. The growth fraction", J. Natl. Cancer Inst. 28:1015-1029.
14. Nilausen, K. and H. Green (1965) "Reversible arrest of growth in  $G_1$  of an established fibroblast line (3T3)", Exp. Cell Res. 40:166-168.
15. Pardee, A. B. (1974) "A restriction point for control of normal animal cell proliferation", Proc. Natl. Acad. Sci. USA 71:1286-1290.
16. Paul, D. (1973) "Quiescent SV40 virus transformed 3T3 cells in culture", Biochem. and Biophys. Res. Commun. 53:745-753.
17. Pohjanpelto, P. and A. Raina (1972) "Identification of a growth factor produced by human fibroblasts in vitro as putrescine", Nature New Biol. 235:247-249.
18. Prescott, D. M. (1968) "Regulation of cell reproduction", Cancer Res. 28:1815-1820.
19. Rothfels, K. H. and L. Siminovitch (1958) "An air-drying technique for flattening chromosomes in mammalian cells grown in vitro", Stain Technology 33:73-77.
20. Rubin, H. (1972) "Inhibition of DNA synthesis in animal cells by ethylene diamine tetraacetate, and its reversal by zinc", Proc. Natl. Acad. Sci. USA 69:712-716.
21. Smith, J. A. and L. Martin (1973) "Do cells cycle?", Proc. Natl. Acad. Sci. USA 70:1263-1267.

22. Stoker, M. G. P. (1972) "Tumour viruses and the sociology of fibroblasts", Proc. R. Soc. Lond. B. 181:1-17.
23. Studzinski, G. P. and J. F. Gierthy (1973) "Selective inhibition of the cell cycle of cultured human diploid fibroblasts by aminonucleoside of puromycin", J. Cell. Physiol. 81:71-84.
24. Tobey, R. A. and K. D. Ley (1970) "Regulation of initiation of DNA synthesis in Chinese hamster cells. I. Production of stable, reversible G<sub>1</sub>-arrested populations in suspension culture", J. Cell Biol. 46:151-157.
25. Todaro, G. J., G. K. Lazar and H. Green (1965) "The initiation of cell division in a contact-inhibited mammalian cell line", J. Cell and Comp. Physiol. 66:325-334.
26. Trujillo, T. T. and M. A. Van Dilla (1972) "Adaptation of the fluorescent feulgen reaction to cells in suspension for flow microfluorometry", Acta Cytol. 16:26-30.
27. Vogt, M. and R. Dulbecco (1963) "Steps in the neoplastic transformation of hamster embryo cells by polyoma virus", Proc. Natl. Acad. Sci. USA 49:171-179.

Table I.  
Effect of Serum on the Population Doubling  
Time of Balb 3T3 A31 HYF and FNE

| % Serum | Doubling Time (hr)* |     |
|---------|---------------------|-----|
|         | Balb 3T3 A31 HYF    | FNE |
| 1.0     | -                   | 51  |
| 2.0     | -                   | 29  |
| 4.0     | -                   | 20  |
| 6.0     | -                   | 16  |
| 8.0     | 16                  | 16  |
| 10.0    | 16                  | 16  |
| 16.0    | 16                  | 16  |
| 20.0    | 16                  | -   |

\* The doubling times were computed from the cell counts using a least squares fitting program (see text).

Table II.

Effect of Serum on the Cell Cycle Distribution of Balb 3T3 A31 HYF

| % Serum | G <sub>1</sub>  |                             | S               |                             | G <sub>2</sub> +M |                             |
|---------|-----------------|-----------------------------|-----------------|-----------------------------|-------------------|-----------------------------|
|         | % of Population | Average Residence Time (hr) | % of Population | Average Residence Time (hr) | % of Population   | Average Residence Time (hr) |
| 0.4     | 96.9            | -                           | 0               | -                           | 3.1               | -                           |
| 1.0     | 93.7            | -                           | 1.7             | -                           | 4.6               | -                           |
| 2.0     | 95.6            | -                           | 0               | -                           | 4.4               | -                           |
| 4.0     | 80.1            | -                           | 6.1             | -                           | 12.8              | -                           |
| 6.0     | 59.5            | -                           | 18.0            | -                           | 22.4              | -                           |
| 8.0     | 50.9            | 6.8                         | 24.1            | 4.0                         | 25.0              | 5.2                         |
| 10.0    | 48.6            | 6.4                         | 28.9            | 4.9                         | 22.5              | 4.7                         |
| 16.0    | 42.8            | 5.6                         | 31.0            | 5.0                         | 26.2              | 5.4                         |
| 20.0    | 38.6            | 5.0                         | 38.2            | 6.2                         | 23.2              | 4.8                         |

Table III.

Effect of Serum on the Cell Cycle Distribution of FNE

| % Serum | $G_1$           |                             | S               |                             | $G_2+M$         |                             |
|---------|-----------------|-----------------------------|-----------------|-----------------------------|-----------------|-----------------------------|
|         | % of Population | Average Residence Time (hr) | % of Population | Average Residence Time (hr) | % of Population | Average Residence Time (hr) |
| 0.1     | 34.6            | -                           | 9.6             | -                           | 55.9            | -                           |
| 0.4     | 49.8            | -                           | 10.0            | -                           | 40.2            | -                           |
| 1.0     | 49.6            | 21.0                        | 22.6            | 12.0                        | 27.8            | 18.0                        |
| 2.0     | 47.7            | 11.4                        | 28.1            | 8.5                         | 24.2            | 9.1                         |
| 4.0     | 56.6            | 9.6                         | 22.3            | 4.9                         | 22.2            | 5.5                         |
| 6.0     | 45.3            | 5.9                         | 25.9            | 4.2                         | 28.8            | 5.9                         |
| 8.0     | 49.1            | 6.6                         | 27.5            | 4.5                         | 23.5            | 4.9                         |
| 10.0    | 50.0            | 6.6                         | 24.7            | 4.2                         | 25.3            | 5.2                         |
| 16.0    | 46.8            | 6.2                         | 27.1            | 4.5                         | 26.1            | 5.3                         |

FIGURE LEGENDS


Figure 1. Idealized DNA histogram obtained by analyzing a non-synchronous population.

Figure 2. Growth of Balb 3T3 A31 HYF and FNE in medium containing different concentrations of new born calf serum.

Figure 3. DNA histograms of populations of Balb 3T3 A31 HYF harvested two days after seeding in medium containing the indicated serum concentration.

Figure 4. DNA histograms of populations of FNE harvested two days after seeding in medium containing the indicated serum concentration.


Figure 5. DNA histograms of populations of FNE harvested six days after seeding in medium containing the indicated serum concentration.



XBL752-5090

Fig. 1






XBL756-5305

Fig. 2


CELL CYCLE DISTRIBUTIONS OF BALB 3T3 A31  
HYF DURING LOGARITHMIC GROWTH



XBL756-5313


Fig. 3

CELL CYCLE DISTRIBUTIONS OF  
FNE DURING LOGARITHMIC GROWTH



XBL756-5314

Fig. 4

CELL CYCLE DISTRIBUTIONS OF FNE  
IN STATIONARY PHASE

XBL756-5315

Fig. 5

U S G O 4 3 0 4 3 7 7

**LEGAL NOTICE**

*This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.*

TECHNICAL INFORMATION DIVISION  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720