UCLA UCLA Previously Published Works

Title

State-of-the-Science Data and Methods Need to Guide Place-Based Efforts to Reduce Air Pollution Inequity.

Permalink

https://escholarship.org/uc/item/4bt2s7mj

Journal Environmental Health Perspectives, 131(12)

Authors

Gohlke, Julia Harris, Maria Roy, Ananya <u>et al.</u>

Publication Date

2023-12-01

DOI

10.1289/EHP13063

Peer reviewed

State-of-the-Science Data and Methods Need to Guide Place-Based Efforts to Reduce Air Pollution Inequity

Julia M. Gohlke,^{1,2} Maria H. Harris,¹ Ananya Roy,¹ Tammy M. Thompson,¹ Mindi DePaola,¹ Ramón A. Alvarez,¹ Susan C. Anenberg,³ Joshua S. Apte,^{4,5} Mary Angelique G. Demetillo,⁶ Isabella M. Dressel,⁶ Gaige H. Kerr,³ Julian D. Marshall,⁷ Aileen E. Nowlan,¹ Regan F. Patterson,⁸ Sally E. Pusede,⁶ Veronica A. Southerland,^{1,3} and Sarah A. Vogel¹

¹Environmental Defense Fund, Washington, District of Columbia, USA

²Department of Population Health Sciences, Virginia Tech, Blacksburg, Virginia, USA

³Department of Environmental and Occupational Health, George Washington University, Washington, District of Columbia, USA

⁴Department of Civil and Environmental Engineering, University of California, Berkeley, Berkeley, California, USA

⁵School of Public Health, University of California, Berkeley, Berkeley, California, USA

⁶Department of Environmental Sciences, University of Virginia, Charlottesville, Virginia, USA

⁷Department of Civil and Environmental Engineering, University of Washington, Seattle, Washington, USA

⁸Department of Civil and Environmental Engineering, University of California, Los Angeles, Los Angeles, California, USA

BACKGROUND: Recently enacted environmental justice policies in the United States at the state and federal level emphasize addressing place-based inequities, including persistent disparities in air pollution exposure and associated health impacts. Advances in air quality measurement, models, and analytic methods have demonstrated the importance of finer-scale data and analysis in accurately quantifying the extent of inequity in intraurban pollution exposure, although the necessary degree of spatial resolution remains a complex and context-dependent question.

OBJECTIVE: The objectives of this commentary were to *a*) discuss ways to maximize and evaluate the effectiveness of efforts to reduce air pollution disparities, and *b*) argue that environmental regulators must employ improved methods to project, measure, and track the distributional impacts of new policies at finer geographic and temporal scales.

DISCUSSION: The historic federal investments from the Inflation Reduction Act, the Infrastructure Investment and Jobs Act, and the Biden Administration's commitment to Justice40 present an unprecedented opportunity to advance climate and energy policies that deliver real reductions in pollution-related health inequities. In our opinion, scientists, advocates, policymakers, and implementing agencies must work together to harness critical advances in air quality measurements, models, and analytic methods to ensure success. https://doi.org/10.1289/EHP13063

Introduction

Twenty-three states and the federal government have enacted environmental justice policies since the 1990s, but only recently have significant resources been allocated at both state and federal levels. Currently, five states (California, Colorado, Illinois, New York, and Washington) and the Biden Administration's Justice40 Initiative¹ directly designate a minimum percentage investment allocation to prioritized locations (Climate and Economic Justice screening tool (https://screeningtool.geoplatform. gov), ClimateXchange (https://climate-xchange.org/dashboard/), Colorado Department of Public Health and Environment (https:// teeo-cdphe.shinyapps.io/COEnviroScreen_English/), Illinois Power Agency,² California Office of Environmental Health Hazard Assessment,³ New York State Climate Justice Working Group,⁴ and Washington State Department of Health⁵). (Table 1; see also "Place-based environmental justice policies with investment or resource allocations" in the Supplemental Material). To define which locations qualify under these environmental justice investment allocation programs, a range of socioeconomic, demographic, and environmental variables at the census tract or

Supplemental Material is available online (https://doi.org/10.1289/EHP13063).

Authors declare they have nothing to disclose beyond funding sources disclosed in the Acknowledgments section.

census block group scale are currently being used (Table 1; see also "Place-based environmental justice policies with investment or resource allocations" in the Supplemental Material). Notably, the Biden Administration has allocated tens of billions of dollars via the Justice40 Initiative,¹ which calls for 40% of federal program benefits to be directed to geographically defined "disadvantaged communities" (Table 1; see also "Place-based environmental justice policies with investment or resource allocations" in the Supplemental Material). The Inflation Reduction Act (IRA) allocates \$296 million for investments in air monitoring specifically.⁶ In addition, for the first time, the federal government is inviting the use of air pollution data to guide investment of >\$30 billion in funds focused on climate pollution reduction, advanced industrials, community revitalization, and other pollution mitigation programs.⁶ These new place-based prioritizations, combined with federal climate and clean energy funding through the Infrastructure Investment and Jobs Act (IIJA),⁷ present an unprecedented opportunity to address longstanding disparities in air pollution exposure and impacts that have persisted under prior regulatory structures (e.g., the Clean Air Act). We assert that to maximize and evaluate the effectiveness of these efforts, federal and state agencies will require improved methods to measure, model, project, and track the distributional impacts of new policies. Hence, there is a pressing need for scientists, advocates, policymakers, and implementing agencies to collaboratively develop new data collection and analytic frameworks to aid the implementation of place-based policies. Here, we offer recommendations informed by relevant recent research on criteria air pollutants leveraging satellite remote sensing, new monitoring technologies, hybrid models, and fine-scale health impact assessment approaches to a) characterize past and current inequities in air pollution exposure, b) model expected health equity benefits of past and potential future air pollution policies, and c) empirically determine whether expected air quality and health benefits are achieved following policy implementation.

Address correspondence to Julia M. Gohlke, Environmental Defense Fund, 555 12th St NW #400, Washington, DC 20004 USA. Telephone: (202) 572-3513. Email: jgohlke@edf.org

Received 23 March 2023; Revised 19 November 2023; Accepted 27 November 2023; Published 18 December 2023.

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehpsubmissions@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Table 1. Environmental ju	istice policies	with place-based	allocations or	investment	provisions
---------------------------	-----------------	------------------	----------------	------------	------------

			Qualifying criteria				
Policy (date)	Allocation provision	Spatial unit	Climate burden	Pollution burden	Health burden	Proportion non-White/ Hispanic	Proportion low income
California Climate Investment Act (2016)	At least 25% ^a	Census tract	_	\checkmark	\checkmark	_	1
Colorado Modernize the Public Utilities Commission Act (2021)	At least 40% renewable energy	Census block group	\checkmark	\checkmark	1	1	1
Illinois Climate and Equitable Jobs Act (2021)	25% solar funding, 45% EV funding	Census block group	—	\checkmark	—	\checkmark	1
New York Climate Leadership and Community Protection Act (2019)	35%, with a goal of 40%	Census tract	1	5	\checkmark	1	1
Washington Climate Commitment Act (2021)	35%, with a goal of 40%, 10% to Indian tribes	Census tract	—	\checkmark	1	1	1
United States Justice40 Executive Order (2020)	40% for qualifying federal programs	Census tract	1	\checkmark	\checkmark		1

Note: Table is based on Climate and Economic Justice screening tool (https://screeningtool.geoplatform.gov), ClimateXchange (https://climate-xchange.org/dashboard/), Colorado Department of Public Health and Environment (https://teeo-cdphe.shinyapps.io/COEnviroScreen_English/), Illinois Power Agency,² California Office of Environmental Health Hazard Assessment,³ New York State Climate Justice Working Group,⁴ and Washington State Department of Health.⁵ —, not applicable; EV, electric vehicle. ^a10% to low-income communities, individuals, and households.

Discussion

Characterizing Inequities in Air Pollution Exposures

Advances in air pollution modeling and satellite-derived measurements have enabled progress in characterizing criteria air pollutants at spatial resolutions relevant to current environmental justice place-based policies (e.g., census tract). Most analyses find significantly higher pollution concentrations in areas with higher proportions of residents identifying as Black, Asian, or Hispanic; reporting low income; or residing in an area subjected to historic patterns of discrimination.^{8–14} Generally, intraurban differences in air pollutant concentrations are greater for primary (i.e., directly emitted) pollutants, such as nitrogen dioxide (NO₂) and ultrafine particles, compared with secondary pollutants (i.e., which are formed from chemical reactions), such as ozone and a portion of particulate matter $\leq 2.5 \ \mu m$ in aerodynamic diameter $(PM_{2.5})$.^{11,14,15} Research into the spatial resolution needed to accurately quantify pollution inequities suggests that countyscale analyses can underestimate the magnitude of total U.S. PM_{2.5} and NO₂ exposure disparities by race and ethnicity compared with those employing finer-scale data aggregation (census tract, block group, and block)¹⁶ but that the resolution of current satellite measurements (e.g., 20–60 km²) can quantify a large (although perhaps incomplete) fraction of intraurban NO₂ inequality.^{8,17,18} National, state, regional, and city government agencies, along with private companies, now have access to funding (and are subject to mandates) that welcome or require a focus on remedying health inequities. There is great potential to leverage recent research and novel data to inform strategies in reducing intraurban and regional pollution disparities.

Modeling to Evaluate Health Equity Impacts of Pollution Mitigation Policies

Recent research has also advanced modeling frameworks to predict expected benefits of policies aimed at reducing race, ethnicity, and income inequities. To estimate health burdens associated with air pollution exposure within a given geography and the expected benefits of exposure reductions, health impact assessments generally use area-level (e.g., county, census tract, census block group) exposure, baseline disease rates, and concentration–response functions derived from previous epidemiological studies. Values for each of these quantities can vary across geographies and population subgroups, driven by environmental and social determinants of health.

Inequities in baseline disease burden are a result of many processes.¹⁹⁻²¹ As one example, current and historical discriminatory practices based on presenting race and ethnicity can lead to inequities in health burden via chronic social stress, in addition to causing air pollution exposure differences via segregation.^{22,23} A recent study has demonstrated the use of fine-scale baseline disease rates in identifying overburdened neighborhoods with overlapping high exposures and poor health.²⁴ An assessment of the San Francisco Bay Area (California) using census block group mortality rates yielded 15% higher spatially aggregated estimates of pollutant-attributable mortality rates, as compared with the application of county baseline disease rates.²⁵ Despite their potential importance in health impact assessment, highly resolved baseline disease rates or individual-level air pollution exposure data are rarely publicly available, in part because of concerns around identifiability; instead, for baseline disease rates, many health impact assessments rely on coarsely resolved data (e.g., county-level or coarser) or modeled estimates at a finer resolution (e.g., census tract or finer).

Recent analyses using a fine-scale (1-km²) gridded exposure data set²⁶ aggregated to the census tract, concentration-response functions derived from a longitudinal study of Medicare recipients,²⁷ and fine-scale (census tract) incidence data estimated PM_{2.5}-attributable mortality for Americans >65 years of age, and reported mortality rates three times higher for Black Americans compared with white Americans.^{28,29} These studies applied racespecific concentration-response functions to represent a range of race-based disparities and discriminatory processes that result in health burden inequities. The race-specific concentration-response functions used in the above studies were derived from an underlying epidemiological study with exposure estimates at the ZIP code level,²⁷ which could lead to mischaracterization due to exposure variation within ZIP codes. In addition, nonlinear concentrationresponse relationships may be important to consider in accounting for the known drivers of disparities.³⁰

To effectively implement place-based policies in support of environmental justice, we believe it will be critical to identify specific emission sources and sectors contributing to air quality inequities with the highest spatial precision that is feasible. In addition, current evidence using reduced-form air pollution models—developed to reduce the computational costs of model

runs-suggests the magnitude of disparities and relative contributions of different sectors can vary across U.S. Regions, as well as urban and rural areas.³¹ Therefore, we suggest that the emission sources that need to be reduced to maximally reduce exposure inequalities may differ by location. Fine-scale (e.g., census tract or higher resolution) source attribution may need to reflect weather patterns, built infrastructure, and local sources³²; current regulatory modeling typically operates at broader spatial scales (county or state) and without fine-scale local information. Source attribution at the hyperlocal scale is currently limited both by input data (e.g., emissions inventories, meteorological fields) and, for conventional models, by the spatial precision of computationally tractable models.³³ Air pollution-relevant emission inventories based on data provided through the U.S. Environmental Protection Agency (EPA) National Emissions Inventory (NEI)³⁴ for the continental United States are available at 12-, 4-, and 1-km² resolution.^{12,31,35–38}

Researchers have made advances in identifying census block or finer exposure inequities associated with different sources, such as restaurants³⁹ and traffic-related air pollution.^{8,11,15,40-44} Other research has studied inequities associated with other sectors of the economy, such as electricity generation^{45,46} and interstate freight.⁴⁷ Using fine-scale air quality change estimates, Wang et al.⁴⁸ and Dressel et al.¹⁷ suggested that targeted, location-specific reduction strategies are needed to reduce current racial and ethnic PM_{2.5} and NO₂ exposure inequalities, with Wang et al.⁴⁹ demonstrating the potential of reduced-form modeling to inform selection of locations for emission reductions. In our view, such location-specific regulatory approaches would complement the main current approaches to air pollution regulation, such as the National Ambient Air Quality Standards and sector-specific emission-reduction technology requirements.

In summary, we believe characterizing within-neighborhood ambient air pollution concentrations and health metrics will enable *a*) more refined characterization of exposure in epidemiological analyses used for defining concentration–response functions, *b*) improved exposure and baseline disease rate inputs for health impact assessments, and *c*) enhanced attribution to emissions sources. The end result would be to better prioritize policies that could substantially reduce air pollution-mediated health inequities (e.g., utility investments in electric vehicle charging infrastructure).

Monitoring Policy Effectiveness

Although the existing federal regulatory monitoring network is essential to develop and evaluate clean air policies and standards, it does not provide geographic coverage adequate to resolve neighborhood-scale pollution variability.^{50,51} Expanding innovative approaches to monitoring will help track and verify progress [i.e., accountability research (see Boogaard et al.⁵²)] in reducing air pollution exposure inequities. For example, the impact of closures of industrial facilities on air pollution and associated health outcomes was quantified using an ensemble model that incorporates land-use and satellite-derived variables to estimate weekly ambient PM2.5 concentrations in each ZIP code before and after facility closure.⁵³ Recent research has demonstrated the potential utility of fine-scale satellite atmospheric composition measurements for evaluating the impacts of policy changes on exposure disparities, such as quantifying changes in the magnitude of racial/ethnic disparities in NO2 concentrations in U.S. urban areas associated with reductions in traffic during the early months of the COVID-19 pandemic,¹⁰ evaluating the effect of diesel emission control strategies on air pollution inequity,^{8,54} and examining the localized air quality impacts of freeway rerouting.55 Studies also show that diesel emission control technologies reduce but do not eliminate disparities.^{8,56} Other research has used empirical models that incorporate regulatory monitor measurements and satellite observations to quantify changes in the magnitude of national racial/ethnic and socioeconomic air pollution disparities over multiple decades.^{9,11}

From our perspective, further advancement in methods for monitoring fine-scale changes in pollution concentrations over time (e.g., mobile monitoring,⁴⁸ low-cost sensors⁵⁷) will assist in tracking and confirmation of the impact of place-based policy interventions on reducing air pollution-related health disparities. Characterizing uncertainty (e.g., data quality, accuracy, representativeness) also must be a key priority.⁵⁸ For example, hourly, daily, and weekly temporal variation in emissions, as well as data on residents' movements through space, can refine analyses of exposure inequities.⁵⁹ In addition, movements of residents over yearly timescales should be considered when tracking effectiveness given that lessons from other place-based programs, such as hazardous waste clean-ups and allocation of education resources, indicate gentrification processes could reduce long-term benefits accrued to the original residents.^{60–62}

Recommendations

We believe that to ensure the effectiveness of future environmental and climate policies in addressing longstanding and persistent air pollution inequities, scientists, advocates, and environmental regulators must collaboratively develop new data collection and analytic frameworks to aid the design and implementation of placebased policies [Figure 1 (inspired by a California Air Resources Board white paper⁶³)]. Specifically, environmental regulators must prioritize projecting and tracking the distributional impacts of new policies at appropriate geographic and temporal scales. The U.S. EPA Scientific Advisory Board recently emphasized this need in calling for the U.S. EPA to "develop a strategy for systematic, quantitative evaluation of the environmental justice impacts of ... regulations," as part of their recent regulatory reviews of science supporting the U.S. EPA's proposed rule on control of air pollution from new motor vehicle standards.⁶⁴ Novel monitoring and modeling technologies have the potential to greatly aid the effort to characterize and abate air pollution sources contributing to health disparities. To meet that potential, regulators will need to update and expand regulatory frameworks to incorporate new data. For example, a recent analysis in California demonstrated how satellite-derived estimates can inform air quality management by screening locations not represented within the regulatory monitoring network.⁶⁵ Recent public comments from a coalition of environmental and public health advocates emphasized that the U.S. "EPA must amend its monitoring network requirements to ensure adequate monitoring of air pollution in at-risk communities, including communities of color and low-income communities" and called on the U.S. EPA to "issue guidance as soon as possible to ensure that data collected. . . using new technologies and hybrid approaches may factor into regulatory decisionmaking."⁶⁶ We recommend that regulators leverage insights from novel data sources, including community-led monitoring supported by new U.S. EPA funding, to improve and expand the regulatory monitoring network and inform regulatory decisionmaking that alleviates pollution disparities.

The scientific community must work closely with regulators, advocates, and impacted communities to advance data collection and analysis strategies that match the goals of environmental and climate justice policies, integrating diverse data streams using accessible reporting methods while also acknowledging and characterizing uncertainty as accurately as possible.⁵⁸ We assert that collaboration by governments, scientists, and advocates will be critical to address key technical, governance, and ethical questions and challenges, including how to enable broad access and




Figure 1. Diagram illustrating relationships between data sources, models, collaboration for synthesis, and policy and communication. Central circles represent components of an integrated data analysis and policy framework to maximize reduction in air pollution-related health inequities. Lighter colored rectangles are examples of observational data sets, models across health and environmental sciences, and relevant policy and communication outcomes. This figure was modified from a figure by the California Air Resources Board.⁶³

understanding of an increasingly complex data ecosystem, synthesize data from multiple sources into information useful to policymakers and community leaders, and ensure feedback mechanisms that incorporate local lived experience. Critically, this collaborative work must be grounded in the principles of environmental justice,67 antiracism,68,69 and community science,⁷⁰ including centering the priorities of impacted communities and ensuring that community representatives serve in decision-making roles. These efforts can build on work already underway, including the collaborative work of the communitybased organization Comite Civico del Valle and government and academic partners to deploy the Imperial County Community Air Monitoring Network⁷¹ and the activities of the National Aeronautics and Space Administration Health and Air Quality Applied Sciences Team to advance the use of satellite data for environmental justice (https://haqast.org/tiger-teams/), and can leverage new resources, including funding through the U.S. EPA's Environmental Justice Collaborative Problem Solving Cooperative Agreement Program (https://www.epa. gov/environmentaljustice/environmental-justice-collaborativeproblem-solving-cooperative-agreement-5), and the Environmental Justice Thriving Communities Technical Assistance Centers (https://www.epa.gov/environmentaljustice/environmental-justicethriving-communities-technical-assistance-centers).

With federal resources made available via the IIJA and IRA, we believe it is now more important than ever to advance the collection and analysis of fine-scale spatial and temporal environmental and health data to support the design, implementation, and evaluation of policies that reduce inequities in air pollution exposure and associated health impacts.

Acknowledgments

Author contributions: Conceptualization: J.M.G., M.H.H., A.R., T.M.T., and M.D.; Methodology: J.M.G., M.H.H., A.R., T.M.T., and M.D.; Investigation: J.M.G., M.H.H., A.R., T.M.T., M.D., S.C.A., J.S.A., M.A.G.D., I.M.D., G.H.K., J.D.M., A.E.N., R.F.P., S.E.P., and V.A.S.; Visualization: J.S.A., A.E.N., S.A.V., M.H.H., and J.M.G.; Funding acquisition: R.A.A. and S.A.V.; Project Administration: J.M.G., M.H.H., M.D., R.A.A., and S.A.V.; Writing—original draft: J.M.G., M.H.H., A.R., and T.M.T.; Writing—review and editing: J.M.G., M.H.H., A.R., T.M.T., M.D., R.A.A., S.C.A., J.S.A., M.A.G.D., I.M.D., G.H.K., J.D.M., A.E.N., R.F.P., S.E.P., V.A.S., and S.A.V.

This manuscript grew from discussions during a year-long virtual seminar series (2021–2022) on air quality and health inequities organized by Environmental Defense Fund. We acknowledge contributions from participants in the seminar series. We also acknowledge authors of the California Air Resources Board white paper⁶³ that inspired Figure 1.

Funding was provided to authors by the following organizations: Valhalla Foundation (J.M.G., M.H.H., A.R., T.M.T., M.D., R.A.A., A.E.N., and S.A.V.), Quadrivium Foundation (J.M.G., M.H.H., A.R., T.M.T., M.D., R.A.A., A.E.N., and S.A.V.), University of Virginia College Science Scholars Program (I.M.D.), University of Virginia Double Hoo Award (I.M.D.), National Aeronautics and Space Administration New (Early Career) Investigator Program in Earth Science (80NSSC21K0935) to S.E.P., National Science Foundation CAREER Award (AGS 2047150) to S.E.P. Data and materials availability: All data are available in the main text or the Supplemental Material.

References

- White House. 2022. FACT SHEET: Biden-Harris Administration holds Justice40 Week of Action to highlight historic investments in overburdened and underserved communities. Published online 23 May 2023. https://www.whitehouse. gov/briefing-room/statements-releases/2022/05/23/fact-sheet-biden-harrisadministration-holds-justice40-week-of-action-to-highlight-historic-investmentsin-overburdened-and-underserved-communities/ [accessed 1 November 2023].
- Illinois Power Agency. 2022. 2022 Long-term Renewable Resources Procurement Plan. Final Plan. https://ipa.illinois.gov/content/dam/soi/en/web/ipa/documents/ 2022-long-term-plan-23-august.pdf [accessed 15 January 2023].
- California Office of Environmental Health Hazard Assessment. 2022. SB535 Disadvantaged Communities, 2022 Update. https://oehha.ca.gov/calenviroscreen/sb535 [accessed 15 January 2023].
- New York State Climate Justice Working Group. 2022. Disadvantaged Communities Criteria. https://climate.ny.gov/resources/climate-justice-workinggroup/#disadvantaged-communities-map [accessed 15 January 2023].

- Washington State Department of Health. 2022. Washington Environmental Health Disparities Map, Version 2.0, Technical Report. https://doh.wa.gov/sites/ default/files/2022-07/311-011-EHD-Map-Tech-Report_0.pdf?uid=63e13f2febb79 [accessed 15 January 2023].
- Nowlan A. 2022. Historic investments in air quality monitoring can give communities a voice in clean air solutions. Published 8 September 2022. https:// globalcleanair.org/health/historic-investments-in-air-quality-monitoring-can-givecommunities-a-voice-in-clean-air-solutions/ [accessed 19 July 2023].
- White House. 2023. A Guidebook to the Bipartisan Infrastructure Law. https:// www.whitehouse.gov/build/guidebook/ [accessed 13 November 2023].
- Demetillo MAG, Harkins C, McDonald BC, Chodrow PS, Sun K, Pusede SE. 2021. Space-based observational constraints on NO₂ air pollution inequality from diesel traffic in major US cities. Geophys Res Lett 48(17):e2021GL094333, https://doi.org/10.1029/2021GL094333.
- Jbaily A, Zhou X, Liu J, Lee TH, Kamareddine L, Verguet S, et al. 2022. Air pollution exposure disparities across US population and income groups. Nature 601(7892):228–233, PMID: 35022594, https://doi.org/10.1038/s41586-021-04190-y.
- Kerr GH, Goldberg DL, Anenberg SC. 2021. COVID-19 pandemic reveals persistent disparities in nitrogen dioxide pollution. Proc Natl Acad Sci USA 118(30): e2022409118, PMID: 34285070, https://doi.org/10.1073/pnas.2022409118.
- Liu J, Clark LP, Bechle MJ, Hajat A, Kim SY, Robinson AL, et al. 2021. Disparities in air pollution exposure in the United States by race/ethnicity and income, 1990–2010. Environ Health Perspect 129(12):127005, PMID: 34908495, https://doi.org/10.1289/EHP8584.
- Tessum CW, Paolella DA, Chambliss SE, Apte JS, Hill JD, Marshall JD. 2021. PM_{2.5} polluters disproportionately and systemically affect people of color in the United States. Sci Adv 7(18):eabf4491, PMID: 33910895, https://doi.org/10. 1126/sciadv.abf4491.
- Lane HM, Morello-Frosch R, Marshall JD, Apte JS. 2022. Historical redlining is associated with present-day air pollution disparities in U.S. cities. Environ Sci Technol Lett 9(4):345–350, PMID: 35434171, https://doi.org/10.1021/acs.estlett.1c01012.
- Bramble K, Blanco MN, Doubleday A, Gassett AJ, Hajat A, Marshall JD, et al. 2023. Exposure disparities by income, race and ethnicity, and historic redlining grade in the greater Seattle area for ultrafine particles and other air pollutants. Environ Health Perspect 131(7):077004, PMID: 37404015, https://doi.org/10.1289/ EHP11662.
- Saha PK, Presto AA, Hankey S, Marshall JD, Robinson AL. 2022. Racial-ethnic exposure disparities to airborne ultrafine particles in the United States. Environ Res Lett 17(10):104047, https://doi.org/10.1088/1748-9326/ac95af.
- Clark LP, Harris MH, Apte JS, Marshall JD. 2022. National and intraurban air pollution exposure disparity estimates in the United States: impact of dataaggregation spatial scale. Environ Sci Technol Lett 9(9):786–791, PMID: 36118958, https://doi.org/10.1021/acs.estlett.2c00403.
- Dressel IM, Demetillo MAG, Judd LM, Janz SJ, Fields KP, Sun K, et al. 2022. Daily satellite observations of nitrogen dioxide air pollution inequality in New York City, New York and Newark, New Jersey: evaluation and application. Environ Sci Technol 56(22):15298–15311, PMID: 36224708, https://doi.org/10. 1021/acs.est.2c02828.
- Demetillo MAG, Navarro A, Knowles KK, Fields KP, Geddes JA, Nowlan CR, et al. 2020. Observing nitrogen dioxide air pollution inequality using high-spatialresolution remote sensing measurements in Houston, Texas. Environ Sci Technol 54(16):9882–9895, PMID: 32806912, https://doi.org/10.1021/acs.est.0c01864.
- Braveman P, Gottlieb L. 2014. The social determinants of health: it's time to consider the causes of the causes. Public Health Rep 129(suppl 2):19–31, PMID: 24385661, https://doi.org/10.1177/00333549141291S206.
- Homan P, Brown TH, King B. 2021. Structural intersectionality as a new direction for health disparities research. J Health Soc Behav 62(3):350–370, PMID: 34355603, https://doi.org/10.1177/00221465211032947.
- Jeffries N, Zaslavsky AM, Diez Roux AV, Creswell JW, Palmer RC, Gregorich SE, et al. 2019. Methodological approaches to understanding causes of health disparities. Am J Public Health 109(S1):S28–S33, PMID: 30699015, https://doi.org/10.2105/ AJPH.2018.304843.
- Gee GC, Payne-Sturges DC. 2004. Environmental health disparities: a framework integrating psychosocial and environmental concepts. Environ Health Perspect 112(17):1645–1653, PMID: 15579407, https://doi.org/10.1289/ehp.7074.
- Payne-Sturges DC, Cory-Slechta DA, Puett RC, Thomas SB, Hammond R, Hovmand PS. 2021. Defining and intervening on cumulative environmental neurodevelopmental risks: introducing a complex systems approach. Environ Health Perspect 129(3):035001, PMID: 33688743, https://doi.org/10.1289/EHP7333.
- Castillo MD, Kinney PL, Southerland V, Arno CA, Crawford K, van Donkelaar A, et al. 2021. Estimating intra-urban inequities in PM_{2.5}-attributable health impacts: a case study for Washington, DC. Geohealth 5(11):e2021GH000431, PMID: 34765851, https://doi.org/10.1029/2021GH000431.
- 25. Southerland VA, Anenberg SC, Harris M, Apte J, Hystad P, van Donkelaar A, et al. 2021. Assessing the distribution of air pollution health risks within cities: a neighborhood-scale analysis leveraging high-resolution data sets in the Bay

Area, California. Environ Health Perspect 129(3):037006, PMID: 33787320, https://doi.org/10.1289/EHP7679.

- Di Q, Amini H, Shi L, Kloog I, Silvern R, Kelly J, et al. 2019. An ensemblebased model of PM_{2.5} concentration across the contiguous United States with high spatiotemporal resolution. Environ Int 130:104909, PMID: 31272018, https://doi.org/10.1016/j.envint.2019.104909.
- Di Q, Wang Y, Zanobetti A, Wang Y, Koutrakis P, Choirat C, et al. 2017. Air pollution and mortality in the Medicare population. N Engl J Med 376(26):2513–2522, PMID: 28657878, https://doi.org/10.1056/NEJMoa1702747.
- Industrial Economics, Incorporated. 2022. Analysis of PM_{2.5}-Related Health Burdens Under Current and Alternative NAAQS. https://globalcleanair.org/ files/2022/05/Analysis-of-PM2.5-Related-Health-Burdens-Under-Current-and-Alternative-NAAQS.pdf [accessed 1 February 2023].
- Spiller E, Proville J, Roy A, Muller NZ. 2021. Mortality risk from PM_{2.5}: a comparison of modeling approaches to identify disparities across racial/ethnic groups in policy outcomes. Environ Health Perspect 129(12):127004, PMID: 34878311, https://doi.org/10.1289/EHP9001.
- Daouda M, Henneman L, Goldsmith J, Kioumourtzoglou MA, Casey JA. 2022. Racial/ethnic disparities in nationwide PM_{2.5} concentrations: perils of assuming a linear relationship. Environ Health Perspect 130(7):077701, PMID: 35857400, https://doi.org/10.1289/EHP11048.
- Tessum CW, Apte JS, Goodkind AL, Muller NZ, Mullins KA, Paolella DA, et al. 2019. Inequity in consumption of goods and services adds to racial-ethnic disparities in air pollution exposure. Proc Natl Acad Sci USA 116(13):6001–6006, PMID: 30858319, https://doi.org/10.1073/pnas.1818859116.
- Lateb M, Meroney RN, Yataghene M, Fellouah H, Saleh F, Boufadel MC. 2016. On the use of numerical modelling for near-field pollutant dispersion in urban environments—a review. Environ Pollut 208(pt A):271–283, PMID: 26282585, https://doi.org/10.1016/j.envpol.2015.07.039.
- Russell A, Dennis R. 2000. NARSTO critical review of photochemical models and modeling. Atmos Environ 34(12–14):2283–2324, https://doi.org/10.1016/ S1352-2310(99)00468-9.
- U.S. EPA (U.S. Environmental Protection Agency). 2015. Air Emissions Inventories. Published 11 March 2015. https://www.epa.gov/air-emissions-inventories/nationalemissions-inventory-nei [accessed 13 November 2023].
- Goodkind AL, Tessum CW, Coggins JS, Hill JD, Marshall JD. 2019. Fine-scale damage estimates of particulate matter air pollution reveal opportunities for location-specific mitigation of emissions. Proc Natl Acad Sci USA 116(18):8775– 8780, PMID: 30962364, https://doi.org/10.1073/pnas.1816102116.
- Harkins C, McDonald BC, Henze DK, Wiedinmyer C. 2021. A fuel-based method for updating mobile source emissions during the COVID-19 pandemic. Environ Res Lett 16(6):065018, https://doi.org/10.1088/1748-9326/ac0660.
- Li M, McDonald BC, McKeen SA, Eskes H, Levelt P, Francoeur C, et al. 2021. Assessment of updated fuel-based emissions inventories over the contiguous United States using TROPOMI NO₂ retrievals. J Geophys Res 126(24): e2021JD035484.
- Ma S, Tong DQ. 2022. Neighborhood emission mapping operation (NEMO): a 1km anthropogenic emission dataset in the United States. Sci Data 9(1):680, PMID: 36351966, https://doi.org/10.1038/s41597-022-01790-9.
- Shah RU, Robinson ES, Gu P, Apte JS, Marshall JD, Robinson AL, et al. 2020. Socio-economic disparities in exposure to urban restaurant emissions are larger than for traffic. Environ Res Lett 15(11):114039, https://doi.org/10.1088/ 1748-9326/abbc92.
- Clark LP, Millet DB, Marshall JD. 2014. National patterns in environmental injustice and inequality: outdoor NO₂ air pollution in the United States. PLoS One 9(4):e94431, PMID: 24736569, https://doi.org/10.1371/journal.pone.0094431.
- Clark LP, Millet DB, Marshall JD. 2017. Changes in transportation-related air pollution exposures by race-ethnicity and socioeconomic status: outdoor nitrogen dioxide in the United States in 2000 and 2010. Environ Health Perspect 125(9):097012, PMID: 28930515, https://doi.org/10.1289/EHP959.
- Chambliss SE, Pinon CPR, Messier KP, LaFranchi B, Upperman CR, Lunden MM, et al. 2021. Local- and regional-scale racial and ethnic disparities in air pollution determined by long-term mobile monitoring. Proc Natl Acad Sci USA 118(37):e2109249118, PMID: 34493674, https://doi.org/10.1073/pnas.2109249118.
- Rowangould GM. 2013. A census of the US near-roadway population: public health and environmental justice considerations. Transp Res D Transp Environ 25:59–67, https://doi.org/10.1016/j.trd.2013.08.003.
- Nguyen NP, Marshall JD. 2018. Impact, efficiency, inequality, and injustice of urban air pollution: variability by emission location. Environ Res Lett 13(2):024002, https://doi.org/10.1088/1748-9326/aa9cb5.
- Thind MPS, Tessum CW, Azevedo IL, Marshall JD. 2019. Fine particulate air pollution from electricity generation in the US: health impacts by race, income, and geography. Environ Sci Technol 53(23):14010–14019, PMID: 31746196, https://doi.org/10.1021/acs.est.9b02527.
- Luo Q, Copeland B, Garcia-Menendez F, Johnson JX. 2022. Diverse pathways for power sector decarbonization in Texas yield health cobenefits but fail to

alleviate air pollution exposure inequities. Environ Sci Technol 56(18):13274–13283, PMID: 36070515, https://doi.org/10.1021/acs.est.2c00881.

- Thind MPS, Tessum CW, Marshall JD. 2022. Environmental health, racial/ethnic health disparity, and climate impacts of inter-regional freight transport in the United States. Environ Sci Technol 57(2):884–895, PMID: 36580637, https://doi.org/ 10.1021/acs.est.2c03646.
- Wang Y, Apte JS, Hill JD, Ivey CE, Patterson RF, Robinson AL, et al. 2022. Location-specific strategies for eliminating US national racial-ethnic PM_{2.5} exposure inequality. Proc Natl Acad Sci USA 119(44):e2205548119, PMID: 36279443, https://doi.org/10.1073/pnas.2205548119.
- Wang Y, Apte JS, Hill JD, Ivey CE, Johnson D, Min E, et al. 2023. Air quality policy should quantify effects on disparities. Science 381(6655):272–274, PMID: 37471550, https://doi.org/10.1126/science.adg9931.
- deSouza P, Kinney PL. 2021. On the distribution of low-cost PM_{2.5} sensors in the US: demographic and air quality associations. J Expo Sci Environ Epidemiol 31(3):514–524, PMID: 33958706, https://doi.org/10.1038/s41370-021-00328-2.
- Sullivan D, Krupnick A. 2018. Using Satellite Data to Fill the Gaps in the US Air Pollution Monitoring Network. Resources for the Future. https://www.rff.org/ publications/working-papers/using-satellite-data-to-fill-the-gaps-in-the-us-air-pollutionmonitoring-network/ [accessed 4 July 2023].
- Boogaard H, van Erp AM, Walker KD, Shaikh R. 2017. Accountability studies on air pollution and health: the HEI experience. Curr Environ Health Rep 4(4):514– 522, PMID: 28988407, https://doi.org/10.1007/s40572-017-0161-0.
- Chen C, Ilango SD, Henneman LRF, Casey JA, Benmarhnia T. 2023. The local impacts of coal and oil power plant retirements on air pollution and cardiorespiratory health in California: an application of generalized synthetic control method. Environ Res 226:115626, PMID: 36907346, https://doi.org/10.1016/j. envres.2023.115626.
- Patterson RF, Harley RA. 2021. Effects of diesel engine emission controls on environmental equity and justice. Environ Justice 14(5):360–371, https://doi.org/ 10.1089/env.2020.0078.
- Patterson RF, Harley RA. 2019. Effects of freeway rerouting and boulevard replacement on air pollution exposure and neighborhood attributes. Int J Environ Res Public Health 16(21):4072, PMID: 31652720, https://doi.org/10.3390/ ijerph16214072.
- Marshall JD, Swor KR, Nguyen NP. 2014. Prioritizing environmental justice and equality: diesel emissions in Southern California. Environ Sci Technol 48(7):4063–4068, PMID: 24559220, https://doi.org/10.1021/es405167f.
- Do K, Yu H, Velasquez J, Grell-Brisk M, Smith H, Ivey CE. 2021. A data-driven approach for characterizing community scale air pollution exposure disparities in inland Southern California. J Aerosol Sci 152:105704, https://doi.org/10.1016/j. jaerosci.2020.105704.
- National Academies of Sciences, Engineering, and Medicine. 2022. Communities, Climate Change, and Health Equity—State-Level Implementation: Proceedings of a Workshop—in Brief. Washington, DC: National Academies Press.
- Park YM, Kwan MP. 2017. Individual exposure estimates may be erroneous when spatiotemporal variability of air pollution and human mobility are ignored. Health Place 43:85–94, PMID: 27914271, https://doi.org/10.1016/j.healthplace.2016.10.002.

- Acolin A, Crowder K, Decter-Frain A, Hajat A, Hall M. 2022. Gentrification, mobility, and exposure to contextual determinants of health. Hous Policy Debate 33(1):194–223, PMID: 37200539, https://doi.org/10.1080/10511482.2022.2099937.
- Loh J, Loh TH. 2020. How we define 'need' for place-based policy reveals where poverty and race intersect. https://www.brookings.edu/research/howwe-define-need-for-place-based-policy-reveals-where-poverty-and-raceintersect/ [accessed 6 September 2022].
- Cassidy AW, Hill EL, Ma L. 2022. Who benefits from hazardous waste cleanups? Evidence from the housing market. NBER Working Paper No. w30661. https://doi.org/10.2139/ssrn.4282511.
- 63. Cohen RC, Preble CV, Apte JS, Kirchstetter TW, Arnett M. 2023. An integrated framework to guide improvements in air quality at community, urban, and regional scales: Recommendations for the future of the air quality ecosystem in California. https://ww2.arb.ca.gov/integrated-framework-guide-improvements-air-quality-community-urban-and-regional-scales [accessed 13 November 2023].
- U.S. EPA. 2022. Science Advisory Board Regulatory Review Report of Science Supporting EPA Decisions for the Proposed Rule: Control of Air Pollution from New Motor Vehicles: Heavy-Duty Engine and Vehicle Standards (RIN 2060-AU41). EPA-SAB-23-001. https://sab.epa.gov/ords/sab/f?p=114:18:9476967133120:::RP,18: P18_ID:2625 [accessed 15 February 2023].
- Gladson L, Garcia N, Bi J, Liu Y, Lee HJ, Cromar K. 2022. Evaluating the utility of high-resolution spatiotemporal air pollution data in estimating local PM_{2.5} exposures in California from 2015–2018. Atmosphere (Basel) 13(1):85, https://doi.org/10. 3390/atmos13010085.
- U.S. EPA. 2023. Comment submitted by Appalachian Mountain Club et al. to EPA Apr 3, 2023 EPA-HQ-OAR-2015-0072-1543. Published online 3 April 2023. https://www.regulations.gov/comment/EPA-HQ-OAR-2015-0072-2233 [accessed 5 July 2023].
- Van Horne YO, Alcala CS, Peltier RE, Quintana PJE, Seto E, Gonzales M, et al. 2023. An applied environmental justice framework for exposure science. J Expo Sci Environ Epidemiol 33(1):1–11, PMID: 35260805, https://doi.org/10.1038/ s41370-022-00422-z.
- Payne-Sturges DC, Gee GC, Cory-Slechta DA. 2021. Confronting racism in environmental health sciences: moving the science forward for eliminating racial inequities. Environ Health Perspect 129(5):055002, PMID: 33945300, https://doi.org/10. 1289/EHP8186.
- Wilkins D, Schulz AJ. 2023. Antiracist research and practice for environmental health: implications for community engagement. Environ Health Perspect 131(5):055002, PMID: 37224068, https://doi.org/10.1289/EHP11384.
- Pandya R, Aurbach EL, Burns K, Chalk AM. 2023. Recommendations for an NSF Convergence Accelerator Track on Community Science A Community Science Report of an NSF-funded Convergence Accelerator Workshop Facilitated by the American Geophysical Union. ESS Open Archive. Preprint posted online March 1 2023, https://doi.org/10.22541/essoar.167768122.22544063/v1.
- English PB, Olmedo L, Bejarano E, Lugo H, Murillo E, Seto E, et al. 2017. The Imperial County Community Air Monitoring Network: a model for communitybased environmental monitoring for public health action. Environ Health Perspect 125(7):074501, PMID: 28886604, https://doi.org/10.1289/EHP1772.