

Lawrence Berkeley National Laboratory

Recent Work

Title

K+n CHARGE EXCHANGE AND TIE , REGGE TRAJECTORY

Permalink

<https://escholarship.org/uc/item/4bw3z0xm>

Authors

Rarita, William
Schwarzschild, Bertram M.

Publication Date

1967-05-17

University of California
Ernest O. Lawrence
Radiation Laboratory

K^+_n CHARGE EXCHANGE AND THE ρ^1 REGGE TRAJECTORY

William Rarita and Bertram M. Schwarzschild

May 17, 1967

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to The Physical Review

UCRL-17572 Rev.
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

K_n^+ CHARGE EXCHANGE AND THE ρ' REGGE TRAJECTORY

William Rarita and Bertram M. Schwarzschild

May 17, 1967

K^+n CHARGE EXCHANGE AND
THE ρ' REGGE TRAJECTORY*

William Rarita[†]

Department of Physics
University of California
Berkeley, California

and

Bertram M. Schwarzschild[‡]

Lawrence Radiation Laboratory
University of California
Berkeley, California

May 17, 1967

ABSTRACT

Finding that the Regge exchange of ρ and A_2 fails to describe our K^+n charge-exchange data at 2.3 GeV/c (the highest energy available to date) when a simultaneous fit is attempted with higher energy data on K^+p and π^-p charge exchange and $\pi^-p \rightarrow \eta^0 n$, we introduce a second, lower lying ρ type trajectory (ρ'). This also provides a possible mechanism for the puzzling π^-p charge-exchange polarization. We find that we are then able simultaneously to fit all these data (including the polarization), together with related total cross-section differences up to 20 GeV/c, with a ρ' whose spin 1 mass is 1.0 GeV [perhaps the $\delta(965)$?] and whose $t=0$ intercept, 1.1 units of angular momentum below the ρ , agrees roughly with the ρ' proposed by Høgaasen and Fischer to describe forward $\bar{p}p$ and np charge exchange, where the ρ , A_2 model also fails. Our ρ and A_2 trajectories turn out essentially traditional. In the fit we permit only small SU_3 breaking between the $K\bar{K}$ and $\pi\pi$ (or $\eta^0\pi$) couplings to the trajectories. We further constrain the fit to obey the

sum rule of Igi and Matsuda. In fitting our K^+ data at 2.3 GeV/c, we include a deuteron correction, and employ exact Legendre functions rather than the high-energy asymptotic Regge forms. We offer predictions for higher energy K^+ charge exchange.

INTRODUCTION

Considerable success has previously been achieved in fitting the high-energy differential cross sections for the reactions:

$$\pi^- p \rightarrow \pi^0 n \quad (1)$$

$$K^- p \rightarrow \bar{K}^0 n \quad (2)$$

$$\pi^- p \rightarrow \eta^0 n \quad (3)$$

in the peripheral region $[-t \lesssim 1(\text{GeV}/c)^2]$, by the t -channel Regge exchange of the $\rho(C^-)$ and $A_2(C^+)$ trajectories.^{1,2} Together with

$$K^+ n \rightarrow K^0 p, \quad (4)$$

these constitute all the isotopically independent reactions

$$PS8 + N \rightarrow PS8 + N \quad (PS8 = \text{Pseudoscalar Octet}, N = \text{Nucleon}) \quad (5)$$

which require $I = 1$ exchange in the t channel. For t -channel exchange, reaction (4) differs from (2) only by a relative sign change in the C^+ and C^- exchange amplitudes. Thus, if the (ρ, A_2) model is adequate, the previous fits ought to determine high-energy $K^+ n$ peripheral charge exchange.

We therefore compared $K^+ n$ charge-exchange data at 2.3 GeV/c (the highest energy available to date) of one of the authors (BMS)³ to the previous fits of the other (WR)¹, and attempted a simultaneous fit of data on all four reactions to the assumption of only ρ and A_2 Regge exchange in the peripheral region. The attempt was in part motivated by the correct prediction of this model that, in contrast to $K^- p$, the $K^+ n$ charge exchange ought to show $|\text{Re } f(t=0)/\text{Im } f(t=0)| \gg 1$. Despite correction

for the deuteron effect in (4), and use of exact Legendre function forms because of the relatively low energy of our K^+ data, we found that the model always gave a cross section only about half as large as that observed for K^+n charge exchange. A kinder fate would have been expected considering the relative absence of resonant activity in the K^+ nucleon system above about 1.3 GeV/c.

The model has other difficulties. It predicts, contrary to fact, zero polarization for π^-p charge exchange. Further, Högaasen and Fischer find that it fails to describe the energy dependence of forward pn and $\bar{p}p$ charge exchange.⁴ There seems to be a simultaneous plausible remedy for all three difficulties, namely, the exchange of a second, lower lying, ρ' trajectory having the same quantum numbers as the ρ . Lying well below the ρ trajectory, its relative contribution to the differential cross sections would decrease rapidly with increasing energy, thus explaining the earlier fits which ignored it.

Adding this ρ' trajectory to the C^- exchange amplitude, which we then subject to the constraint of a recent superconvergent sum rule of Igi and Matsuda,⁵ we have achieved a good simultaneous fit to extensive data on processes (1) through (4), together with related total cross-section differences and π^-p charge-exchange polarization. This fit involves only small SU_3 symmetry breaking, the ρ' being assumed to belong to an octet. The resulting ρ' trajectory is consistent with the intercept found by Högaasen and with the $\delta(965 \text{ MeV})$ meson at $\alpha = 1$.

FORMALISM

All the processes (5) (see Fig. 1) require that the t-channel exchanged object satisfy $C = P = (-1)^J = G(-1)^I$. Anticipating Reggeization and SU_3 , we separate the $I = 1$ exchange into two parts:

$$C^+(P^+, G^- \text{ J even}) \text{ and } C^-(P^-, G^+, \text{ J odd}).$$

Clearly, for reaction (1) only the C^- exchange is possible, while for (3) we have only C^+ . The $K^\pm N$ charge exchanges admit both. Therefore we define the t-channel helicity nonflip amplitudes

$$A(K^- p \rightarrow \bar{K}^0 n) = {}_{\kappa} A^+ + {}_{\kappa} A^- \quad (6)$$

$$A(\pi^- p \rightarrow \pi^0 n) = {}_{\pi} A^- \quad (7)$$

$$A(\pi^- p \rightarrow \eta^0 n) = {}_{\pi} A^+, \quad (8)$$

giving

$$A(K^+ n \rightarrow K^0 p) = {}_{\kappa} A^+ - {}_{\kappa} A^- \quad (9)$$

with similar formulae for the helicity flip amplitude B. The superscript sign refers to both charge conjugation and J parity (or signature) $(-1)^J = C$.

Now let each amplitude be a sum over contributing Regge trajectories, e. g.,

$${}_{\pi} A^- = \sum_i {}_{\pi} A_i^- \text{ etc.} \quad (10)$$

Then we have from the factorization theorem:

$${}_{\pi} A_i / {}_{\kappa} A_i = {}_{\pi} B_i / {}_{\kappa} B_i \equiv F_i(t) \quad (11)$$

If SU_3 is unbroken, and all contributing trajectories are octet members, we have

$$F_i(t) = \begin{cases} F^+ = (2/3)^{1/2} \text{ for } C_i = +1, \\ F^- = -\sqrt{2} \text{ for } C_i = -1 \end{cases} \quad (12)$$

independent of t .

The experimentally observed quantities are given in terms of the helicity amplitudes by:

$$\frac{d\sigma}{dt} = \frac{1}{\pi} \left(\frac{1}{4p}\right)^2 \left\{ \left(1 - \frac{t}{4M^2}\right) |A|^2 - \frac{t}{4M^2} \left[\frac{s + P^2}{1 - (t/4M^2)} - s \right] |B|^2 \right\} \quad (13)$$

$$\sigma_T(K^\pm p) - \sigma_T(K^\pm n) = \text{Im}(\kappa A^+ \mp \kappa A^-)_{t=0}/p \quad (14)$$

$$\sigma_T(\pi^+ p) - \sigma_T(\pi^- p) = \sqrt{2} \text{Im}(\pi A^-)_{t=0}/p \quad (15)$$

$$P \frac{d\sigma}{dt} (\pi^- p \rightarrow \pi^0 n) =$$

$$\frac{1}{\pi} \left(\frac{1}{4p^2}\right) \left[\left(1 - \frac{t}{4M^2}\right) \right]^{1/2} \left\{ \frac{-t}{4M^2} \left[\frac{s + p^2}{1 - (t/4M^2)} - s \right] \right\}^{1/2} \pi_{\pi} A^- \times \pi_{\pi} B^- \quad (16)$$

where M is the nucleon mass, s and t are the invariant squares of energy and momentum transfer, p is the incoming lab momentum, and P is the $\pi^- p$ charge-exchange polarization. \underline{A} and \underline{B} are vectors in the complex plane.

There is only one reasonably well established trajectory for each of C^+ and C^- , namely the A_2 and ρ . As discussed above, however, they do not seem to suffice for all the processes under consideration. Including now the speculative ρ' , we parametrize the various high-energy asymptotic Regge helicity amplitudes as follows:

$$\kappa A^- = \kappa A_\rho + \kappa A_{\rho'}, \text{ and } \kappa A^+ = \kappa A_{A_2} \equiv \kappa A_R \quad (10')$$

with similar relations for the κB , πA , and πB amplitudes.

$$\kappa A_i^\pm = -C_i(t) (\alpha_i + 1) \frac{(e^{-i\pi\alpha_i \pm 1})}{\sin \pi\alpha_i} \left(\frac{E}{E_0}\right)^{\alpha_i} \quad (17)$$

$$\kappa B_i^\pm = -D(t) \alpha_i (\alpha_i + 1) \frac{(e^{-i\pi\alpha_i \pm 1})}{\sin \pi\alpha_i} \left(\frac{E}{E_0}\right)^{\alpha_i - 1},$$

where E/E_0 is the lab energy of the incoming meson in GeV.

We take

$$\begin{aligned} C_\rho(t) &= C_\rho^0 (1 + C_\rho^1 t) & D_\rho(t) &= D_\rho^0 \exp(D_\rho^1 t) \\ C_{\rho'}(t) &= C_{\rho'}^0 \exp(C_{\rho'}^1 t) & D_{\rho'}(t) &= D_{\rho'}^0 \exp(D_{\rho'}^1 t) \\ C_R(t) &= \alpha_R C_R^0 \exp(C_R^1 t) & D_R(t) &= \alpha_R D_R^0 \exp(D_R^1 t) \end{aligned} \quad (17')$$

$$\alpha_i^0(t) = \alpha_i^0 + \alpha_i^1 t \quad (18)$$

$$F_i(t) = F_i^0 \exp(F_i^1 t). \quad (11')$$

The distribution of the factors of α in Eqs. (17) implies a specific mechanism for the required vanishing of flip residue functions and the "ghost killing" for even-signature amplitudes at $\alpha=0$. We assume that the "nonsense" vertices (see Fig. 2) each provide a factor $\sqrt{\alpha}$ for all exchanged trajectories, and that in the case of even-trajectory exchange, every vertex provides an additional factor $\sqrt{\alpha}$. This is the so-called Chew ghost-killing mechanism, with all trajectories "choosing sense." Alternate mechanisms have been suggested, but there is some evidence in favor of our choice.⁶

For $C_\rho(t)$, we chose originally the form $C^0 \{ [G+1] \exp(C^1 t) - G \}$ to provide a possible mechanism for the crossover of the $\pi^+ p$ and $\pi^- p$ elastic differential cross sections via a sign change in $C_\rho(t)$. The fitting program always chose the case $G \gg 1 \gg C^1$. Hence the form in (17').

Table I shows the data to be fitted. Of the 24 parameters in Eqs. (17') to (11') varied to fit these data, one degree of fitting freedom is lost via the constraint of the Igi sum rule discussed below, which relates α_ρ^0 , α_ρ^0 , C_ρ^0 , C_ρ^0 , F_ρ^0 , and F_ρ^0 , in an equation of constraint. We have also constrained the six SU₃ breaking parameters F_i^0 and F_i^1 to give symmetry breaking of less than 25%. Thus we are left effectively with 17 free parameters and 6 restricted parameters.

SPECIAL TREATMENT OF LOW-ENERGY K⁺N DATA

The K⁺n charge-exchange data at 2.3 GeV/c require special treatment, (a) because of the relatively low energy for the applicability of the high-energy asymptotic forms (17), and (b) because the target neutron is bound in a deuteron.

A. Explicit Legendre Function Formulation

The high-energy Regge behavior ($\sim E^\alpha$) of the amplitudes comes from the asymptotic behavior of the Legendre functions:

$$\left. \begin{array}{l} \alpha > -1/2, P_\alpha(w) \\ \alpha < +1/2, M_\alpha(w) \equiv \frac{-\tan\pi\alpha}{\pi} Q_{-\alpha-1}(w) \end{array} \right\} \xrightarrow{w \rightarrow \infty} \frac{(2w)^\alpha}{\sqrt{\pi}} \frac{\Gamma(\alpha + \frac{1}{2})}{\Gamma(\alpha + 1)} \quad (19)$$

where P and Q are the Legendre functions of the first and second kind. Following a suggestion of Read et al.⁷ based on Mandelstam's extension⁸ of the Regge formalism to $\alpha < -1/2$, we will use P_α for $\alpha \geq 0$ and M_α for $\alpha < 0$, denoting this generically by L_α .

Retreating from asymptotically high energies, we may write the nonflip amplitudes more generally as

$$A_i^\pm = -a(t) \left(\frac{-qq'}{ME_0} \right)^\alpha (\alpha + 1/2) L_\alpha(w) \frac{(e^{-i\pi\alpha_{\pm 1}})}{\sin \pi\alpha} \quad (20)$$

where q , q' , and $w = -\cos \theta_t$ are the (nonphysical) momenta and scattering angle cosine in the t channel. These are given by

$$\begin{aligned} t &= 4(q^2 + M^2) = 4(q'^2 + m^2) \\ -s &= q^2 + q'^2 - 2qq' \cos \theta_t \end{aligned} \quad (21)$$

giving

$$w = \frac{2(s - M^2 - m^2) + t}{(4M^2 - t)^{1/2} (4m^2 - t)^{1/2}} \quad (22)$$

$m = (E^2 - p^2)^{1/2}$ is the meson mass. $(-qq'/ME_0)^\alpha = \{[(4M^2 - t)(4m^2 - t)]^{1/2}/4ME_0\}^\alpha$ is factored out of the residue function $a(t)$ to cancel the anomalous t -channel threshold singularities in w . From (19) we have then

$$A_i^\pm \xrightarrow{E \rightarrow \infty} -a(t) \left(\frac{2E}{E_0} \right)^\alpha \frac{(\alpha + 1/2)}{\sqrt{\pi}} \frac{\Gamma(\alpha + \frac{1}{2})}{\Gamma(\alpha + 1)} \frac{(e^{-i\pi\alpha_{\pm 1}})}{\sin \pi\alpha} \quad (23)$$

Comparing (23) with (17) we have finally in terms of the fitted parameters,

$$\kappa A_i^\pm = -C_i(t) (\alpha_i + 1) \frac{\Gamma(\alpha_i + 1)}{\Gamma(\alpha_i + \frac{1}{2})} \frac{\sqrt{\pi}}{2\alpha_i} \left(\frac{-qq'}{ME_0} \right)^{\alpha_i} L_{\alpha_i}(w) \frac{(e^{-i\pi\alpha_{i\pm 1}})}{\sin \pi\alpha_i} \quad (20')$$

Similarly we write the general form for the helicity flip amplitudes

$$B_i^\pm = -b(t) \left(\frac{-qq'}{ME_0} \right)^{\alpha-1} (\alpha + \frac{1}{2}) \frac{d}{dw} L_\alpha(w) \frac{(e^{-i\pi\alpha_{\pm 1}})}{\sin \pi\alpha} \quad (24)$$

Invoking the general property $d/dw L_\alpha(w) = \alpha(w L_\alpha - L_{\alpha-1})/(w^2 - 1)$, we get

$$B_i^\pm \xrightarrow{w \rightarrow \infty} -b(t) \left(\frac{2E}{E_0}\right)^{\alpha-1} \frac{2\alpha(\alpha+\frac{1}{2})}{\sqrt{\pi}} \frac{\Gamma(\alpha+\frac{1}{2})}{\Gamma(\alpha+1)} \frac{(e^{-i\pi\alpha_{i\pm 1}})}{\sin \pi\alpha}, \quad (25)$$

which by comparison with (17) gives

$$\kappa B_i^\pm = -D_i(t) (\alpha_i + 1) \frac{\Gamma(\alpha_i + 1)}{\Gamma(\alpha_i + \frac{1}{2})} \frac{\sqrt{\pi}}{2\alpha_i} \left(\frac{-qq'}{ME_0}\right)^{\alpha_i-1} \frac{d}{dw} L_{\alpha_i}(w) \left(\frac{e^{-i\pi\alpha_{i\pm 1}}}{\sin \pi\alpha_i}\right). \quad (24')$$

The factor $[\Gamma(\alpha+1)]^{-1}$ in (19) serves to cancel the unwanted poles in the signature factors for negative α . In our formalism we have effectively replaced it by $(\alpha+1)$. So long as we have $\alpha^+ > -2$ and $\alpha^- > -3$ (which turns out to be the case), the empirical factors $C(t)$ and $D(t)$ can make up the difference.

We replace Eqs. (17) by (20') and (24') when fitting the K^+ charge-exchange and total cross-section difference data at 2.3 GeV/c. To all the rest of our data, which are above 5 GeV/c, we apply the asymptotic forms.

B. Deuteron Correction

We must now express the observed $K^+ d \rightarrow K^0 p(p)$ distribution $(d\sigma/dt)_d$ in terms of the free neutron $K^+ n$ charge-exchange cross section $(d\sigma/dt)_n$, and thus in terms of the Regge amplitudes. The data $(d\sigma/dt)_d$ had been determined by attributing to each K^0 the lab momentum p' it would have if its observed direction \hat{p}' had resulted from a collision with a stationary free neutron.³ Then the impulse and closure approximation gives⁹:

$$\left(\frac{d\sigma}{dt}\right)_d = \left(\frac{d\sigma}{dt}\right)_n \frac{[1 - H + R(1 - H/3)]}{(1 + R)}, \quad (26)$$

where

$$R(t) = \frac{(\frac{d\sigma}{dt})_n^{\text{spin flip}}}{(\frac{d\sigma}{dt})_n^{\text{nospin flip}}}$$

and

$$H(t) = \int d^3 \underline{r} |\psi_d(\underline{r})|^2 \exp[i(\hat{p}' p' - p) \cdot \underline{r}], \quad (27)$$

$\psi_d(\underline{r})$ being the Hulthén deuteron wave function.

For $t=0$ we have $H=1$ and $R=0$, causing $(d\sigma/dt)_d$ to vanish in the forward direction whether or not the two-body cross section vanishes. With increasing $-t$, H approaches zero, falling to about 0.1 at $t = -0.13$ GeV/c, causing $(d\sigma/dt)_d$ to approach $(d\sigma/dt)_n$.

It remains to make the connection between R and the t -channel helicity amplitudes A, B . $R = f_2^2/f_1^2$ where f_1 and f_2 are, in conventional notation, given by $f(\theta) = f_1 + f_2 \sigma \cdot \hat{n}$. Now the s -channel helicity amplitudes g_1 and g_2 are given by $f = g_1 + g_2 \sigma \cdot \hat{k}_f \sigma \cdot \hat{k}_i$. Comparing the two representations and using well known properties of the Pauli matrices σ , we get

$$R = \left| \frac{g_2 \sin \theta}{g_1 + g_2 \cos \theta} \right|^2, \quad (28)$$

where θ is the c.m. scattering angle. Finally, having followed the formalism of Singh¹⁰ for the t -channel helicity amplitudes, we have

$$g_1 = \frac{(k^2 + M^2)^{1/2} \pm M}{8\pi\sqrt{s}} [\pm A' + (\sqrt{s} \mp M)B] \quad (29)$$

$$A = A' + \frac{E + (t/4M)}{1 - (t/4M^2)} B ,$$

k being the c. m. momentum.

In summary then, the K^+n charge-exchange data $(d\sigma/dt)_d$ are fitted to Eq. (26) in which the factor $(d\sigma/dt)_n$ is given by the Regge two-body formula (13). Note that the free neutron cross section, especially near the forward direction, cannot be extracted from the deuterium data without the aid of a model which gives $R(t)$.

A SUPERCONVERGENT SUM RULE

To test the validity of additional Regge poles (and cuts) with the quantum numbers of the ρ , proposed to explain the π^-p charge-exchange polarization, Igi and Matsuda have obtained a superconvergent sum rule.⁵ From a dispersion relation for that part of the $t=0$ amplitude which vanishes at infinity faster than E^{-1} (i. e., $\alpha < -1$), they get the sum rule

$$4\pi f^2 = \frac{1}{2\pi} \int_{E=m}^{\infty} \{p[\sigma_T(\pi^+p) - \sigma_T(\pi^-p)] - \sqrt{2} \sum_i \beta_i L_{\alpha_i^0}(\frac{E}{m})\} dE, \quad (30)$$

where $\sum \beta L$ is the sum, over contributing singularities with $\alpha^0 > -1$ (in our case ρ and ρ'), of the imaginary parts of the forward Regge π^-p charge-exchange scattering amplitudes. $f^2 = 0.081$ is the πN coupling constant squared.

$$\beta_i L_{\alpha_i^0} \left(\frac{E}{M} \right) \xrightarrow{E \rightarrow \infty} \text{Im}_{\pi} A_i(t=0) \text{ asymptotic}$$

giving

$$\beta_i = F_i^0 C_i^0 (\alpha_i^0 + 1) \frac{\Gamma(\alpha_i^0 + 1)}{\Gamma(\alpha_i^0 + \frac{1}{2})} \sqrt{\pi} \left(\frac{m}{2} \right)^{\alpha_i^0}. \quad (30')$$

The integrand of (30) vanishes at energies sufficiently high that all contributions other than ρ and ρ' become negligible [cf. optical theorem, Eq. (15)]. We take the integral up to $39m$ (≈ 5.5 GeV/c), using Igi's numerical determination of $\int p \Delta \sigma_T dE$. Then Eqs. (30) relate C_ρ^0 , $C_{\rho'}^0$, α_ρ^0 , $\alpha_{\rho'}^0$, F_ρ^0 , and $F_{\rho'}^0$, in an equation of constraint, which we impose upon the fitting.

RESULTS

The fitted parameters resulting from a least-squares fit to the data are given in Table II. In Figs. 3 through 9 the resulting theoretical curves are shown, superimposed upon the data. For 194 data points we have a χ^2 of 191. The ρ and A_2 trajectories turn out essentially traditional, i. e., not unlike the results of the usual fits to the high-energy cross sections without a ρ' . The ρ' intercept, -0.48 , is reasonably consistent with Högaasen's determination, -0.6 , from the $\bar{p}p$ and pn data.⁴ The ρ' slope, $1.44/(\text{GeV})^2$, gives a mass, at $\alpha=1$, of 1.01 GeV, suggesting the $\delta(965)$, about which little is known except that it is an isovector. Relative to the ρ , the ρ' contribution near the forward direction is given roughly by

$$\text{Im}A_{\rho'} / \text{Im}A_{\rho} \approx e^{3.7t/E} \Delta\alpha$$

$$\text{Im}B_{\rho'} / \text{Im}B_{\rho} \approx 3e^{0.3t/E} \Delta\alpha$$

$$\text{Re}A_{\rho} \approx \text{Im}A_{\rho'}; \text{Re}A_{\rho'} \approx -\text{Im}A_{\rho}$$

where E is in GeV, and $\Delta\alpha \equiv \alpha_{\rho'} - \alpha_{\rho}$.

For the high-energy cross-section data we see that the ρ' plays only a small role. In Fig. 3 we have the usual dip in the π^-p charge-exchange cross section at $t \approx -0.6$, independent of energy, due to $\alpha_{\rho} = 0$. Our "ghost killing and sense choosing" mechanism predicts an analogous dip in $\pi^-p \rightarrow \eta^0 n$ for $\alpha_R(t \approx -0.9) = 0$. Figure 5 shows that our data do not extend to sufficiently large momentum transfer to serve as evidence in this matter. The "nonsense choosing" mechanism of Gell-Mann would not require such a dip for even-signature trajectories.⁶ Note that our η production data measure only etas decaying to two photons. Therefore, to arrive at the SU_3 breaking parameter F_R^0 , one needs to know the branching ratio for $\eta \rightarrow 2\gamma$. Using¹¹ $B(\eta \rightarrow 2\gamma) = 0.303$ we find F_R^0 differing from the unbroken F^+ for the A_2 by only 1%. For ρ and ρ' we have ten and twenty percent SU_3 breaking in F_i^0 . In each case the symmetry breaking is slightly greater for $t \neq 0$ due to the nonvanishing F_i^1 . The zero of $C_{\rho}(t)$ occurs at $t = -0.34$, roughly the first inflection point in the π^-p charge-exchange cross sections.

In the K^+N system at 2.3 GeV/c (Figs. 6 and 7), we see the ρ' asserting itself. In the forward direction κA_{ρ} and κA_R are, roughly speaking, equal, and lie at $\pi/4$ and $3\pi/4$ in the complex plane. Thus their imaginary parts subtract and real parts add for K^+ charge exchange (vice versa for K^-). The ρ' does not significantly alter the traditional

result that the forward $K^+(K^-)$ amplitude is predominantly real (imaginary). We find $\text{Re}/\text{Im } f_0(K^+) = +7.7$. But when we include the $\Delta\sigma_T(K^+N)$ datum at 2.3 GeV/c and apply the optical theorem (Eq. 14), we see the ρ' at work. At high energies the cancellation of $\text{Im}(A_{\kappa R} A_{\kappa \rho}^*)_{t=0}$ results in $\Delta\sigma_T(K^+N) \approx 0$, as seen in Fig. 7. This trend would continue down to 2.3 but for the ρ' term, $\sim -1/E$, which emerges at lower energies and causes $\Delta\sigma_T$ to increase negatively. This is required by the datum at 2.3. The error bar here is given by the amplitude of the small wiggles (presumably not a t-channel Regge effect) in the $\Delta\sigma_T(K^+N)$ data in this region.¹² The quoted experimental error is considerably smaller.

To see the contribution of the flip amplitudes to $d\sigma/dt$ (Eq. 13), we write for small t.

$$\frac{\left\{ \frac{-t}{4M^2} \left[\frac{s+p^2}{1 - (t/4M^2)} \right]^{-s} \right\}^{1/2} B_i}{(1 - t/4M^2)^{1/2} A_i} \approx \sqrt{-t} \frac{D_i^0}{C_i^0} \frac{\alpha_i^0}{2} \exp[D_i^1 - C_i^1 + (\alpha'/\alpha^0)_i] t.$$

For each trajectory, the t-channel helicity flip contribution becomes comparable to the nonflip at $-t \lesssim 0.1 \text{ (GeV)}^2$. This rapid rise from zero in the forward direction is responsible for the initial rise in each of the different cross sections. In the case of K^+n charge exchange at 2.3 (GeV/c), ρ' adds significantly and positively to the flip amplitude, producing a considerable initial rise and increasing the cross section in the peak region by about a factor of two. Without the ρ' , the theoretical curve in this region had stubbornly remained a factor of two below the data.

Figure 6 also shows the improvement in the fit at larger angles (smaller $-\cos\theta_t$) due to the use of the exact Legendre functions rather

than the high-energy asymptotic expressions at this energy. Near the forward direction, where the deuteron effect is significant, the free neutron cross section is also shown. Without knowing R from the fitted Regge parameters, one could not say to what extent the forward dip in the data reflects the two-body cross section rather than the forward vanishing required by the deuteron effect.

For the π^-p charge-exchange polarization (Fig. 8), the ρ' is of course, in our model, the sine qua non. For $-t < 0.34$, where our $C_\rho(t)$ goes through zero, we have

$$A_{\pi^-}^- \times B_{\pi^-}^- = \sin \frac{\pi}{2} (\alpha_\rho - \alpha_{\rho'}) (|A_{\pi^- \rho}| |B_{\pi^- \rho'}| - |A_{\pi^- \rho'}| |B_{\pi^- \rho}|).$$

The data require, and we find, $A_{\rho'} B_{\rho} > A_{\rho} B_{\rho'}$, giving positive polarization. The approximate orthogonality of the ρ and ρ' amplitudes for small t gives roughly maximal polarization, given the magnitudes and signs of the amplitudes. If the polarization is in fact due to a ρ' trajectory lying about one unit of angular momentum below the ρ , we have perforce a $\sim 1/E$ falloff in the polarization. The data, being quite uncertain, are consistent with this, but are also consistent with an energy-independent polarization. Better polarization data would constitute a severe test of our model. The unsightly high shoulder in the theoretical curves at the smallest t data emerged upon imposition of the sum rule constraint. Previously the two data around $-t = 0.03$ had been better fit. The good fit for $\Delta\sigma_T(\pi p)$, shown in Fig. 4, guarantees the convergence of the sum rule integral (Eq. 30) above 5 GeV/c. The ρ' term turns out to make 1/15th the contribution of the ρ term to this integral.

Finally, in Fig. 10 are shown some higher energy K^+n charge-exchange cross-section predictions from our fitted parameters. As the ρ' contribution wanes with increasing energy, the forward turnover diminishes, but, as in the case of K^-p charge exchange (Fig. 9), some turnover persists at high energies. For comparison with experiment, the deuteron effect, which causes the observed cross section to vanish at $t=0$, is also shown. Except for the 9.5 GeV/c prediction, where the difference has become negligible, exact Legendre functions were used here in place of the asymptotic forms.

POSTSCRIPT

Our Regge fit gives for the real part of the forward K^+n charge-exchange scattering amplitude at 2.3 GeV/c

$$k \operatorname{Re} f(t=0) = -0.69 .$$

Note that, unlike the imaginary part, this is not a directly observable quantity in deuterium. In a Phys. Rev. Letter just published [18, 801 (1967)], A. A. Carter, using forward K^+N dispersion relations, calculates for this quantity the value -0.70 ± 0.01 .

ACKNOWLEDGMENTS

We wish to thank Dr. Charles Chiu for his helpfulness in several phases of this work, and Prof. Geoffrey Chew for calling the Igi sum rule to our attention. Rarita thanks Prof. Burton Moyer for the hospitality of the Physics Department, Berkeley. Schwarzschild wishes especially to thank Prof. Gerson Goldhaber, who together with his wife Sulamith, of

blessed memory, gave guidance and forbearance in generous measure during his graduate studies, of which this paper represents the completion.

Table I. Data.

| <u>Reaction</u> | <u>Lab momenta (GeV/c) and references</u> |
|--|---|
| | <u>Differential cross section</u> |
| $K^+n \rightarrow K^0p$ | 2.3 ^a |
| $K^-p \rightarrow \bar{K}^0n$ | (5, 7, 9.5) ^b |
| $\pi^-p \rightarrow \pi^0n$ | (5.9, 9.8, 13.3, 18.2) ^c |
| $\pi^-p \rightarrow \eta^0n \times B(\eta \rightarrow 2\gamma)$ | (5.9, 9.8, 13.3, 18.2) ^{d, e, f} |
| | <u>Total cross-section differences</u> |
| $\sigma_T(K^+p) - \sigma_T(K^+n)$ | 2.3, ^g (6, 8, 10, 12, 14, 16, 18, 20) ^h |
| $\sigma_T(K^-p) - \sigma_T(K^-n)$ | (6, 10, 12, 14, 16, 18) ^h |
| $\sigma_T(\pi^-p) - \sigma_T(\pi^+p)$ | (5, 5.2, 5.4, 5.6, 5.8, 6.0, 6.2, 6.4) ⁱ (8, 10, 12, 14, 16, 18, 20) ^h |
| | <u>Polarization</u> |
| $\pi^-p \rightarrow \pi^0n$ | (5.9, 11.2) ^j |
| <p>a. I. Butterworth, J. Brown, G. Goldhaber, S. Goldhaber, A. Hirata, J. Kadyk, B. Schwarzschild, and G. Trilling, Phys. Rev. Letters <u>15</u>, 734 (1965).</p> <p>b. P. Astbury, G. Brautti, G. Finocchiaro, A. Michelini, K. Terwilliger, D. Websdale, C. West, P. Zanella, W. Beuch, W. Fischer, B. Gobbi, M. Peppin, E. Polgar, C. Verkerk, and M. Pouchon, CERN Preprint 66/1057/5, 1966.</p> <p>c. Saclay-Orsay Collaboration: A. Stirling, P. Sonderegger, J. Kirz, P. Falk-Vairant, O. Guisan, C. Bruneton, P. Borgeaud, M. Yvert, J. Guillaud, C. Caverzasio, and B. Amblard, Phys. Rev. Letters <u>14</u>, 763 (1965); Phys. Letters <u>20</u>, 75 (1966).</p> <p>d. Saclay-Orsay Collaboration (see Ref. c), Phys. Letters <u>18</u>, 200 (1965).</p> <p>e. Branching ratio ($\eta \rightarrow 2\gamma/\eta \rightarrow$ all neutrals) = 0.416, as given in G. DiGiugno, R. Querzoli, G. Triose, F. Vanoli, M. Giorgi, and P. Schiavon, Phys. Rev. Letters <u>16</u>, 767 (1966).</p> <p>f. Branching ratio ($\eta \rightarrow$ all neutrals) = 0.729, as given in A. Rosenfeld et al., Rev. Mod. Phys. <u>39</u>, 1 (1967).</p> <p>g. T. Kycia, Bull. Am. Phys. Soc., April 1967 (JB2).</p> <p>h. W. Galbraith, E. Jenkins, T. Kycia, B. Leontic, R. H. Phillips, A. Read, and R. Rubinstein, Phys. Rev. <u>138</u>, B913 (1965).</p> <p>i. A. Citron, W. Galbraith, T. Kycia, B. Leontic, R. H. Phillips, A. Rousset, and P. Sharp, Phys. Rev. <u>144</u>, 1101 (1966).</p> <p>j. P. Bonamy, P. Borgeaud, C. Bruneton, P. Falk-Vairant, O. Guisan, P. Sonderegger, C. Caverzasio, J. Guillaud, J. Schneider, M. Yvert, I. Mannelli, F. Sergiampietri, and L. Vincelli, Phys. Letters <u>23</u>, 501 (1966).</p> | |

Table II. Fitted parameters.

| | Trajectories | | KN residue parameters | | | | $\pi N/\text{KN } \text{SU}_3$ breaking | |
|------------|--------------|------------|-----------------------|---------------|------------------------------|------------------------------|---|------------------------------|
| | α^0 | α^1 | C^0 (mb×GeV) | D^0 (mb) | C^1 (GeV) ⁻² | D^1 (GeV) ⁻² | F^0/F^\pm | F^1 (GeV) ⁻² |
| $\rho(-)$ | 0.58 | 0.92 | 1.30 | 22.7 | 2.92 | 0.26 | 1.10 | -0.006 |
| $\rho'(-)$ | -0.48 | 1.44 | 5.02 | -264 | 4.4 | 2.95 | 0.80 | +0.20 |
| $A_2(+)$ | 0.37 | 0.41 | 5.50 | -116 | 0.42 | 0.66 | 1.01 | -0.07 |

Footnotes and References

*Work done under auspices of the U. S. Atomic Energy Commission.

†Visiting scientist.

‡This paper was accepted by the University of California, Berkeley, as part of the Ph. D. dissertation.

1. R. J. N. Phillips and W. Rarita, Phys. Rev. 139, B1336 (1965);
Phys. Rev. Letters 15, 807 (1965).
2. F. Arbab and C. B. Chiu, Phys. Rev. 147, 1045 (1966).
3. Table I, Ref. a.
4. H. Högaasen and W. Fischer, Phys. Letters 22, 516 (1966).
5. K. Igi and S. Matsuda, Phys. Rev. Letters 18, 625 (1967).
6. F. Arbab, N. Bali, and J. Dash, Ambiguities in the Phenomenological Determination of Regge Pole Parameters, Phys. Rev. (to be published).
7. A. Read, J. Orear, and H. Bethe, Nuovo Cimento 29, 1051 (1963).
8. S. Mandelstam, Ann. Phys. (N. Y.) 19, 254 (1959).
9. B. M. Schwarzschild, Lawrence Radiation Laboratory, University of California, Berkeley, Thesis (UCRL-17572, Appendix B).
10. V. Singh, Phys. Rev. 129, 1889 (1963). We have interchanged Singh's A and A'.
11. Table I, Refs. e and f.
12. Table I, Ref. g.

FIGURE CAPTIONS

Fig. 1. Reactions (1) through (4), for which we investigate peripheral scattering data, constitute a complete, isotopically independent, set of the reactions (pseudoscalar octet + nucleon \rightarrow pseudoscalar octet + nucleon) which require isovector exchange in the t (peripheral) channel.

Fig. 2. If one thinks of $\alpha(t)$ as the spin of the exchanged "particle" for a given t , then the t -channel helicity flip vertex above violates angular momentum conservation when $\alpha = 0$. We assume that each such "nonsense" vertex contributes a factor $\sqrt{\alpha}$ to the t -channel helicity flip amplitude.

Fig. 3. π^-p charge-exchange differential cross sections, incoming lab momenta from 5.9 to 18.2 GeV/c. Data from Saclay-Orsay Collaboration, Stirling et al. (Table I, ref. c). Solid curves are our Regge fits, Table II parameters.

Fig. 4. πp total cross-section difference from 5 to 20 GeV/c. Data from Galbraith et al., Brookhaven (Table I, Refs. h and i). Solid curve is our Regge fit, Table II parameters.

Fig. 5. $\pi^-p \rightarrow \eta^0 n$ differential cross sections reduced by branching ratio $B(\eta^0 \rightarrow 2\gamma)$, which is the only mode observed in these data, Stirling et al., Saclay-Orsay Collaboration (Table I, Ref. d). Solid curves are our Regge fits, Table II parameters. To make contact with the KN normalization parameters via SU_3 , we use for the branching ratio 0.303 (Table I, Refs. e and f).

Fig. 6. K^+n charge exchange at 2.3 GeV/c, the highest energy data available to date (Butterworth, Goldhaber, Schwarzschild, et al.,

Fig. 6 (Cont.)


Berkeley, Table I, Ref. a). In the forward region we show the Regge fit, with deuteron correction, to the deuterium data, as well as the free neutron cross section deduced from the fitted parameters, Table II. At larger angles one sees the correction due to the use of exact Legendre functions, as well as the fit using the high-energy asymptotic forms.

Fig. 7. K^+N and K^-N total cross-section differences, 6 to 20 GeV/c. $\sigma_T(K^-p) - \sigma_T(K^-n) > 0$, and $\sigma_T(K^+p) - \sigma_T(K^+n) < 0$. Data of Galbraith et al., Brookhaven (Table I, Ref. h). We include also the K^+ datum at 2.3, for which we indicate here (and use in fitting) an uncertainty given by the amplitude of the small wiggles of $\Delta\sigma_T(K^+)$ in this region, the quoted experimental error being considerably smaller (Kycia, Brookhaven, Table I, Ref. g). The solid curves are our Regge fits, Table II parameters.

Fig. 8. π^-p charge-exchange polarization, 5.9 GeV/c (solid error bars and fitted curve) and 11.2 GeV/c (dashed bars and curve). Data of Bonamy et al., Saclay-Orsay-Pisa Collaboration (Table I, Ref. j).

Fig. 9. K^-p charge-exchange differential cross sections, incoming lab momenta 5 to 9.5 GeV/c. Data of Astbury et al., CERN-ETH Zürich (Table I, Ref. b). Solid curves are our Regge fits, Table II parameters.


Fig. 10. Predictions of higher energy K^+n charge exchange, from the fitted parameters, Table II. Up to 5 GeV/c we use here the exact Legendre functions. The dashed curves near $t=0$ show the predicted $K^+d \rightarrow K^0p(p)$, i. e., the deuteron correction.



XBL673-2445


Fig. 1

A "nonsense" Vertex
(small arrows indicate helicities)




XBL673-2438

Fig. 2



XBL673-2440

Fig. 3



XBL673-2437

Fig. 4






Fig. 5




XBL673-2439

Fig. 6



XBL673-2436

Fig. 7



XBL673-2435

Fig. 8





Fig. 9



XBL675-3203

Fig. 10

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

