Lawrence Berkeley National Laboratory

Recent Work

Title

A Method for Quantifying the Acute Health Impacts of Residential Non-Biological Exposure Via Inhalation

Permalink

<https://escholarship.org/uc/item/4bz7p67s>

Authors

Logue, Jennifer
Sherman, Max
Singer, Bret

Publication Date

2014-08-01

A Method for Quantifying the Acute Health Impacts of Residential Non-Biological Exposure Via Inhalation

J.M. Logue, M.H. Sherman, B.C. Singer

Environmental Energy Technologies Division

August 2014

Funding was provided by the U.S. Dept. of Energy Building Technologies Program, Office of Energy Efficiency and Renewable Energy under DOE Contract No. DE-AC02-05CH11231; by the U.S. Dept. of Housing and Urban Development Office of Healthy Homes and Lead Hazard Control through Interagency Agreement I-PHI-01070, and by the California Energy Commission through Contract 500-08-061.

Disclaimer

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof, or The Regents of the University of California.

Ernest Orlando Lawrence Berkeley National Laboratory is an equal opportunity employer.

SUMMARY

INTRODUCTION The inability to monetize the health costs of acute exposures in homes and the benefits of various control options is a barrier to justifying policies and approaches that can reduce exposure and improve health.

METHODS We synthesized relationships between short-term outdoor concentration changes and health outcomes to estimate the health impacts of short-term in-home exposures. Damage and cost impacts of specific health outcomes were taken from the literature. We assessed the impact of vented and non-vented residential natural gas cooking burners on Southern California occupants for two pollutants (NO₂ and CO).

RESULTS Despite only looking at the impact of two pollutants on acute exposure-related health outcomes, the annual health benefits of using venting range hoods exceed the costs.

CONCLUSIONS The established methodology will provide a useful tool for quantifying the costs of acute exposures in homes and will allow for identification of cost effective methods for reducing exposures.

IMPLICATIONS Acute exposures in homes can have substantial impacts on the health of occupants especially for those in an already compromised state of health. Range hoods have the potential to significantly reduce acute exposures associated with cooking as well as reduce chronic exposure that result from aggregate cooking emissions. This report quantified the costs and benefits of mitigating two pollutants (NO₂ and CO) associated with gas cooking and indicated that range hoods are cost effective based on those two pollutants alone. It is expected that particle emissions could have a much larger effect than NO₂ and CO. Particle emissions are associated with food cooking and not just fuel usage. The benefits of removing cooking-related particles increase the value for gas cooking appliances and provide value for cooking with electric appliances as well.

CITATION Logue, J.M; Sherman, M.H; Singer, B.C.; (2014) *A method for quantifying the acute health impacts of residential non-biological exposures via inhalation*. Indoor Air Conference, Hong Kong, China, 7-12 July 2014

INTRODUCTION

Previous work at LBNL has developed a methodology for assessing and quantifying the impacts of long term (chronic) health impacts (Logue et al. 2012). The previous methodology was used to determine the chronic health impacts of indoor exposure on occupants of the housing stock. This paper is intended to present a complementary methodology for assessing costs of acute health impacts. Homes contain a variety of short-term episodic sources that impact indoor concentrations on timescales of a few hours to a few days. These short-term spikes in concentration may have a minimal effect on long-term concentrations but a significant effect on concentrations on shorter timescales. The objective of this work was to develop a method to identify illnesses associated with specific exposures, quantify the associated illness rates, quantify the health burden in Disability Adjusted Life Years (DALYs), and estimate the resulting costs of health care.

Initial assessments of the relationship between extremely high exposures during air pollution episodes and illness were easily quantifiable using graphical methods. As outdoor concentrations decreased due to increased regulation, statistical tools became necessary to characterize these relationships (Carracedo-Martinez et al. 2010). Initial large-scale assessments of population impacts of changes in outdoor pollutant concentrations on health predominantly used Poisson regression-based assessments, either parametric or nonparametric, that linked changes in outdoor concentrations to changes in population health outcomes. In recent years, case-crossover (CCO) assessments have increased in popularity because they eliminate problems with confounding and selecting correct degrees of freedom. Using these approaches, a large literature exists linking changes in outdoor concentrations with changes in population health outcomes. These studies, in conjunction with estimates of changes in home concentration resulting from changes in outdoor concentrations, allowed us to develop relationships between indoor concentrations and health outcomes.

In order to apply the methodology for acute exposures due to specific sources, we needed to determine time-resolved exposure concentrations for occupants of the US housing stock due to episodic and intermittent sources. LBNL recently developed a Population Impact Assessment Framework that allows for assessment of minute-by-minute concentrations due to specific sources in a representative subset of homes (Logue et al. 2013). This modeling framework allows us to determine the impact of sources on concentrations over a variety of timescales relevant to acute exposures. This document presents initial results for quantifying the health damage and health costs associated with exposure to nitrogen dioxide (NO₂) and carbon monoxide (CO) emitted by natural gas cooking burners in California homes.

METHODOLOGIES

For this analysis, we compiled and reviewed studies of the acute health impacts for NO₂ and CO. We included studies examining the US population as well as those in other countries with similar lifestyles. These studies estimate the change in a given morbidity/mortality outcome within a studied population, as a function of short-term (acute) changes in outdoor concentrations. Distributions of relative risks (RRs) and odds ratios (ORs), the two “measures of effect”, were used in these studies to describe the change in the probability of an outcome per change in exposure. To identify relevant CCO-based studies, we used the ISI Web of Knowledge database with combinations of the keywords: "case-crossover", "air pollution", "acute", "health", "United States", "nitrogen dioxide", and "carbon monoxide". We also considered the relationships between changes in exposure and changes in health outcomes included in the Cost Benefit Analysis of the Clean Air Act (EPA 1999). The search resulted in 11 studies that covered 15 acute health outcomes. Table 1 presents the top 5 relationships that

drove the damage and health costs for NO₂ and CO in this analysis. The remaining studies did not have an appreciable effect on study outcomes.

In order to translate RR and OR data into changes in incidence of outcomes as a function of changes in exposure, we needed to select concentration response (C-R) functions. For the relationships used in the EPA Cost Benefit Analysis of the Clean Air Act (1999), a specific C-R function is specified. For the literature-derived CCO relationships, we used the log linear concentration response function:

$$\Delta Incidence = -[y_o * (\exp(-\beta \Delta C_{exposure}) - 1)] * population \quad (1)$$

where *population* is the population exposed, $\Delta C_{exposure}$ is the absolute change in exposure concentration, and y_o is the baseline incidence of illness in the population, Table 2. Health outcome-specific values for y_o are listed in Table 2. The β value is traditionally determined from the relationship of the change in relative risk per change in outdoor concentration in each study, $\Delta C_{outdoor}$, Equation 2.

$$\beta_{indoor\ exposure} = \frac{\ln(RR)}{\Delta C_{outdoors}} \quad (2)$$

In order to use the selected concentration-response relationship, Equation 1, for studies that reported *OR* values, we used the relationship developed by Zhang and Yu (1998) to translate *OR* to *RR*.

Table 1: Case-crossover and Poisson distribution acute health impact studies included in analysis. Min age and max age are the age range for which the relationship is valid.

<i>References</i>	<i>Pollutants</i>	<i>Outcome</i>	<i>Location(s)</i>	<i>Time frame</i>	<i>Min Age</i>	<i>Max Age</i>
(Burnett et al. 1999)	NO ₂ , CO	HA: Ischemic Heart Disease	Toronto, CA, USA	1980-1994	NA	NA
(Burnett et al. 1997)	NO ₂	HA: All Respiratory	Toronto, CA, USA	Summers 1992-1994	NA	NA
(Dennekamp et al. 2010)	CO	OHCA: All Cardiac	Melbourne, Australia	2003-2006	35	NA
(Mustafic et al. 2012)	CO, NO ₂	MI	Various (Meta Analysis)	1988-2011	NA	NA
(Wellenius et al. 2005)	CO, NO ₂	Stroke: Ischemic	9 US Cities nationwide	1986-1999	65	NA

All respiratory includes asthma, COPD, lung cancer, pneumonia, and tuberculosis. HA=hospital admission, OCHA=out of hospital cardiac arrest, COPD=chronic obstructive pulmonary disease, MI=myocardial infarction, NA= not applicable.

Table 2: Disease prevalence rates in United States and likelihood of mortality.

<i>Outcome</i>	<i>Y_o</i>	<i>Reference</i>	<i>Mortality rate</i>	<i>Reference</i>
HA: All respiratory	1.85 E-5	(ALA 2008; Wier et al. 2011; ALA 2012; CDC 2012; AAAA&I 2013)	19.7%	(Dalal et al. 2011; ALA 2012; CDC 2012; Joynt et al. 2013; AAAA&I 2013)
HA: Ischemic Heart Disease	3.69 E-6	(Murphy et al. 2013)	89.7%	(Murphy et al. 2013)
MI	3.76 E-5	(ALA 2008)	18.5%	(Joynt et al. 2013)
OHCA: All Cardiac	2.57 E-6	(Roger et al. 2011)	90.4%	(CDC 2011)
Stroke: Ischemic	5.41 E-6	(Roger et al. 2012)	25.0%	(Hankey 2003)

The available studies that look at the relationship between exposure and health compare changes in outdoor concentrations to changes in health outcomes. US residents spend more than 90% of their time indoors and more than 70% in their homes (Klepeis et al. 2001). Since the indoor environment provides protection from outdoor-generated pollutants, both the concentrations indoors and the concentrations people are actually exposed to will be lower than the concentrations measured outdoors. Using β values derived from the outdoor concentrations measured in these studies to assess relationships of these pollutants with health outcome may underestimate the impact of indoor exposures on health. For this reason, we attempted to determine the change in indoor concentrations resulting from the reported change in outdoor concentration for each study when possible so that we can use Equation 3 to calculate a β value for use with indoor exposures. This approach does not consider the increased variability in indoor levels for any specific outdoor level and may introduce bias into the effects estimates.

$$\beta_{indoor\ exposure} = \frac{\ln(RR)}{\Delta C_{indoor}} \quad (3)$$

Chen et al. (2012) looked at the influence of city-by-city variations on the impact of outdoor concentration on indoor home concentrations and estimates of mortality resulting from short-term changes in outdoor concentration. Chen et al. found, for PM10, strong associations between changes in city-specific derived mortality coefficients and changes in indoor exposure. Since we are interested in the impacts of indoor exposures on health, we used the same method as Chen et al. to determine the equivalent change in indoor concentrations associated with changes in outdoor concentrations for the city in which each of the studies was conducted. We could only do this for studies conducted in the US and Canada. For Canadian studies we used the data from the closest US city. Studies conducted outside the US where not corrected and likely underestimate health impacts.

Several authors have determined the DALYs lost per incidence of specific diseases using the preeminent work of Murray and Lopez (Murray and Lopez 1996a; Murray and Lopez 1996b). Multiplying disease incidence by a “DALY factor” yields total DALYs lost.

$$DALYs = (\partial DALYs / \partial Disease\ Incidence) * Disease\ Incidence \quad (4)$$

Equation 4 uses a partial derivative in recognition that DALY losses are incrementally impacted by causes other than disease. The total burden of disease in a community can be calculated as the aggregate, across all diseases, of DALY factors multiplied by disease incidence rates. One major question in public health costing is whether acute exposure related deaths result in a substantial loss of life or merely accelerate the death of those in an already frail state of health who would have died soon anyway. Advancing death by only a few days is often referred to as “mortality displacement” or “harvesting”. Several studies have addressed the harvesting question using distributed lag models (DLM) (Zanobetti et al. 2002; Dominici et al. 2003; Roberts and Switzer 2004; Murray and Lipfert 2012). One issue with using existing DLM approaches is that the only option for exit from the frail population is death, and there are only two life states: healthy and frail. As Zanobetti and Schwartz (2008) point out, for certain health outcomes there is also the option of getting healthier. They specifically reference myocardial infarction and pneumonia, which, if survived, usually lead to a recovery and re-entry to the healthy pool, or at least a healthier state than the frail pool.

In order to determine the DALYs associated with each of these outcomes, we divided the outcomes into three groups: 1) those who were hospitalized but survived (independent of the

reason for hospitalization), 2) those who were hospitalized due to chronic health issues and died, and 3) those who were hospitalized due to a one-time event not related to a chronic health issue and died. For those in the first group, we assumed the event did not have a long term impact on occupant health and assigned a DALY loss of 4 per 10,000 hospitalizations based on Lyvovsky (2000). This assumes that any loss of life or life quality is attributed to an underlying chronic health issue and not a particular acute event. For those in the second group, we assume that that death results in a short-term mortality shift but did not result in a substantial loss of life. For those with chronic health issues that die, we will assume that they are part of the frail population with life expectancies of 11.8 days to 102 days, representing the range of values found in the literature for populations identified as frail or having the shortened expected life span (Manton et al. 1993; Murray and Nelson 2000; Murray and Lipfert 2012). When assigning DALYs to each incidence, we assumed a log normal distribution with a 95th percentile range of 2.4 to 31 days or 0.0065 to 0.084 DALYs with a central estimate of 0.023 DALYs/incidence. For the third group, those that died due to one-time events (stroke, cardiac arrest or myocardial infarction), DALYs assigned are based on the estimated life expectancy if the patient had survived the event. The DALYs assignments per health outcome are specified in Table 3, and the survivability of hospital admissions for each health outcome is included in Table 2. Medical costs for each health outcome are summarized in Table 3.

Table 3: Treatment cost and DALYs lost for each health outcome.

<i>Outcome</i>	<i>Cost of Treatment</i>	<i>Reference</i>	<i>DALYs Lost (If patient dies)</i>	<i>Reference</i>
HA: All Respiratory	\$2,521	MEPS	88.8% [0.023], 11.2% [1.2]	(Manton et al. 1993; Murray and Nelson 2000; Dick et al. 2012; Murray and Lipfert 2012)
HA: Ischemic Heart Disease	\$3,159	MEPS	0.023	(Manton et al. 1993; Murray and Nelson 2000; Murray and Lipfert 2012)
MI	\$15,631	(Azoulay et al. 1999)	5	(Goldberg et al. 1998)
OHCA: All Cardiac	\$3,159	MEPS	5	(Cobbe et al. 1996)
Stroke: Ischemic	\$9,526	(Russo and Andrews 2008)	2	(May et al. 1994)

As an initial application of this acute assessment methodology, we assessed the impact of select natural gas cooking pollutants on acute exposure related health outcomes for the population living in Southern California (SoCal) homes. The two pollutants analyzed were nitrogen dioxide (NO₂) and carbon monoxide (CO). A previous study that used the PIAM framework assessed the minute-by-minute concentration impacts of natural gas cooking on 6,969 representative SoCal homes that reported natural gas cooking in the home at least once a week and on the 19,465 occupants of those homes (Logue et al. 2013). We used the modeled occupant exposure concentrations to assess the impacts of cooking with no range hood, and the benefit of cooking using venting range hoods during all cooking events on the exposure of the occupants over the time frames of exposure specified by the aggregated studies. Logue et al. (2013) assumed an average range hoods capture efficiency (CE) of 55%.

RESULTS AND DISCUSSION

We used the studies included in Tables 1 through 3 to determine the population impacts on health outcomes and the impact of those outcomes on DALYs lost and medical costs. We calculated the incidence of disease and DALYs per incidence of disease for each C-R function

for each occupant using a Monte Carlo approach. We repeated this process for 5000 iterations. The resulting distribution of total DALYs lost and medical cost for the population derived from the 5000 iterations per modeled occupant was used to report the median and 95th percentile confidence interval of the total disease burden.

Across the SoCal population, per 100,00 people weekly in winter, exposure to gas cooking related pollutants (NO₂ and CO) if no range hood is used results in an estimated 0.38 (95%CI: 0.38-0.39) DALYs lost due to health impacts and \$18,850 (95%CI: \$18,800-18,950) in medical expenditures. Over the course of a year, this translate to a loss of 19.2 DALYs (95%CI: 18.7-19.6) and a medical cost of \$943,200 (95%CI: \$938,200-948,200) per 100,000 SoCal occupants of homes that cook with natural gas. The DALYs lost due to acute exposures calculated here for NO₂ and CO exceed, on an annual basis, the estimates for DALYs lost due to NO₂ and CO chronic exposures in the average US home (Logue et al. 2012).

If homes use range hoods during all cooking events with average capture efficiency (55%), the health impact of cooking is reduced to an annual DALYs loss of 8.9 (95%CI: 8.7 - 9.2) and a medical cost of \$458,200 (95%CI: \$455,200 - 461,200) per 100,000 SoCal occupants of homes that cook with natural gas. Using the assumed cost of \$100,000 per DALY lost (Logue et al. 2012), the total cost saved by using standard range hoods during all cooking events due to reductions in NO₂ and CO exposure is estimated to save \$1,430,000-1,580,000 per 100,000 SoCal occupants annually. The calculated benefits of range hood use only apply to homes with gas cooking burners that also have venting range hoods.

Logue and Singer (2013) have previously reported on the annual energy cost of using venting range hoods. The results for IECC climate zone 3C, the climate zone that contains the majority of the SoCal region, indicated that using currently available venting range hoods during all cooking events would result in an energy cost to consumers of \$385,000 to \$729,000 per 100,000 SoCal occupants. The study also found that the cost could be reduced by 23% though increases in system efficiency.

CONCLUSIONS

This analysis only estimated the cost of acute health impacts from two pollutants (NO₂ and CO). Several other pollutants, particularly PM_{2.5}, will have a significant impact on health and health related costs on acute and chronic time scales due to cooking emissions (both from natural gas cooking burners and food preparation). Despite only looking at the impact of two pollutants on acute exposure related health outcomes, the annual health benefits of using venting range hoods exceed the annual energy cost. Further analysis is needed to determine the impact of other pollutants and to determine installation costs for venting range hoods in homes based on home characteristics.

ACKNOWLEDGEMENT

Funding was provided by the U.S. Dept. of Energy Building Technologies Program, Office of Energy Efficiency and Renewable Energy under DOE Contract No. DE-AC02-05CH11231; by the U.S. Dept. of Housing and Urban Development Office of Healthy Homes and Lead Hazard Control through Interagency Agreement I-PHI-01070, and by the California Energy Commission through Contract 500-08-061.

REFERENCES

AAAA&I.2013 "Asthma Statistics." Retrieved July 22, 2013, 2013, from <http://www.aaaai.org/about-the-aaaai/newsroom/asthma-statistics.aspx>.

- ALA.2008. Lung Disease Data: 2008. Washington DC, American Lung Association http://www.lung.org/assets/documents/publications/lung-disease-data/LDD_2008.pdf
- ALA.2012. Trends in Lung Cancer Morbidity and Mortality, American Lung Association, Epidemiology and Statistics Unit <http://www.lung.org/finding-cures/our-research/trend-reports/lc-trend-report.pdf>
- Azoulay A, Eisenberg M and Pilote L.1999. "Cost of treatment of acute myocardial infarction in the United States and Canada." *Journal of the American College of Cardiology* **33**(2 SUPPL. A): 307A.
- Burnett RT, Cakmak S, Brook JR and Krewski D.1997. "The role of particulate size and chemistry in the association between summertime ambient air pollution and hospitalization for cardiorespiratory diseases." *Environmental Health Perspectives* **105**(6): 614-620.
- Burnett RT, Smith-Doiron M, Stieb D, Cakmak S and Brook JR.1999. "Effects of particulate and gaseous air pollution on cardiorespiratory hospitalizations." *Archives of Environmental Health* **54**(2): 130-139.
- Carracedo-Martinez E, Taracido M, Tobias A, Saez M and Figueiras A.2010. "Case-Crossover Analysis of Air Pollution Health Effects: A Systematic Review of Methodology and Application." *Environmental Health Perspectives* **118**(8): 1173-1182.
- CDC.2011. Out-of-Hospital Cardiac Arrest Surveillance — Cardiac Arrest Registry to Enhance Survival (CARES), United States, October 1, 2005–December 31, 2010. Atlanta, GA, Center for Disease Control and Prevention, US Department of Health and Human Services. **60**:8 <http://www.cdc.gov/mmwr/pdf/ss/ss6008.pdf>
- CDC.2012. Reported Tuberculosis in the United States. Atlanta, GA, U.S. Department of Health and Human Services, CDC, <http://www.cdc.gov/tb/statistics/reports/2011/pdf/report2011.pdf>
- Chen C, Zhao B and Weschler CJ.2012. "Indoor Exposure to "Outdoor PM10" Assessing Its Influence on the Relationship Between PM10 and Short-term Mortality in US Cities." *Epidemiology* **23**(6): 870-878.
- Cobbe SM, Dalziel K, Ford I and Marsden AK.1996. "Survival of 1476 patients initially resuscitated from out of hospital cardiac arrest." *BMJ* **312**: 1633-1637.
- Dalal AA, Shah M, D'Souza AO and Rane P.2011. "Costs of COPD exacerbations in the emergency department and inpatient setting." *Respiratory Medicine* **105**(3): 454-460.
- Dennekamp M et al.2010. "Outdoor Air Pollution as a Trigger for Out-of-hospital Cardiac Arrests." *Epidemiology* **21**(4): 494-500.
- Dick A et al.2012. "Long-term survival and healthcare utilization outcomes attributable to sepsis and pneumonia." *BMC Health Services Research* **12**(1): 1-10.
- Dominici F, McDermott A, Zeger SL and Samet JM.2003. "Airborne particulate matter and mortality: Timescale effects in four US cities." *American Journal of Epidemiology* **157**(12): 1055-1065.
- EPA.1999. The Benefits and Costs of the Clean Air Act 1990 to 2010. Washington, D.C., Environmental Protection Agency:EPA-410-R-99-001 <http://www.epa.gov/air/sect812/1990-2010/fullrept.pdf>
- Goldberg RJ et al.1998. "Age-related trends in short- and long-term survival after acute myocardial infarction: a 20-year population-based perspective (1975–1995)." *The American Journal of Cardiology* **82**(11): 1311-1317.
- Hankey GJ.2003. "Long-term outcome after ischaemic stroke/transient ischaemic attack." *Cerebrovascular Diseases* **16**: 14-19.
- Joynt KE, Orav E and Jha AK.2013. "Mortality rates for medicare beneficiaries admitted to critical access and non–critical access hospitals, 2002-2010." *JAMA* **309**(13): 1379-1387.

- Klepeis NE et al.2001. "The National Human Activity Pattern Survey (NHAPS): A Resource for Assessing Exposure to Environmental Pollutants." Journal of Exposure Analysis and Environmental Epidemiology **11**: 231-252.
- Logue JM, Klepeis NE, Lobscheid AB and Singer BC.2013. "Pollutant Exposures from Natural Gas Cooking Burners: A Simulation-Based Assessment for Southern California." Under Review by Environmental Health Perspectives.
- Logue JM, Price PN, Sherman MH and Singer BC.2012. "A Method to Estimate the Chronic Health Impact of Air Pollutants in US Residences." Environmental Health Perspectives **120**(2): 216-222.
- Logue JM and Singer BC.2013. "Energy Impacts of Effective Range Hood Use for All U.S. Residential Cooking." under review by HVAC&R.
- Lvovsky K, Huges G, Maddison D, Ostro B and Pearce D.2000. Environmental costs of fossil fuels: a rapid assessment method with application to six cities. Washington, D.C., The World Bank Environment Department: The World Bank Environment Department Papers, No.78
- Manton KG, Stallard E and Liu K.1993. Chapter 7: Frailty and Forecasts of Active Life Expectancy in the United States. Springer Series in Statistics. Dietz, K, Gail, M, Krickeberg, K et al. New York, NY.
- May DS, Casper ML, Croft JB and Giles WH.1994. "Trends in survival after stroke among Medicare beneficiaries." Stroke **28**(8): 1617-1622.
- Murphy SL, Xu JQ and Kochanek KD.2013. Deaths: Final Data for 2010. National Vital Statistics Reports, Division of Vital Statistics. **61**Division of Vital Statistics
- Murray CJ and Lipfert FW.2012. "A new time-series methodology for estimating relationships between elderly frailty, remaining life expectancy, and ambient air quality." Inhalation Toxicology **24**(2): 89-98.
- Murray CJ and Nelson CR.2000. "State-space modeling of the relationship between air quality and mortality." Journal of the Air & Waste Management Association **50**(7): 1075-1080.
- Murray CL and Lopez AD.1996a. The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020. Cambridge, MA, Harvard University.
- Murray CL and Lopez AD.1996b. Global health statistics: A compendium of incidence, prevalence, and mortality estimates for over 200 conditions. Global burden of disease and injury series. Cambridge, MA, Harvard University. **II**: 906.
- Mustafic H et al.2012. "Main Air Pollutants and Myocardial Infarction A Systematic Review and Meta-analysis." Jama-Journal of the American Medical Association **307**(7): 713-721.
- Roberts S and Switzer P.2004. "Mortality displacement and distributed lag models." Inhalation Toxicology **16**(14): 879-888.
- Roger VL et al.2012. "Heart Disease and Stroke Statistics—2012 Update: A Report From the American Heart Association." Circulation **125**(1): e2-220.
- Roger VL et al.2011. "Heart Disease and Stroke Statistics-2011 Update A Report From the American Heart Association." Circulation **123**(4): E18-E209.
- Russo CA and Andrews RM.2008. Hospital Stays for Stroke and Other Cerebrovascular Diseases, 2005. Healthcare Cost and Utilization Project. Rockville, MD, Agency for Healthcare Research and Quality Statistical Brief #51 <http://www.hcup-us.ahrq.gov/reports/statbriefs/sb51.pdf>
- Wellenius GA, Schwartz J and Mittleman MA.2005. "Air pollution and hospital admissions for ischemic and hemorrhagic stroke among medicare beneficiaries." Stroke **36**(12): 2549-2553.

- Wier LM, Elixhauser A, Pfuntner A and Au DH.2011. Overview of Hospitalizations among Patients with COPD, 2008. Rockville, MD, Agency for Healthcare Research and Quality:Statistical Brief #106 <http://www.hcup-us.ahrq.gov/reports/statbriefs/sb106.pdf>
- Zanobetti A and Schwartz J.2008. "Mortality displacement in the association of ozone with mortality - An analysis of 48 cities in the United States." American Journal of Respiratory and Critical Care Medicine **177**(2): 184-189.
- Zanobetti A et al.2002. "The temporal pattern of mortality responses to air pollution: A multicity assessment of mortality displacement." Epidemiology **13**(1): 87-93.
- Zhang J and Yu KF.1998. "What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes." Jama-Journal of the American Medical Association **280**(19): 1690-1691.