

UCLA

UCLA Previously Published Works

Title

The Relationship Between Family Socioeconomic Status and Adolescent Sleep and Diurnal Cortisol.

Permalink

<https://escholarship.org/uc/item/4c6429fg>

Journal

Psychosomatic Medicine, 84(7)

Authors

Seeman, Teresa
Fuligni, Andrew
Rocha, Sarah
et al.

Publication Date

2022-09-01

DOI

10.1097/PSY.0000000000001104

Peer reviewed



HHS Public Access

Author manuscript

Psychosom Med. Author manuscript; available in PMC 2023 September 01.

Published in final edited form as:

Psychosom Med. 2022 September 01; 84(7): 848–855. doi:10.1097/PSY.0000000000001104.

The relationship between family socioeconomic status and adolescent sleep and diurnal cortisol

Sarah Rocha, M.A.,

Department of Psychology, University of California, Los Angeles

David M. Almeida, Ph.D.,

Department of Human Development and Family Studies, Pennsylvania State University, University Park, PA

Jessica J. Chiang, Ph.D.,

Department of Psychology, Georgetown University, Address: 306N White-Gravenor Hall, 37th and O streets NW, Washington, D.C., 20057

Steve W. Cole, Ph.D.,

Cousins Center for Psychoneuroimmunology, Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, Address: 11-934 Factor Building, Los Angeles CA 90095-1678

Michael R. Irwin, M.D.,

Department of Psychology, Cousins Center for Psychoneuroimmunology, Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, Address: 300 UCLA Medical Plaza, #3109, Los Angeles, CA 90095

Teresa Seeman, Ph.D.,

Division of Geriatrics, David Geffen School of Medicine, University of California, Los Angeles, Address: 10945 Le Conte Ave, Suite 2339, Los Angeles, CA 90095-1687

Andrew J. Fuligni, Ph.D.

Department of Psychology, Cousins Center for Psychoneuroimmunology, Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, Address: 760 Westwood Plaza, Box 62, Los Angeles, CA 90095

Abstract

Objective: To investigate the associations between indices of family socioeconomic status and sleep during adolescence and to examine whether measures of hypothalamic-pituitary-adrenal (HPA) axis functioning mediate the observed associations.

Address for Correspondence: Sarah Rocha, M.A., Department of Psychology, University of California, Los Angeles, Address: 2311 Franz Hall, Los Angeles, CA 90095, Phone: 480-7103-719; rocha@ucla.edu; Fax: 310-794-6297.

Conflicts of Interest and Source Funding: The study was supported by funding from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (R01-HD062547); the University of California, Los Angeles California Center for Population Research, which was supported by the National Institute of Child Health and Human Development (R24-HD041022); and the University of California, Los Angeles Older Americans Independence Center, which was supported by the National Institute of Aging (P30-AG017265 and P30-AG028748). The content does not necessarily represent the official views of the National Institutes of Health and its separate institutes. All authors declare no conflict of interest.

Methods: A total of 350 ethnically diverse adolescents (57% female; mean age_{wave1}=16.4, SD=0.7 years) completed a three-wave longitudinal study in which sleep and cortisol data were collected at two-year time intervals. Sleep duration, latency, and variability was assessed via actigraphy over a period of eight days per study wave. Salivary cortisol was collected across three days per study wave to assess cortisol diurnal slope, area under the curve and the cortisol awakening response. Adolescents' caregivers reported their education levels, family income, and economic hardship.

Results: A greater family income-to-needs ratio was associated with longer adolescent sleep duration ($b = 2.90, p = .023$), whereas greater parental education was associated with shorter sleep duration ($b = -3.70, p = .030$), less sleep latency ($b = -0.74, p = .016$), and less variability across days ($b = -2.06, p = .010$). Diurnal cortisol slope statistically mediated the association of parental education with sleep duration ($b = -0.48, 95\% \text{ CI } [-1.099, -0.042]$), but not the association of income-to-needs ratio with sleep duration.

Conclusion: Findings suggest parental education and family resources may have unique impacts upon sleep and HPA axis functioning during the period of adolescence. Future research is needed to examine family and behavioral factors that may underlie SES associations with adolescent sleep and HPA axis functioning.

Keywords

sleep; adolescence; socioeconomic status; diurnal cortisol; actigraphy

Introduction

Scholars have pointed to sleep as a potential source of disparities in education and health that can emerge during adolescence and extend into adulthood (1,2). Sleep plays a critical role in adolescent development, consistently exhibiting associations with learning (3), mood (4), and mental and physical health (5,6). Epidemiologic research suggests that socioeconomic status (SES) may be related to sleep, with higher-SES individuals showing more positive sleep outcomes (7,8). However, the relationship between SES and sleep during adolescence is still somewhat uncertain. While some studies have found disparities in sleep duration between high- and low-SES adolescents (8–10), null associations are also common (11–13).

Inconsistency in the literature on SES-sleep associations may be due to the diverse ways SES and sleep can be operationalized. Sleep is often measured in terms of duration; however, other components of sleep may also have implications for health. Consistency in bedtime routines has been often cited as a necessary component for good sleep hygiene, and increasing evidence suggests that adolescents with greater variability in nightly sleep duration may be at risk for negative health outcomes (14,15). Sleep latency, or the time it takes for adolescents to fall asleep, can also be an indicator of sleep issues. A short sleep latency may imply feelings of sleepiness, while a long sleep latency can suggest difficulties falling asleep (16). Assessing multiple components of sleep behavior is needed to form a more complete picture of the relationship between SES and sleep, as single indicators of sleep do not capture the multiple dimensions of sleep quality.

Furthermore, SES is a multifaceted construct (17) and use of single or composite measures of SES may not fully capture the unique associations different SES measures may have with sleep outcomes. For example, common measures of SES, such as family income and parental education, have the potential to impact youths' sleep in varying ways. Low-income families may be less able to provide a home environment conducive to sleep; previous literature suggests that aspects of the home environment such as noise, light and family chaos contribute to income associations with youths' sleep quality (13,18). Parental education, on the other hand, maybe be associated with parenting practices that influence youths' sleep. Previous literature suggests that mothers with lower educational attainment are less likely to have consistent bedtime routines for their children (19). Alternatively, highly educated parents may have greater academic expectations for their adolescents (20) and greater time spent in engaging in academic activities such as homework can come at the cost of youths' sleep (21). Finally, measures of economic strain, such as perceived hardship and difficulty making ends meet, have been related to sleepiness during adolescence indirectly via family chaos (18). Examining multiple measures of SES simultaneously and estimating their associations with different components of sleep behavior (e.g., duration, latency, variability) can help bring clarity to the literature's inconclusive associations between SES and youths' sleep.

Furthermore, there is still much to be understood about the mechanisms linking SES indices with adolescent sleep, particularly potential physiologic systems. One physiologic system that has shown strong associations with both SES and sleep is the hypothalamic-pituitary-adrenal (HPA) axis. The HPA axis reacts to stressful events, releasing the hormone cortisol (22), and significant evidence suggests that HPA axis activity also is associated with chronic stressors like poverty (23). Cortisol is released by the HPA axis in a typical fashion wherein circulating levels are high in the morning, peak shortly after wake, and then decline gradually across the day until bedtime (24). Consistent literature has linked SES with the pattern of the decline in circulating cortisol across the day, wherein high-SES is associated with a steeper decline (25,26). Deviations from typical HPA axis functioning, such as a less steep or "blunted" pattern of diurnal secretion, may indicate dysregulation, and this pattern of cortisol secretion is associated with numerous adverse mental and physical health outcomes (27). SES has also been linked with the total cortisol output over the day, measured as the area under the curve (AUC); however, the direction of the association has been mixed (25,28). Similarly mixed associations have been found in regard to the initial surge in cortisol secretion following wake, known as the cortisol awakening response (CAR) (26). Adding to this complexity, both large and small values of CAR may be indicative of dysregulation (29). Thus, SES appears to be relevant to typical HPA axis functioning, though associations may most consistently be found in relation to the slope of the decline in cortisol across the day.

Given the circadian rhythm of cortisol secretion and its relationship with sleep-wake patterns (24), scholars have investigated whether alterations of typical HPA axis functioning are related to sleep disruption. Evidence suggests there is likely a reciprocal relationship between cortisol secretion and sleep patterns (30,31). For example, a study of adolescents found that longer sleep duration was predictive of a steeper decline in cortisol levels the next day, and a steeper decline predicted greater hours of sleep the following day (32). If

altered HPA-axis functioning disrupts sleep, this could position it as a potential pathway by which socioeconomic status influences sleep patterns. Considering evidence for a reciprocal relation between diurnal cortisol and sleep duration, altered HPA axis functioning could induce a cascading effect in which initial dysregulation is exacerbated by ongoing sleep difficulties.

The goals of this study were the following: 1) to examine the association between SES and sleep using multiple indicators of SES and 2) to determine whether HPA axis functioning acts as a statistical mediator of the association between SES and sleep. We focused on parental education, family income-to-needs ratio (INR), and parental economic hardship (EH) as our indicators of SES. We predicted that adolescents from families with greater INR, higher parental education, and less parental EH would have longer sleep duration and less daily variability in their sleep. Given that both short and long sleep latency can indicate sleep difficulties, we predicted that SES would be related to latency in either direction. Next, we predicted that HPA axis functioning would mediate the association between SES and sleep variables, such that higher-SES would be related to better sleep outcomes indirectly through a steeper cortisol diurnal slope. We also explored the mediating role of the CAR and AUC, but due their less consistent associations with SES, we predicted that mediation would occur through either smaller or higher values of AUC and CAR.

Methods

Participants:

Participants were 350 adolescents recruited from four public high schools in the Los Angeles area who took part in a three-wave longitudinal study with their primary caregivers (88% mothers, 8% fathers, 4% others) in which data were collected on daily experiences at 2-year time intervals. Adolescents were majority female (57%) and ethnically diverse (42% Latinx, 30% European American, 22% Asian American and 6% identifying with another ethnicity). Forty-one percent of adolescents had at least one parent born outside the US and 2% did not provide information on caregiver country of origin. In the first wave, which took place between October 2011 and June 2012, a total of 316 adolescents (mean age = 16.40, $SD=0.74$) agreed to participate in the study (see Table 1 for descriptive statistics). In the following wave of data collection, conducted in October 2013 to June 2014, 214 of the original participants returned, and 34 new, grade-matched participants were added to the sample to account for attrition, for a total of 248 participants (mean age=18.31, $SD=0.77$). In the final wave of data collection (October 2015-August 2016), 180 participants who had participated in one or both of the previous two waves (mean age = 20.29, $SD=0.74$) returned. Given that a subset of adolescents was recruited at the second wave, study completion was determined by the percent of possible waves in which an adolescent collected some sleep or cortisol data. Study completion varied by adolescent ethnicity, $F(3, 346) = 5.408$ $p=.001$, with Asian adolescents completing the fewest waves of data collection. Additionally, female adolescents were more likely to complete all possible waves of data collection $r(347) = .11$, $p=.05$, as were those with higher family incomes, $r(306)=.14$, $p=.01$, higher parental education, $r(346) = .14$, $p = .01$, and those whose parents were born in the US, $r(340) = .11$, $p = .03$.

Procedures:

Adolescents and their caretakers were recruited via flyers and presentations at school. Those interested in participating in the study were contacted by phone and written consent was obtained in-person during the first study visit at participants' homes. Adolescents and their caregivers reported demographic information and completed surveys during the initial visit. At each wave of data collection, adolescents were instructed to wear an actigraph watch (Micro Motionlogger Sleep Watch, Ambulatory Monitoring, Inc.; Ardsley, NY) for eight nights to record their sleep. Additionally, for the first three nights of each study wave, adolescents were asked to collect saliva samples to measure diurnal cortisol at designated timepoints throughout the day. Adolescents were compensated after each wave of data collection, receiving \$50 after the first wave, \$75 after the second, and \$125 after the third. To further incentivize retention, adolescents were given 2 movie theater passes for completing the daily protocols correctly. All study procedures were approved by the University of California, Los Angeles Institutional Review Board.

Measures:**SES Variables:**

Parent Education: Caregivers reported the highest level of education they achieved (1 = some elementary school; 2 = completed elementary school; 3 = some junior-high school; 4 = completed junior-high school; 5 = some high school; 6 = graduated from high school; 7 = trade or vocational school; 8 = some college; 9 = graduated from college; 10 = some medical, law, or graduate school; 11 = graduated from medical, law, or graduate school) and, if applicable, the highest level of education of the adolescents' other caregiver. Responses for both caregivers were averaged for one composite score of parental education. Education reported at study entry (i.e., Wave 1 or Wave 2) was used in all analyses.

Income-to-Needs Ratio: At each study wave, adolescents' primary caregivers reported their total household income over the past year, and how many people lived in their household. Given the greater fluctuation in income as compared to parental education, we used income at each wave to predict the sleep and cortisol parameters at the same wave. Each family's income-to-needs ratio (INR) was computed by dividing total reported household income over the U.S. Department of Health and Human Services' federal poverty threshold by family size. For waves one through three, the 2012, 2014 and 2016 guidelines were used, respectively. An INR = 1 represents a family living at the poverty level. Descriptive statistics for family INR across study waves are reported in Table 1.

Parental Economic Hardship: A modified version of Conger et al.'s (33) well-established questionnaire for measuring family economic pressure was completed by the primary caregiver at each wave. The questionnaire was broken down into 3 subsections concerning their ability to: 1) pay bills 2) save and 3) afford material items over the past 3 months. First, caregivers were asked to rate their difficulty paying bills on a 4-point Likert scale with endpoints: "no difficulty at all" to "a great deal of difficulty." Second, caregivers were asked to estimate their savings on a 4-point scale from "with more than enough money left over" to "very short on money." Third, caregivers were provided with a list of 7 affirmative

statements on their ability to purchase essential material needs (e.g., food, clothing, medical care) and rated their agreement on a 4-point scale from “not at all true” to “very true”. Responses to the 3-subsections are averaged to create a composite score of perceived economic hardship (18,33). Mean parental economic hardship (EH) scores are reported in Table 1.

Sleep Variables: Sleep variables were based on adolescents’ sleep actigraphy data. Adolescents were instructed to wear the actigraph watch on their non-dominant hand and to keep it on for the duration of the time they spent in bed at night. An event marker was pressed to indicate when adolescents turned off the lights to go to bed, when they woke up, and when/if they got out of bed during the night. Data were scored using software package Action 4 (Ambulatory Monitoring, Inc.; Ardsley, NY). Sleep statistics for each night were calculated using one-minute epochs and the Sadeh actigraph scoring algorithm, which has been validated and used in studies with children and adolescents (15,34). Time in bed began when the adolescent pressed the event marker indicating lights out and ended when pressed again the next morning. Sleep onset began after at least 3 consecutive minutes of sleep were scored, and sleep offset was determined after the last five or more consecutive minutes of sleep. A total of 343 participants completed sleep actigraphy measurement during at least one study wave, 206 completed actigraphy for at least two waves, and 101 completed actigraphy for all three waves. During data collection periods, adolescents on average wore the watch for 6.19 nights. Descriptive statistics on sleep parameters are reported in Table 1.

Sleep Duration (minutes): Nightly sleep duration was computed as the total amount of time spent in bed that was scored as sleep for each 8 days of sleep assessment. Sleep duration was averaged across the 8 days to determine mean weekly sleep duration.

Sleep Latency (minutes): Time to fall asleep (latency) was computed as the minutes between the start of time in bed (as determined by event marker press) and sleep onset (as determined by actigraphy scoring), representing the time it took for adolescents to fall asleep.

Daily Sleep Variability (minutes): Variability was computed to represent daily-level variations in sleep duration. To do so, we computed the absolute value of the difference between each individual day’s sleep duration and the average sleep duration for the week.

Salivary Cortisol: At each study wave, adolescents collected five saliva samples using Salivettes (Sarstedt, Nümbrecht, Germany) at designated time points (wake, 15-min post-wake, 30-min post-wake, dinner, and bedtime) for three consecutive days. Text messages were sent prior to each collection time to encourage compliance. Adolescents were asked to refrain from drinking, brushing teeth, or using tobacco products 30 min prior to saliva collection. Participants used electronic time stampers (Dymo Corporation, Stamford, CT) and stamping booklets to record sampling time and completion. Samples were stored in adolescents’ household refrigerators until they were picked up by research staff one to three weeks later. Research staff stored the collected samples at -80°C prior to assay. Saliva samples were shipped to the Laboratory of Biological Psychology at the Technical University of Dresden, Germany where they were assayed for cortisol

concentrations (nmol/L) using high-sensitivity chemiluminescence-immunoassays (IBL International, Hamburg, Germany). In the first wave of data collection, 308 adolescents collected saliva during the three-day period. In the following two waves, 221 and 158 adolescents, respectively, collected saliva during the three-day study period. Adolescents collected an average of 4.63 of the five daily saliva samples at each wave.

We computed three indices of HPA axis functioning: diurnal cortisol slope, the CAR, and AUC. Daily-level slope, CAR, and AUC values were averaged across the three days to characterize the typical diurnal patterns of adolescents and minimize the influence of daily-level fluctuations in cortisol levels. Sample values greater than 60nmol/L (2.1% of the total samples provided) were removed from the sample to reduce the impact of outliers on diurnal profiles.

Diurnal Slope: Diurnal slope was computed by subtracting the bedtime cortisol concentration from the waking cortisol concentration and dividing the value by the hours passed between the wake and bed samples (35,36). Average diurnal slope was computed during at least one study wave for N=334 adolescents, during at least two waves for N=216 adolescents, and N=110 for all three waves.

Cortisol Awakening Response: The cortisol awakening response, CAR, was computed by subtracting the salivary cortisol concentration from the sample taken 30-minutes after wake from the cortisol concentration of the wake sample. The CAR can be affected by the timing of saliva collection (37) and so daily-level CAR values were flagged where the wake and +30 minute sample were taken <15 minutes apart (N= 25) or >60 minutes apart (N= 24). As a sensitivity check, all analyses with the CAR were repeated wherein flagged wake and +30 min samples were removed from the dataset. Average CAR was computed during at least one study wave for N=343 adolescents, during at least two waves for N=221 adolescents, and N=119 for all three waves.

Area Under the Curve: To measure total cortisol output, we computed area under the curve (AUC) with respect to ground using the trapezoid equation established by Preussner and colleagues (38). Daily-level AUC was computed for each day across the three-day period where participants collected all five saliva sample. Average AUC was computed during at least one study wave for N=328 adolescents, during at least two waves for N=208 adolescents, and N=94 for all three waves.

Analytic Strategy

Using STATA 16.1, multilevel models were run to estimate the associations between SES variables and sleep. Study waves (level 1) were nested within participants (level 2). These models allowed for the inclusion of adolescents with missing data, such that adolescents were included in the analytic dataset if they participated in at least one wave of data collection. Individual observations within adolescents were dropped if missing information on age, gender, ethnicity, immigration status or sleep actigraphy data.

To assess the unique contribution of each SES characteristic on sleep outcomes, each multilevel model included all three SES variables: parental education, family INR, and

parental EH). Parental education (a level 2 or person-varying SES variable, given the lack of change in parental education over time) was grand-mean centered. Family INR and parental EH (level 1 or wave-varying SES variables) were also grand-mean centered, pooling the model estimates across within- and between-person associations. Separate multilevel models were run for each sleep outcome variable: duration, latency, and variability. Model covariates included age, gender (ref. = female), ethnicity (ref.=Latinx), parents' immigration status (ref = U.S. born), and average wake time (minutes). Age was centered around the youngest age in the sample (14.5). Grand-mean centered sleep duration was added as an additional covariate for the model examining sleep variability as the dependent variable, given the associations between less sleep duration and greater variability (39). The final analytical sample for the multilevel models included N=585 observations from 311 adolescents.

Next, we repeated analyses with the addition of three SES by age interaction terms to the model (one for each SES predictor variable) to investigate whether the relationship between SES and sleep varied as a function of age.

Finally, for models which demonstrated a significant association between SES and sleep, we estimated mediation by indices of HPA axis functioning. Separate mediation models were run to examine mediation by diurnal slope, the CAR, and AUC. The framework outlined by Krull and colleagues (2016) was used to run multilevel mediation models. Multilevel estimation offers an advantage over single-level procedures for assessing mediation when there are nested data (40). Confidence intervals for the indirect effects were obtained using percentile-based bootstrapping with 10,000 replications. Model covariates remained the same as above. The final analytical sample for the mediation models included N=562 observations from 311 adolescents.

Results:

Descriptive Statistics:

Table 1 presents descriptive statistics for age and socioeconomic information by study wave, as well as descriptive statistics for sleep and diurnal cortisol variables. Across study waves adolescents' average sleep duration decreased and their daily sleep variability became greater (Table 1). Bivariate correlation coefficients between key study variables at study entry are presented in Table 2. Parental education was associated with lower sleep latency, $r(291) = -.12$, $p = .04$. None of the other indicators of SES were associated with sleep parameters in a bivariate manner.

SES-Sleep Associations:

As shown in Table 3, multi-level models indicated that a higher family income-to-needs ratio was significantly associated with longer adolescent sleep duration ($b = 2.90$, $p = .023$), but not latency or variability. In contrast, greater parental education was significantly associated with shorter sleep duration ($b = -3.70$, $p = .030$), shorter sleep latency ($b = -0.74$, $p = .016$), and less daily variability in duration ($b = -2.06$, $p = .010$), the last when additionally controlling for average sleep duration. Lastly, parental EH was not associated

with adolescent sleep, and follow-up analyses suggested that associations between SES sleep parameters did not vary by age.

Mediation Analyses:

Parental Education: The observed associations of parent education with adolescent sleep duration and sleep variability were tested for mediation by diurnal cortisol indices. As shown in Figure 1, mediation analyses suggested that greater parental education was significantly associated with a flatter (less negative) diurnal cortisol slope ($b = 0.05$, $p = 0.008$). A flatter diurnal slope, in turn, was associated with shorter sleep duration ($b = -9.74$, $p = .009$), and the indirect effect was significant ($b = -0.48$, 95% CI [-1.099, -0.042]), suggesting parental education was related to shorter sleep duration indirectly via a flatter diurnal slope. However, diurnal cortisol slope was not a significant mediator for the association of parental education with sleep variability ($b = 0.24$, 95% CI [-0.006, 0.567]), or latency ($b = -0.05$, 95% CI [-0.140, 0.016]). Additionally, neither the CAR nor AUC were associated with parent education and neither emerged as significant statistical mediators.

Income-to-Needs Ratio: A greater family INR was significantly associated with a steeper (more negative) diurnal cortisol slope ($b = -0.03$, $p = 0.033$). However, the indirect effect was non-significant, indicating that diurnal cortisol slope did not mediate the association between family INR and sleep duration ($b = 0.29$, 95% CI [-0.005, 0.711]). Neither the CAR nor AUC were associated with family INR and neither statistically mediated the association between family INR and sleep duration.

Discussion

The present study examined the association between socioeconomic status and sleep and the mediating role of HPA axis functioning in a longitudinal study of adolescents. Study results suggest that the relationship between socioeconomic status and sleep during adolescence is complex. As hypothesized, higher family INR was associated with greater minutes of sleep duration. Greater parental education was associated with less sleep variability and shorter sleep latency. Contrary to our hypothesis, parental education was also associated with fewer minutes of sleep duration. Finally, evidence was found supporting HPA axis functioning, particularly a flatter (less negative) diurnal cortisol slope, as a statistical mediator of the high-parental education association with shorter sleep duration. In contrast, diurnal cortisol slope was not found to statistically mediate the association between family INR and sleep duration.

Although the association of greater parental education with shorter sleep is inconsistent with findings from a large previous epidemiologic study in adolescents (11), that study relied on a less precise measure of duration (i.e., self-reported frequency of receiving <7 hours of sleep per night). In contrast, actigraphy-based studies in adolescent populations have not found an association between parental education and sleep duration (11,41). Our finding that parental education is associated with less sleep variability is consistent with actigraphy based work (11), suggesting that youth with more highly educated parents may have more consistent bedtime routines. While less variability in nightly sleep duration is generally thought to

be a positive sleep outcome (42), it may also indicate that these youth are sleeping less on a consistent basis. Likewise, the observed association of parental education with shorter sleep latency may reflect greater feelings of sleepiness for those with more highly educated parents (16). Taken together, these findings suggest that higher parental education may be related to insufficient sleep during adolescence. Information on the types of activities youth engage in instead of sleeping can help clarify why higher parental education might be associated with less sleep. School or academic-related obligations are common contributors to insufficient sleep during adolescence (43). Enrollment in advanced coursework may vary by SES and has been associated with reduced sleep among high-school aged adolescents (44). Higher SES youth may also be more likely to participate in extracurricular activities (45) and these obligations can similarly come at the cost of sleep (46). Future research should consider collecting information on academic and extracurricular obligations to assess whether they may explain parental education associations with sleep.

Notably, we also found that HPA axis functioning, as indicated by a flatter diurnal slope, mediated the association between higher parental education and shorter sleep duration. A potential interpretation of this finding could be that chronic stressors associated with higher parental education are able to indirectly influence sleep through dysregulation of the stress-response system. While research is limited on the relationship between parental education and diurnal cortisol in adolescence, a recent study of first year college students found that students whose parents had an academic degree had higher levels of hair cortisol concentrations than those whose parents did not have an academic degree (47). The authors of the study suggested that youth with educated parents may feel greater pressure to succeed academically, potentially pursuing academic studies that are too difficult for them. High academic demands in turn can contribute to youths' feelings of stress (48), which may be reflected in diurnal cortisol rhythms. However, this study does not have sufficient data to confirm whether parental education was associated with differences in academic demands or participation in advanced coursework. Further research is needed to examine the relationship between academic-related stressors and diurnal cortisol during adolescence and assess if these factors may explain part of the association between parental education and sleep behavior.

The relationship between income-based measures of SES and sleep duration during adolescence is somewhat inconsistent in previous research, with studies showing both a positive association, no association, or a positive association only for children but not adolescents (9,11,12,41). Relatively few studies have examined the relationship between income and actigraphy-based measures of sleep during adolescence using actigraphy (10,11,13,41) and our study contributes evidence to the literature suggesting that family financial resources may impact how long adolescents are sleeping. In terms of HPA activity, we did find a significant association between family INR and diurnal cortisol slope, such that higher family INR was related to a steeper decline in cortisol levels across the day. This finding is consistent with previous literature and may reflect theory that youth with fewer economic resources show a more blunted pattern of diurnal cortisol secretion as a result of exposure to chronic stressors. Diurnal cortisol, however, did not explain the association between family INR and sleep duration, suggesting the role of other factors. Household stressors, such as family chaos, can potentially explain part of the association between

income and sleep quality in adolescents (18). Feelings of stress or worries that occur before bedtime have also been shown to disrupt low-SES children's sleep (13). Alternatively, aspects of the home sleep environment, such as noise and light levels, could be stressors that contribute to low-income youths' sleep problems (13,49)

The present study had several strengths. To our knowledge, this is the first study to assess associations of multiple measures of SES simultaneously with sleep and cortisol during adolescence. This study provides a step forward in understanding physiologic systems that may underly SES differences in sleep by providing evidence that altered cortisol secretion may explain part of the association between parental education and sleep. Next, this study benefits from its use of sleep actigraphy, which estimates sleep from participants' movement. Much of the literature on SES-disparities in adolescent sleep has relied on subjective or self-reported sleep (8,12,18), which is often only loosely correlated with objectively-measured sleep behavior (50). Actigraphy-based sleep assessment eliminates the potential reporting bias of survey-based measures of sleep and provides a minimally invasive alternative to the gold standard of sleep assessment, polysomnography. Finally, the broad age-span of the study allowed for an examination of the association between SES and sleep over the transition from adolescence to young adulthood, an understudied developmental period.

The study also had several limitations. Study results may be affected by attrition bias as higher-SES adolescents (i.e., those with higher family incomes and greater parental education) were more likely to complete all study waves. Further completion differences by gender, ethnicity, and parent immigration status may have also affected results and might limit generalizability of findings. Additionally, all measures of SES used in this study were parent-report, which may not encompass adolescents' perceptions of their socioeconomic status. Previous research suggests that adolescent-perceived financial strain is associated with physical and mental health outcomes even after controlling for parent SES measures (51). The lack of information regarding youths' perceived economic hardship is a limitation of this study and may provide context for the null associations of parent economic hardship with adolescent sleep behavior. Future research assessing socioeconomic differences during adolescence should consider including questions regarding youths' perceptions of economic hardship, which may offer additional insight into the potentially complex relationship that SES has with sleep during this developmental period. Finally, and importantly, our study design cannot establish causal relationships between SES, HPA axis activity, and sleep. Previous research suggests that adolescents moving into poverty may be at risk for shorter sleep duration, while those maintaining their poverty status do not show changes in sleep (52). Additional longitudinal research following youth from childhood into adulthood is needed to capture SES trajectories and examine whether HPA axis dysregulation is an underlying mechanism linking changes in SES with sleep disruption.

Conclusion

This study finds evidence that differing measures of socioeconomic status have divergent relationships with sleep during late adolescence. Consistent with previously observed patterns, family income-to-needs ratio was associated with longer sleep duration. By

contrast, parental education was associated with shorter sleep duration, shorter sleep latency, and less sleep variability. Findings suggested parental education was linked with sleep duration via flatter diurnal cortisol slopes. As such, parental education may be reflective a complex system of behavioral or micro-cultural processes that induce distinctive patterns of stress in adolescence that manifest as disturbances in sleep and HPA axis functioning. These results highlight the importance of assessing multiple SES indicators when studying sleep behavior, with particular care for how indicators might operate distinctively in the lives of adolescents versus adults.

Abbreviations:

SES	Socioeconomic status
INR	Income-to-needs ratio
EH	Economic hardship
CAR	Cortisol awakening response
AUC	Area under the curve

References

1. Levy DJ, Heissel JA, Richeson JA, Adam EK. Psychological and biological responses to race-based social stress as pathways to disparities in educational outcomes. *American Psychologist*. 2016;71. [PubMed: 26766766]
2. Jackson CL, Redline S, Emmons KM. Sleep as a Potential Fundamental Contributor to Disparities in Cardiovascular Health. *Annual Review of Public Health*. 2015;
3. Dewald JF, Meijer AM, Oort FJ, Kerkhof GA, Bögels SM. The influence of sleep quality, sleep duration and sleepiness on school performance in children and adolescents: A meta-analytic review. *Sleep Medicine Reviews*. 2010.
4. Short MA, Louca M. Sleep deprivation leads to mood deficits in healthy adolescents. *Sleep Medicine*. 2015;
5. Roberts RE, Duong HT. The Prospective Association between Sleep Deprivation and Depression among Adolescents. *Sleep*. 2014;
6. Owens J, Au R, Carskadon M, Millman R, Wolfson A, Braverman PK, Adelman WP, Breuner CC, Levine DA, Marcell A v., Murray PJ, O'Brien RF. Insufficient sleep in adolescents and young adults: An update on causes and consequences. *Pediatrics*. 2014;
7. Grandner MA, Williams NJ, Knutson KL, Roberts D, Jean-Louis G. Sleep disparity, race/ethnicity, and socioeconomic position. *Sleep Medicine*. 2016.
8. Keyes KM, Maslowsky J, Hamilton A, Schulenberg J. The great sleep recession: Changes in sleep duration among US adolescents, 1991–2012. *Pediatrics*. 2015;
9. Hysing M, Petrie KJ, Bøe T, Lallukka T, Sivertsen B. The social gradient of sleep in adolescence: results from the youth@hordaland survey. *European journal of public health*. 2016;
10. Marco CA, Wolfson AR, Sparling M, Azuaje A. Family Socioeconomic Status and Sleep Patterns of Young Adolescents. *Behavioral Sleep Medicine*. 2011;
11. Schmeer KK, Tarrence J, Browning CR, Calder CA, Ford JL, Boettner B. Family contexts and sleep during adolescence. *SSM - Population Health*. 2019;
12. Jarrin DC, McGrath JJ, Quon EC. Objective and subjective socioeconomic gradients exist for sleep in children and adolescents. *Health Psychology*. 2014;
13. Bagley EJ, Kelly RJ, Buckhalt JA, El-Sheikh M. What keeps low-SES children from sleeping well: The role of presleep worries and sleep environment. *Sleep Medicine*. 2015;

14. He F, Bixler EO, Berg A, Imamura Kawasawa Y, Vgontzas AN, Fernandez-Mendoza J, Yanosky J, Liao D. Habitual sleep variability, not sleep duration, is associated with caloric intake in adolescents. *Sleep Medicine*. 2015;
15. Park H, Tsai KM, Dahl RE, Irwin MR, McCreath H, Seeman TE, Fuligni AJ. Sleep and inflammation during adolescence. *Psychosomatic Medicine*. 2016;
16. Banks S, Dinges DF. Behavioral and physiological consequences of sleep restriction. *Journal of Clinical Sleep Medicine*. 2007.
17. Duncan GJ, Magnuson KA. Off with hollingshead: Socioeconomic resources, parenting, and child development. In: *Socioeconomic Status, Parenting, and Child Development*. 2014.
18. Philbrook LE, Saini EK, Fuller-Rowell TE, Buckhalt JA, El-Sheikh M. Socioeconomic status and sleep in adolescence: The role of family chaos. *Journal of Family Psychology*. 2020;
19. Lee S, Hale L, Berger LM, Buxton OM. Maternal Perceived Work Schedule Flexibility Predicts Child Sleep Mediated by Bedtime Routines. *Journal of Child and Family Studies*. 2019;
20. Davis-Kean PE. The influence of parent education and family income on child achievement: The indirect role of parental expectations and the home environment. *Journal of Family Psychology*. 2005;19.
21. Zhang J, Li AM, Fok TF, Wing YK. Roles of Parental Sleep/Wake Patterns, Socioeconomic Status, and Daytime Activities in the Sleep/Wake Patterns of Children. *Journal of Pediatrics*. 2010;156.
22. Dickerson SS, Kemeny ME. Acute stressors and cortisol responses: A theoretical integration and synthesis of laboratory research. Vol. 130, *Psychological Bulletin*. 2004.
23. Lupien SJ, King S, Meaney MJ, McEwen BS. Can poverty get under your skin? Basal cortisol levels and cognitive function in children from low and high socioeconomic status. *Development and Psychopathology*. 2001;13. [PubMed: 11346048]
24. Buckley TM, Schatzberg AF. Review: On the interactions of the hypothalamic-pituitary-adrenal (HPA) axis and sleep: Normal HPA axis activity and circadian rhythm, exemplary sleep disorders. Vol. 90, *Journal of Clinical Endocrinology and Metabolism*. 2005.
25. Desantis AS, Kuzawa CW, Adam EK. Developmental origins of flatter cortisol rhythms: Socioeconomic status and adult cortisol activity. *American Journal of Human Biology*. 2015;
26. Dowd JB, Simanek AM, Aiello AE. Socio-economic status, cortisol and allostatic load: A review of the literature. *International Journal of Epidemiology*. 2009.
27. Adam EK, Quinn ME, Tavernier R, McQuillan MT, Dahlke KA, Gilbert KE. Diurnal cortisol slopes and mental and physical health outcomes: A systematic review and meta-analysis. *Psychoneuroendocrinology*. 2017.
28. Cohen S, Doyle WJ, Baum A. Socioeconomic status is associated with stress hormones. *Psychosomatic Medicine*. 2006;68.
29. Chida Y, Steptoe A. Cortisol awakening response and psychosocial factors: A systematic review and meta-analysis. *Biological Psychology*. 2009.
30. Gillin JC, Jacobs LS, Fram DH, Snyder F. Acute effect of a glucocorticoid on normal human sleep. *Nature*. 1972;
31. Kumari M, Badrick E, Ferrie J, Perski A, Marmot M, Chandola T. Self-reported sleep duration and sleep disturbance are independently associated with cortisol secretion in the Whitehall II study. *Journal of Clinical Endocrinology and Metabolism*. 2009;
32. Zeiders KH, Doane LD, Adam EK. Reciprocal relations between objectively measured sleep patterns and diurnal cortisol rhythms in late adolescence. *Journal of Adolescent Health*. 2011;
33. Conger RD, Conger KJ, Elder GH, Lorenz FO, Simons RL, Whitbeck LB. A Family Process Model of Economic Hardship and Adjustment of Early Adolescent Boys. *Child Development*. 1992;
34. Acebo C, Sadeh A, Seifer R, Tzischinsky O, Hafer A, Carskadon MA. Sleep/wake patterns derived from activity monitoring and maternal report for healthy 1- to 5-year-old children. *Sleep*. 2005;
35. Kuhlman KR, Chiang JJ, Bower JE, Irwin MR, Seeman TE, McCreath HE, Almeida DM, Dahl RE, Fuligni AJ. Sleep problems in adolescence are prospectively linked to later depressive symptoms via the cortisol awakening response. *Development and Psychopathology*. 2020;32.

36. Adam EK, Doane LD, Zinbarg RE, Mineka S, Craske MG, Griffith JW. Prospective prediction of major depressive disorder from cortisol awakening responses in adolescence. *Psychoneuroendocrinology*. 2010;35.
37. Stalder T, Kirschbaum C, Kudielka BM, Adam EK, Pruessner JC, Wüst S, Dockray S, Smyth N, Evans P, Hellhammer DH, Miller R, Wetherell MA, Lupien SJ, Clow A. Assessment of the cortisol awakening response: Expert consensus guidelines. Vol. 63, *Psychoneuroendocrinology*. 2016.
38. Pruessner JC, Kirschbaum C, Meinlschmid G, Hellhammer DH. Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology*. 2003;28.
39. Becker SP, Sidol CA, van Dyk TR, Epstein JN, Beebe DW. Intraindividual variability of sleep/wake patterns in relation to child and adolescent functioning: A systematic review. Vol. 34, *Sleep Medicine Reviews*. 2017.
40. Krull JL, Cheong J, Fritz MS, MacKinnon DP. Moderation and mediation in interindividual longitudinal analysis. In *developmental psychopathology*. Chichester, United Kingdom: John Wiley & Sons, Ltd; 2016;16–18. Available at: 10.1002/9781119125556.devpsy121. Accessed May 3, 2021.
41. Moore M, Kirchner HL, Drotar D, Johnson N, Rosen C, Redline S. Correlates of adolescent sleep time and variability in sleep time: The role of individual and health related characteristics. *Sleep Medicine*. 2011;12. [PubMed: 21145280]
42. Chaput JP, Dutil C, Featherstone R, Ross R, Giangregorio L, Saunders TJ, Janssen I, Poitras VJ, Kho ME, Ross-White A, Zankar S, Carrier J. Sleep timing, sleep consistency, and health in adults: a systematic review. *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*. 2020;45. [PubMed: 31121100]
43. Adam EK, Snell EK, Pendry P. Sleep timing and quantity in ecological and family context: A nationally representative time-diary study. *Journal of Family Psychology*. 2007;21.
44. Jin Q, Shi Q. A Comparison of the Number of Hours of Sleep in High School Students Who Took Advanced Placement and/or College Courses and Those Who Did Not. *The Journal of School Nursing*. 2008;24.
45. McNeal RB. High school extracurricular activities: Closed structures and stratifying patterns of participation. *Journal of Educational Research*. 1998;91.
46. Gaarde J, Hoyt LT, Ozer EJ, Maslowsky J, Deardorff J, Kyauk CK. So Much to Do Before I Sleep: Investigating Adolescent-Perceived Barriers and Facilitators to Sleep. *Youth and Society*. 2020;52.
47. Bertrams A, Minkley N. Preliminary Indications That First Semester Students From Academic Households Exhibit Higher Hair Cortisol Concentrations Than Their Peers From Nonacademic Households. *Frontiers in Psychiatry*. 2020;11. [PubMed: 32116834]
48. Suldo SM, Shaunessy E, Hardesty R. Relationships among stress, coping, and mental health in high-achieving high school students. *Psychology in the Schools*. 2008;45.
49. Doane LD, Breitenstein RS, Beekman C, Clifford S, Smith TJ, Lemery-Chalfant K. Early Life Socioeconomic Disparities in Children's Sleep: The Mediating Role of the Current Home Environment. *Journal of Youth and Adolescence*. 2019;
50. Aili K, Åström-Paulsson S, Stoetzer U, Svartengren M, Hillert L. Reliability of actigraphy and subjective sleep measurements in adults: The design of sleep assessments. *Journal of Clinical Sleep Medicine*. 2017;
51. Quon EC, McGrath JJ. Subjective socioeconomic status and adolescent health: a meta-analysis. Vol. 33, *Health psychology: official journal of the Division of Health Psychology, American Psychological Association*. 2014.
52. Sivertsen B, Bøe T, Skogen JC, Petrie KJ, Hysing M. Moving into poverty during childhood is associated with later sleep problems. *Sleep Medicine*. 2017;




Figure 1: Parental education and sleep duration: mediation by diurnal cortisol slope
 Mediation pathway of the association between parent education and sleep duration. Models control for wake time, age, gender, ethnicity, family INR, parent EH, parent immigration. Beta coefficients are provided with standard errors. Significant indirect, direct, and total pathways are denoted with an asterisk if the bootstrapped confidence interval of the beta coefficient did not include zero.

Table 1.

Descriptive statistics for sociodemographic, sleep, and diurnal cortisol variables

Variable	Wave 1 (n = 311)		Wave 2 (n = 223)		Wave 3 (n = 159)	
	Range	Mean (SD)	Range	Mean (SD)	Range	Mean (SD)
Age	14.50–20.50	16.39 (0.74)	14.50–22.17	18.31 (0.77)	16.50–22.09	20.29 (0.74)
Parental Education	1–11	7.18 (1.87)	1–11	7.42 (1.96)	1–11	7.44 (2.05)
Family INR	0–14.32	2.80 (2.13)	0–17.69	3.58 (2.74)	0–14.73	3.82 (2.83)
Parental EH	1–3.95	2.15 (0.76)	1–4	2.09 (0.79)	1–4	2.02 (0.73)
Duration (min)	177.4–603.0	447.7 (57.84)	221.4–575.4	446.5 (64.23)	161.0–592.4	430.77 (75.14)
Latency (min)	0–57.50	10.38 (9.85)	0–95.00	10.10 (11.31)	0–64.00	11.30 (11.83)
Variability (min)	0.75–175.33	57.13 (28.99)	0.75–166.46	60.28 (29.61)	0.75–164.05	62.03 (35.89)
Diurnal Slope	–2.58–0.22	–1.01 (0.44)	–3.34–1.01	–1.26 (0.63)	–4.75–0.67	–1.49 (0.96)
CAR	–22.16–37.25	4.73 (9.19)	–24.64–35.58	4.47 (10.81)	–29.51–50.77	4.18 (13.05)
AUC	32.98–499.64	173.07 (70.76)	28.70–531.05	218.64 (87.00)	17.02–710.02	225.76 (112.92)

SD = standard deviation. n = number of adolescents completing at least some measurement of sleep or cortisol during study wave

Table 2

Bivariate correlations between key study variables, as measured at study entry

Variable	1	2	3	4	5	6	7	8	9	10	11	12
1. Age												
2. Female	0											
3. US-Born Parent	-.04	-.19**										
4. Parent Education	-.13*	-.11*	.44**									
5. Family INR	.03	-.06	.39**	.49**								
6. Parental EH	.02	.08	-.20**	-.28**	-.43**							
7. CAR	-.17**	.19**	.01	.05	-.04	-.10						
8. Diurnal Slope	-.13*	-.05	-.06	.09	-.05	-.07	.44**					
9. AUC	.06	.21**	.11	.01	.05	-.09	.51**	-.18**				
10. Sleep Duration	-.16**	.16*	.05	.01	.08	0.04	.05	-.09	-.12			
11. Sleep Latency	-.10	-.06	-.11	-.12*	-.08	.05	-.14*	-.12*	-.06	-.06		
12. Sleep Variability	.04	.09	.05	.03	0	-.03	.02	.12*	-.08	-.11	.05	
13. Wake Time	-.02	.03	-.07	.03	.03	.01	0.07	.07	-.11	.38**	.15*	.31**

Results of bivariate correlations between study variables.

* p < .05

** p < .01

*** p < .001

Table 3.

Multi-level associations between SES and sleep

Variable	Duration	Latency	Daily Variability
	b (SE)	b (SE)	b (SE)
Intercept	495.01 (7.54) ***	11.80 (1.38) ***	71.04 (3.93) ***
Age	-12.21 (1.46) ***	-10 (0.27)	-1.14 (0.82)
Parent EH	4.53 (3.64)	0.57 (0.67)	0.44 (1.78)
Family INR	2.90 (1.28) *	0.12 (0.23)	0.12 (0.62)
Parent Education	-3.70 (1.75) *	-0.74 (0.30) *	-2.06 (0.80) **
Male	-24.07 (5.38) ***	0.60 (0.97)	-7.86 (2.56) **
Asian American	-18.55 (7.89) *	-1.41 (1.43)	3.88 (3.71)
European American	9.78 (7.05)	-2.92 (1.28) *	-1.76 (3.29)
Other Ethnicity	7.43 (11.80)	2.23 (2.13)	-5.18 (5.47)
Immigrant Parent	-2.21 (7.14)	-0.04 (1.28)	-9.82 (3.28) **
Wake Time	0.34 (0.03) ***	0.02 (0.01) **	0.13 (0.02) ***

Results of multilevel models predicting sleep outcomes from SES indices, controlling for age, gender, ethnicity, parent immigration, and wake time. Age was centered at 14.5. Parent EH, family INR, parental education, and wake time were grand-mean centered. Reference variables were female, Latinx, and adolescents of US-born parents.

* p < .05

**

p < .01

p < .001.