

UC Davis

UC Davis Previously Published Works

Title

Imidazolidine-2,4,5- and pirimidine-2,4,6-triones - New primary pharmacophore for soluble epoxide hydrolase inhibitors with enhanced water solubility

Permalink

<https://escholarship.org/uc/item/4c74q533>

Journal

Bioorganic & Medicinal Chemistry Letters, 30(3)

ISSN

0960-894X

Authors

Burmistrov, Vladimir
Morisseau, Christophe
D'yachenko, Vladimir
[et al.](#)

Publication Date

2020-02-01

DOI

10.1016/j.bmcl.2019.126908

Peer reviewed



Published in final edited form as:

Bioorg Med Chem Lett. 2020 February 01; 30(3): 126908. doi:10.1016/j.bmcl.2019.126908.

Imidazolidine-2,4,5- and pirimidine-2,4,6-triones – new primary pharmacophore for soluble epoxide hydrolase inhibitors with enhanced water solubility.

Vladimir Burmistrov^{a,b}, Christophe Morisseau^a, Vladimir D'yachenko^b, Dmitry Karlov^{c,d,e}, Gennady M. Butov^b, Bruce D. Hammock^{a,*}

^aDepartment of Entomology and Nematology, and Comprehensive Cancer Center, University of California, Davis, CA, 95616, USA

^bDepartment of Chemistry, Technology and Equipment of Chemical Industry, Volzhsky Polytechnic Institute (branch) Volgograd State Technical University, Volzhsky, 404121, Russia

^cSkolkovo Institute of Science and Technology, Skolkovo Innovation Center, Moscow, 143026, Russia

^dDepartment of Chemistry, Lomonosov Moscow State University, Moscow, 119991, Russia

^eInstitute of Physiologically Active Compounds, Russian Academy of Sciences, Chernogolovka, Moscow Region, 142432, Russia

Abstract

A series of inhibitors of the soluble epoxide hydrolase (sEH) containing imidazolidine-2,4,5-trione or pirimidine-2,4,6-trione has been synthesized. Inhibition potency of the described compounds ranges from 8.4 μ M to 0.4 nM. The tested compounds possess higher water solubility than their preceding ureas. Molecular docking indicates new bond between the triones and the active site of sEH that in part explain the observed potency of the new pharmacophores. While less potent than the corresponding ureas, the modifications of urea group reported herein yield compounds with higher water solubility, thus permitting easier formulation.

Abstract

Imidazolidine-2,4,5-triones were systematically studied as sEH inhibitors.

21 adamantyl imidazolidine-2,4,5- and pirimidine-2,4,6-triones were synthesized.

Molecular docking highlights possible new bonds with the enzyme.

*To whom correspondence should be addressed: Bruce D. Hammock, Department of Entomology, University of California at Davis, One Shields Avenue, Davis, CA 95616. Tel: 530 752 7519; Fax: 530-751-1537; bdhammock@ucdavis.edu.


Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Proposed that triones can act as a prodrugs releasing more potent ureas.

Graphical Abstract



Keywords

soluble epoxide hydrolase; epoxyeicosatrienoic acids; inhibitor; adamantane; urea; imidazolidine-2,4,5-trione; pirimidine-2,4,6-trione

The human soluble epoxide hydrolase (sEH) is involved in the metabolism of arachidonic acid epoxides and other natural epoxy-fatty acids,¹ which have numerous, largely beneficial, biological activities.² Through the addition of a water molecule, sEH converts epoxides into corresponding vicinal diols thus affecting inflammatory processes, pain and other disease states.² Thereby inhibition of sEH could be beneficial in treatment of many renal, cardiovascular and neuronal diseases.^{3,4} Although thousands of various sEH inhibitors (sEHI) were synthesized over the last two decades⁵⁻⁷, they have limited solubility, making them hard to formulate, as well as limited bioavailability, especially toward the CNS where sEH is emerging as a potential target for neurological diseases.⁸ Toward improving water solubility and metabolic stability, herein, we changed the most common sEHI primary pharmacophore, an urea group, with imidazolidine-2,4,5-trione or pirimidine-2,4,6-trione groups and investigate the effects of such substitution on the potency and properties of the resulting compounds.

Urea-type sEH inhibitors bearing either adamantyl or aromatic moiety as lipophilic group both possess poor water solubility,^{9,10} that could be explained by the intermolecular interactions between urea molecules.^{11,12} Unfortunately the same hydrogen bond acceptors (HBAs) and hydrogen bond donors (HBDs) from the urea group are essential for sEH inhibition.¹³ However, we proposed that certain changes in the urea-group can improve water solubility while sustaining potency on a decent (nanomolar) level. Thus, we converted urea group in known sEH inhibitors into imidazolidine-2,4,5-trione or pirimidine-2,4,6-trione, because additional HBAs should enhance water solubility and prevent intermolecular interactions between urea molecules.

To synthesize imidazolidine-2,4,5-triones, the reaction of ureas with oxalyl chloride. Reaction was carried out in anhydrous THF for 2h with reflux (Scheme 1).^{14,15}

In most cases imidazolidine-2,4,5-triones are significantly less active than the corresponding ureas, with loss of potency between 2.3 and 6,000 folds, except for compound **1c**, which is

as potent as the corresponding urea (Table 1). Imidazolidine-2,4,5-trione **1k** (8.4 μ M, 700-folds less active than the preceding urea) is the least active among compounds **1a-k**. Presence of 5 HBAs in the molecule probably prevents it from entering the active site of the sEH. However, for compound **1c** its activity equals activity of corresponding urea.

As expected, in most cases, the triones yielded more water soluble than the corresponding urea. Compound **1b** is 40-fold more soluble than its corresponding urea (*t*-TUCB). However, in some cases (**1c** and **1k**) triones were less soluble than its preceding ureas. In addition, melting points of triones were up to 101°C less than those for the corresponding ureas except for compounds **1a** and **1b**. Because in some conditions, triones can degrade back to the original ureas, the better physical properties of triones can be enhanced their formulation as pro-drugs of urea-based sEHI.

It was previously shown that diureas, which contain two urea groups linked with aliphatic spacer are very potent sEHI.⁹ Molecular docking suggests that the high potency of these compounds is due to the binding of the second urea group with Ser374 of the sEH. We used diureas with various linkers between adamantane and urea fragments to synthesize corresponding di-imidazolidine-2,4,5-triones **2a-g** and tri-imidazolidine-2,4,5-trione **2h** (Scheme 2).

Data (Table 2) shows that the di-imidazolidine-2,4,5-triones **2a-h** possess significantly lower melting points than those for corresponding ureas. Compounds **2f** and **2g** are liquid at room temperature, which suggests that their melting points are at least 193 and 167°C lower. The ureas are known to form intermolecular hydrogen bonds resulting in a brick wall fashion crystalline lattice. Thus, the strong decrease in melting point observed, coupled with the higher molecular weight of the imidazolidine-2,4,5-triones, illustrates a lower number of intermolecular interactions in the trione crystals. Therefore, the up to 115-fold decrease in inhibitory activity (except of compound **2d** which is 2.2-fold more active than its preceding urea) of imidazolidine-2,4,5-triones is somewhat not surprising. As observed for the single triones (Table 1), the di-triones are also 16- and 9-fold more soluble in water than their corresponding diureas.

To rationalize the activity and solubility trends in the current data sets of sEH inhibitors we performed a classic QSAR study based on fragment descriptors and regularized linear regression. The obtained model is satisfactory and can explain some features of the data set and the calculated F-statistics values confirmed the significance of the constructed models and the quality of the constructed models is satisfactory. The fragment descriptors which showed non-zero coefficients in the descriptor selection procedure are shown at Fig. 1 (A) (activity). The only descriptors which demonstrate suitable significance level in a t-test are *frag7* ($p = 0.07$) and *frag10* ($p = 0.002$). Their influence may be interpreted in the following way: *frag7* can be found in compounds which contain N-adamantylurea fragments which is easily accommodated in the sEH active site and form hydrogen bonds with Asp335 and Tyr383, while unsubstituted N-adamantyl-imidazolidine-trione containing compounds contain *frag10* and, according to the docking results, have an alternative binding mode to the enzyme. Asp335 can exist in two different conformations in sEH active site: the most stable is highlighted by red color (Fig 2 (A)) it forms two hydrogen bonds with the backbone

amides of Trp336 and Gly266 and the other one is shown at Fig 2 (A) with ordinary color scheme. The latter one is observed for the binding mode of compounds **2a** where one of carbonyl oxygens forms hydrogen bonds with the backbone amides of Trp336 and Gly266 substituting Asp335 side chain. Thus, the number of hydrogen bonds between the enzyme and the ligand are roughly the same for **2a** and its urea derivative **2a***. The urea derivative forms two hydrogen bonds with Asp335 while one of the carbonyl oxygens of **2a** forms two hydrogen bonds with the backbone amide groups of Trp336 and Phe267.

Finally, we synthesized two pyrimidine-2,4,6-triones **3a** and **3b** (Scheme 3) by substituting oxalyl chloride with malonyl chloride in the above described reaction.

Compound **3a** is more active than compound **3b** (Table 3), confirming that a single methylene spacer between the adamantane and the primary pharmacophore lead to increase of the inhibitory activity.⁹ Both pyrimidine-2,4,6-triones **3a** and **3b** show higher inhibitory activity against sEH when compared to the corresponding imidazolidine-2,4,5-triones **1e** and **1d** (Table 1), but they are up to 100-fold less active than the corresponding ureas. Pyrimidine-2,4,6-triones **3a** and **3b** are 21 and 10-fold more soluble than the ureas and approximately 4-fold more soluble than the corresponding imidazolidine-2,4,5-triones **1d** and **1e**. Because, like the imidazolidine-2,4,5-triones in some conditions, the pyrimidine-2,4,6-triones can degrade back to the original ureas, the better physical properties of the triones can enhanced their formulation as pro-drugs of urea-based sEHI.

Series of imidazolidine-2,4,5-triones and pyrimidine-2,4,6-triones were synthesized and investigated for their inhibitory activity against sEH as well as their physical properties. Synthesized compounds are less active than its preceding ureas but possess higher water solubility and lower melting point. Molecular docking with the new compounds highlights possible new bonds with the enzyme that could be used to increase potency. In addition, the new pharmacophore yield molecules that should easier to formulate, and that could be used as pro-drugs for urea-based inhibitor of sEH.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by Russian Fund for Basic Research (grant number 18-43-343002), Ministry of Education and Science of the Russian Federation (base part of state assignment for 2017-2019; project no. 4.7491.2017/BCh), National Institute of Environmental Health Sciences (NIEHS) grant R35 ES030443, and NIEHS Superfund Research Program grant P42 ES004699.

References and Notes

1. Arand M; Grant DF; Beetham JK; Friedberg T; Oesch F; Hammock BD FEBS Lett. 1994, 338, 251. [PubMed: 8307189]
2. Imig JD; Zhao X; Zaharis CZ; Olearczyk JJ; Pollock DM; Newman JW; Kim IH; Watanabe T; Hammock BD Hypertension. 2005, 46, 975. [PubMed: 16157792]
3. Fleming I; Rueben A; Popp R; Fisslthaler B; Schrodt S; Sander A; Haendeler J; Falck JR; Morisseau C; Hammock BD; Busse R Arterioscler. Thromb. Vasc. Biol 2007, 27, 2612. [PubMed: 17872452]

4. Imig JD Expert Opin. Drug Metab. Toxicol 2008, 4, 165. [PubMed: 18248310]
5. Hwang SH; Tsai HJ; Liu JY; Morisseau C; Hammock BD J. Med. Chem 2007, 50, 3825. [PubMed: 17616115]
6. Morisseau C; Goodrow MH; Newman JW; Wheelock CE; Dowdy DL; Hammock BD Biochem. Pharm 2002, 63,1599. [PubMed: 12007563]
7. Burmistrov V; Morisseau C; Harris TR; Butov G; Hammock BD Bioorg. Chem 2018, 76, 510. [PubMed: 29310082]
8. Zarriello S; Tuazon JP; Corey S; Schimmel S; Rajani M; Gorsky A; Incontri D; Hammock BD; Borlongan CV Prog. Neurobiol 2018, 172, 23. [PubMed: 30447256]
9. Burmistrov V; Morisseau C; Lee KSS; Shihadih DS; Harris TR; Butov GM; Hammock BD Bioorg. Med. Chem. Lett, 2014, 24, 2193. [PubMed: 24685540]
10. Hwang SH; Weckler AT; Zhang G; Morisseau C; Nguyen LV; Fu SH; Hammock BD Bioorg. Med. Chem. Lett, 2013, 23, 3732. [PubMed: 23726028]
11. Burmistrov V; Morisseau C; D'yachenko V; Rybakov VB; Butov GM; Hammock BD J. Fluor. Chem 2019, 220, 48.
12. Custelcean R Chem. Commun 2008, 295.
13. Kodani SD; Bhakta S; Hwang SH; Pakhomova S; Newcomer ME; Morisseau C; Hammock BD Bioorg. Med. Chem. Lett 2018, 28, 762. [PubMed: 29366648]
14. D'yachenko VS; Burmistrov VV; Nishi K; Kim I-H; Butov GM Chemistry of Heterocyclic Compounds. 2017, 53(10), 1080.
15. Kratky M; Mandikova J; Trejtnar F; Buchta V; Stola ikova J; Vinšova J Chem. Pap2015, 69, 1108.
16. Wagner K; Inceoglu B; Dong H; Yang J; Hwang SH; Jones P; Morisseau C; Hammock BD Eur. J. Pharm 2013, 700, 93.
17. Ulu A; Appt SE; Morisseau C; Hwang SH; Jones PD; Rose TE; Dong H; Lango J; Yang J; Tsai HJ; Miyabe C; Fortenbach C; Adams MR; Hammock BD British J. Pharm 2012, 165, 1401.
18. Jones PD; Wolf NM; Morisseau C; Whetstone P; Hock B; Hammock BD Anal. Biochem 2005, 343, 66. [PubMed: 15963942]
19. Burmistrov VV; Rasskazova EV; D'yachenko VS; Vernigora AA; Butov GM Russ. J. Org. Chem 2019, 55, 1229.




Figure 1.

(A) Molecular fragments have non-zero coefficients in the final regression equation for pIC₅₀. Central atom of each fragment is highlighted by blue color, the aliphatic carbon atoms are labeled by grey color while the aromatic ones are colored with yellow; (B) The final prediction results of the final model, the possible outlier is shown in red color; (C) The structural formula of the possible outlier.







Figure 2.
(A) The binding mode of the compound **2a**. Black points show the hydrogen bonds which compound **2a** maintain with protein residues. The green circle labels atoms which are in close proximity to each other. (B) The superposition of the docked structured of **2a** and its urea derivative in the binding site.

**Scheme 1.**

Reagents and conditions: *a*. Oxalyl chloride (1.2 eq.), THF, 66 °C, 2 h.

**Scheme 2.**












Reagents and conditions: *a*. Oxalyl chloride (1.2 eq.), THF, 66 °C, 2 h.












**Scheme 3.**

Reagents and conditions: a. Malonyl chloride (1.2 eq.), THF, 66 °C, 2 h.

Table 1

IC₅₀ values and some physicochemical properties for imidazolidine-2,4,5-triones **1a-k** and their corresponding ureas.

#	Structure	mp (°C)	Solubility (μM) ^a	Human sEH IC ₅₀ (nM) ^b
1a		212–214	280±10	86.8
1a*		194–195 ¹⁹	100±5	4.5
1b		295–296	210±10	7.7
1b* t-TUCB		244–273 ⁵	5 ¹⁶	1±0.1 ¹⁷
1c		205–207	250±10	0.4
1c*		241–243 ⁷	500±25 ⁷	0.4 ⁷
1d		155–158	250±10	6127
1d*		196–197 ¹¹	85±5 ¹¹	1.0 ¹¹
1e		90–91	350±10	310.6
1e*		191–192 ¹¹	65±5 ¹¹	0.7 ¹¹
1f		159–160	900±25	466.4

#	Structure	mp (°C)	Solubility (μM) ^a	Human sEH IC ₅₀ (nM) ^b
1f*		172–173 ¹¹	55±5 ¹¹	55.6 ¹¹
1g		129–130	200±10	370.7
1g*		181–182 ¹¹	65±5 ¹¹	3.6 ¹¹
1h		144–145	250±10	160.4
1h*		183–184 ¹¹	85±5 ¹¹	94.2 ¹¹
1i		260–263	100±5	22.8
1i*		266–268 ⁷	20±5 ⁷	9.6 ⁷
1j		153–155	100±5	216.1
1j*		194–196 ⁷	40±5 ⁷	13.8 ⁷
1k		185–186	150±10	8417
1k*		273–274	1750±25	12.0



^aSolubilities were measured in sodium phosphate buffer (pH 7.4, 0.1 M) containing 1% of DMSO.

^bDetermined via a kinetic fluorescent assay. Results are means of three separate experiments.¹⁸

Table 2

IC₅₀ values and some physicochemical properties for imidazolidine-2,4,5-triones **2a-h** and its corresponding preceding ureas.

#	Structure	mp (°C)	Solubility (μM) ^a	Human sEH IC ₅₀ (nM) ^b
2a		217–220	120±5	1.6
2a*		228–230 ⁹	25±5 ⁹	0.6 ⁹
2b		127–130	110±5	3.8
2b*		256–258 ⁹	30±5 ⁹	0.9 ⁹
2c		118–121	90±5	8.4
2c*		210–213 ⁹	n/a	2.3 ⁹
2d		122–125	100±5	649.6
2d*		130–131 ⁹	n/a	1442.6 ⁹
2e		39–40	900±25	294.4
2e*		229–230 ⁷	100±10 ⁷	3.4 ⁷
2f		n/a ^c	1300±25	92.0
2f*		192–193 ⁷	80±10 ⁷	0.8 ⁷
2g		n/a ^c	600±10	438.1
2g*		166–167 ⁷	75±5 ⁷	1.5 ⁷

#	Structure	mp (°C)	Solubility (μM) ^a	Human sEH IC ₅₀ (nM) ^b
2h		167–168	650±25	80.3
2h*		222–223	70±5	6.7

^aSolubilities were measured in sodium phosphate buffer (pH 7.4, 0.1 M) containing 1% of DMSO.

^bDetermined via a kinetic fluorescent assay. Results are means of three separate experiments.¹⁸

^cLiquid compounds. Do not crystallize at -20°C.





Table 3.

The obtained regression equations for activity and solubility

Outcome type	Equation	Q ²	R ²	RMSE _{cv}	F-value
Activity (pIC ₅₀)	$pIC_{50} = 7.386 - 0.174616 * counts_frag1 - 0.069048 * counts_frag2 - 0.043058 * counts_frag3 + 0.002409 * counts_frag4 + 0.015568 * counts_frag5 + 0.106981 * counts_frag6 + 0.107737 * counts_frag7 + 0.113848 * counts_frag8 + 0.177734 * counts_frag9 + 0.196090 * counts_frag10$	0.46	0.67	0.85	5.72
	descriptor	Std. error	t-value	p-value	
	intercept	0.81	9.01	< 10 ⁻⁶	
	frag1	0.19	-1.5565	0.12	
	frag2	0.16	-0.6346	0.53	
	frag3	0.09	-0.7240	0.47	
	frag4	0.03	0.2218	0.83	
	frag5	0.07	0.8327	0.41	
	frag6	0.11	0.9326	0.36	
	frag7	0.08	1.8443	0.07	
	frag8	0.22	1.1164	0.27	
	frag9	0.20	0.6367	0.53	
	frag10	0.08	3.2223	0.002	

Table 4

IC₅₀ values and some physicochemical properties for pyrimidine-2,4,6-triones **3a** and **3b** and its corresponding preceding ureas.

#	Structure	mp (°C)	Solubility (μM) ^a	Human sEH IC ₅₀ (nM) ^b
3a		90–91	1400±50	61.0
1a*		191–192 ¹¹	65±5 ¹¹	0.7 ¹¹
3b		115–116	900±50	210.4
1b*		196–197 ¹¹	85±5 ¹¹	1.0 ¹¹

^aSolubilities were measured in sodium phosphate buffer (pH 7.4, 0.1 M) containing 1% of DMSO.

^bDetermined via a kinetic fluorescent assay. Results are means of three separate experiments.¹⁸