UC Irvine

UC Irvine Previously Published Works

Title

Better cognitive control of emotional information is associated with reduced pro-inflammatory cytokine reactivity to emotional stress.

Permalink

<https://escholarship.org/uc/item/4cw651fq>

Journal

Stress (Amsterdam, Netherlands), 19(1)

ISSN

1025-3890

Authors

Shields, Grant S
Kuchenbecker, Shari Young
Pressman, Sarah D
et al.

Publication Date

2016

DOI

10.3109/10253890.2015.1121983

Peer reviewed

Stress

The International Journal on the Biology of Stress

ISSN: 1025-3890 (Print) 1607-8888 (Online) Journal homepage: <http://www.tandfonline.com/loi/ists20>

Better Cognitive Control of Emotional Information is Associated with Reduced Pro-Inflammatory Cytokine Reactivity to Emotional Stress

Grant S. Shields, Shari Young Kuchenbecker, Sarah D. Pressman, Kenneth D. Sumida & George M. Slavich

To cite this article: Grant S. Shields, Shari Young Kuchenbecker, Sarah D. Pressman, Kenneth D. Sumida & George M. Slavich (2015): Better Cognitive Control of Emotional Information is Associated with Reduced Pro-Inflammatory Cytokine Reactivity to Emotional Stress, *Stress*, DOI: [10.3109/10253890.2015.1121983](https://doi.org/10.3109/10253890.2015.1121983)

To link to this article: <http://dx.doi.org/10.3109/10253890.2015.1121983>

Accepted author version posted online: 19 Nov 2015.

Submit your article to this journal [↗](#)

Article views: 6

View related articles [↗](#)

View Crossmark data [↗](#)

Full Terms & Conditions of access and use can be found at
<http://www.tandfonline.com/action/journalInformation?journalCode=ists20>

Better Cognitive Control of Emotional Information is Associated with Reduced Pro-Inflammatory Cytokine Reactivity to Emotional Stress

Grant S. Shields, Shari Young Kuchenbecker, Sarah D. Pressman, Kenneth D. Sumida, and George M. Slavich

doi: 10.3109/10253890.2015.1121983

Abstract

Stress is strongly associated with several mental and physical health problems that involve inflammation, including asthma, cardiovascular disease, certain types of cancer, and depression. It has been hypothesized that better cognitive control of emotional information may lead to reduced inflammatory reactivity to stress and thus better health, but to date no studies have examined whether differences in cognitive control predict pro-inflammatory cytokine responses to stress. To address this issue, we conducted a laboratory-based experimental study in which we randomly assigned healthy young-adult females to either an acute emotional stress (emotionally evocative video) or no-stress (control video) condition. Salivary levels of the key pro-inflammatory cytokines IL-1 β , IL-6, and IL-8

were measured before and after the experimental manipulation, and following the last cytokine sample, we assessed participants' cognitive control of emotional information using an emotional Stroop task. We also assessed participants' cortisol levels before and after the manipulation to verify that documented effects were specific to cytokines and not simply due to increased nonwater salivary output. As hypothesized, the emotional stressor triggered significant increases in IL-1 β , IL-6, and IL-8. Moreover, even in fully adjusted models, better cognitive control following the emotional (but not control) video predicted less pronounced cytokine responses to that stressor. In contrast, no effects were observed for cortisol. These data thus indicate that better cognitive control specifically following an emotional stressor is uniquely associated with less pronounced pro-inflammatory cytokine reactivity to such stress. These findings may therefore help explain why superior cognitive control portends better health over the lifespan.

© 2015 Taylor & Francis. This provisional PDF corresponds to the article as it appeared upon acceptance. Fully formatted PDF and full text (HTML) versions will be made available soon.

DISCLAIMER: The ideas and opinions expressed in the journal's *Just Accepted* articles do not necessarily reflect those of Taylor & Francis (the Publisher), the Editors or the journal. The Publisher does not assume any responsibility for any injury and/or damage to persons or property arising from or related to any use of the material contained in these articles. The reader is advised to check the appropriate medical literature and the product information currently provided by the manufacturer of each drug to be administered to verify the dosages, the method and duration of administration, and contraindications. It is the responsibility of the treating physician or other health care professional, relying on his or her independent experience and knowledge of the patient, to determine drug dosages and the best treatment for the patient. *Just Accepted* articles have undergone full scientific review but none of the additional editorial preparation, such as copyediting, typesetting, and proofreading, as have articles published in the traditional manner. There may, therefore, be errors in *Just Accepted* articles that will be corrected in the final print and final online version of the article. Any use of the *Just Accepted* articles is subject to the express understanding that the papers have not yet gone through the full quality control process prior to publication.

**Better Cognitive Control of Emotional Information is Associated with
Reduced Pro-Inflammatory Cytokine Reactivity to Emotional Stress**

Grant S. Shields ¹, Shari Young Kuchenbecker ², Sarah D. Pressman ³,
Kenneth D. Sumida ⁴, and George M. Slavich ⁵

¹ Department of Psychology, University of California, Davis, CA, USA

² Department of Psychology, Chapman University, Orange, CA, USA

³ Department of Psychology and Social Behavior, University of California, Irvine, USA

⁴ Crean College of Health & Behavioral Sciences, Chapman University, Orange, CA, USA

⁵ Cousins Center for Psychoneuroimmunology and Department of Psychiatry and
Biobehavioral Sciences, University of California, Los Angeles, CA, USA

Correspondence: George M. Slavich, Ph.D., Cousins Center for Psychoneuroimmunology,
University of California, Los Angeles, UCLA Medical Plaza 300, Room 3156, Los Angeles, CA
90095-7076, USA. Tel: +1 310-825-2576. E-mail address: gslavich@mednet.ucla.edu

Article Type: Short Communication

Abstract: 250 words

Body: 2,625

References: 23

Figures: 1

Tables: 2

Abstract

Stress is strongly associated with several mental and physical health problems that involve inflammation, including asthma, cardiovascular disease, certain types of cancer, and depression. It has been hypothesized that better cognitive control of emotional information may lead to reduced inflammatory reactivity to stress and thus better health, but to date no studies have examined whether differences in cognitive control predict pro-inflammatory cytokine responses to stress. To address this issue, we conducted a laboratory-based experimental study in which we randomly assigned healthy young-adult females to either an acute emotional stress (emotionally evocative video) or no-stress (control video) condition. Salivary levels of the key pro-inflammatory cytokines IL-1 β , IL-6, and IL-8 were measured before and after the experimental manipulation, and following the last cytokine sample, we assessed participants' cognitive control of emotional information using an emotional Stroop task. We also assessed participants' cortisol levels before and after the manipulation to verify that documented effects were specific to cytokines and not simply due to increased nonwater salivary output. As hypothesized, the emotional stressor triggered significant increases in IL-1 β , IL-6, and IL-8. Moreover, even in fully adjusted models, better cognitive control following the emotional (but not control) video predicted less pronounced cytokine responses to that stressor. In contrast, no effects were observed for cortisol. These data thus indicate that better cognitive control specifically following an emotional stressor is uniquely associated with less pronounced pro-inflammatory cytokine reactivity to such stress. These findings may therefore help explain why superior cognitive control portends better health over the lifespan.

Keywords: life stress, cognition, emotion, Stroop, inflammation, cytokine, cortisol, risk, health, disease

Introduction

A large body of research has implicated psychological stress in the onset or progression of several disorders, including asthma, cardiovascular disease, certain cancers, and depression (Dienes et al., 2006; Slavich & Irwin, 2014). Because inflammation is involved in many different health outcomes, recent research has examined the role that pro-inflammatory cytokines play in linking stress with poor health (Cohen et al., 2012). Stress can trigger increased inflammatory activity at the molecular level (Slavich & Cole, 2013), and persistent elevations in inflammation can lead to a systemic inflammatory state that promotes disease (Cohen et al., 2012; Rohleder et al., 2010). Despite these associations, many people who experience stress do not develop inflammation-related health problems. Differences in how individuals biologically respond to stress might help explain health disparities following stress, but the factors that predict individual differences in cytokine reactivity to stress remain poorly understood.

One factor hypothesized to moderate biological reactivity to stress involves the extent to which individuals can cognitively control their thoughts and attention during stressful circumstances (Compton et al., 2013). Better cognitive control has been associated with less emotional reactivity to stress (Compton et al., 2011), and emotional responses to stress have in turn been found to predict individuals' pro-inflammatory cytokine reactivity to stress (Denson et al., 2009; Moons & Shields, 2015). One resulting possibility is that cognitive control, especially of emotional information, may influence cytokine responses to stress. To date, however, no studies have examined whether individuals' ability to cognitively control emotional information predicts their pro-inflammatory cytokine reactivity to emotional stress.

To address this issue, we randomly assigned healthy young-adult women to watch either an emotional stress-inducing or non-stress inducing video, before and after which we assessed salivary levels of three cytokines—namely, interleukin (IL)-1 β , IL-6, and IL-8. Salivary cytokine

are ideal for examining stress-related inflammatory reactivity because they are clinically relevant, and increase reliably and on a much faster timescale than cytokines in blood (Slavish et al., 2015). In addition, we characterized participants' cognitive control abilities using an emotional Stroop task. Consistent with prior research (Slavich & Irwin, 2014), we hypothesized that participants in the emotional stress-induction condition would exhibit greater pro-inflammatory cytokine responses than participants in the control group. We further hypothesized that better cognitive control of emotional information would be associated with less pronounced cytokine reactivity to the emotional stress-inducing (but not control) video, based on research showing that better cognitive control specifically within a stressful context predicts reduced reactivity to stress (Shields et al., 2015).

Lastly, to examine the specificity of these effects, we assessed participants' cortisol reactivity to the stress-inducing or non-stress inducing video. This permitted us to ensure that the emotional stressor specifically triggered increases in pro-inflammatory cytokines, which would be expected given that this stressor did not involve characteristics—specifically, uncontrollability and social-evaluative threat—that prior research has shown are required for a cortisol response (Dickerson & Kemeny, 2004). Assessing cortisol also enabled us to verify that individual differences in cognitive control of emotional information were specifically related to differences in inflammatory responding and not nonwater salivary output more generally.

Methods

Participants and Procedure

Participants were 37 healthy young-adult women from 18-22 years-old ($M_{\text{age}}=19.19$), sampled from a university community. Women taking hormone medication over the previous three months were excluded. Most participants ($N=21$) were in the luteal phase of their menstrual cycle. Participants' average body mass index (BMI) was 23.07 ($SD=5.07$). Oral health was controlled by ensuring participants' compliance with instructions on proper oral hygiene during the 48 hours before the visit.

Participants arrived between 12pm and 6pm for a one-hour study visit, and first completed brief screening measures.¹ They then provided baseline saliva samples for cytokine and cortisol measurement. Next, participants were randomly assigned to watch either a stress-inducing or non-stress inducing video (see below). A second saliva sample was collected approximately 10 minutes after the onset of the video, and participants then reported their state negative affect. Finally, participants completed the Stroop and were debriefed. The emotional stress manipulation was placed at the beginning of the study, prior to acclimation, to increase its stressfulness. The emotional stress-inducing video was not expected to influence performance on the cognitive task, given prior research showing that non-specific negative mood inductions do not impair cognitive control (Baumeister et al., 1998). All procedures adhered to American Psychological Association ethical principles and received Chapman University Institutional Review Board approval.

Emotional Stress Induction

Participants randomly assigned to the emotional stress-induction condition watched a 4-minute video of a 2-day-old crying infant being circumcised. Stimuli of this nature reliably induce an immune response (Schaller et al., 2010). In contrast, participants in the control

¹ Time of arrival did not influence the results.

condition watched a length-matched, non-emotional video. As described below, we evaluated the success of this stress manipulation by assessing participants' negative affect post-video, using the Positive and Negative Affect Scale (Watson et al., 1988). Reliability for this measure was excellent ($\alpha=.91$).

Cytokine and Cortisol Assays

We assessed participants' cytokine and cortisol levels immediately before and approximately 10 minutes after the video. We focused on the cytokines IL-1 β , IL-6, and IL-8 because they respond to emotional stressors and are implicated in the pathophysiology of several diseases. Saliva samples were obtained using a passive drool method. Samples were then immediately placed in a dry-ice filled container and transported to a -80°C freezer. Cytokines were measured using high-sensitivity multiplex immunoassay kits manufactured by R&D Systems (Minneapolis, MN), which have a minimal detectable dose of 0.04 pg/mL for IL-8, 0.08 pg/mL for IL-1 β , and 0.14 pg/mL for IL-6. The salivary cytokine levels we obtained were similar to those observed in prior studies (i.e., Byrne et al., 2013), and these levels tend to be higher than those obtained from blood (Slavish et al., 2015). Cortisol was measured using ELISA assay kits manufactured by R&D Systems (Minneapolis, MN), which have a minimum detectable dose of 0.16 ng/mL.

Cognitive Control

Cognitive control of emotional information was assessed using an emotional Stroop task. Participants viewed 108 Ekman male happy, sad, or angry expression faces with a superimposed happy, sad, or angry word (4 words per emotion). Thirty-six face/words plus an additional 12 control blurred faces were presented in each block (48 trials per block), with three blocks in total. Each trial began with the presentation of an emotional word overlaid on a face. Participants ignored the face and reported the emotion of the word. After participants responded, the face/word disappeared. The inter-trial interval varied from 1000-2000ms.

A cognitive control score was calculated for each participant by subtracting latencies to correctly label words on a control blurred/indistinguishable face from latencies to correctly label words on an angry face. We selected angry faces because anger produces the biggest emotion-related interference on such tasks (Gotlib et al., 2004). However, using happy or sad faces as the interfering faces did not alter the findings. Higher scores on this task indicate worse cognitive control of emotional information.

Analytic Strategy

Variables with significant skew (IL-1 β , IL-6, IL-8, cortisol, and negative affect) were log-transformed for analyses. To test the hypothesis that the emotional stress-inducing video would increase pro-inflammatory cytokine activity, and that differences in cognitive control would predict these responses, we conducted an Emotional Stroop $\times$ 2 (Condition: Stress vs. Control) $\times$ 3 (Cytokine Type: IL-1 β , IL-6, IL-8) mixed-model ANOVA using Condition as a between-subjects factor, Cytokine as a within-subjects factor, Emotional Stroop as a continuous variable, and post-video cytokines as the outcome. These analyses controlled for participants' age, BMI, hormonal cycle, and baseline cytokine levels. The covariates were chosen to avoid confounding the relation of biological reactivity and cognitive control with extraneous variables, but importantly, the results did not differ when these covariates were excluded. Analyses of cortisol were conducted using an ANOVA with the same covariates and factors, except that cortisol replaced the cytokine variables. Degrees of freedom in mixed-models were estimated using the Satterwaite approximation, which relaxes assumptions of homogeneity but entails that the degrees of freedom often contain non-integer numbers. All reported means and standard errors are least-squares means and their respective standard errors.

Results

Preliminary Analyses

Participants in the emotional stress induction and control groups did not differ on measures of age, BMI, menstrual cycle phase, or baseline levels of IL-1 β , IL-6, IL-8, or cortisol (p s>.17). Importantly, however, participants in the emotional stress condition reported more post-video negative affect ($M=2.82$, $SE=0.07$) than those in the control condition ($M=2.51$, $SE=0.08$), $t(34)=2.866$, $p=.007$, indicating that the experimental manipulation was successful. Finally, consistent with research showing no effects of acute negative mood inductions on Stroop performance, participants' cognitive control scores did not differ between conditions, $t(34)=0.115$, $p=.909$, which is critical for ensuring independence between the factors under study.

Primary Analyses

The mixed-model ANOVA used to test the main hypotheses revealed a significant main effect of Condition on cytokine reactivity, $F(1,25.7)=4.75$, $p=.039$, $\eta^2_{\text{partial}}=.156$, and a non-significant Condition $\times$ Cytokine Type interaction, $p=.816$ (see Table 1). These results indicate that cytokine reactivity differed significantly for participants in the two conditions and that this effect was similar for all three cytokines (see Table 2). Indeed, as hypothesized, participants in the emotional stress condition had greater cytokine reactivity ($M=3.74$, $SE=0.09$) than participants in the control group ($M=3.35$, $SE=0.12$), $t(27.6)=2.585$, $p=.015$. Therefore, the emotional stress induction was successful in triggering increased inflammatory activity relative to the control task.

Next, we examined the joint influence of Condition and participants' Emotional Stroop scores in predicting their pro-inflammatory cytokine levels. These analyses revealed a significant Condition $\times$ Emotional Stroop interaction effect, $F(1,25.7)=8.00$, $p=.009$, $\eta^2_{\text{partial}}=.237$, indicating that participants' cytokine reactivity depended on both their experimental condition and their ability to exert cognitive control over emotional information.

In addition, we found that the three-way Condition $\times$ Emotional Stroop $\times$ Cytokine Type interaction was non-significant, $p=.334$, indicating that these effects did not differ for the three cytokines assessed (See Figure 1).

To understand this interaction between Condition and Emotional Stroop, we conducted analyses of least-squares means. As hypothesized, among participants demonstrating better cognitive control (i.e., interference costs 1 *SD* below the mean), those in the emotional stress condition did not exhibit a greater cytokine response ($M=3.62$, $SE=0.13$) relative to those in the control condition ($M=3.72$, $SE=0.17$), $t(27.6)=-0.485$, $p=.631$. Conversely, among participants demonstrating average (i.e., interference costs at the mean) or worse cognitive control (i.e., interference costs 1*SD* above the mean), those in the emotional stress condition exhibited a significantly greater cytokine response than those in the control condition, $ps \leq .01$. In sum, therefore, individuals exhibiting better cognitive control of emotional information had no or only a negligible cytokine response to the emotional stressor, whereas those exhibiting average or poor cognitive control had a typical cytokine response to the stressor.

To verify that these results represented a specific link between cognitive control and cytokine reactivity and that they were not due to a general increase in secretory output from salivary glands, we focused next on cortisol reactivity, which was not expected to differ by condition given that the emotional stressor did not include characteristics required for a cortisol response—namely, uncontrollability and social-evaluative threat. As hypothesized, participants in the emotional stress condition did not exhibit greater cortisol reactivity than participants in the control condition, $F(1,30)=0.76$, $p=.390$ (Table 2). We also examined whether individual differences in cognitive control were associated with differences in cortisol responding or whether there was a Condition $\times$ Emotional Stroop interaction effect in the prediction of cortisol responding, but there were not, $Fs < 1.14$, $ps > .295$. Therefore, differences in cognitive control of

emotional information appear to be specifically related to differences in cytokine reactivity to emotional stress and not salivary output or salivary stress hormone reactivity more generally.

Discussion

Psychological stress is a well-known trigger of pro-inflammatory cytokine activity, but few studies have examined factors that predict individual differences in these reactions. We addressed this issue by showing that individuals' ability to exert cognitive control over emotional information was strongly associated with their salivary cytokine reactivity to a brief emotional stressor. These effects were specific to participants in the emotional stress condition and were robust when adjusting for participants' baseline cytokine levels, age, BMI, and menstrual cycle phase. Moreover, the emotional stressor did not trigger increases in cortisol, and there were no associations between participants' cognitive control characteristics and their cortisol dynamics. As such, we conclude that better cognitive control of emotional information specifically predicts reduced inflammatory reactivity to stress in young-adult women. These effects may thus help explain why superior cognitive control is associated with better health.

Although no studies have examined associations between cognitive control and inflammatory reactivity to stress, one study has investigated links between cognitive control and cortisol reactivity. Specifically, Compton et al. (2013) found that better cognitive control during a cognitive stressor predicted lesser cortisol reactivity. We did not observe such associations here, but we also did not expect these effects given that the emotional stressor we employed did not involve uncontrollability or social-evaluative threat, which are required for cortisol reactivity (Dickerson & Kemeny, 2004).

What mechanism might underlie salivary cytokine increases in the timeframe assessed here? We can only speculate, but the observed increases may represent a stress-induced redistribution of immune system resources. Saliva contains high concentrations of cytokines that

are part of a complex network that is involved in innate and adaptive immunity (Fábián et al., 2012). During stress, cytokines are redistributed closer to sites of potential injury or infection (e.g., the mouth) to protect the host against possible viruses or bacteria (Dhabhar & McEwen, 1996). This redistribution is partially mediated by sympathetic activity, which is extremely rapid and could thus cause increases in cytokines within the timeframe assessed here (Bosch et al., 2005). Notably, this is not the first study to document increases in salivary cytokines approximately 10 minutes after onset of a stressor (see Minetto et al., 2005).

Several limitations should be noted. First, the sample was relatively small, homogenous, and female. Additional research is thus needed to examine the generalizability of these effects. Second, research has demonstrated the validity of oral measures of cytokine activity (Slavish et al., 2015), but confirmatory results using other sampling procedures would be beneficial. Third, we examined only one type of stress—emotional stress. Because other stressors (e.g., uncontrollable, socially evaluative stressors) elicit different biological responses and may be preferentially associated with mental health outcomes such as learned helplessness (Grahn et al., 1999), future research should study relations between cognitive control, cytokines, and other types of stress. Finally, although the stress manipulation was experimental, all of the associations with cognitive control were correlational. Future studies should thus examine whether cytokine reactivity differs after manipulating cognitive control.

Several strengths are also noteworthy. First, all of the results held across the three cytokines that were measured, providing a within-study replication of the effects. Second, we employed the emotional Stroop task, which is a gold-standard measure of cognitive control. Third, the effect sizes obtained in this study indicated that cognitive control accounted for a large amount of variance (i.e., 23.7%) in stress-induced changes in cytokines, even after controlling for covariates. Finally, as a result, the data may elucidate a biological mechanism

underlying previously observed associations between cognitive control and health (Shields et al., 2015).

In conclusion, we found that individuals who demonstrate better cognitive control over emotional information exhibit less pronounced salivary cytokine responses to an emotional stressor. Although stress increases risk for several inflammation-related disease conditions that represent the leading causes of death in the United States today (Slavich, 2015), many individuals who experience major stressors do not get ill. The present data are important in this context as they highlight a novel mechanism—specifically, cognitive control—that could potentially be modified to reduce inflammatory reactivity and improve health. Additional research should examine the generalizability of these findings and elucidate neurocognitive and biological mechanisms that link cognitive control with cytokine reactivity and health.

Acknowledgements

Preparation of this report was supported by National Institute of Mental Health grant K08 MH103443 and a Society in Science – Branco Weiss Fellowship to George Slavich. We thank Le'Quan Jackson, Brendan Everett, Cecilia Lau, and Jared Celniker for helping conduct the study; Naveen Jonathan at the Chapman University Frances Smith Center for Individual and Family Therapy for providing rooms for subject running; and David Barrow at the UNC Cytokine and Biomarker Analysis Facility for conducting the cytokine assays. Shari Young Kuchenbecker is now at R.W. Research, Inc. and at the Western Positive Psychology Association.

Declaration of interest

There are no conflicts of interest that could have inappropriately influenced the study or the reporting of the findings.

References

- Baumeister RF, Bratslavsky E, Muraven M, Tice DM. (1998). Ego depletion: is the active self a limited resource? *J Pers Soc Psychol* 74:1252-1265.
<http://dx.doi.org/10.1037/0022-3514.74.5.1252>
- Bosch JA., Berntson GG, Cacioppo JT, Marucha PT. (2005). Differential mobilization of functionally distinct natural killer subsets during acute psychologic stress. *Psychosom Med* 67:366-375.
- Byrne ML, O'Brien-Simpson NM, Reynolds EC, Walsh KA, Laughton K, Waloszek JM, . . . Allen NB. (2013). Acute phase protein and cytokine levels in serum and saliva: a comparison of detectable levels and correlations in a depressed and healthy adolescent sample. *Brain Behav Immun* 34:164-175. <http://dx.doi.org/10.1016/j.bbi.2013.08.010>
- Cohen S, Janicki-Deverts D, Doyle WJ, Miller GE, Frank E, Rabin BS, Turner RB. (2012). Chronic stress, glucocorticoid receptor resistance, inflammation, and disease risk. *Proc Natl Acad Sci USA* 109:5995-5999.
<http://dx.doi.org/10.1073/pnas.1118355109>
- Compton RJ, Arnstein D, Freedman G, Dainer-Best J, Liss A, Robinson MD. (2011). Neural and behavioral measures of error-related cognitive control predict daily coping with stress. *Emotion* 11:379-390. <http://dx.doi.org/10.1037/a0021776>
- Compton RJ, Hofheimer J, Kazinka R. (2013). Stress regulation and cognitive control: evidence relating cortisol reactivity and neural responses to errors. *Cogn Affect Behav Neurosci* 13:152-163. <http://dx.doi.org/10.3758/s13415-012-0126-6>
- Denson TF, Spanovic M, Miller N. (2009). Cognitive appraisals and emotions predict cortisol and immune responses: a meta-analysis of acute laboratory social stressors and emotion inductions. *Psychol Bull* 135:823-853. <http://dx.doi.org/10.1037/a0016909>

- Dhabhar FS, McEwen BS. (1996). Stress-induced enhancement of antigen-specific cell-mediated immunity. *J Immunol* 156:2608-2615.
- Dickerson SS, Kemeny ME. (2004). Acute stressors and cortisol responses: a theoretical integration and synthesis of laboratory research. *Psychol Bull* 130:355-391.
<http://dx.doi.org/10.1037/0033-2909.130.3.355>
- Dienes KA, Hammen CL, Henry RM, Cohen AN, Daley S. (2006). The stress sensitization hypothesis: understanding the course of bipolar disorder. *J Affect Disord* 95:43-49.
<http://dx.doi.org/10.1016/j.jad.2006.04.009>
- Fábián TK, Hermann P, Beck A, Fejérdy P, Fábián G. (2012). Salivary defense proteins: their network and role in innate and acquired oral immunity. *Int J Mol Sci* 13:4295-4320.
<http://dx.doi.org/10.3390/ijms13044295>
- Gotlib IH, Krasnoperova E, Yue DN, Joormann J. (2004). Attentional biases for negative interpersonal stimuli in clinical depression. *J Abnorm Psychol* 113:127-135.
<http://dx.doi.org/10.1037/0021-843X.113.1.121>
- Grahn RE, Will MJ, Hammack SE, Maswood S, McQueen MB, Watkins LR, Maier SF. (1999). Activation of serotonin-immunoreactive cells in the dorsal raphe nucleus in rats exposed to an uncontrollable stressor. *Brain Res*, 826:35-43.
[http://dx.doi.org/10.1016/S0006-8993\(99\)01208-1](http://dx.doi.org/10.1016/S0006-8993(99)01208-1)
- Minetto M, Rainoldi A, Gazzoni M, Terzolo M, Borrión P, Termine A, ... Paccotti P. (2005). Differential responses of serum and salivary interleukin-6 to acute strenuous exercise. *Eur J Appl Physiol* 93:679-686. <http://dx.doi.org/10.1007/s00421-004-1241-z>
- Moons WG, Shields GS. (2015). Anxiety, not anger, induces inflammatory activity: an avoidance/approach model of immune system activation. *Emotion* 15:463-476.
<http://dx.doi.org/10.1037/emo0000055>

- Rohleder N, Wolf JM, Wolf OT. (2010). Glucocorticoid sensitivity of cognitive and inflammatory processes in depression and posttraumatic stress disorder. *Neurosci Biobehav Rev* 35:104-114. <http://dx.doi.org/10.1016/j.neubiorev.2009.12.003>
- Schaller M, Miller GE, Gervais WM, Yager S, Chen E. (2010). Mere visual perception of other people's disease symptoms facilitates a more aggressive immune response. *Psychol Sci* 21:649-652. <http://dx.doi.org/10.1177/0956797610368064>
- Shields GS, Moons WG, Slavich GM. (2015). Better executive function following acute stress is associated with less perceived stress and fewer life stress-related physical health complaints in young adulthood. Manuscript in preparation.
- Slavich GM. (2015). Understanding inflammation, its regulation, and relevance for health: a top scientific and public priority. *Brain Behav Immun* 45:13-14. <http://dx.doi.org/10.1016/j.bbi.2014.10.012>
- Slavich GM, Cole SW. (2013). The emerging field of human social genomics. *Clin Psychol Sci* 1:331-348. <http://dx.doi.org/10.1177/2167702613478594>
- Slavich GM, Irwin MR. (2014). From stress to inflammation and major depressive disorder: a social signal transduction theory of depression. *Psychol Bull* 140:774-815. <http://dx.doi.org/10.1037/a0035302>
- Slavish DC, Graham-Engeland JE, Smyth JM, Engeland CG. (2015). Salivary markers of inflammation in response to acute stress. *Brain Behav Immun* 44:253-269. <http://dx.doi.org/10.1016/j.bbi.2014.08.008>
- Watson D, Clark LA, Tellegen A. (1988). Development and validation of brief measures of positive and negative affect: the PANAS scales. *J Pers Soc Psychol* 54:1063-1070. <http://dx.doi.org/10.1037/0022-3514.54.6.1063>

Table 1

Type III SSs Mixed-Model ANOVA Table from Primary Analysis

Variable	MSE	df_1	df_2	F	p
Baseline Cytokines	378.23	1	84.9	359.21	<.001
Condition	0.21	1	25.7	4.75	.039
Emotional Stroop	0.00	1	25.8	2.05	.164
Cytokine Type	0.42	2	69.3	7.06	.002
Body Mass Index	0.00	1	25.7	1.11	.303
Age	0.12	1	26.0	1.85	.186
Phase of the menstrual cycle	0.00	1	26.2	0.13	.726
Condition × Emotional Stroop	0.44	1	25.7	8.00	.009
Condition × Cytokine Type	0.01	2	59.7	0.20	.816
Emotional Stroop × Cytokine Type	0.04	2	59.1	0.80	.455
Condition × Emotional Stroop × Cytokine Type	0.06	2	59.4	1.12	.334

Note: Significant effects are presented in boldface.

Table 2

Descriptive Statistics of Natural Log-Transformed Cytokine and Cortisol Levels by Experimental Condition

Variable	Pre-Video Mean (SE)	Post-Video Mean (SE)	Correlation Pre- to Post-Video
Control Condition			
Interleukin-1 β (ln)	4.30 (0.27)	4.06 (0.23)	.92
Interleukin-6 (ln)	0.93 (0.19)	0.80 (0.17)	.86
Interleukin-8 (ln)	5.69 (0.21)	5.61 (0.18)	.82
Cortisol (ln)	1.19 (0.20)	1.04 (0.21)	.94
Emotional Stressor Condition			
Interleukin-1 β (ln)	4.14 (0.24)	4.24 (0.20)	.88
Interleukin-6 (ln)	1.29 (0.16)	1.43 (0.15)	.61
Interleukin-8 (ln)	5.52 (0.19)	5.70 (0.16)	.80
Cortisol (ln)	1.14 (0.18)	0.92 (0.18)	.87

Note: All cytokine and cortisol values were natural log transformed to correct for skew.

Figure Caption

Figure 1. Moderation of pro-inflammatory cytokine responses to acute stress by cognitive control of emotional information for participants randomly assigned to an acute emotional stress (emotionally evocative video) versus no-stress (control video) condition. Across the three cytokines measured (i.e., IL-1 β , IL-6, and IL-8), better cognitive control of emotional information was associated with less pronounced cytokine reactivity for participants randomly assigned to the emotional stress (but not control) condition, even after adjusting for several covariates—specifically, participants' baseline cytokine levels, age, BMI, and menstrual cycle phase. The bar graphs present raw data (pg/mL) for illustrative purposes, while analyses, as well as the indicators of significance on the bar graphs, controlled for covariates and used the natural log transformation of each cytokine to correct for skew.