

# UC Davis

## UC Davis Previously Published Works

### Title

Adenylyl Cyclase 6 Mediates Inhibition of TNF in the Inflammatory Reflex

### Permalink

<https://escholarship.org/uc/item/4cz7992d>

### Journal

Frontiers in Immunology, 9(NOV)

### ISSN

1664-3224

### Authors

Tarnawski, Laura

Reardon, Colin

Caravaca, April S

et al.

### Publication Date

2018

### DOI

10.3389/fimmu.2018.02648

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



# Adenylyl Cyclase 6 Mediates Inhibition of TNF in the Inflammatory Reflex

Laura Tarnawski<sup>1\*</sup>, Colin Reardon<sup>2</sup>, April S. Caravaca<sup>1</sup>, Mauricio Rosas-Ballina<sup>3</sup>, Michael W. Tusche<sup>4</sup>, Anna R. Drake<sup>5</sup>, LaQueta K. Hudson<sup>3</sup>, William M. Hanes<sup>3</sup>, Jian Hua Li<sup>3</sup>, William R. Parrish<sup>3</sup>, Kaie Ojamaa<sup>6</sup>, Yousef Al-Abed<sup>3</sup>, Michael Faltys<sup>5</sup>, Valentin A. Pavlov<sup>3</sup>, Ulf Andersson<sup>3,7</sup>, Sangeeta S. Chavan<sup>3</sup>, Yaakov A. Levine<sup>5</sup>, Tak W. Mak<sup>4</sup>, Kevin J. Tracey<sup>3</sup> and Peder S. Olofsson<sup>1,3</sup>

<sup>1</sup> Department of Medicine, Center for Bioelectronic Medicine, Center for Molecular Medicine, Solna, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden, <sup>2</sup> Department of Anatomy, Physiology & Cell Biology, University of California, Davis, Davis, CA, United States, <sup>3</sup> Laboratory of Biomedical Science, The Feinstein Institute for Medical Research, Northwell Health, Manhasset, NY, United States, <sup>4</sup> The Campbell Family Institute for Breast Cancer Research, University Health Network, Toronto, ON, Canada, <sup>5</sup> SetPoint Medical, Valencia, CA, United States, <sup>6</sup> Department of Biomedical Sciences, New York Institute of Technology College of Osteopathic Medicine, Old Westbury, NY, United States, <sup>7</sup> Department of Women's and Children's Health, Karolinska Institutet, Stockholm, Sweden

## OPEN ACCESS

### Edited by:

Christoph Thiemermann,  
Queen Mary University of London,  
United Kingdom

### Reviewed by:

Sergio Iván Valdés-Ferrer,  
Instituto Nacional de Ciencias  
Médicas y Nutrición Salvador Zubirán  
(INCMNSZ), Mexico  
Sina Maren Coldewey,  
Universitätsklinikum Jena, Germany

### \*Correspondence:

Laura Tarnawski  
laura.tarnawski@ki.se

### Specialty section:

This article was submitted to  
Inflammation,  
a section of the journal  
Frontiers in Immunology

**Received:** 12 September 2018

**Accepted:** 26 October 2018

**Published:** 27 November 2018

### Citation:

Tarnawski L, Reardon C, Caravaca AS, Rosas-Ballina M, Tusche MW, Drake AR, Hudson LK, Hanes WM, Li JH, Parrish WR, Ojamaa K, Al-Abed Y, Faltys M, Pavlov VA, Andersson U, Chavan SS, Levine YA, Mak TW, Tracey KJ and Olofsson PS (2018) Adenylyl Cyclase 6 Mediates Inhibition of TNF in the Inflammatory Reflex. *Front. Immunol.* 9:2648. doi: 10.3389/fimmu.2018.02648

Macrophage cytokine production is regulated by neural signals, for example in the inflammatory reflex. Signals in the vagus and splenic nerves are relayed by choline acetyltransferase<sup>+</sup> T cells that release acetylcholine, the cognate ligand for alpha7 nicotinic acetylcholine subunit-containing receptors ( $\alpha 7nAChR$ ), and suppress TNF release in macrophages. Here, we observed that electrical vagus nerve stimulation with a duration of 0.1–60 s significantly reduced systemic TNF release in experimental endotoxemia. This suppression of TNF was sustained for more than 24 h, but abolished in mice deficient in the  $\alpha 7nAChR$  subunit. Exposure of primary human macrophages and murine RAW 264.7 macrophage-like cells to selective ligands for  $\alpha 7nAChR$  for 1 h *in vitro* attenuated TNF production for up to 24 h in response to endotoxin. Pharmacological inhibition of adenylyl cyclase (AC) and knockdown of adenylyl cyclase 6 (AC6) or c-FOS abolished cholinergic suppression of endotoxin-induced TNF release. These findings indicate that action potentials in the inflammatory reflex trigger a change in macrophage behavior that requires AC and phosphorylation of the cAMP response element binding protein (CREB). These observations further our mechanistic understanding of neural regulation of inflammation and may have implications for development of bioelectronic medicine treatment of inflammatory diseases.

**Keywords:** inflammatory reflex, choline acetyltransferase, acetylcholine, adenylyl cyclase 6, sustained TNF inhibition, vagus nerve stimulation/VNS,  $\alpha 7nAChR$

## INTRODUCTION

Infection and injury activate immune cells to produce cytokines and other factors that mediate inflammation. Macrophage function is critical for protection from infection and injury. Animals rendered deficient in macrophages, or subjected to genetic modifications that prevent cytokine production, have a diminished inflammatory response. During inflammation, the cellular response can be sustained for hours or days. Recent discoveries reveal that the responsiveness of innate immune cells when re-encountering pathogens is modulated by transcription

factors and epigenetic reprogramming (1). However, the mechanisms initiating these changes that regulate inflammation are not fully understood.

Glucocorticoids, anti-inflammatory cytokines, immunoresolvents, and other humoral anti-inflammatory agents regulate immune responses and the resolution of inflammation, e.g. by inhibiting macrophage production of pro-inflammatory cytokines (2). Neural reflex circuits also regulate macrophage release of pro-inflammatory cytokines (3–6). Electrical activation of the vagus nerve (milliseconds to minutes) can significantly reduce endotoxin-induced release of Tumor Necrosis Factor (TNF), IL-1, IL-6, and HMGB1 (7, 8). Data from experimental disease models and clinical studies using vagus nerve stimulation (VNS) to treat rheumatoid arthritis and Crohn's disease indicate that transient electrical stimulation of the vagus nerve results in a prolonged alleviation of inflammation and disease severity (9–12). However, the mechanism underlying this sustained reduction of cytokine production in response to transient nerve activation has been unknown.

It was previously established that action potentials in the vagus nerve culminate in activation of the  $\alpha 7$  nicotinic acetyl choline receptor subunit ( $\alpha 7$ nAChR) on macrophages (4, 13–15). The molecular signal that completes this inhibitory reflex requires an intact splenic nerve, and choline acetyltransferase<sup>+</sup> (ChAT<sup>+</sup>) T cells that release acetylcholine (ACh), the cognate ligand for the  $\alpha 7$ nAChR (5). Several mechanisms by which nAChRs control inflammatory responses have been proposed, including activation of JAK2/STAT3 and inhibition of NF $\kappa$ B (16–19). The  $\alpha 7$ nAChR is an ion channel in neurons (20) but reportedly, the ion flux itself is not sufficient to mediate inhibition of TNF release (21), suggesting involvement of other molecular components.

In neurons, signal transduction via  $\alpha 7$ nAChR increases cyclic adenosine monophosphate (cAMP) levels and phosphorylation of the transcription factor cAMP response element binding protein (CREB) (22), a pathway that is essential for memory (23). In macrophages, increased cAMP promotes intracellular accumulation of the transcription factor c-FOS, and promotes suppression of inflammatory cytokine production (24). Interestingly, CREB has been implicated in cAMP-induced inhibition of NF $\kappa$ B-mediated transcription of TNF (25). Accordingly, we hypothesized that a cAMP and CREB-mediated increase in *c-Fos* expression underlies the development of persistently attenuated macrophage responses after VNS.

Here, we observed that VNS  $\geq 24$  h prior to an inflammatory insult reduced endotoxin induced TNF release through a mechanism that required the  $\alpha 7$ nAChR. Cholinergic inhibition of TNF in macrophages was abolished by inhibition of adenylyl cyclase, CREB phosphorylation, and expression of *c-Fos*.


## RESULTS

### Vagus Nerve Stimulation Inhibits Cytokine Release for More Than 24 h

To study the kinetics of TNF release in endotoxemia following VNS, we subjected rats to vagus nerve stimulation (VNS) for 60 s at 10 Hz, or sham surgery. The rats were then injected

intraperitoneally with bacterial endotoxin (lipopolysaccharide, LPS) at various time points after VNS. Serum TNF levels were significantly lower in VNS-treated rats compared to sham treated animals when endotoxin was administered 0, 2, 24, and 48 h after stimulation (Figure 1A). The VNS-mediated inhibition of endotoxin-induced TNF was not significant at 72 h after nerve stimulation.

Next, mice were treated with 60 s of VNS at 10 Hz, or sham surgery, and subjected to endotoxemia 24 h after treatment. Endotoxin-induced TNF levels were significantly lower in VNS-treated mice as compared to sham treated animals (Figure 1B). We previously observed that a single pulse, equivalent to 0.1 s stimulation, is sufficient to significantly reduce TNF release in



experimental endotoxemia (8). To investigate if a single pulse confers a sustained reduction in TNF release, mice treated with 0.1 s of VNS were subjected to endotoxemia, 24 h after treatment. Endotoxin-induced TNF levels were significantly lower in VNS treated mice compared to sham treated animals (Figure 1B).

Together, these results suggest that a short activation of the inflammatory reflex reduces TNF production during endotoxemia for  $\geq 24$  h.

### $\alpha 7$ nAChR Is Required for the Sustained Inhibition of Cytokine Release *in vivo*


Prior work established that the  $\alpha 7$  nicotinic acetylcholine receptor subunit ( $\alpha 7$ nAChR) expressed by macrophages is required for the functional integrity of the inflammatory reflex (14, 15, 26). To study the role of  $\alpha 7$ nAChR in the persistent inhibition by VNS of TNF release in endotoxemia, we next studied  $\alpha 7$ nAChR deficient and wild type mice. Endotoxin-induced TNF was significantly reduced at 24 h after VNS in wild type mice (Figure 1C, left panel), but not in  $\alpha 7$ nAChR deficient mice (Figure 1C, right panel). Thus,  $\alpha 7$ nAChR is required for the persistent inhibition of TNF release by transient electrical stimulation of the vagus nerve.

### Exposure of Macrophages to Selective $\alpha 7$ nAChR Agonists Increases Intracellular cAMP Levels

$\alpha 7$ nAChR subunits form ion channels on neuronal cells (20). Although ion flux has been observed in macrophages exposed to cholinergic agonists, this effect is reportedly unlikely to affect cytokine production (21).

In neurons,  $\alpha 7$ nAChR can interact with adenylyl cyclase 6 (AC6) and promote cAMP formation (27, 28). To investigate the possibility that a similar interaction occurs in macrophages, bone marrow derived macrophages from  $\alpha 7$ nAChR  $^{+/+}$  and  $\alpha 7$ nAChR  $^{-/-}$  mice, respectively, were exposed *in vitro* to the selective  $\alpha 7$ nAChR agonist, *N*-(3*R*)-1-Azabicyclo[2.2.2]oct-3-yl-2,3-dihydro-1,4-benzodioxin-6-carboxamide fumarate (PHA568487) (29). Addition of PHA568487 significantly increased cAMP levels in  $\alpha 7$ nAChR  $^{+/+}$ , but not in  $\alpha 7$ nAChR  $^{-/-}$  derived macrophages (Figure 2A), indicating that  $\alpha 7$ nAChR signaling regulates cAMP in murine macrophages.

To study the mechanism of cholinergic control of TNF release we utilized macrophage-like RAW 264.7 cells. First, we investigated whether cholinergic activation confers prolonged inhibition of endotoxin-induced TNF release in RAW 264.7 cells. Cells were exposed to 10  $\mu$ M ACh or vehicle for 1 h, and then washed and re-suspended in fresh medium. Cells were then rested for 24 h and subsequently exposed to endotoxin for 4 h. TNF levels in culture medium were significantly lower in cultures treated with ACh (Figure 2B) compared to cultures treated with vehicle. To investigate whether blocking of cAMP synthesis prevents cholinergic inhibition of endotoxin-induced TNF expression, RAW 264.7 cells were cultured in the presence of the adenylyl cyclase-inhibitor [cis-N-(2-Phenylcyclopentyl)azacyclotridec-1-en-2-amine.HCl] (MDL




12,330A) and exposed to the selective  $\alpha 7$ nAChR agonist choline for 30 min. The cells were then incubated with endotoxin for 3 h and TNF mRNA was measured in cell lysates by qPCR. Choline-mediated suppression of TNF mRNA was significantly reduced in the presence of MDL 12,330A (Figure 2C). Based on the observations that  $\alpha 7$ nAChR can interact with AC6 (27)

and that blocking cAMP synthesis reduced cholinergic inhibition of endotoxin-induced TNF expression, RAW 264.7 cells were transfected with siRNA targeting AC6 or scrambled siRNA. AC6 protein knock-down was confirmed by Western blot (Figure 2D, inset). Transfected cells were subsequently incubated with choline and cultured in the presence of endotoxin for 3 h. Knock down of AC6 cells abolished the choline-dependent reduction of TNF mRNA (Figure 2D), indicating that AC6 is required for the  $\alpha 7$ nAChR-mediated reduction of endotoxin-induced TNF mRNA. Together, these data indicate that cholinergic inhibition of endotoxin-induced TNF expression requires AC6-dependent cAMP production.

### c-Fos Is Essential for $\alpha 7$ nAChR-Mediated Inhibition of TNF

It was previously established that increased cAMP levels promote expression and phosphorylation of c-FOS, which regulates macrophage TNF expression (24). Accordingly, we measured c-FOS protein in endotoxin-exposed RAW 264.7 cells in the presence of choline or vehicle (Figure 3A). Addition of choline augmented the production of c-FOS, supporting involvement of c-FOS in  $\alpha 7$ nAChR-mediated regulation of TNF. Based on these observations, we hypothesized that increased c-Fos expression in response to selective  $\alpha 7$ nAChR agonists contributes to the cholinergic inhibition of TNF production in macrophages. To test this, RAW 264.7 cells were transfected with siRNA targeting c-Fos or scrambled siRNA. Transfected cells were subsequently incubated with choline and cultured in the presence of endotoxin for 3 h. In cells transfected with scrambled siRNA, choline significantly inhibited endotoxin-induced TNF. In contrast, the inhibitory effect on endotoxin-mediated TNF expression was completely abolished in cells transfected with siRNA targeting c-Fos (Figure 3B). These data demonstrate that c-Fos is required for cholinergic suppression of endotoxin-induced TNF expression.




### Phosphorylation Defective CREB Prevents $\alpha 7$ nAChR-Mediated Inhibition of TNF

In neurons, signal transduction via  $\alpha 7$ nAChR leads to increased cAMP levels and phosphorylation of the transcription factor CREB (22), a pathway that is essential for long-term retention of information, i.e., memory (23). CREB has also been implicated in cAMP-induced inhibition of NF $\kappa$ B-mediated transcription of TNF (25) and activation of CREB is known to promote expression of c-Fos (30, 31). RAW 264.7 cells and human macrophages cultured in the presence of ACh or vehicle showed increased CREB phosphorylation in cultures exposed to ACh as measured by western blot (data not shown). Accordingly, we hypothesized that CREB phosphorylation underlies sustained cholinergic reduction of endotoxin-induced TNF release in macrophages. To study this, we first established whether cholinergic stimulation of human differentiated macrophages confers sustained inhibition of TNF release. Human macrophages *in vitro* were exposed to ACh or vehicle for 1 h, the ACh washed away, and the cells rested for 24 h. Subsequently, cultures were exposed to endotoxin for 4 h and TNF levels were measured in culture medium. ACh exposure significantly reduced endotoxin-induced TNF release to the medium 24 h after ACh exposure (Figure 4A), demonstrating that ACh exposure confers persistent reduction in endotoxin-induced TNF release in human macrophages. Next, human macrophages were transduced with a phosphorylation defective CREB (ACREB), using an engineered adenovirus (32). TNF production by human primary macrophages transduced with control virus was significantly inhibited at 24 h after exposure to the selective  $\alpha 7$ nAChR agonist 3-(2,4-dimethoxybenzylidene)anabaseine (GTS21; Figure 4B). However, GTS21 failed to elicit prolonged deactivation of macrophages transduced with the ACREB containing adenovirus (Figure 4B), indicating that CREB phosphorylation is required for the sustained inhibition of cytokine release.

### DISCUSSION

The results here indicate that electrical stimulation of the vagus nerve for 0.1–60 s reduces endotoxin-induced TNF release for  $\geq 24$  h by a mechanism that requires the  $\alpha 7$ nAChR subunit. Exposure of macrophages to selective  $\alpha 7$ nAChR agonists for 1 h resulted in prolonged inhibition of endotoxin-induced TNF release. The effect required adenylyl cyclase, CREB phosphorylation and c-FOS. Thus, we observe that macrophages retain information entrained by electrical signals in the vagus nerve.

VNS pulse trains shorter than 10 min significantly reduces inflammation and release of pro-inflammatory cytokines in experimental models of inflammation and clinical trials (9, 11, 12, 15). In the present study, one minute of VNS attenuated the systemic response to endotoxin for  $\geq 24$  h in mice and rats *in vivo*. For the acute effect of VNS on endotoxin-induced TNF release, the  $\alpha 7$ nAChR subunit is required (14, 15). The prolonged inhibition of endotoxin-induced TNF release observed here also required the  $\alpha 7$ nAChR subunit, as VNS




**FIGURE 4 |** Functional CREB mediates persistent inhibition of TNF. **(A)** Human macrophages were incubated *in vitro* with acetylcholine (ACh), washed and rested for 24 h, and subsequently exposed to endotoxin (LPS) for 4 h. TNF was measured in the culture media using ELISA. **(B)** Primary human macrophages were transduced with adenovirus engineered to encode a dominant negative phosphorylation defective CREB (ACREB), or a GFP-expressing control adenovirus. After viral transduction, cells were stimulated with the selective  $\alpha 7$ nAChR agonist GTS21, washed, and cultured for 24 h before exposure to endotoxin. TNF was measured in culture media by ELISA. Mean TNF levels  $\pm$  SEM in cells transduced with control virus or the phosphorylation defective ACREB-expressing virus (ACREB) are shown. \* $p < 0.05$ .

failed to reduce endotoxin-induced TNF release 24 h after stimulation in  $\alpha 7$ nAChR<sup>-/-</sup> mice. Priming of macrophages with a cholinergic agonist resulted in significant attenuation of TNF production 24 h after agonist removal, revealing a mechanism that can persist for hours after a transient nerve signal.

Cholinergic ligands are known to reduce NF $\kappa$ B activation and TNF release in macrophages (18, 19). de Jonge and colleagues concluded that nicotine increases phosphorylation of STAT3 and reduced TNF release from peritoneal macrophages without affecting TNF transcription (16). Others have shown that increased cAMP levels attenuate TNF expression through upregulation of *c-Fos* and direct inhibition of NF $\kappa$ B binding to the TNF promoter (24, 33). The cholinergic inhibition of TNF mRNA expression in the present study was abrogated by the adenylyl cyclase inhibitor MDL 12,330A. A mechanistic role for cAMP in this context was further supported by our observation that knock-down of AC6 and *c-FOS*, respectively, abolished cholinergic attenuation of TNF transcription. Accordingly, these results indicate that cholinergic inhibition of TNF transcription and perhaps also release likely involves cAMP- and *c-Fos*-mediated transcriptional changes.

The  $\alpha 7$ nAChR subunit has been implicated in long term neuronal memory formation (23), a process in which the



**FIGURE 5 |** Proposed model of  $\alpha 7$ nAChR mediated prolonged regulation of TNF in macrophages. Activation of  $\alpha 7$ nAChR promotes activity of adenylyl cyclase 6, which increases production of cAMP. Increased intracellular levels of cAMP promotes phosphorylation of CREB, which enhances expression of *c-Fos*. *c-Fos*, in turn, is involved in NF $\kappa$ B mediated inhibition of TNF transcription. In this way,  $\alpha 7$ nAChR activation can inhibit endotoxin-induced TNF production.

transcription factor CREB also plays a key role (22). CREB was furthermore reported to modulate cytokine production in macrophage-like cells (34) and has been implicated in cAMP-mediated regulation of *c-Fos* expression (25). In the present study, functional inactivation of CREB abolished the cholinergic inhibition of TNF release in macrophages. Based on this observation, we propose the possibility that  $\alpha 7$ nAChR activation elicits CREB-dependent transcriptional changes in macrophages, which reduces endotoxin-induced TNF release (Figure 5).

Taken together, the results here identify mechanistic elements involved in the sustained reduction of TNF release after electrical vagus nerve stimulation and show that the cholinergic neurotransmitter acetylcholine can potentially confer a prolonged change in macrophage activation behavior. The resulting inhibition of TNF may attenuate excessive systemic inflammation.

In conclusion, our findings indicate that macrophages retain information from cholinergic signals. This mechanism requires the  $\alpha 7$ nAChR subunit, AC6, cAMP synthesis, phosphorylation of CREB, and expression of *c-Fos*. This data improves our understanding of the inflammatory reflex and potentially opens new avenues for therapeutic advancements in treatment of excessive inflammation.

## MATERIALS AND METHODS

### Ethical Statement

Experimental *in vivo* protocols were approved by the Institutional Animal Care and Use Committee (IACUC) at the Feinstein Institute for Medical Research, Northwell Health, which follows the National Institutes of Health guidelines for the ethical treatment of animals. Animals were housed under

a standard 12 h light/dark cycle and acclimatized for at least 1 week before conducting experiments. Water and regular rodent chow were available *ad libitum*. Human PBMCs from blood donors derive from completely anonymized, excess leukocytes, which do not require ethical approval.

## Vagus Nerve Stimulation

Male Sprague-Dawley rats were anesthetized using ketamine (100 mg/kg) and xylazine (10 mg/kg; IP). The left carotid sheath containing the vagus nerve was isolated and stimulated for 60 s at 1.0 mA, 10 Hz frequency, 0.25 ms biphasic pulse using a bipolar electrode (MicroProbes for Life Science). In sham-operated animals, the vagus nerve was visualized but not manipulated.

Male C57Bl/6, wild type and  $\alpha 7nAChR$  deficient, mice were anesthetized with ketamine (100 mg/kg) and xylazine (10 mg/kg), IP. A midline cervical incision was made, the salivary glands separated, and the left carotid sheath containing the vagus nerve was isolated between the sternomastoid and sternohyoid muscles and secured within a silastic covered bipolar platinum/iridium electrode (MicroProbes for Life Science). Charge-balanced biphasic stimulation was delivered at 0.75 mA, 0.20 ms pulse width, 10 Hz for 60 s by a custom stimulator (SetPoint Medical). In some animals, the nerve was stimulated with a single biphasic pulse, 0.1 s, at 1.0 mA. In sham-operated animals the vagus nerve was visualized but not manipulated. Skin was closed with surgical staples and the mice recovered.

## Rodent Endotoxemia

At various times post-VNS and sham-VNS (0, 2, 24, 48, and 72 h in rat, 24 h in mouse) VNS, endotoxin (0111:B4; Sigma-Aldrich) was injected intraperitoneally (IP; 1.5 mg/kg in rats; 8 mg/kg in mice). Rodents were euthanized after 90 min and serum TNF concentrations were measured using ELISA (R&D Systems).

## Macrophage cultures

RAW264.7 cells were incubated with 0 or 10  $\mu M$  acetylcholine in serum-free Opti-Mem media (Gibco) containing 1 mM pyridostigmine bromide (Sigma-Aldrich), for 60 min. Media was collected 4 h after endotoxin (10 ng/ml LPS) exposure and analyzed for TNF by ELISA (R&D Systems).

Human blood was collected from healthy subjects. Peripheral blood mononuclear cells (PBMC) were collected and monocytes isolated using the Monocyte Isolation Kit II (Miltenyi Biotec). Human monocyte and macrophage cultures were exposed to 10  $\mu M$  acetylcholine chloride (Sigma-Aldrich) in serum-free Opti-Mem media (Gibco) for 60 min in the presence of 1 mM pyridostigmine bromide. Cultures were washed to remove the cholinergic agonist and challenged with endotoxin in serum-free Opti-Mem media (Gibco) 24 h later. Media was collected 4 h after endotoxin addition and analyzed for TNF by ELISA (R&D Systems).

## cAMP and Adenylyl Cyclase Assay

Bone-marrow derived macrophages from wild type ( $n = 2$ ) and  $\alpha 7nAChR^{-/-}$  ( $n = 3$ ) mice were obtained by flushing

the femurs and tibia with sterile PBS. After 6 days of culture in the presence of 20 ng/ml M-CSF (PeproTech), cells were scraped, re-plated, and exposed to the  $\alpha 7nAChR$  selective agonist PHA568487 (Tocris) for 15 min. Wells were then washed in ice cold PBS, and cell lysis buffer (R&D systems) added on dry ice. Competitive cAMP ELISAs were conducted according to manufacturer instructions (R&D systems). Samples were freeze-thawed twice and centrifuged before assay. Protein concentration was determined by the Bradford assay (Bio-Rad) and used for normalization of cAMP values.

RAW264.7 cells were incubated with the indicated concentrations of [cis-N-(2-Phenylcyclopentyl)azacyclotridec-1-en-2-amine.HCl] (MDL 12,330A; Sigma Aldrich) for 30 min, followed by 50 mM choline chloride for 30 min. Endotoxin was added for 3 h. RNA was then isolated, cDNA generated, and analyzed by qPCR. Data shown (normalized to GAPDH) are representative of three (3) independent experiments and expressed as fold increase relative to unstimulated controls.

## siRNA Transfection

siRNA (125 pmol, Dharmacon) directed against adenylyl cyclase 6 (M-043463-01) or c-Fos (L-041157-00), or scrambled siRNA (D-001206-14-20), were transfected into RAW264.7 cells. 72 h later, cells were incubated with 50 mM choline for 30 min, followed by 100 ng/ml LPS for 3 h. RNA levels were analyzed by qPCR. Data shown (normalized to GAPDH) are representative of three independent experiments and expressed as fold increase relative to unstimulated controls.

## Western Blot

Protein was extracted from cell lysates and Western blot was performed using the anti-AC5/6 (sc-590, Santa Cruz Biotechnology), anti-c-Fos (sc-52, Santa Cruz Biotechnology), goat-anti-human CREB (AF2989, R&D Systems), rabbit anti-phospho-CREB (AF2510, R&D Systems), and anti-tubulin (05-661, Millipore) antibodies.

## ACREB Virus Transduction

Human macrophage cultures were transduced (Lipofectamine<sup>®</sup> 2000, ThermoFisher) with a replication deficient adenovirus (35) encoding green fluorescent protein (GFP) and a phosphorylation defective CREB (ACREB) (32) or a control virus encoding GFP. Three days after exposure to virus, cells were incubated with 100  $\mu M$  3-[(3E)-3-[(2,4-dimethoxyphenyl)methylidene]-5,6-dihydro-4H-pyridin-2-yl]pyridine (GTS21), a stable and selective  $\alpha 7nAChR$  agonist (36, 37), for 2 h, washed, and then cultured for 24 h. Endotoxin was added to the cultures for 4 h, then media collected for measurement of TNF by ELISA (R&D systems).

## Statistical Analysis

Differences between experimental groups were evaluated using two-tailed Student's *t*-test or ANOVA followed by Tukey

*post-hoc* analysis as appropriate.  $p < 0.05$  was considered significant.

## AUTHOR CONTRIBUTIONS

LT: project development, manuscript writing, and manuscript editing. PO and KT: project development, data collection, data analysis, manuscript writing, and manuscript editing. CR, MR-B, MT, AD, LH, WH, JL, WP, KO, YA-A, ME, VP, UA, SC, YL, TM, and KT: project development, data collection, data analysis, and manuscript editing. AC: manuscript writing and manuscript editing.

## REFERENCES

1. Netea MG, Joosten LAB, Latz E, Mills KHG, Natoli G, Stunnenberg HG, et al. Trained immunity: a program of innate immune memory in health and disease. *Science* (2016) 352:aaf1098. doi: 10.1126/science.aaf1098
2. Nathan C. Points of control in inflammation. *Nature* (2002) 420:846–52. doi: 10.1038/nature01320
3. Tracey KJ. Reflex control of immunity. *Nat Rev Immunol.* (2009) 9:418–28. doi: 10.1038/nri2566
4. Rosas-Ballina M, Ochani M, Parrish WR, Ochani K, Harris YT, Huston JM, et al. Splenic nerve is required for cholinergic antiinflammatory pathway control of TNF in endotoxemia. *Proc Natl Acad Sci USA.* (2008) 105:11008–13. doi: 10.1073/pnas.0803237105
5. Rosas-Ballina M, Olofsson PS, Ochani M, Valdés-Ferrer SI, Levine YA, Reardon C, et al. Acetylcholine-synthesizing T cells relay neural signals in a vagus nerve circuit. *Science* (2011) 334:98–101. doi: 10.1126/science.120985
6. Mirakaj V, Dalli J, Granja T, Rosenberger P, Serhan CN. Vagus nerve controls resolution and pro-resolving mediators of inflammation. *J Exp Med.* (2014) 211:1037–48. doi: 10.1084/jem.20132103
7. Huston JM, Gallowitsch-Puerta M, Ochani M, Ochani K, Yuan R, Rosas-Ballina M, et al. Transcutaneous vagus nerve stimulation reduces serum high mobility group box 1 levels and improves survival in murine sepsis\*. *Crit Care Med.* (2007) 35:2762–8. doi: 10.1097/01.CCM.0000288102.15975.BA
8. Olofsson PS, Levine YA, Caravaca A, Chavan SS, Pavlov VA, Faltys M, et al. Single-pulse and unidirectional electrical activation of the cervical vagus nerve reduces TNF in endotoxemia. *Bioelectron Med.* (2015)37–42. doi: 10.15424/bioelectronmed.2015.00006
9. Steinberg BE, Sundman E, Terrando N, Eriksson LI, Olofsson PS. Neural control of inflammation: implications for perioperative and critical care. *Anesthesiology* (2016) 124:1174–89. doi: 10.1097/ALN.0000000000001083
10. Levine YA, Koopman FA, Faltys M, Caravaca A, Bendele A, Zitnik R, et al. Neurostimulation of the cholinergic anti-inflammatory pathway ameliorates disease in rat collagen-induced arthritis. *PLoS ONE* (2014) 9:e104530. doi: 10.1371/journal.pone.0104530
11. Koopman FA, Chavan SS, Miljko S, Grazio S, Sokolovic S, Schuurman PR, et al. Vagus nerve stimulation inhibits cytokine production and attenuates disease severity in rheumatoid arthritis. *Proc Natl Acad Sci USA.* (2016) 113:8284–9. doi: 10.1073/pnas.1605635113
12. Bonaz B, Sinniger V, Hoffmann D, Clarençon D, Mathieu N, Dantzer C, et al. Chronic vagus nerve stimulation in Crohn's disease: a 6-month follow-up pilot study. *Neurogastroenterol Motil.* (2016) 28:948–53. doi: 10.1111/nmo.12792
13. Tracey KJ. The inflammatory reflex. *Nature* (2002) 420:853–9. doi: 10.1038/nature01321
14. Wang H, Yu M, Ochani M, Amella CA, Tanovic M, Susarla S, et al. Nicotinic acetylcholine receptor  $\alpha 7$  subunit is an essential regulator of inflammation. *Nature* (2002) 421:384–8. doi: 10.1038/nature01339
15. Olofsson PS, Katz DA, Rosas-Ballina M, Levine YA, Ochani M, Valdés-Ferrer SI, et al.  $\alpha 7$  nicotinic acetylcholine receptor ( $\alpha 7$ nAChR) expression in bone

## FUNDING

GCRC MO1 RR018535 NCRR (KT), Wenner-Gren Foundation (PO), Knut and Alice Wallenberg's foundation (PO), Heart-Lung Foundation (PO), Swedish research council (PO) and SetPoint Medical (YL).

## ACKNOWLEDGMENTS

We thank Dr. Mark Montminy for providing the ACREB adenovirus.

- marrow-derived non-T cells is required for the inflammatory reflex. *Mol Med.* (2012) 18:539–43. doi: 10.2119/molmed.2011.00405
16. Jonge WJ de, Zanden EP van der, The FO, Bijlsma ME, Westerloo DJ van, Bennink RJ, et al. Stimulation of the vagus nerve attenuates macrophage activation by activating the Jak2-STAT3 signaling pathway. *Nat Immunol.* (2005) 6:844. doi: 10.1038/ni1229
  17. Wang H, Liao H, Ochani M, Justiniani M, Lin X, Yang L, et al. Cholinergic agonists inhibit HMGB1 release and improve survival in experimental sepsis. *Nat Med.* (2004) 10:1216–21. doi: 10.1038/nm1124
  18. Rosas-Ballina M, Goldstein RS, Gallowitsch-Puerta M, Yang L, Valdés-Ferrer SI, Patel NB, et al. The selective  $\alpha 7$  agonist GTS-21 attenuates cytokine production in human whole blood and human monocytes activated by ligands for TLR2, TLR3, TLR4, TLR9, and RAGE. *Mol Med.* (2009) 15:195–202. doi: 10.2119/molmed.2009.00039
  19. Parrish WR, Rosas-Ballina M, Gallowitsch-Puerta M, Ochani M, Ochani K, Yang L-H, et al. Modulation of TNF release by choline requires  $\alpha 7$  subunit nicotinic acetylcholine receptor-mediated signaling. *Mol Med.* (2008) 14:567–74. doi: 10.2119/2008-00079.Parrish
  20. Shen J, Yakel JL. Nicotinic acetylcholine receptor-mediated calcium signaling in the nervous system. *Acta Pharmacol Sin.* (2009) 30:673–80. doi: 10.1038/aps.2009.64
  21. Báez-Pagán CA, Delgado-Vélez M, Lasalde-Dominicci JA. Activation of the macrophage  $\alpha 7$  nicotinic acetylcholine receptor and control of inflammation HHS public access. *J Neuroimmune Pharmacol.* (2015) 10:468–76. doi: 10.1007/s11481-015-9601-5
  22. Huang YY, Nguyen PV, Abel T, Kandel ER. Long-lasting forms of synaptic potentiation in the mammalian hippocampus. *Learn Mem* (1996) 3:74–85.
  23. Buccafusco JJ, Letchworth SR, Bencherif M, Lippello PM. Long-lasting cognitive improvement with nicotinic receptor agonists: Mechanisms of pharmacokinetic-pharmacodynamic discordance. *Trends Pharmacol Sci.* (2005) 26:352–60. doi: 10.1016/j.tips.2005.05.007
  24. Koga K, Takaesu G, Yoshida R, Nakaya M, Kobayashi T, Kinjo I, et al. Cyclic adenosine monophosphate suppresses the transcription of proinflammatory cytokines via the phosphorylated c-fos protein. *Immunity* (2009) 30:372–83. doi: 10.1016/j.immuni.2008.12.021
  25. Parry GC, Mackman N. Role of cyclic AMP response element-binding protein in cyclic AMP inhibition of NF-kappaB-mediated transcription. *J Immunol.* (1997) 159:5450–6.
  26. Olofsson PS, Rosas-Ballina M, Levine YA, Tracey KJ. Rethinking inflammation: neural circuits in the regulation of immunity. *Immunol Rev.* (2012) 248:188–204. doi: 10.1111/j.1600-065X.2012.01138.x
  27. Couturier S, Bertrand D, Matter JM, Hernandez MC, Bertrand S, Millar N, et al. A neuronal nicotinic acetylcholine receptor subunit ( $\alpha 7$ ) is developmentally regulated and forms a homo-oligomeric channel blocked by  $\alpha$ -BTX. *Neuron* (1990) 5:847–56. doi: 10.1016/0896-6273(90)90344-F
  28. Oshikawa J, Toya Y, Fujita T, Egawa M, Kawabe J, Umemura S, et al. Nicotinic acetylcholine receptor  $\alpha 7$  regulates cAMP signal within lipid rafts. *Am J Physiol Cell Physiol.* (2003) 285:C567–74. doi: 10.1152/ajpcell.00422.2002



29. Walker DP, Wishka DG, Piotrowski DW, Jia S, Reitz SC, Yates KM, et al. Design, synthesis, structure-activity relationship, and in vivo activity of azabicyclic aryl amides as  $\alpha 7$  nicotinic acetylcholine receptor agonists. *Bioorg Med Chem.* (2006) 14:8219–48. doi: 10.1016/j.bmc.2006.09.019
30. Boutillier AL, Barthel F, Roberts JL, Loeffler JP. Beta-adrenergic stimulation of cFOS via protein kinase A is mediated by cAMP regulatory element binding protein (CREB)-dependent and tissue-specific CREB-independent mechanisms in corticotrope cells. *J Biol Chem.* (1992) 267:23520–6.
31. Impey S, McCorkle SR, Cha-Molstad H, Dwyer JM, Yochum GS, Boss JM, et al. Defining the CREB regulon: a genome-wide analysis of transcription factor regulatory regions. *Cell* (2004) 119:1041–54. doi: 10.1016/j.cell.2004.10.032
32. Herzig S, Long F, Jhala US, Hedrick S, Quinn R, Bauer A, et al. CREB regulates hepatic gluconeogenesis through the coactivator PGC-1. *Nature* (2001) 413:179–83. doi: 10.1038/35093131
33. Cai W, Du A, Feng K, Zhao X, Qian L, Ostrom RS, et al. Adenylyl Cyclase 6 activation negatively regulates TLR4 signaling through lipid raft-mediated endocytosis. *J Immunol.* (2013) 191:6093–100. doi: 10.4049/jimmunol.1301912
34. Avni D, Ernst O, Philosoph A, Zor T. Role of CREB in modulation of TNF $\alpha$  and IL-10 expression in LPS-stimulated RAW264.7 macrophages. *Mol Immunol.* (2010) 47:1396–403. doi: 10.1016/j.molimm.2010.02.015
35. Foxwell B, Browne K, Bondeson J, Clarke C, de Martin R, Brennan F, et al. Efficient adenoviral infection with IkappaB alpha reveals that macrophage tumor necrosis factor alpha production in rheumatoid arthritis is NF-kappaB dependent. *Proc Natl Acad Sci USA.* (1998) 95:8211–5.
36. Mahnir V, Lin B, Prokai-Tatrai K, Kem WR. Pharmacokinetics and urinary excretion of DMXBA (GTS-21), a compound enhancing cognition. *Biopharm Drug Dispos.* (1998) 19:147–51.
37. Papke RL, Porter Papke JK. Comparative pharmacology of rat and human  $\alpha 7$  nAChR conducted with net charge analysis. *Br J Pharmacol.* (2002) 137:49–61. doi: 10.1038/sj.bjp.0704833

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Tarnawski, Reardon, Caravaca, Rosas-Ballina, Tusche, Drake, Hudson, Hanes, Li, Parrish, Ojamaa, Al-Abed, Faltys, Pavlov, Andersson, Chavan, Levine, Mak, Tracey and Olofsson. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.